



## BIOMEDICAL SCIENCES

# $\beta$ -Cyclodextrins alter the energy metabolism-related enzyme activities in rats

FRANCINE R. IANISKI, AMANDA L. DE OLIVEIRA, NATHANA JAMILLE MEZZOMO, ITIANE D. DE FRANCESCHI, GUILHERME M. DO CARMO, CAMILA R. CREMONESE, MATHEUS D. BALDISSERA, JENIFER P. ZANON, JENIFER KOLLING, JORDANA DANIELA FRIEDERICH, IVANA Z. DA SILVA, JANICE L. GIONGO, LUCIANE R. FEKSA, RODRIGO A. VAUCHER, CLOVIS MILTON D. WANNMACHER & VIRGINIA C. RECH

**Abstract:** Although widely used in medicine, separation technology, and other fields, the effects of cyclodextrins on the activities of phosphoryl transfer enzymes have not been previously evaluated. *In vivo* studies evaluated the function of cyclodextrins as active compounds. Despite the use of cyclodextrins as active compounds, the effects of cyclodextrins on hepatic and renal tissues remain to be fully elucidated. The primary objective of this study was to evaluate the effects of  $\beta$ -cyclodextrins, methyl- $\beta$ -cyclodextrin (M- $\beta$ -cyclodextrins), and (2-hydroxypropyl)- $\beta$ -cyclodextrin (HP- $\beta$ -cyclodextrins) on enzyme activities regulating the maintenance of energy homeostasis in the kidney and liver tissues in relation to toxicity. Serum levels of liver and kidney markers were measured, and oxidative stress parameters were assessed. After 60-day treatments, we observed that the administration of  $\beta$ -cyclodextrins and M- $\beta$ -cyclodextrins inhibited the hepatic activity of pyruvate kinase, an irreversible enzyme within the glycolytic pathway. Additionally, administration of HP- $\beta$ -cyclodextrins inhibited creatine kinase activity and increased the total sulfhydryl content in kidneys. Here, we demonstrated for the first time that  $\beta$ -cyclodextrins, M- $\beta$ -cyclodextrins, and HP- $\beta$ -cyclodextrins cause bioenergetic dysfunction in renal and hepatic tissues. These findings suggest that understanding the balance between cyclodextrins' efficacy and adverse effects is essential for better accepting their use in medicine.

**Key words:** creatine kinase, adenylate kinase, pyruvate kinase, oxidative stress, biochemical, blood serum.

## INTRODUCTION

The  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrins (CDs) are cyclic oligosaccharide molecules. These molecules possess nanometric cavities and are composed of 6 – 8 D-glucose units (Souza et al. 2016). These cavities are hydrophobic and interact with drugs that are poorly soluble in water. This reaction forms inclusion complexes that improve the biocompatibility of drugs and their use within biological environments. This property promotes the sustained release of drugs (Yang

et al. 2013) and allows for the lodging of organic compounds. Hydroxyls in CDs cause the external environment to become hydrophilic (Souza et al. 2016).

For these attributes, CDs have been widely used for applications in the pharmaceutical field, the pesticide, foodstuffs, toilet articles, and textile processing industries, and for supramolecular host-guest chemistry and molecular encapsulation (Gao et al. 2006, Mura 2020, Tang et al. 2020). CDs can interact

with cholesterol and phospholipids in the biomembranes of the gastrointestinal tract. They can also pass through the stomach and small intestine but are degraded in the intestinal colon. This characteristic is associated with the potential adaptability of these drug carriers (Uekama et al. 2006).

$\beta$ CDs possessing either methyl (M- $\beta$ CD) or hydroxypropyl (HP- $\beta$ CD) groups have been used in pharmaceutical preparations, as toxicity studies have shown that these CDs are practically non-toxic after oral administration (Mura 2020). Nishihira et al. (2017) demonstrated that resveratrol complexed with HP- $\beta$ -CD did not cause any significant differences in hematological and biochemical parameters. Therefore, CDs can carry hydrophobic molecules in an aqueous environment, act as non-toxic and biocompatible carriers, and are widely used throughout the pharmaceutical field (Monteil et al. 2017, Souza et al. 2018).

However, the effects of CDs on the activities of phosphoryl transfer enzymes, pyruvate kinase (PK), creatine kinase (CK), and adenylate kinase (AK) have yet to be previously evaluated. The coupling between the processes of consumption and production of cellular ATP must be tightly balanced, and this coupling is fundamental for various metabolic processes within living organisms and cell signaling (Rech et al. 2018). The enzymes PK, CK, and AK allow this connection between cell compartments by transferring the  $\gamma$ -phosphoryl group of ATP from mitochondria to the cytosol (Dzeja & Terzic 2003). Based on this, these enzymes are crucial for maintaining cellular energy homeostasis in most mammalian tissues. Imbalances between the flow and distribution of cellular energy have been associated with cardiovascular and neurodegenerative diseases, and these imbalances can influence the uncontrolled growth and metastatic potential of tumor cells

(Dzeja & Terzic 2003, 2009). A decrease in the functions of these enzymes may be associated with a crucial role in the physiopathology of neurodegeneration and the subsequent loss of neurons (Bortoluzzi et al. 2014). Consequently, the study of the activities of these enzymes is critical to corroborate reports that CDs are not toxic to living organisms.

In addition to the phosphotransfer network system, oxidative stress is associated with several diseases, including neurodegenerative processes. Oxidative stress is an imbalance in relation to the pro-oxidant side of pro-oxidant/antioxidant homeostasis that occurs in several human diseases, including diabetes, chronic renal failure, and respiratory distress syndrome (Dalle-Donne et al. 2003). Yalcin et al. (2016) demonstrated that using HP- $\beta$ CD microspheres decreased lipid peroxidation levels and the production of reactive oxygen species in a model of A $\beta$  (1-42)-induced neurotoxicity. Considering the importance of CDs as drug carriers or as active compounds, we evaluate the effect of  $\beta$ CD administration on energy parameters and oxidative stress in hepatic and renal tissues to better understand their mechanisms of action.

## MATERIALS AND METHODS

### Animals and Reagents

Thirty-two male Wistar rats (sixty-day-old) were obtained from the Central Animal House of the Federal University of Santa Maria and used in the experiments. The animals were maintained on a 12:12 h light/dark cycle in a room acclimatized at constant temperature ( $22 \pm 1^\circ\text{C}$ ). The rats were provided free access to water and commercial chow (Guabi, RS, Brazil). The Ethics Committee approved the experimental protocol for Animal Research of the Franciscan University, Santa Maria, Brazil, under protocol number 010/2011 and followed the 'Principles

of Laboratory Animal Care' (NIH publication 85-23, revised 1996). All chemicals, including  $\beta$ CD, M- $\beta$ CD, and HP- $\beta$ CD, were purchased from Sigma (St. Louis, MO, USA).

### Treatment of Rats

The animals were randomly divided into four groups of eight rats, each that included groups I (Control), II ( $\beta$ CD), III (M- $\beta$ CD), and IV (HP- $\beta$ CD). The animals received intraperitoneal treatments of saline solution (0.9% NaCl), 66 mg  $\beta$ CD/kg (body weight), M- $\beta$ CD/kg (body weight), and HP- $\beta$ CD/kg (body weight), respectively, for 60 days. These doses were chosen based on previous Bellringer et al. (1995) data. The intraperitoneal route was chosen according to Trautwein et al. (1999), as some CDs interfered with the absorption of certain nutrients.

### Preparation of Hepatic and Renal Tissues

After sixty days of treatment, the animals were euthanized by decapitation. The liver and kidneys were quickly removed, dissected, and frozen at  $-80\text{ }^{\circ}\text{C}$  for no more than a week until the determination of energy metabolism and oxidative stress parameters.

Kidneys and liver were homogenized to evaluate the parameters of the phosphoryl transfer network (1:10, w/v with a Potter-Elvehjem glass homogenizer) in SET buffer (0.32 M sucrose, 1 mM EGTA, 10 mM Tris-HCl, pH 7.4). The homogenate was centrifuged at  $800 \times g$  for 10 min at  $4\text{ }^{\circ}\text{C}$  to discard nuclei and cell debris, and the pellet was discarded. The supernatant that contained a suspension of mixed and preserved organelles, including mitochondria, was then separated and fractionated. An aliquot of the supernatant was used for AK activity determination. The pellet was discarded, and the remaining supernatant was centrifuged at  $10,000 \times g$  for 15 min at  $4\text{ }^{\circ}\text{C}$ .

The supernatant of this second centrifugation corresponded to the cytosolic fraction. It was collected to determine PK and cytosolic CK activities (CK<sub>cyt</sub>). The pellet, containing mitochondria, myelin, synaptosomes, and membrane fragments, was washed twice with the same Tris-sucrose isotonic buffer and then resuspended in 100 mM  $\text{MgSO}_4$ -Trizma buffer (pH 7.5) for determination of mitochondrial CK activity (CK<sub>mit</sub>). The samples were stored for no more than a week at  $-80\text{ }^{\circ}\text{C}$ . The other half of the liver and the other kidneys were weighed and homogenized (1:10, w/v) in 20 mM sodium phosphate buffer (pH 7.4) containing 140 mM KCl. Homogenates were centrifuged at  $800 \times g$  for 10 min at  $4\text{ }^{\circ}\text{C}$ , and the pellet was discarded. The supernatant containing a suspension of mixed and preserved organelles, including mitochondria, was separated and used for the various oxidative stress analyses.

### Determination of energy parameters

The activities of CK, PK, and AK enzymes were determined in liver and kidney homogenates according to the method of Hughes (1962), Leong et al. (1981), and Dzeja et al. (1999), respectively. The results were expressed as  $\beta$ mol of creatine formed per min per mg of protein for CK, as  $\beta$ mol of pyruvate formed per min per mg of protein for PK, and as  $\beta$ mol of ATP formed per min per mg of protein for AK.

### 2'7' Dihydrodichlorofluorescein oxidation assay (H<sub>2</sub>DCF-DA)

The levels of reactive species were evaluated according to LeBel et al. (1992) using reduced 2', 7'-dihydrodichlorofluorescein diacetate (H<sub>2</sub>DCF-DA). H<sub>2</sub>DCF-DA is enzymatically hydrolyzed to H<sub>2</sub>DCF by intracellular esterases. H<sub>2</sub>DCF is oxidized to DCF by reactive oxygen species (ROS) or reactive nitrogen species (RNS) present in the samples. The intensity of the DCF fluorescence

corresponds to the proportion of reactive species present within the sample. Fluorescence was measured using excitation and emission wavelengths of 480 and 535 nm, respectively. The results are expressed as nmol of DCF. mg protein<sup>-1</sup>.

### **Thiobarbituric acid-reactive substances (TBARS)**

Thiobarbituric acid-reactive substances were determined according to the method described by Ohkawa et al. (1979), and this method essentially evaluates malondialdehyde (MDA), a lipid peroxidation product. This attack on the lipids primarily occurs due to hydroxyl radicals ( $\bullet\text{OH}$ ) derived from  $\text{H}_2\text{O}_2$  by the iron-catalyzed Fenton reaction or the Haber–Weiss reaction (Kehrer 2000). Each sample was assessed at 532 nm using a spectrophotometer. The results are expressed as nmol of MDA. mg protein<sup>-1</sup>.

### **Determination of carbonyls**

Carbonyl protein content was determined using the method described by Reznick & Packer (1994). The oxidation of amino acids forms carbonyls in proteins or adducts formed along with lipoperoxidation products such as 4-hydroxy-2-nonenal (HNE). Each sample was measured at 370 nm, and total carbonylation was calculated using a molar extinction coefficient of 22,000  $\text{M}^{-1} \text{cm}^{-1}$  as described by Levine et al. (1990). The results are expressed as nmol of carbonyls. mg protein<sup>-1</sup>.

### **Total sulfhydryl content**

The sulfhydryl assays are based on the method of Aksenov & Markesbery (2001), where the reduction of 5, 50-dithio-bis (2-nitrobenzoic acid) (DTNB) by thiols generates a yellow derivative (TNB) with absorption that can be measured spectrophotometrically at 412 nm.

The results are expressed as nmol of TNB per mg of protein.

### **Protein determination**

The protein content of the homogenates was determined according to the method of Lowry et al. (1951) using bovine serum albumin as the standard.

### **Serum Biochemical Parameters**

Blood samples were carefully collected and centrifuged at  $900 \times g$  for 15 min. The serum was removed and stored at  $-80^\circ\text{C}$  until analysis. Glucose, triglycerides (TG), total cholesterol, low-density lipoprotein (LDL), high-density lipoprotein (HDL), very low-density lipoprotein (VLDL), and total protein were measured by an automated analyzer (Vitros 250, Ortho-Clinical Diagnostics, Rochester, NY) using Johnson and Johnson kits that incorporated a chemical drying method. Aspartate aminotransferase (AST), alanine aminotransferase (ALT), urea, creatinine, uric acid, lipase, albumin, gamma-glutamyl transferase, phosphatase alkaline, iron, phosphate, sodium, potassium, calcium, and magnesium were also measured.

### **Statistical analysis**

Data were analyzed by repeated ANOVA tests followed by the Tukey test when the F values were significant. Differences between groups were rated as significant at  $p < 0.05$ . All analyses were performed using SPSS software.

## **RESULTS**

After treatment for 60 days, the animals were initially evaluated to assess the effects of the intraperitoneal administrations of the CDs on the activities of the enzymes AK, PK, and CK, all of which are important enzymes involved in the phosphotransfer network. Subsequently, we

studied the effects of the treatments on serum biochemical parameters.

### Creatine kinase activity

First, we observed that CKcyt activity in kidney tissue was not significantly different among the groups ( $p > 0.05$ ) (Table I). CKmit activity in the kidney tissue of rats [F (3, 22) = 5.07;  $p < 0.05$ ] exhibited a significant decrease in the HP- $\beta$ CD group compared to that of the control group (Table I). In contrast, CKmit activity was not altered by  $\beta$ CD or by M- $\beta$ CD ( $p > 0.05$ ) (Table I).

### Pyruvate kinase activity

PK activity in the kidney tissue of rats was not significantly different among the groups ( $p > 0.05$ ) (Table I). However, the activity of this enzyme in the liver was significantly reduced [F (3.23) = 7.11;  $p < 0.05$ ] in the  $\beta$ CD and M- $\beta$ CD groups compared to that in the co-control group (Table I).

### Adenylate kinase activity

AK activity in the kidney and liver tissues of rats was not significantly different among the groups ( $p > 0.05$ ) (Table I).

### 2',7'-Dihydrodichlorofluorescein (DCFH) oxidation levels

The generation of reactive species as evaluated by 2',7'-dichlorofluorescein production was not significantly different among the groups ( $p > 0.05$ ) in the kidney tissue of rats (Table II).

DCFH oxidation levels in the liver were significantly increased in response to M- $\beta$ CD and HP- $\beta$ CD administration [F (3, 25) = 4.822;  $p < 0.05$ ] in rats; however, these levels were not increased in response to  $\beta$ CD (Table II).

### Lipid peroxidation

Next, we evaluated the effects of CDs on MDA levels. We observed that this metabolite did not exhibit any significant differences among the groups ( $p > 0.05$ ) in the kidney and liver tissues of rats (Table II).

### Carbonyl formation

Carbonyl formation in the kidney and liver tissues of the rats was not significantly different among the groups ( $p > 0.05$ ) (Table II).

### Total sulfhydryl content

Total sulfhydryl content in the liver was not altered by CD administration ( $p > 0.05$ ) compared to levels in the control group (Table II).

**Table I. Effect of intraperitoneal administration of  $\beta$ CD, M- $\beta$ CD, and HP- $\beta$ CD on cytosolic creatine kinase (CKcyt), mitochondrial creatine kinase (CKmit) activity in kidney tissue and adenylate kinase (AK) pyruvate kinase (PK) activity in kidney and liver tissues from rats.**

Tissues	Enzyme activity	Control	$\beta$ CD	M- $\beta$ CD	HP- $\beta$ CD
Kidney	CKcyt	0.53 $\pm$ 0.08	0.5 $\pm$ 0.08	0.48 $\pm$ 0.1	0.46 $\pm$ 0.07
	CKmit	0.38 $\pm$ 0.03	0.33 $\pm$ 0.03	0.35 $\pm$ 0.08	0.28 $\pm$ 0.02*
	AK	0.13 $\pm$ 0.02	0.14 $\pm$ 0.02	0.14 $\pm$ 0.02	0.16 $\pm$ 0.01
	PK	6.50 $\pm$ 0.88	7.78 $\pm$ 0.84	7.55 $\pm$ 0.85	6.77 $\pm$ 0.35
Liver	AK	0.33 $\pm$ 0.02	0.33 $\pm$ 0.05	0.34 $\pm$ 0.03	0.34 $\pm$ 0.07
	PK	15.7 $\pm$ 3.2	9.8 $\pm$ 3.3*	10.3 $\pm$ 2.1*	13.7 $\pm$ 3.6

Values are mean  $\pm$  standard deviation for six to eight independent experiments (animals) performed in triplicate and are expressed as  $\beta$ mol creatine per min per mg of protein for CK; performed in duplicate and are expressed as  $\beta$ mol of ATP per min per mg of protein for AK; performed in duplicate and are expressed as  $\beta$ mol of pyruvate per min per mg of protein for PK. \* $p < 0.05$ , compared to the Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M- $\beta$ CD: methyl; HP- $\beta$ CD: hydroxypropyl.

II). In contrast, total sulfhydryl content was significantly increased in the kidney following HP- $\beta$ CD administration [F (3, 23) = 87.488,  $p < 0.001$ ] in rats; however, this content was not increased in response to  $\beta$ CD or M- $\beta$ CD (Table II).

### Serum Biochemical Parameters

The results for serum biochemical parameters are shown in Table III. AST levels were significantly increased by M- $\beta$ CD and HP- $\beta$ CD administration in the serum [F (3, 26) = 7.950;  $p < 0.05$ ] of rats, and these levels were not increased in response to  $\beta$ CD. ALT, lipase, gamma-glutamyl transferase, and glucose levels were not significantly different among groups.

BCD or HP- $\beta$ CD administration did not alter the serum's albumin, total protein, and phosphatase alkaline. However, the M- $\beta$ CD administration significantly increased albumin [F (3, 27) = 6.259;  $p < 0.05$ ], total protein [F (3, 28) = 4.606;  $p < 0.05$ ], and phosphatase alkaline [F (3, 22) = 7.467;  $p < 0.05$ ] levels in the serum of rats.

The analysis of the serum electrolyte concentrations shows (Tables III and IV) a significant reduction in the concentration of iron in response to  $\beta$ CD, M- $\beta$ CD, and

HP- $\beta$ CD administration [F (3,23) = 19.682;  $p < 0.05$ ] in the serum of rats. However, only M- $\beta$ CD administration resulted in a significant increase in sodium [F (3, 28) = 5.759;  $p < 0.05$ ], calcium [F (3, 28) = 5.505;  $p < 0.05$ ], magnesium [F (3, 28) = 4.810;  $p < 0.01$ ] and phosphate [F (3,28) = 2.755;  $p > 0,05$ ] in the serum of rats, but not by  $\beta$ CD or HP- $\beta$ CD.

Renal function (Table V) was evaluated according to urea, creatine, and uric acid levels. We observed a significant increase in urea [F (3, 28) = 3.161;  $p < 0.05$ ] and creatinine [F (3, 28) = 3.400;  $p < 0.05$ ] concentrations following  $\beta$ CD administration in the serum of rats, and this was not observed in response to M- $\beta$ CD or HP- $\beta$ CD administration.

We then analyzed the lipid profiles (Table VI). We found that M- $\beta$ CD administration increased LDL levels [F (3, 24) = 7.929;  $p < 0.05$ ], while  $\beta$ CD or HP- $\beta$ CD administration did not alter this parameter. Moreover, other parameters, including total cholesterol, HDL, VLDL, and TG, were not significantly different among the groups in the serum of rats.

**Table II.** *In vivo* effect of  $\beta$ CD, M- $\beta$ CD and HP- $\beta$ CD on the oxidative stress parameters from kidney and liver of rats.

Tissues	Tests	Control	$\beta$ CD	M- $\beta$ CD	HP- $\beta$ CD
Kidney	Sulfhydryl content	18.4 $\pm$ 4.9	22.2 $\pm$ 1,7	24.4 $\pm$ 7.5	60.9 $\pm$ 5.4*
	DCF	9.4 $\pm$ 0.7	9.7 $\pm$ 1.1	9.8 $\pm$ 0.6	9.3 $\pm$ 1.0
	TBARS	1887 $\pm$ 354	1918 $\pm$ 252	2064 $\pm$ 203	1958 $\pm$ 262
	Carbonylation protein	6.4 $\pm$ 1.6	5.7 $\pm$ 0.4	7.8 $\pm$ 1.8	6.2 $\pm$ 2.3
Liver	Sulfhydryl content	28.6 $\pm$ 3.8	33.7 $\pm$ 5,4	27 $\pm$ 4.1	35.1 $\pm$ 8.6
	DCF	4.3 $\pm$ 0.2	4.6 $\pm$ 0.2	4.7 $\pm$ 0.3*	4.7 $\pm$ 0.2*
	TBARS	1134 $\pm$ 88	1252 $\pm$ 72	1239 $\pm$ 86	1192 $\pm$ 105
	Carbonylation protein	5.0 $\pm$ 16	4.8 $\pm$ 0.8	5.5 $\pm$ 1.7	5.7 $\pm$ 0.5

Values are means  $\pm$  standard deviation for six to eight independent experiments (animals) per group, performed in triplicate. \* $p < 0.05$ , compared to Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M- $\beta$ CD: methyl; HP- $\beta$ CD: hydroxypropyl; DCF: dichlorofluorescein; TBARS: thiobarbituric acid-reactive substances.

**Table III.** *In vivo* effect of βCD, M-βCD and HP-βCD on the serum biochemical parameters from rats.

Tests	Control	βCD	M-βCD	HP-βCD
AST (U/L)	247 ± 58	267 ± 29	518 ± 248*	550 ± 121*
ALT (U/L)	69.3 ± 9.8	70.8 ± 11	68.7 ± 7.6	70.6 ± 4.7
Albumin (g/dL)	2.7 ± 0.1	3.0 ± 0.2	3.2 ± 0.3*	3.0 ± 0.2
Gamma glutamyl transferase (U/L)	5.0 ± 0.0	5.0 ± 0.0	5.0 ± 0.0	5.3 ± 0.7
Glucose (mg/mL)	101 ± 5	104 ± 8	102 ± 12	100 ± 11
Total protein (g/dL)	6.1 ± 0.3	6.2 ± 0.3	6.7 ± 0.3*	6.3 ± 0.4
Lipase (U/L)	88 ± 19	67 ± 21	99 ± 26	97 ± 26
Phosphatase alkaline (U/L)	137 ± 25	124 ± 17	197 ± 47*	165 ± 19

Values are means ± standard deviation for six to eight independent experiments (animals) performed in triplicate. \* $p < 0.05$ , compared to Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M-βCD: methyl; HP-βCD: hydroxypropyl; AST: aspartate aminotransferase; ALT: alanine aminotransferase.

**Table IV.** *In vivo* effect of βCD, M-βCD and HP-βCD on the serum electrolyte concentrations from rats.

Tests	Control	βCD	M-βCD	HP-βCD
Sodium (mg/dL)	141 ± 5	144 ± 4	155 ± 12*	144 ± 3
Potassium (mmol/L)	6.7 ± 0.7	7.3 ± 0.4	7.4 ± 0.7	7.1 ± 0.3
Iron (mg/dL)	226 ± 31	143 ± 16*	192 ± 16*	149 ± 26*
Calcium (mg/dL)	9.6 ± 0.3	9.6 ± 0.3	10.5 ± 0.8*	9.8 ± 0.3
Magnesium (mg/dL)	2.1 ± 0.2	2.2 ± 0.2	2.4 ± 0.2*	2.1 ± 0.2
Phosphate (mg/dL)	8.3 ± 1.0	8.7 ± 0.9	9.3 ± 0.7*	8.6 ± 0.6

Values are means ± standard deviation for six to eight independent experiments (animals) performed in triplicate. \* $p < 0.05$ , compared to Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M-βCD: methyl; HP-βCD: hydroxypropyl.

## DISCUSSION

The effects of βCD and M-βCD on bioenergetic dysfunction in rats' renal and hepatic tissues have been characterized here for the first time. Our data revealed that βCD and M-βCD administration inhibited hepatic PK activity. Administration of HP-βCD inhibited CK activity and increased the total sulfhydryl content in the kidney. In general, these findings may suggest that understanding the balance between the efficacy and adverse effects of using CDs is fundamental for better accepting their use in medicine.

The literature has strong evidence that CDs are virtually non-toxic to the biological

environment (Loftsson & Duchene 2007, Nishihira et al. 2017). However, when associating drugs with CDs, it is expected that there is no toxicity on the part of the carrier and that possible toxicity of the associated drug is minimized through this association. Thus, there is an immense need for further studies related to the possible toxicity of CDs. Many mechanisms from the action of cyclodextrins in the biological environment will still be reported in future research, enriching our understanding of them.

It is also important to evaluate the bioenergetic function in tissues. Cell viability is highly dependent on ATP required for the function of ATPase enzymes responsible for many cellular processes, including intracellular

**Table V.** *In vivo* effect of  $\beta$ CD, M- $\beta$ CD and HP- $\beta$ CD on the measurements of renal function in rats.

Tests	Control	$\beta$ CD	M- $\beta$ CD	HP- $\beta$ CD
Urea (mg/dL)	36.7 $\pm$ 5	45.5 $\pm$ 6.6*	41.3 $\pm$ 6.3	41.5 $\pm$ 4.3
Uric acid (mg/dL)	1.3 $\pm$ 0.3	1.5 $\pm$ 0.2	1.5 $\pm$ 0.3	1.4 $\pm$ 0.3
Creatinine (mg/dL)	0.5 $\pm$ 0.1	0.6 $\pm$ 0.1*	0.56 $\pm$ 0.1	0.56 $\pm$ 0.1

Values are means  $\pm$  standard deviation for six to eight independent experiments (animals) performed in triplicate. \* $p < 0.05$ , compared to Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M- $\beta$ CD: methyl; HP- $\beta$ CD: hydroxypropyl.

and extracellular electrolyte and biomolecule homeostasis (Wallimann et al. 1992, Rech et al. 2018). The creatine kinase/phosphocreatine system (CK/PCr system) is responsible for the rapid replacement of ATP (Wallimann et al. 1992, Rech et al. 2018) in tissues or cells that require high levels of energy (Machado et al. 2016).

From the analysis of the CK system, it was observed that there are distinct CK isoenzymes that are compartmentalized specifically in areas that produce (mitochondria) or use (cytosol) ATP (Wallimann et al. 1992). In this study, HP- $\beta$ CD inhibited CKmit activity in the kidneys of rats. Inhibition of CKmit enzyme activity can cause energy loss, ADP accumulation, and excess intracellular calcium, ultimately leading to oxidative stress (Machado et al. 2016). Some oxidative damage biomarkers include protein oxidation (indexed by protein carbonyls) and lipid peroxidation products (Dalle-Donne et al. 2003). In the interim, it is important to determine if the activities of thiol-containing enzymes such as PK, CK, and AK can change in response to oxidative stress (Rech et al. 2008, De Franceschi et al. 2013). In this study, it was observed that the administration of HP- $\beta$ CD caused an increase in total sulfhydryl levels but did not affect the levels of TBARS, DCFH oxidation, and protein carbonylation in the kidney, indicating that HP- $\beta$ CD did not induce oxidative stress in the rat kidney. Based on this, it can be concluded that there is a possibility that a relationship exists between the increase in total sulfhydryl levels and the reduction of CKmit activity. We also

observed that  $\beta$ CD and M- $\beta$ CD administration did not alter CK activity in the kidney.

In liver tissue, the administration of HP- $\beta$ CD increased the oxidation of DCFH but did not alter the activity of the enzymes PK and AK. In contrast, the administration of M- $\beta$ CD, in addition to increasing the oxidation of DCFH due to an increase in oxidative species, also reduced the activity of PK. Here, it must be emphasized that PK activity is the primary route for producing pyruvate in the liver, as demonstrated by Burgess et al. (2008). This reference highlights the importance of analyzing this route, and the results of this study shed light on the possible effects of the use of CDs on bioenergetic dysfunction. Moreover, PK activity in the liver was also reduced by  $\beta$ CD administration; however, this treatment did not affect any of the oxidative stress parameters evaluated. The observed inhibited activity of PK implies a reduction in the amount of cellular pyruvate, an important antioxidant agent (Figueiredo et al. 2009), and a reduction in the ATP cell pool (Rech et al. 2018) that can induce cell death (Gabardo et al. 2015).

According to the proposed experimental design, it was observed that there was no induction of oxidative stress in response to the administration of  $\beta$ CD, M- $\beta$ CD, or HP- $\beta$ CD in animals compared to that observed in the control group. This result highlights the non-toxic characteristics of CDs concerning the evaluated parameters. Our findings regarding the non-toxicity of CDs in biological environments are corroborated by the study performed by



**Table VI.** *In vivo* effect of  $\beta$ CD, M- $\beta$ CD and HP- $\beta$ CD on the lipids measurements in rats.

Tests	Control	$\beta$ CD	M- $\beta$ CD	HP- $\beta$ CD
Cholesterol (mg/mL)	83 $\pm$ 13	86 $\pm$ 13	99 $\pm$ 14	85 $\pm$ 11
VLDL (mg/mL)	9.9 $\pm$ 1.2	9.1 $\pm$ 1.9	9.1 $\pm$ 1.2	8.9 $\pm$ 1.2
LDL (mg/mL)	26.3 $\pm$ 5.3	23.4 $\pm$ 4.7	36.4 $\pm$ 7.2*	24.2 $\pm$ 4.9
HDL (mg/mL)	48.8 $\pm$ 5.5	51.3 $\pm$ 3.9	55.8 $\pm$ 6.6	52 $\pm$ 6.0
TG (mg/mL)	52 $\pm$ 8	46 $\pm$ 9	46 $\pm$ 6	45 $\pm$ 6

Values are means  $\pm$  standard deviation for six to eight independent experiments (animals) performed in triplicate. \* $p < 0.05$ , compared to Control group (ANOVA followed by Tukey test). CD: cyclodextrins; M- $\beta$ CD: methyl; HP- $\beta$ CD: hydroxypropyl; VLDL: very low-density lipoprotein; LDL: low-density lipoprotein; HDL: high-density lipoprotein; TG: triglycerides.

Nishihira et al. (2017). Although the DCs affected the activities of some enzymes involved in energy metabolism, no correlation between these effects and the evaluated parameters of oxidative stress was observed.

From the analysis of bioenergetic functions, it can be determined that there is a significant relationship between CK and AK enzymes. Both CK and AK are closely associated in such a manner that when the activity of one enzyme is reduced, the activity of the other enzyme is enhanced (Dzeja & Terzic 2009, Figueiredo et al. 2009). The AK enzyme catalyzes the reversible reaction of nucleotide phosphoryl exchange among ATP, ADP, and AMP. Thus, this enzyme transfers the phosphoryl group from ATP, produced in mitochondria, to the cytosol, where it is consumed. However, no interaction between CK and AK activities was found in this study.

Furthermore, it was observed that the administration of M- $\beta$ CD and HP- $\beta$ CD significantly increased the serum levels of AST. According to Mello et al. (2012), an increase in serum AST levels indicates the induction of chronic hepatocellular damage. Despite this, it is generally observed that hepatic changes cause variations in AST levels accompanied by increased ALT levels (Spanish Society of Clinical Biochemistry and Molecular Pathology 2012).

Additionally, in this study, it was observed that the administration of M- $\beta$ CD significantly

increased the serum levels of albumin, total proteins, and alkaline phosphatase. It is widely established that serum albumin is present in much greater amounts in the blood than other proteins (Chandrasekaran et al. 2016). It should be considered that the increase in the serum albumin level might be related to the increase in the total protein. It is well established that in liver disorders, serum albumin levels decrease; however, it is also established that these levels increase in response to dehydration. These results, together with increased sodium, calcium, magnesium, and phosphate levels, are characteristic of dehydration rather than liver damage caused by M- $\beta$ CD. In agreement with Moe (2008), it has been observed that most patients with renal disease in conjunction with hyperphosphatemia and hypermagnesemia exhibit increased gastrointestinal absorption. Furthermore, magnesium, phosphate, and calcium are essential for many biological and cellular functions. Thus, it must be noted that maintaining the homeostasis of these ions is a central role performed by the kidneys (Blaine et al. 2015).

Among the findings of the present study, it was observed that treatment with  $\beta$ CD significantly increased urea and creatinine levels in the rats' serum. These parameters are associated with liver damage. Mosher & Thompson (2002) observed liver damage when

performing experiments that included the oral administration of  $\beta$ CD. However, it was also observed in the present study that  $\beta$ CD did not alter AST and ALT levels. Thus, it is advisable to perform further studies to elucidate possible tissue alterations that the administration of  $\beta$ CD may cause.

A significant reduction in iron levels was observed based on the analysis of serum electrolyte concentrations. In response to  $\beta$ CD and HP- $\beta$ CD treatments, the reduction was approximately 40% and around 15% for M- $\beta$ CD. It is common knowledge that iron is a facilitator of reversible redox reactions, and this element is, therefore, essential for life (O'Brien 2011). From our study, we observed that the administration of M- $\beta$ CD increased the iron levels in rats. It is known that excess iron can lead to the production of reactive oxygen species (ROS) that can damage cell structures. To prevent this harmful process, homeostasis is necessary to avoid tissue damage (Grotto 2008). In addition to the analysis of iron, another study highlighted the effects of CDs on blood components such as LDL. LDL comprises an important group of plasma lipoproteins that transport fats (mainly cholesterol) to cells. In this study, the administration of M- $\beta$ CD increased the serum levels of LDL without affecting the levels of total cholesterol, VLDL, HDL, and TG. However, an *in vitro* study found that CDs could induce lipid depletion of native LDL (Ao et al. 2016).

Oxidative stress occurs because of the overproduction of ROS and a failure in the function of antioxidant molecules (Dalle-Donne et al. 2003). Thiol compounds are reducing agents that contain a sulfhydryl or thiol group (-SH), and they are found in free form or linked to proteins in the biological environment. In this study, the total thiol content was increased 3-fold in the kidneys of animals treated with HP- $\beta$ CD. This significant increase was not reflected

in DCFH, TBARS, and carbonylation protein measurements. However,  $\beta$ CD and M- $\beta$ CD did not alter the parameters tested in this tissue. Based on the findings from renal tissue, the increase in serum levels of creatinine and urea in animals treated with  $\beta$ CD does not correlate with the findings of energy metabolism and oxidative stress.

The hepatic tissue exhibited an increase in the oxidation of DCFH in response to the administration of M- $\beta$ CD and HP- $\beta$ CD, indicating an increase in free radicals. The other parameters related to oxidative stress were not affected in this tissue. Inhibition of hepatic PK may be associated with increased reactive oxygen species and DCFH oxidation in response to M- $\beta$ CD. However, this analysis cannot inhibit the enzyme influenced by  $\beta$ CD. Similar to the results reported by Bellringer et al. (1995), these results indicate a possible kidney injury caused by  $\beta$ CD. However, these results do not rule out a possible liver injury even without altering ALT, gamma-glutamyl transferase, or alkaline phosphatase (indicators of liver injury). This possibility can be considered in this tissue based on the observation that the enzyme PK is inhibited, and this typically occurs in the presence of gluconeogenesis. We also assessed pancreatic function by measuring the lipase enzymes and glycemia, which were not affected by the treatments.

These results demonstrate that the  $\beta$ CDs used in this study did not affect most of the evaluated oxidative stress parameters. It is possible that CDs can be modulated by mechanisms other than oxidative stress. CDs have been widely used as drug carriers. Regardless, understanding the balance between the efficacy and adverse effects of using CDs is fundamental for a better acceptance of the clinical use of these compounds.

## Acknowledgments

We are sincerely grateful to Dc. Renata P. Raffin by HP- $\beta$ CD, and Dc. Paula Angélica Roratto by technical support. The financial support by Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Universidade Franciscana is gratefully acknowledged.

## REFERENCES

- AKSENOV MY & MARKESBERY WR. 2001. Changes in thiol content and expression of glutathione redox system genes in the hippocampus and cerebellum in Alzheimer's disease. *Neurosci Lett* 302: 141-145.
- AO M, GANA C, SHAO W, ZHOU X & CHENA Y. 2016. Effects of cyclodextrins on the structure of LDL and its susceptibility to copper-induced oxidation. *Eur J Pharm Sci* 25: 183-189.
- BELLINGER ME, SMITH TG, READ R, GOPINATH C & OLIVIERI P. 1995.  $\beta$ -cyclodextrin: 52-week toxicity studies in the rat and dog. *Food Chem Toxicol* 33: 367-376.
- BLAINE J, CHONCHOL M & LEVI M. 2015. Renal control of calcium, phosphate, and magnesium homeostasis. *Clin J Am Soc Nephrol* 10: 1257-1272.
- BORTOLUZZI VT, DE FRANCESCHI ID, RIEGER E & WANNMACHER CMD. 2014. Co-administration of creatine plus pyruvate prevents the effects of phenylalanine administration to female rats during pregnancy and lactation on enzymes activity of energy metabolism in cerebral cortex and hippocampus of the offspring. *Neurochem Res* 39: 1594-1602.
- BURGESS SC, IIZUKA K, JEOUNG NH, HARRIS RA, KASHIWAYA Y, VEECH RL, KITAZUME T & UYEDA K. 2008. Carbohydrate-response element-binding protein deletion alters substrate utilization producing an energy-deficient liver. *J Biol Chem* 283: 1670-1678.
- CHANDRASEKARAN S, SUDHA N, PREMNATH D & ENOCH IVMV. 2016. Binding of 6-methyl-3-phenyliminomethyl-4H-chromen-4-one with bovine serum albumin in free and  $\beta$ -cyclodextrin-complexed forms: modulation of the binding by  $\beta$ -cyclodextrin. *J Mol Liq* 222: 398-406.
- DALLE-DONNE I, ROSSI R, GIUSTARINI D, MILZANI A & COLOMBO R. 2003. Protein carbonyl groups as biomarkers of oxidative stress. *Clin Chim Acta* 329: 23-38.
- DE FRANCESCHI ID, RIEGER E, VARGAS AP, ROJAS DB, CAMPOS AG, RECH VC, FEKSA LR & WANNMACHER CMD. 2013. Effect of leucine administration to female rats during pregnancy and lactation on oxidative stress and enzymes activities of phosphoryltransfer network in cerebral cortex and hippocampus of the offspring. *Neurochem Res* 38: 632-643.
- DZEJA PP & TERZIC A. 2003. Phosphotransfer networks and cellular energetics. *J Exp Biol* 206: 2039-2047.
- DZEJA PP & TERZIC A. 2009. Adenylate kinase and amp signaling networks: metabolic monitoring, signal communication and body energy sensing. *Int J Mol Sci* 10: 1729-1772.
- DZEJA PP, VITKEVICIUS KT, REDFIELD MM, BURNETT JC & TERZIC A. 1999. Adenylate kinase-catalyzed phosphotransfer in the myocardium: increased contribution in heart failure. *Circ Res* 84: 1137-1143.
- FIGUEIREDO VC, FEKSA LR & WANNMACHER CMD. 2009. Cysteamine prevents inhibition of adenylate kinase caused by cystine in rat brain cortex. *Metab Brain Dis* 24: 373-381.
- GABARDO T, PERIPOLLI CM, DE ANDRADE RB, GEMELLI T, LIMA JDO, OLIVEIRA AS, MEDEIROS NS, WANNMACHER C, DANI C & FUNCHAL C. 2015. Assessment of changes in energy metabolism parameters provoked by carbon tetrachloride in Wistar rats and the protective effect of white grape juice. *Toxicol Rep* 2: 645-653.
- GAO Y, ZHAO X, DONG B, ZHENG L, LI N & ZHANG S. 2006. Inclusion complexes of  $\alpha$ -cyclodextrin with ionic liquid surfactants. *J Phys Chem B* 110: 8576-8581.
- GROTTO HZW. 2008. Metabolismo do ferro: uma revisão sobre os principais mecanismos envolvidos em sua homeostase. *Rev Bras Hematol Hemoter* 30: 390-397.
- HUGHES BP. 1962. A method for the estimation of serum creatine kinase and its use in comparing creatine kinase and aldolase activity in normal and pathological sera. *Clin Chim Acta* 7: 597-603.
- KEHRER JP. 2000. The Haber-Weiss reaction and mechanisms of toxicity. *Toxicol* 149: 43-50.
- LEBEL CP, ISCHIROPOULOS H & BONDY SC. 1992. Evaluation of the probe 2 $\beta$ ,7 $\beta$  dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chem Res Toxicol* 5: 227-231.
- LEONG SF, LAI JC, LIM L, CLARK JB. 1981. Energy-metabolising enzymes in brain regions of adult and aging rats. *J Neurochem* 37: 1548-1556.
- LEVINE RL, GARLAND D, OLIVER CN, AMICI I, CLIMENT AG, LENZ BW, AHN BW, SHALTIEL S & STADTMAN ER. 1990. Determination of carbonyl content in oxidatively modified proteins. *Meth Enzymol* 186: 464-478.

- LOFTSSON T & DUCHENE D. 2007. Cyclodextrins and their pharmaceutical applications. *Int J Pharm* 329: 1-11.
- LOWRY OH, ROSEBROUGH NJ, FARR AL & RANDALL RJ. 1951. Protein measurement with the Folin phenol reagent. *J Biol Chem* 193: 265-275.
- MACHADO FS, KUO J, WOHLBERG MF, FRUSCIANTE MR, FREITAS M, OLIVEIRA AS, ANDRADE RB, WANNMACHER CMD, DANI C & FUNCHAL C. 2016. Subchronic treatment with açai frozen pulp prevents the brain oxidative damage in rats with acute liver failure. *Metab Brain Dis* 31: 1427-1434.
- MELLO A, MEDEIROS MC, LACERDA DS, ANDRADE RB, GEMELLI T, GUERRA RB, WANNMACHER CMD, GOMEZ R & FUNCHAL C. 2012. Toxicological evaluation of chronic exposure to the organochalcogen 3-methyl-1-phenyl-2-(phenylseleno) oct-2-en-1-one in male rats. *Food Chem Toxicol* 50: 2450-2455.
- MOE SM. 2008. Disorders involving calcium, phosphorus, and magnesium. *Prim Care: Clinics Office Practice* 35: 215-237.
- MONTEIL M, LECOUEVEY M, LANDY D, RUELLAN S & MALLARD I. 2017. Cyclodextrins: A promising drug delivery vehicle for bisphosphonate. *Carbohydr Polym* 156: 285-293.
- MOSHER G & THOMPSON DO. 2002. Complexation and cyclodextrins. In Swaebrick J, Boylan J C. *Encyclopedia Pharm Technol* 1: 531-558.
- MURA P. 2020. Advantages of the combined use of cyclodextrins and nanocarriers in drug delivery: A review *Int J Pharm* 579: 119-181.
- NISHIHARA VSK ET AL. 2017. Resveratrol inclusion complex with  $\beta$ -cyclodextrin (rcd): characterization and evaluation of toxicity in wistar rats. *Int J Pharm Pharm Sci* 1: 27-33.
- O'BRIEN TX. 2011. Iron metabolism, anemia, and heart failure. *J Am Coll Cardiol* 58: 1252-1253.
- OHKAWA H, OHISHI N & YAGI K. 1979. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 95: 351-358.
- RECH VC, FEKSA LR, FLECK RM, ATHAYDES GA, DORNELLES PK, RODRIGUES-JUNIOR V & WANNMACHER CMD. 2008. Cysteamine prevents inhibition of thiol-containing enzymes caused by cystine or cystine dimethylester loading in rat brain cortex. *Metab Brain Dis* 23: 133-145.
- RECH VC, MEZZOMO NJ, ATHAYDES GA, FEKSA LR, FIGUEIREDO VC, KESSLER A, FRANCESCHI ID & WANNMACHER CMD. 2018. Thiol/disulfide status regulates the activity of thiol-containing kinases related to energy homeostasis in rat kidney. *An Acad Bras Cienc* 90: 99-108. <https://doi.org/10.1590/0001-3765201720160348>.
- REZNICK AZ & PACKER L. 1994. Oxidative damage to proteins: spectrophotometric method for carbonyl assay. *Meth Enzymol* 233: 357-363.
- SOUZA ACA ET AL. 2018. The inclusion complex of carvacrol and  $\beta$ -cyclodextrin reduces acute skeletal muscle inflammation and nociception in rats. *Pharm Reports* 70: 1139-1145.
- SOUZA TM, CORDEIRO RFB, VIANA M, AGUIAR LCS, DE SENNA LF, MALTA LFB & DÉLIA E. 2016. Inclusion compounds of dibenzylthiourea with hydroxypropylated-cyclodextrins for corrosion protection of carbon steel in acidic medium. *J Mol Struct* 1125: 331-339.
- SPANISH SOCIETY OF CLINICAL BIOCHEMISTRY AND MOLECULAR PATHOLOGY. 2012. Available from: <http://www.seqc.es/article/articleview/330/1/170>.
- TANG W, ZOU C, DA C, CAO Y & PENG H. 2020. A review on the recent development of cyclodextrin-based materials used in oilfield applications. *Carbohydrate Polymers* 240: 116-321.
- TRAUTWEIN EA, FORGBERT K, RIECKHOFF D & ERBERSDOBLER HF. 1999. Impact of L-cyclodextrin and resistant starch on bile acid metabolism and fecal steroid excretion in regard to their hypolipidemic action in hamsters. *Biochim Biophys Acta* 1437: 1-12.
- UEKAMA K, HIRAYAMA F & HIDETOSHI A. 2006. Recent aspect of cyclodextrin-based drug delivery system. *J Incl Phenom Macrocycl Chem* 56: 3-8.
- WALLIMANN T, WYSS M, BRDICZKA D, NICOLAY K & EPPENBERGER HM. 1992. Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: the 'phosphocreatine circuit' for cellular energy homeostasis. *Biochem J* 281: 21-40.
- YALCIN A, SODDU E, BAYRAKDAR ET, UYANIKGIL Y, KANIT L, ARMAGAN G, RASSU G, GAVINI E & GIUNCHEDI P. 2016. Neuroprotective effects of engineered polymeric nasal microspheres containing hydroxypropyl-b-cyclodextrin on b-amyloid (1-42) induced toxicity. *J Pharm Sci* 105: 2372-2380.
- YANG L, MA S, ZHOU S, CHEN W, YUAN M & YIN Y. 2013. Preparation and characterization of inclusion complexes of naringenin with  $\beta$ -cyclodextrin or its derivative. *Carbohydr Polym* 98: 861-869.

**How to cite**

IANISKI FR ET AL. 2023. β-Cyclodextrins alter the energy metabolism-related enzyme activities in rats. *An Acad Bras Cienc* 95: e20201783. DOI 10.1590/0001-3765202320201783.

*Manuscript received on November 18, 2020; accepted for publication on June 17, 2021*

**FRANCINE R. IANISKI<sup>1,2</sup>**

<https://orcid.org/0000-0002-8801-6322>

**AMANDA L. DE OLIVEIRA<sup>1#</sup>**

<https://orcid.org/0000-0002-7179-2450>

**NATHANA JAMILLE MEZZOMO<sup>3</sup>**

<https://orcid.org/0000-0001-7855-4343>

**ITIANE D. DE FRANCESCHI<sup>4</sup>**

<https://orcid.org/0000-0003-2093-5236>

**GUILHERME M. DO CARMO<sup>5</sup>**

<https://orcid.org/0000-0002-6907-2200>

**CAMILA R. CREMONESE<sup>1</sup>**

<https://orcid.org/0000-0002-1392-0936>

**MATHEUS D. BALDISSERA<sup>1</sup>**

<https://orcid.org/0000-0002-3280-8528>

**JENIFER P. ZANON<sup>1</sup>**

<https://orcid.org/0000-0002-4288-1307>

**JENIFER KOLLING<sup>1</sup>**

<https://orcid.org/0000-0002-1054-2116>

**JORDANA DANIELA FRIEDERICH<sup>1</sup>**

<https://orcid.org/0000-0002-1393-2136>

**IVANA Z. DA SILVA<sup>1</sup>**

<https://orcid.org/0000-0002-7067-5519>

**JANICE L. GIONGO<sup>6</sup>**

<https://orcid.org/0000-0002-2199-0838>

**LUCIANE R. FEKSA<sup>4</sup>**

<https://orcid.org/0000-0001-7080-0028>

**RODRIGO A. VAUCHER<sup>6</sup>**

<https://orcid.org/0000-0003-3496-7335>

**CLOVIS MILTON D. WANNMACHER<sup>4</sup>**

<https://orcid.org/0000-0002-7826-8416>

**VIRGINIA C. RECH<sup>2</sup>**

<https://orcid.org/0000-0003-1330-8031>

<sup>1</sup>Franciscan University, Nanotechnology Laboratory, Postgraduate Program in Nanoscience, Dos Andradas Street, nº 1614, Block I, 97010-032 Santa Maria, RS, Brazil

<sup>2</sup>Machado de Assis Integrated Colleges, Santos Dumont Street, nº 820, 98780-109 Santa Rosa, RS, Brazil

<sup>3</sup>Federal University of Santa Maria, Biochemistry and Molecular Biology Department, Natural and Exact Sciences Centre, Roraima Avenue, nº 1000, Camobi, 97105-900 Santa Maria, RS, Brazil

<sup>4</sup>Federal University of Rio Grande do Sul, Biochemistry Department, Basic Health Sciences Institute, Ramiro Barcelos Street, nº 2600, 90035-003 Porto Alegre, RS, Brazil

<sup>5</sup>University Center Univel, Pharmacy College, Tito Muffato Avenue, nº 2317, 85806-080 Cascavel, PR, Brazil

<sup>6</sup>Federal University of Pelotas, Postgraduate Program in Biochemistry and Bioprospecting, Capão do Leão Campus, s/n, 96050-500 Pelotas, RS, Brazil

Correspondence to: **Virginia Cielo Rech**

E-mail: [vga.cielo@gmail.com](mailto:vga.cielo@gmail.com)

#The data of this work are part of Amanda L. de Oliveira's master's dissertation

**Authors contributions**

V. C. Rech, A. L. de Oliveira and F. R. Ianiski, and I. Z. da Silva designed the research. A. L. de Oliveira, N. J. Mezzomo, G. M. do Carmo, C. R. Cremonese, M. D. Baldissera, J. P. Zanon, J. Kolling, J. D. Friederich, were involved in rat care and treatment. I. D. de Franceschi, A. L. de Oliveira, L. R. Feksa, and J. L. Giongo performed the measurements of enzymes of phosphoryl transfer network. N. J. Mezzomo, G. M. do Carmo, and R. A. Vaucher performed serum biochemical analyzes. I. D. de Franceschi, A. L. de Oliveira executed the measurements of oxidative stress. V. C. Rech and F. R. Ianiski performed the statistics analysis. V. C. Rech, L. R. Feksa, and C. D. Wannmacher collected and analyzed the data. V. C. Rech and F. R. Ianiski wrote the paper. V. C. Rech, F. R. Ianiski, R. A. Vaucher, I. Z. da Silva, L. R. Feksa, J. L. Giongo, and C. M. D. Wannmacher revised it critically the paper.

