cambridge.org/par

# **Research Article**

**Cite this article:** Hekimoglu O, Elverici C, Kuyucu AC (2023). Predicting climate-driven distribution shifts in *Hyalomma marginatum* (Ixodidae). *Parasitology* 1–11. https://doi.org/ 10.1017/S0031182023000689

Received: 9 April 2023 Revised: 2 June 2023 Accepted: 14 July 2023

#### **Keywords:**

climate change; Crimean-Congo haemorrhagic fever; ecological niche modelling; *Hyalomma marginatum*; MaxEnt; ticks

**Corresponding author:** Arda Cem Kuyucu; Email: ckuyucu@hacettepe.edu.tr

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution and reproduction, provided the original article



is properly cited.

# Predicting climate-driven distribution shifts in *Hyalomma marginatum* (Ixodidae)

Olcay Hekimoglu<sup>1</sup>, Can Elverici<sup>1,2</sup> 💿 and Arda Cem Kuyucu<sup>1</sup> 💿

<sup>1</sup>Biology Department, Hacettepe University, Ankara, Turkey and <sup>2</sup>Biodiversity Institute, University of Kansas, Lawrence, KS, USA

# Abstract

*Hyalomma marginatum* is an important tick species which is the main vector of Crimean-Congo haemorrhagic fever and spotted fever. The species is predominantly distributed in parts of southern Europe, North Africa and West Asia. However, due to ongoing climate change and increasing reports of *H. marginatum* in central and northern Europe, the expansion of this range poses a potential future risk. In this study, an ecological niche modelling approach to model the current and future climatic suitability of *H. marginatum* was followed. Using high-resolution climatic variables from the Chelsa dataset and an updated list of locations for *H. marginatum*, ecological niche models were constructed under current environmental conditions using MaxEnt for both current conditions and future projections under the ssp370 and ssp585 scenarios. Models show that the climatically suitable region for *H. marginatum* matches the current distributional area in the Mediterranean basin and West Asia. When applied to future projections, the models suggest a considerable expansion of *H. marginatum*'s range in the north in Europe as a result of rising temperatures. However, a decline in central Anatolia is also predicted, potentially due to the exacerbation of drought conditions in that region.

# Introduction

Human intervention in ecosystems, including globalization, urbanization, rapid transportation, land use and climate change, has led to a global increase in the range, distribution and transmission rate of numerous pathogens and vector organisms that serve as reservoirs for these pathogens (Kovats *et al.*, 2001; Andersen and Davis, 2017; Carvalho *et al.*, 2017; Semenza and Suk, 2018; Aguilar-Domínguez *et al.*, 2021; Leder *et al.*, 2021). While the most severe impacts of climate change are expected to affect ectotherm populations in the tropics, population growth rates of insects and other arthropods are also anticipated to increase in mid-to-high latitude regions due to the influence of warmer temperatures on growth rates (Deutsch *et al.*, 2008; Bonebrake and Deutsch, 2012; Rocklöv and Dubrow, 2020). In addition to the effects of temperature on the biology of disease-transmitting vector organisms, increased precipitation and extreme precipitation events in some regions may lead to an increase in vector abundance, whereas longer dry periods might increase tick mortality (Githeko *et al.*, 2000; Rocklöv and Dubrow, 2020).

Ticks are the vectors of numerous critical pathogens and pose a significant threat to animal and public health (Jongejan and Uilenberg, 2004). The species Hyalomma marginatum Koch, 1844 presents a substantial health threat as the main vector of Crimean-Congo haemorrhagic fever (CCHF) and also the vector of babesiosis and Rickettsia in the Mediterranean region, Africa and Asia (Ergönül, 2006; Vatansever et al., 2007; Gale et al., 2010; Ros-García et al., 2011; Bonnet et al., 2022; Sultankulova et al., 2022). Effective and sustainable control of ticks, like many arthropod pests and vectors, can be achieved with a solid understanding of the species' biology, ecology and distribution (Bonnet et al., 2022). Knowledge of H. marginatum's current distribution is crucial for projecting areas at risk for CCHF expansion. The Mediterranean basin holds special importance as the main distributional area and reservoir for *H. marginatum*. The Mediterranean is also among the areas projected to be most negatively affected by climate change (Ulbrich et al., 2006; Bardsley and Edwards-Jones, 2007; Newbold et al., 2020). Studies have indicated an increase in CCHF transmission in the Mediterranean basin (Maltezou and Papa, 2010) and a tendency for the northward expansion of CCHF's range in this region (Estrada-Peña and Venzal, 2007; Williams et al., 2015; Andersen and Davis, 2017; Fernández-Ruiz and Estrada-Peña, 2021). Recent reports have shown that the vector of CCHF, H. marginatum Koch, 1844 might have expanded its geographic range and shifted northwards into previously unoccupied areas (Vial et al., 2016; Bah et al., 2022). Furthermore, the expansion of suitable areas has been documented in central Europe and Balkans in addition to Mediterranean countries (Fernández-Ruiz and Estrada-Peña, 2021). Recent detections have highlighted that northern latitudes might be becoming more suitable for H. marginatum activity and survival as a result of climate change; however, most of these are records of individual tick detections and yet there is not sufficient evidence of establishment of northern populations (Duscher et al., 2018; Chitimia-Dobler et al., 2019; Grandi et al., 2020; McGinley et al., 2021).

Thanks to advances in machine-learning algorithms in ecology, such as MaxEnt (Phillips and Schapire, 2004; Elith et al., 2011; Crisci et al., 2012; Merow et al., 2013) and the advent of high-resolution climatic datasets such as Chelsa (Karger et al., 2017) and WorldClim (Fick and Hijmans, 2017), ecological niche modeling ENM has become a fundamental method for investigating possible current and future distributions of diseasetransmitting vectors over the past 2 decades (Chalghaf et al., 2018; Raghavan et al., 2020; Aguilar-Domínguez et al., 2021; Alkishe et al., 2021; Moo-Llanes et al., 2021). Correlative ecological niche models primarily rely on combining georeferenced species records with predictive environmental variables (e.g. climatic, topographic, soil) to build a coefficient matrix representing the organism's multidimensional niche (Warren and Seifert, 2011; Peterson and Soberón, 2012; Sillero and Barbosa, 2021). ENMs have been widely used to predict the potential future distributions of vectors under various global circulation scenarios of climate, in addition to their current potential niche (Aguilar-Domínguez et al., 2021; Alkishe and Peterson, 2022; Wu et al., 2022).

The main objective of this study was to predict climatically suitable areas for *H. marginatum* under both present-day conditions and future climate scenarios. To achieve this aim, bioclimatic parameters for present and near-future scenarios (2011–2040 and 2041–2070) obtained from the Chelsa database and *H. marginatum* locations from the literature and previous field surveys by the authors were used to build ecological niche models with MaxEnt to show suitable areas for present and future projections. The findings of this study will help predict potential new suitable areas in the Mediterranean basin and Europe that *H. marginatum* may colonize in the future and enhance surveil-lance efforts in areas identified as high risk.

#### **Materials and methods**

# Occurrence records

Local data from Turkey were derived from 2 sources: (1) from published studies by first author (O. H.) until 2021, which used both morphological and molecular methods to identify tick samples (Hekimoglu and Ozer, 2017; Hekimoglu et al., 2021) and additional field collected samples from 2018 (n = 13 locations) and 2021(n = 7 locations) which have been identified using 16s rDNA. Morphological identification was conducted using taxonomic keys (Apanaskevich and Horak, 2008; Estrada-Peña et al., 2018). Since H. marginatum populations show morphological variations (Apanaskevich and Horak, 2008) and Hyalomma species have been one of the most misidentified taxa using morphological methods (Estrada-Pena and De La Fuente, 2016), conducting a molecular step is extremely important. For this purpose, mitochondrial 16S rDNA sequences (Mangold et al., 1998) were generated for samples of localities where molecular data were not generated before. In total, 66 geographical points from Turkey were included in the analyses.

To represent the distribution area of the species worldwide, we included geographic locations of *H. marginatum* from different parts of the world from a previous compilation by Estrada-Pena and De La Fuente (2016), composed of literature reviews between 1970 and 2014. Firstly, the geographic information of *H. marginatum* was recorded on a separate data sheet. Secondly, these raw records were cleaned and reduced by removing localities, (1) where tick collection had been conducted from birds since finding ticks on birds does not mean that ticks can establish populations in these areas (Estrada-Peña *et al.*, 2011; Fernández-Ruiz and Estrada-Peña, 2021); and (2) sampling information from Cyprus was removed as molecular data do not confirm that *H. marginatum* exists in Cyprus (Hekimoglu and Ozer, 2017). After these steps, a total of 565 geographic coordinates were obtained.

## Environmental variables

For environmental predictors, Chelsa V2.1 dataset was used (Karger et al., 2017, 2022) which is available at https://chelsaclimate.org, a relatively new high-resolution (30 arcsec,  $\sim$ 1 km) climate dataset that includes additional important microclimatic variables in addition to the counterparts (Chelsa-Bioclim) of popular WorldClim bioclim variables (Hijmans et al., 2005; Fick and Hijmans, 2017). These include variables related to microclimate, water content of air and humidity which are important for H. marginatum (Estrada-Peña et al., 2011; Estrada-Peña, 2023). Furthermore, while the WorldClim dataset used for current distribution predictions uses extrapolations made between 1970 and 2000, the Chelsa dataset uses a dataset created for the period between 1980 and 2010. The additional important variables (Chelsa-Bioclim+) included near surface relative humidity (hurs), vapour pressure deficit (vpd), climate moisture index (cmi), growth degree days above 0°C, net primary productivity (npp) and surface downwelling shortwave radiation (rsds). Growth season temperature (gst) and growth season precipitation (gsp) were also included, which are indicative of H. marginatum, whose activity coincides greatly with the growth season (Estrada-Peña et al., 2011). Bioclim variables 8, 9, 18 and 19 from the Worldclim dataset (Fick and Hijmans, 2017) are reported to have significant spatial artefact problems visible as anomalous discontinuities between neighbouring pixels and thus it is generally advised to remove them before carrying out analyses (Escobar et al., 2014; Escobar, 2020; Aguilar-Domínguez et al., 2021; Alkishe and Peterson, 2022). In this study, these variables were removed from the Chelsa dataset since it was observed that the same artefacts are also present in Chelsa-Bioclim data. All remaining variable rasters were clipped to the area of interest, which is between latitudes  $-20^{\circ}$  and  $60^{\circ}$ and longitudes 20° and 60° WGS84. All geographic computations were carried out with QGIS (QGIS Geographic Information System, 2022), GDAL library in Python (Open Source Geospatial Foundation, 2022) and R version 4.2.2 (R Core Team, 2022).

## Future projections

For future predictions, environmental variables from different global circulation models (GCM) of ssp370 and ssp585 scenarios for CMIP6 (Eyring et al., 2016) were used. 'Shared socioeconomic pathways' (SSPs) complement the 'representative concentration pathways' (RCPs) by involving socio-economic factors such as population growth, urbanization and technological advances, ssp585 represents the worst-case scenario while ssp370 represents a middle way between the worst and more optimistic scenarios (Hausfather, 2020). These bad-to-worst case scenarios were chosen for a more prudent risk prediction as anthropogenic climate risks are becoming much more difficult to manage and complicated all around the world (Kemp et al., 2022). All of the available scenarios for Chelsa dataset were included for a better estimation. These include UKESM1-0-LL (Sellar et al., 2019), MRI-ESM2-0 (Oshima et al., 2020), MPI-ESM1-2 (Mauritsen et al., 2019), IPSL-CM6-LR (Boucher et al., 2020) and GFDL-ESM4 (Dunne et al., 2020) for the Chelsa dataset, and also available at https:// chelsa-climate.org.

# Ecological niche modelling

Before building the models, occurrence records that fell outside environmental raster pixels were removed. In order to reduce spatial autocorrelation, the locations for *H. marginatum* were thinned to a 5 km radius using the spThin package for R (Aiello-Lammens et al., 2015). The resulting occurrence data consisted of 470 locations (Fig. 1) which were split into a training set (70%: n = 335), a preliminary test set (cross-validation set) for evaluating the candidate models (25%: n = 111) and another secondary independent test set for evaluation of final models (5%: n = 24). To simulate the accessible M space (Soberon and Peterson, 2005; Barve et al., 2011) for H. marginatum, a buffered minimum convex polygon with a buffer area of 100 km around the occurrence records was created and environmental variable rasters representing the M space were clipped to this buffer area before building models. Thinning and the creation of the buffer zone were carried out with ellipsenm package for R, available at https://github.com/marlonecobos/ellipsenm (Cobos et al., 2020). Three sets were built by setting correlation thresholds between variables using the vifcor function in the R package usdm (Naimi, 2017). The vifcor function selects the variables with lower variance inflation factors (vif) from correlation pairs to build the sets. The correlation thresholds were 0.9, 0.8 and 0.75, respectively. For the first model dataset sets 2 and 3 were identical, so only set 1 and set 2 were used to build the models. For the Chelsa dataset some of the Bioclim+ variables are not available for future projections (vpd, hurs, rsds and cmi). Thus, 3 different sets were built not including these variables to build a separate second model (model 2) for future projections.

ENMs were built with the maximum entropy algorithm (Phillips and Schapire, 2004) using MaxEnt 3.4.4 (Steven *et al.*, 2021) implemented in the Kuenm package (Cobos *et al.*, 2019) for R. Using the Kuenm package, calibration models were built with all combinations of MaxEnt features L (linear), Q (quadratic), P (product) and H (hinge). All models were repeated with regularization multiplier parameters 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4 and 5. The performance of all these models were evaluated primarily with the partial receiver operating characteristics (pROC) test and an omission threshold rate of 5% (Peterson and Soberón, 2012; Aguilar-Domínguez *et al.*, 2021), secondary evaluations were done with Akaike information

criterion corrected for small sample size (AICc). The selected eventual calibration models were statistically significant, that have omission rates below 5% and with  $\Delta$ AICc values below 2 (Warren and Seifert, 2011; Nuñez-Penichet *et al.*, 2021).

After model calibrations, final models were created with the selected parameters in the resulting calibration models using all the same occurrences of training and testing data with 10 boot-strap replicates. Then, these final models were additionally evaluated with the independent location data (n = 24) that were not used for building and selecting these models.

Map binarization was done by using the average logistic thresholds of maximum training sensitivity plus specificity of 10 replicates (Liu et al., 2013). Consensus maps of the niche models for the future projections were created by taking the average of GCMs. In order to show the effects of model projections under novel future conditions, we transferred models with allowed projections in novel climate conditions (extrapolation with clamping to the GCM scenarios) (Cobos et al., 2019). To see the places where conditions are more extreme compared to the calibration area of the models, a mobility-oriented parity metric (MOP) analysis was carried out and extrapolation risk of transfer regions was calculated with nearest 10% reference (Owens et al., 2013; Alkishe et al., 2020; Flores-López et al., 2022). Because of memory limitations, resolution of calibration area and transfer rasters were changed from 30 s to 2.5 min resolution and MOP analysis were carried out with the mop function included in the kuenm package (Cobos et al., 2019). Again, consensus maps for MOP were created taking the average of all GCMs for each ssp scenario and 4 MOP maps were created in total.

#### **Results**

# Model parameters

Environmental parameters selected based on correlation thresholds and vif for variable sets used in model calibrations are shown in Table 1. In total 420 candidate models were created




Figure 1. All occurrence points of Hyalomma marginatum after cleaning and thinning.

	Model 1		Model 2		
	Set 1	Set 2	Set 1	Set 2	Set 3
Bio3 (isothermality)	$\checkmark$	$\checkmark$	$\checkmark$	$\checkmark$	х
Bio4 (temperature seasonality)	$\checkmark$	x	$\checkmark$	x	х
Bio6 (min. air temp. of the coldest month)	$\checkmark$	х	$\checkmark$	х	х
Bio7 (annual range of air temperature)	$\checkmark$		$\checkmark$	$\checkmark$	$\checkmark$
Bio13 (precipitation amount of the wettest month)	$\checkmark$		$\checkmark$	$\checkmark$	$\checkmark$
Bio14 (precipitation amount of the driest month)	$\checkmark$		$\checkmark$	$\checkmark$	х
Bio15 (precipitation seasonality)	$\checkmark$		$\checkmark$	$\checkmark$	$\checkmark$
Gdd0 (growing degree days heat sum above 0°C)	x	x	$\checkmark$	х	х
Gst (mean temperature of the growing season)	$\checkmark$			$\checkmark$	$\checkmark$
Gsp (precipitation amount of the growing season)	$\checkmark$	x	$\checkmark$	х	х
Net primary productivity (npp)	$\checkmark$	х		х	х
Surface downwelling shortwave radiation (rsds)		x	x	х	х
Climate moisture index (cmi)	$\checkmark$	х	х	х	х

Table 1. Environmental predictors used for different sets in model 1 and model 2 with explanation of the predictors in the first column

for the first model (model 1). All of 420 candidate models were statistically significant (pROC test P < 0.05); however, of these only 82 fulfilled the omission rate criteria of 5% (omission rate = 3.6%). Only one of these met the AICc criteria of  $\leq 2$ ( $\Delta AICc = 0.90$ ). This final selected model is a linear quadratic model with a regularization multiplier parameter of 0.2, which uses variables included in set 1 with a mean area under the curve (AUC) ratio of 1.21. The final evaluation with the independent test set resulted in an omission rate of 8.3%. Although this value is larger than the 5% threshold set for evaluations with the cross-validation sets, the independent test set had a sample size of n = 24, and 8.3% omission rate means that the final model classified 22 of 24 test locations correctly which is a pretty good result for a machine learning algorithm. The percent contribution and permutation importance of 12 predictors are shown in Table 2. Out of these, the most important parameter was cmi with

Model 1		1	Model 2		
Predictor	Percent contribution	Predictor	Percent contribution		
Cmi	33.1	Bio13	31.3		
Bio14	24.2	Bio14	19.4		
Rsds	8.4	Gdd0	13		
Bio6	7	Bio6	8.2		
Bio15	6	Npp	8		
Bio13	5.7	Bio15	7.5		
Gst	4.2	Bio4	4.9		
Npp	4.1	Bio3	4.5		
Bio3	2.1	Gst	2.2		
Bio7	1.8	Bio7	1.1		
Bio4	1.8	Gsp	$\approx 0$		
Gsp	1.5				

a contribution of 33.1%, followed by Bio14 (precipitation amount of the driest month) with a 24.2% contribution and rsds with an 8.4% contribution. Bio6 (minimum air temperature of the coldest month) was the fourth most important predictor (7% contribution).

Out of 630 candidate models for the second Chelsa model (model 2) built for future projection, 625 were statistically significant (pROC test P < 0.05). Out of these, 201 met the 5% omission rate criteria ( $\approx 4.5\%$ ), and only 2 of them met the AICc criteria: a PH (product hinge) model with a regularization multiplier of 2  $(\Delta AICc \approx 0)$ , and a QPH (quadratic product hinge) model with a regularization multiplier of 2 ( $\Delta AICc \approx 1.27$ ). Both of these final calibration models used the set 1 variables created for the second model. Mean AUC values were 1.23 and 1.20 for the product hinge and quadratic product hinge models, respectively. The final evaluation with the independent test set showed omission rates 8.3% and 12.5% for the PH and QPH models, respectively, indicating that the PH model is better for predictions. For the second model, there were 11 predictors. The biggest contributors were Bio13 (precipitation amount of wettest month) with 31.3% followed by Bio14 (19.4% contribution) and Gdd0 (growth degree days above 0 °C) with a 13% contribution. Like model 1, Bio6 was the fourth important predictor for model 2 with 8.2% contribution. Threshold values for creating binary maps were 0.409 and 0.483, respectively.

# Current and future predictions

Suitable areas for present conditions are shown in Fig. 2. According to the first model, the current climatically suitable region for *H. marginatum* stretches out from Iberian Peninsula to Anatolia and the Caspian Sea. This region includes most of the Mediterranean basin. This pattern mostly coincides with the reported locations of *H. marginatum*. The predictions of 2 models are mostly compatible with each other. Compared to the first model, the prediction of the second model for the current distribution shows a wider suitable area in some regions, especially in North Africa and near Hungary. Also, the first model shows wider suitable regions in Trans-Caucasus and Spain compared to the second model. This might be due to the fact that the second model does not include parameters like cmi and rsds which are




Figure 2. Maps of predicted suitable areas from the ENM results. (A) Red areas show the suitable regions under current conditions according to model 1. (B) Green areas show the suitable regions under current conditions for model 2.

present in the first model. On the contrary, the second model includes another important parameter, gdd0, which was eliminated from the first model in the correlation and vif selection procedure.

The predicted changes in suitable areas under GCM scenarios compared to present conditions are also shown in Fig. 3. Future projections under ssp370 scenarios for 2011-2040 show a significant widening of the climatically suitable area. The direction of the expansion is northwards in Europe, westwards in Anatolia and southwards in North African regions. Interestingly, the projections for the 2041-2070 period do not show such widening compared to the 2011-2040 period in eastern Europe and Anatolia; for example, the patches of new barely suitable regions in Baltic regions and Russia are lost in the 2041-2070 scenarios. In addition to that, there are significant declines in some regions in both ssp370 and ssp585 scenarios; these areas of suitability decline are especially distinct in central Anatolia and some areas in Balkans and central Europe. While both ssp370 and ssp585 show similar patterns in new areas for the 2041-2070 period, ssp585 scenarios present a wider new region of low suitability in France and eastern Germany. The results of MOP analysis done for between calibration area and future projections are shown in Fig. 4. High extrapolation risk areas are located relatively far from the current distribution and projected shifting areas of H. marginatum and the regions with high or strict extrapolation start from the north and east of Caspian Sea, Middle East (Arabian Peninsula, Israel and Egypt) and Africa (excluding northwest Africa).

# Discussion

The outputs from the first model show that the main climatic limiting factors for H. marginatum are cmi and precipitation, followed by surface radiation and minimum temperatures. Relative humidity or moisture is crucial for ticks, as they are highly susceptible to cuticular water loss in their off-host state (Knülle and Rudolph, 1982; Benoit and Denlinger, 2010; Estrada-Peña et al., 2012; Requena-García et al., 2017; Leal et al., 2020). The importance of water content of air for H. marginatum has also been in previous habitat suitability demonstrated models (Estrada-Peña et al., 2011; Fernández-Ruiz and Estrada-Peña, 2021). Another important parameter is incoming surface radiation, which is a relatively neglected parameter in ENMs for ticks despite the fact that it is a critical parameter for small arthropods (Davis et al., 2005; Battisti et al., 2013; Barrett and O'Donnell, 2023). Solar radiation might affect many different parameters related to ticks. Firstly, it is the main driver of microclimate temperature, and higher levels of solar radiation might increase tick abundance and questing activity (Kiewra et al., 2014; Del Fabbro et al., 2015). However, due to ultraviolet-B (UVB) radiation, very high levels of solar radiation might also have deleterious effects on small ectotherms like acari (Sakai et al., 2012; Sudo and Osakabe, 2015).

The predicted area for the current conditions in model 1 is mostly compatible with previous habitat suitability models created for *H. marginatum* (Williams *et al.*, 2015; Fernández-Ruiz and Estrada-Peña, 2021). Additionally, the model output shows




Figure 3. Maps of predicted suitable areas for future average of 5 GCM scenarios with differing degrees of loss and gain compared to current conditions for model 2. (A) For ssp370 in the 2011–2040 period. (B) For ssp370 in the 2041–2070 period (C) for ssp585 in the 2011–2040 period (D) for ssp585 in the 2041–2070 period.




Figure 4. Extrapolation risk in 4 GCM projections of *H. marginatum* with MOP10%. Green to black scale shows increasing risk extrapolation where black areas are regions with strict extrapolation. (A) For ssp370 in the 2011–2040. (B) For ssp370 in the 2041–2070 period. (C) For ssp585 in the 2011–2040 period. (D) For ssp585 in the 2041–2070 period.

potential suitable areas in Trans-Caucasus and Iran, especially in northern Iran near the coast of the Caspian Sea, despite that the dataset of locations used for the models did not include any locations from this region. Although no exact coordinates are presented in the literature, CCHF cases and also *H. marginatum* ticks carrying the virus have been reported from this area (Shemshad *et al.*, 2012; Sofizadeh *et al.*, 2014; Telmadarraiy *et al.*, 2015; Sedaghat *et al.*, 2017).

Model 2 output shows a larger climatically suitable area compared to the more restrictive predictions of model 1. Important parameters like moisture index (cmi) and surface radiation (rsds) were left out for this model, as they are not available for future scenarios. Moisture index is replaced by precipitation of the wettest period (Bio13) as the most important parameter in model 2, which was in sixth place in the previous model. Although net precipitation amount is not as strong as water deficit or moisture index as a predictor, this shows the importance of the water balance for climatic suitability. In addition, surface radiation is replaced by gdd0 (growth degree days above 0 °C), which was eliminated from model 1 due to correlation threshold with other parameters. Fulfilling the required degree days is an important necessity for *H. marginatum* populations to establish in a new region (Estrada-Peña, 2023). It has been reported that a yearly accumulated temperature of 3000–4000 °C is necessary for this species (Estrada-Peña et al., 2011); a northern limit roughly coincides with 47°N.

With the increased temperature in recent decades, new reports of *H. marginatum* have also been increasing in Europe (Hornok and Horváth, 2012; Duscher et al., 2018; Bah et al., 2022; Lesiczka et al., 2022). Gillingham et al. (2023), who modelled the probability of survival and establishment of H. marginatum populations depending on cumulative degree days for the United Kingdom, forecasted increased risk in southern UK in the future. Also, the prediction maps for the climatically suitable area for ssp370 and ssp585 scenarios point to a northward expansion in Europe in the future. Depending on the models trained with trends in climate and tick records between 1970 and 2018, Fernández-Ruiz and Estrada-Peña (2021), assumed that the suitable climatic area in Europe would expand while maintaining the original Mediterranean distribution. Additionally, Williams et al. (2015) also predicted a northward expansion under AR5 climate scenarios in the future. Another parameter that should be considered is minimum temperatures, which occupied fourth place in both models 1 and 2. In addition to the above-mentioned effect of temperature (degree days) on development time, overwintering survival is another important factor for H. marginatum, as most adults overwinter in the field (Valcárcel et al., 2020). An increase in the minimum temperatures in the field might increase winter survival and contribute to the ongoing and possible future expansion in the northern limits of H. marginatum populations (Estrada-Peña et al., 2012). Projections for future distributions are compatible with these predictions, showing a very similar pattern of new suitable regions in North Spain, France, the Balkans and western Anatolia. The projections for 2041-2070 point to a continuation of this expansion; however, the projection also points to significant declines in some regions, mostly in central Anatolia and the Balkans. This is most probably due to the decrease in precipitation, as climatic simulations predict that the impact of decreased precipitation will be much higher in the eastern Mediterranean, especially in Turkey (Hoerling et al., 2012; Turkes et al., 2020). It was previously suggested that the northern limit for *H. marginatum* is thought to be mainly determined by temperature, while the southern limit is determined by precipitation and humidity (Gray et al., 2009), and expansion in the north and contraction in the south is an expected outcome in the future.

It also has to be considered that in some regions, the change in suitability might be more complex than this pattern. With projected ongoing climate change, an interchange of climatically suitable and unsuitable areas can be seen in models (Gillingham *et al.*, 2023). A previous model using RCP8.5 scenarios carried out locally for Romania depicted an increase in suitable areas until 2050, followed by a decreasing suitability until 2070 (Domşa *et al.*, 2016), which was also detected by future predictions in the current study (Fig. 3). This rather complex pattern in suitability might be due to the 2 main factors (temperature and precipitation) determining suitability. For instance, present-day models show a highly suitable region for *H. marginatum* in the central and central-north parts of Turkey. Contrasting with the relatively more northward expansion in Europe, future projections indicate that after 2040, this suitable area might shift towards coastal areas where required precipitation would be available compared to the now more arid continental regions.

Assessing the potential distribution range of ticks is important to predict the risk of emergence and re-emergence of tick-borne diseases (Estrada-Peña, 2008; Zhao et al., 2021). In this sense, the interaction between ticks and climate is extremely important, as the survival and biological functions of ticks depend strongly on the external micro-abiotic environment (Gray et al., 2009). It should also be kept in mind that finding a suitable host is one of the most important drivers, which affects the distribution of tick species in an area. On the contrary, H. marginatum is a 2-host species and uses a wide range of vertebrates at different stages of its life cycle; this complex behaviour of this species is another reason for the difficulties in predicting the future distributions of these vectors (Apanaskevich and Horak, 2008; Valcárcel et al., 2020; Bonnet et al., 2022). For instance, an area might be classified as suitable for a species by an ENM in future, but the absence or scarce distribution of suitable hosts would prevent the establishment of populations in the area. Conversely, unexpected novel hosts in new environments would provide the necessary link for dispersal in new environments (Bakkes et al., 2021). Additionally, landscape cover, configuration and habitat fragmentation are other potential factors on the distribution of ticks and transmission of tick-borne diseases (Suzán et al., 2012; Perez et al., 2016). A previous study showed that in Turkey, the number of CCHF cases was associated with high landscape fragmentation and connectivity (Estrada-Pena et al., 2010). On the contrary, Bah et al. (2022) showed that in southern France, sclerophyllous vegetated or sparsely vegetated open natural areas present favourable habitats for H. marginatum rather than more humid or urbanized areas, a similar pattern is also present in Anatolia where the highest distribution is in the centralnorth while H. marginatum is significantly absent both in humid and densely forested Black Sea coast and hot and dry southern regions. Ecological niche models are generally more deterministic models that build the representative fundamental niche using presently available abiotic parameters and locations, then project this assumed niche space to assumed future abiotic parameter data calculated by predictive global circulation simulations (Escobar, 2020; Sillero et al., 2021). Thus, these projections are for predicting the climatically suitable areas in the predicted future under assumed scenarios. However, these kinds of models are among the most useful tools presently available for estimating the potential risk of vector-borne diseases and deciding the necessary precautions (Medlock and Leach, 2015, Ortega-Guzmán et al., 2022). MOP analysis is one of the tools to interpret the potential risk of uncertainty for transfer of models to novel conditions (Owens et al., 2013; Cobos et al., 2019). MOP analysis showed that the current and predicted future distribution areas did not include areas with high risk of extrapolation. Uncertainty maps for H. marginatum are similar to the Europe and North Africa projections of a previous uncertainty analysis of worldwide projections of the distribution of another tick

species (*Rhipicephalus sanguineus*) to RCP scenarios done by Alkishe *et al.* (2020). The main difference is that the present MOP results of projections to ssp scenarios show new and wider areas with high extrapolation in North Africa and Middle East compared to the previous paper by Alkishe *et al.* (2020).

With the ongoing human-induced changes in the biotic and abiotic environment we have been observing a significant increase in dispersal and abundance of H. marginatum and also an increase in the transmission of pathogens carried by H. marginatum. Future distribution projections indicate a significant increase in potential risk due to these factors, so it is important to build both statistical and process-based mechanistic models with new data. Also publishing coordinates is strongly suggested, as it is observed that there are many studies which report species presence without any coordinates that can help future coordination between researchers and provide more reliable models. As H. marginatum is the vector of CCHF, change in predicted distributional range might point out potential new risk areas or widening of current endemic areas. This study may be helpful to forecast new risk areas and therefore to expand awareness and to start well-adapted prevention strategies against CCHF in these areas. Similarly, our models show that the climatically suitable region for H. marginatum matches the current distributional area, which can be interpreted as the need to strengthen and to maintain control measures to CCHF in the future.

**Data availability.** Locations used in this study are available at https://data. mendeley.com/datasets/n8b3kps7nk/1.

Acknowledgements. The numerical calculations reported in this paper were partially performed at TUBITAK ULAKBIM, High Performance and Grid Computing Center (TRUBA resources). We also thank 3 anonymous reviewers for reviewing the paper and for their helpful suggestions.

Author's contributions. O. H. and A. C. K. designed the study; O. H. prepared and cleaned the data; A. C. K. and C. E. carried out the analysis and interpreted the results; A. C. K. and C. E. prepared the tables and maps; A. C. K. prepared the original draft; O. H., A. C. K. and C. E. reviewed and edited the manuscript.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interest. None.

## References

- Aguilar-Domínguez M, Moo-Llanes DA, Sánchez-Montes S, Becker I, Feria-Arroyo TP, de León AP and Romero-Salas D (2021) Potential distribution of *Amblyomma mixtum* (Koch, 1844) in climate change scenarios in the Americas. *Ticks and Tick-borne Diseases* **12**, 101812.
- Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B and Anderson RP (2015) spThin: an R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38, 541–545.
- Alkishe A, Cobos ME, Peterson AT and Samy AM (2020) Recognizing sources of uncertainty in disease vector ecological niche models: An example with the tick Rhipicephalus sanguineus sensu lato. Perspectives in Ecology and Conservation, 18, 91–102.
- Alkishe A and Peterson AT (2022) Climate change influences on the geographic distributional potential of the spotted fever vectors Amblyomma maculatum and Dermacentor andersoni. PeerJ 10, e13279.
- Alkishe A, Raghavan RK and Peterson AT (2021) Likely geographic distributional shifts among medically important tick species and tick-associated diseases under climate change in North America: a review. *Insects* 12, 225.
- Andersen LK and Davis MDP (2017) Climate change and the epidemiology of selected tick-borne and mosquito-borne diseases: update from the International Society of Dermatology Climate Change Task Force. *International Journal of Dermatology* 56, 252–259.
- Apanaskevich DA and Horak IG (2008) The genus Hyalomma Koch, 1844: V. Re-evaluation of the taxonomic rank of taxa comprising the H. (Euhyalomma) marginatum Koch complex of species (Acari: Ixodidae)

with redescription of all parasitic stages and notes on biology. *International Journal of Acarology* **34**, 13–42.

- Bah MT, Grosbois V, Stachurski F, Muñoz F, Duhayon M, Rakotoarivony I, Appelgren A, Calloix C, Noguera L, Mouillaud T, Andary C, Lancelot R, Huber K, Garros C, Leblond A and Vial L (2022) The Crimean–Congo haemorrhagic fever tick vector *Hyalomma marginatum* in the south of France: modelling its distribution and determination of factors influencing its establishment in a newly invaded area. *Transboundary and Emerging Diseases* 69, e2351–e2365.
- Bakkes DK, Ropiquet A, Chitimia-Dobler L, Matloa DE, Apanaskevich DA, Horak IG, Mans BJ and Matthee CA (2021) Adaptive radiation and speciation in *Rhipicephalus ticks*: a medley of novel hosts, nested predator-prey food webs, off-host periods and dispersal along temperature variation gradients. *Molecular Phylogenetics and Evolution* 162, 107178.
- **Bardsley DK and Edwards-Jones G** (2007) Invasive species policy and climate change: social perceptions of environmental change in the Mediterranean. *Environmental Science and Policy* **10**, 230–242.
- **Barrett M and O'Donnell S** (2023) Individual reflectance of solar radiation confers a thermoregulatory benefit to dimorphic male bees (*Centris pallida*) using distinct microclimates. *PLoS ONE* **18**, e0271250.
- Barve N, Barve V, Jiménez-Valverde A, Lira-Noriega A, Maher SP, Peterson AT, Soberón J and Villalobos F (2011) The crucial role of the accessible area in ecological niche modeling and species distribution modeling. *Ecological Modelling* 222, 1810–1819.
- Battisti A, Marini L, Pitacco A and Larsson S (2013) Solar radiation directly affects larval performance of a forest insect. *Ecological Entomology* 38, 553–559.
- Benoit JB and Denlinger DL (2010) Meeting the challenges of on-host and off-host water balance in blood-feeding arthropods. *Journal of Insect Physiology* 56, 1366–1376.
- Bonebrake TC and Deutsch CA (2012) Climate heterogeneity modulates impact of warming on tropical insects. *Ecology* **93**, 449–455.
- Bonnet SI, Vourc'h G, Raffetin A, Falchi A, Figoni J, Fite J, Hoch T, Moutailler S and Quillery E (2022) The control of *Hyalomma* ticks, vectors of the Crimean–Congo hemorrhagic fever virus: where are we now and where are we going? *PLoS Neglected Tropical Diseases* 16, e0010846.
- Boucher O, Servonnat J, Albright AL, Aumont O, Balkanski Y, Bastrikov V, Bekki S, Bonnet R, Bony S, Bopp L, Braconnot P, Brockmann P, Cadule P, Caubel A, Cheruy F, Codron F, Cozic A, Cugnet D, D'Andrea F, Davini P, de Lavergne C, Denvil S, Deshayes J, Devilliers M, Ducharne A, Dufresne JL, Dupont E, Éthé C, Fairhead L, Falletti L, Flavoni S, Foujols MA, Gardoll S, Gastineau G, Ghattas J, Grandpeix JY, Guenet B, Guez LE, Guilyardi E, Guimberteau M, Hauglustaine D, Hourdin F, Idelkadi A, Joussaume S, Kageyama M, Khodri M, Krinner G, Lebas N, Levavasseur G, Lévy C, Li L, Lott F, Lurton T, Luyssaert S, Madec G, Madeleine JB, Maignan F, Marchand M, Marti O, Mellul L, Meurdesoif Y, Mignot J, Musat I, Ottlé C, Peylin P, Planton Y, Polcher J, Rio C, Rochetin N, Rousset C, Sepulchre P, Sima A, Swingedouw D, Thiéblemont R, Traore AK, Vancoppenolle M, Vial J, Vialard J, Viovy N and Vuichard N (2020) Presentation and evaluation of the IPSL-CM6A-LR climate model. Journal of Advances in Modeling Earth Systems 12, e2019MS002010.
- Carvalho BM, Rangel EF and Vale MM (2017) Evaluation of the impacts of climate change on disease vectors through ecological niche modelling. *Bulletin of Entomological Research* **107**, 419–430.
- Chalghaf B, Chemkhi J, Mayala B, Harrabi M, Benie GB, Michael E and Ben Salah A (2018) Ecological niche modeling predicting the potential distribution of *Leishmania* vectors in the Mediterranean basin: impact of climate change. *Parasites and Vectors* 11, 1–9.
- Chitimia-Dobler L, Schaper S, Rieß R, Bitterwolf K, Frangoulidis D, Bestehorn M, Springer A, Oehme R, Drehmann M, Lindau A, Mackenstedt U, Strube C and Dobler G (2019) Imported Hyalomma ticks in Germany in 2018. Parasites and Vectors 12, 1–9.
- Cobos ME, Townsend Peterson A, Barve N and Osorio-Olvera L (2019) Kuenm: an R package for detailed development of ecological niche models using MaxEnt. *PeerJ* 2019, e6281.
- Cobos ME, Osorio-Olvera L, Soberón J, Peterson AT, Barve V and Barve N (2020) ellipsenm: Ecological niche's characterizations using ellipsoids. R. package version 0.34. Available at https://github.com/marlonecobos/ellipsenm
- Crisci C, Ghattas B and Perera G (2012) A review of supervised machine learning algorithms and their applications to ecological data. *Ecological Modelling* 240, 113–122.

- Davis AK, Farrey BD and Altizer S (2005) Variation in thermally induced melanism in monarch butterflies (Lepidoptera: Nymphalidae) from three North American populations. *Journal of Thermal Biology* **30**, 410–421.
- **Del Fabbro S, Gollino S, Zuliani M and Nazzi F** (2015) Investigating the relationship between environmental factors and tick abundance in a small, highly heterogeneous region. *Journal of Vector Ecology* **40**, 107–116.
- Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC and Martin PR (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences* 105, 6668–6672.
- Domşa C, Sándor AD and Mihalca AD (2016) Climate change and species distribution: possible scenarios for thermophilic ticks in Romania. *Geospatial Health* 11, 151–156.
- Dunne JP, Horowitz LW, Adcroft AJ, Ginoux P, Held IM, John JG, Krasting JP, Malyshev S, Naik V, Paulot F, Shevliakova E, Stock CA, Zadeh N, Balaji V, Blanton C, Dunne KA, Dupuis C, Durachta J, Dussin R, Gauthier PPG, Griffies SM, Guo H, Hallberg RW, Harrison M, He J, Hurlin W, McHugh C, Menzel R, Milly PCD, Nikonov S, Paynter DJ, Ploshay J, Radhakrishnan A, Rand K, Reichl BG, Robinson T, Schwarzkopf DM, Sentman LT, Underwood S, Vahlenkamp H, Winton M, Wittenberg AT, Wyman B, Zeng Y and Zhao M (2020) The GFDL Earth System Model Version 4.1 (GFDL-ESM 4.1): overall coupled model description and simulation characteristics. Journal of Advances in Modeling Earth Systems 12, e2019MS002015.
- Duscher GG, Hodžić A, Hufnagl P, Wille-Piazzai W, Schötta AM, Markowicz MA, Estrada-Peña A, Stanek G and Allerberger F (2018) Adult Hyalomma marginatum tick positive for Rickettsia aeschlimannii in Austria, October 2018. Eurosurveillance 23, 1800595.
- Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE and Yates CJ (2011) A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17, 43–57.
- Ergönül Ö (2006) Crimean–Congo haemorrhagic fever. *The Lancet Infectious Diseases* 6, 203–214.
- Escobar LE (2020) Ecological niche modeling: an introduction for veterinarians and epidemiologists. Frontiers in Veterinary Science 7, 519059.
- **Escobar LE, Lira-Noriega A, Medina-Vogel G and Peterson AT** (2014) Potential for spread of the white-nose fungus (*Pseudogymnoascus destructans*) in the Americas: use of Maxent and NicheA to assure strict model transference. *Geospatial Health* **9**, 221–229.
- Estrada-Peña A (2008) Climate, niche, ticks, and models: what they are and how we should interpret them. *Parasitology Research* **103**, 87–95.
- Estrada-Peña A (2023) The climate niche of the invasive tick species Hyalomma marginatum and Hyalomma rupes (Ixodidae) with recommendations for modeling exercises. Experimental and Applied Acarology 89, 1– 20. doi: 10.21203/rs.3.rs-2199505/v1
- Estrada-Pena A and De La Fuente J (2016) Species interactions in occurrence data for a community of tick-transmitted pathogens. *Scientific Data* 3, 1–13.
- Estrada-Peña A and Venzal JM (2007) Climate niches of tick species in the Mediterranean region: modeling of occurrence data, distributional constraints, and impact of climate change. *Journal of Medical Entomology* 44, 1130–1138.
- Estrada-Peña A, Vatansever Z, Gargili A and Ergönul Ö (2010) The trend towards habitat fragmentation is the key factor driving the spread of Crimean–Congo haemorrhagic fever. *Epidemiology & Infection* 138, 1194–1203.
- Estrada-Peña A, Martínez Avilés M and Muñoz Reoyo MJ (2011) A population model to describe the distribution and seasonal dynamics of the tick *Hyalomma marginatum* in the Mediterranean basin. *Transboundary and Emerging Diseases* 58, 213–223.
- Estrada-Peña A, Ayllón N and De La Fuente J (2012) Impact of climate trends on tick-borne pathogen transmission. *Frontiers in Physiology* 3, 64.
- Estrada-Peña A, Mihalca AD and Petney TN (2018) Ticks of Europe and North Africa: a Guide to Species Identification. Cham, Switzerland: Springer. doi: 10.1007/978-3-319-63760-0
- Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ and Taylor KE (2016) Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development* 9, 1937–1958.
- Fernández-Ruiz N and Estrada-Peña A (2021) Towards new horizons: climate trends in Europe increase the environmental suitability for permanent populations of *Hyalomma marginatum* (Ixodidae). *Pathogens (Basel, Switzerland)* **10**, 1–13.

- Fick SE and Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37, 4302–4315.
- Flores-López CA, Moo-Llanes DA, Romero-Figueroa G, Guevara-Carrizales A, López-Ordoñez T, Casas-Martínez M and Samy AM (2022) Potential distributions of the parasite *Trypanosoma cruzi* and its vector *Dipetalogaster maxima* highlight areas at risk of Chagas disease transmission in Baja California Sur, Mexico, under climate change. *Medical and Veterinary Entomology* 36, 469–479.
- Gale P, Estrada-Peña A, Martinez M, Ulrich RG, Wilson A, Capelli G, Phipps P, De La Torre A, Muñoz MJ, Dottori M, Mioulet V and Fooks AR (2010) The feasibility of developing a risk assessment for the impact of climate change on the emergence of Crimean–Congo haemorrhagic fever in livestock in Europe: a review. *Journal of Applied Microbiology* 108, 1859–1870.
- Gillingham EL, Medlock JM, Macintyre H and Phalkey R (2023) Modelling the current and future temperature suitability of the UK for the vector *Hyalomma marginatum* (Acari: Ixodidae). *Ticks and Tick-borne Diseases* 14, 102112.
- Githeko AK, Lindsay SW, Confalonieri UE and Patz JA (2000) Climate change and vector-borne diseases: a regional analysis. *Bulletin of the World Health Organization* 78, 1136–1147.
- Grandi G, Chitimia-Dobler L, Choklikitumnuey P, Strube C, Springer A, Albihn A, Jaenson TGT and Omazic A (2020) First records of adult *Hyalomma marginatum* and *H. rufipes* ticks (Acari: Ixodidae) in Sweden. *Ticks and Tick-borne Diseases* 11, 101403.
- Gray JS, Dautel H, Estrada-Peña A, Kahl O and Lindgren E (2009) Effects of climate change on ticks and tick-borne diseases in Europe. *Interdisciplinary Perspectives on Infectious Diseases* **2009**, 1–12.
- Hausfather Z (2020) CMIP6: The Next Generation of Climate Models Explained-Carbon Brief. Available at https://www.carbonbrief.org/cmip6the-next-generation-of-climate-models-explained/
- Hekimoglu O and Ozer NA (2017) Distribution and phylogeny of *Hyalomma* ticks (Acari: Ixodidae) in Turkey. *Experimental & Applied Acarology* 73, 501–519.
- Hekimoglu O, Sahin MK, Ergan G and Ozer N (2021) A molecular phylogenetic investigation of tick species in Eastern and Southeastern Anatolia. *Ticks and Tick-Borne Diseases* **12**, 101777.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG and Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25, 1965–1978.
- Hoerling M, Eischeid J, Perlwitz J, Quan X, Zhang T and Pegion P (2012) On the increased frequency of Mediterranean drought. *Journal of Climate* **25**, 2146–2161.
- Hornok S and Horváth G (2012) First report of adult *Hyalomma marginatum rufipes* (vector of Crimean–Congo haemorrhagic fever virus) on cattle under a continental climate in Hungary. *Parasites and Vectors* 5, 1–4.
- Jongejan F and Uilenberg G (2004) The global importance of ticks. Parasitology 129, S3-S14.
- Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP and Kessler M (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific Data* 4, 1–20.
- Karger DN, Lange S, Hari C, Reyer CPO, Conrad O, Zimmermann NE and Frieler K (2022) CHELSA-W5E5: daily 1 km meteorological forcing data for climate impact studies. *Earth System Science Data Discussions* 15, 1– 28. doi: 10.5194/essd-2022-367
- Kemp L, Xu C, Depledge J, Ebi KL, Gibbins G, Kohler TA, Rockström J, Scheffer M, Schellnhuber HJ, Steffen W and Lenton TM (2022) Climate endgame: exploring catastrophic climate change scenarios. *Proceedings of the National Academy of Sciences* 119, e2108146119.
- Kiewra D, Kryza M and Szymanowski M (2014) Influence of selected meteorological variables on the questing activity of *Ixodes ricinus* ticks in Lower Silesia, SW Poland. *Journal of Vector Ecology* 39, 138–145.
- Knülle W and Rudolph D (1982) Humidity Relationships and Water Balance of Ticks.In Obenchain FD and Galun R (eds). Physiology of Ticks. Oxford, UK: Pergamon Press. doi: 10.1016/b978-0-08-024937-7.50007-x
- Kovats RS, Campbell-Lendrum DH, McMichael AJ, Woodward A and Cox JSH (2001) Early effects of climate change: do they include changes in vector-borne disease? *Philosophical Transactions of the Royal Society B: Biological Sciences* 356, 1057–1068.
- Leal B, Zamora E, Fuentes A, Thomas DB and Dearth RK (2020) Questing by tick larvae (Acari: Ixodidae): a review of the influences that affect offhost survival. Annals of the Entomological Society of America 113, 425–438.

- Leder K, Openshaw JJ, Allotey P, Ansariadi A, Barker SF, Burge K, Clasen TF, Chown SL, Duffy GA, Faber PA, Fleming G, Forbes AB, French M, Greening C, Henry R, Higginson E, Johnston DW, Lappan R, Lin A, Luby SP, McCarthy D, O'Toole JE, Ramirez-Lovering D, Reidpath DD, Simpson JA, Sinharoy SS, Sweeney R, Taruc RR, Tela A, Turagabeci AR, Wardani J, Wong T and Brown R (2021) Study design, rationale and methods of the revitalising informal settlements and their environments (RISE) study: a cluster randomised controlled trial to evaluate environmental and human health impacts of a water-sensitive intervention in informal settlements in Indonesia and Fiji. *BMJ Open* 11, e042850.
- Lesiczka PM, Daněk O, Modrý D, Hrazdilová K, Votýpka J and Zurek L (2022) A new report of adult Hyalomma marginatum and Hyalomma rufipes in the Czech Republic. Ticks and Tick-borne Diseases 13, 101894.
- Liu C, White M and Newell G (2013) Selecting thresholds for the prediction of species occurrence with presence-only data. *Journal of Biogeography* **40**, 778–789. https://doi.org/10.1111/jbi.12058
- Maltezou HC and Papa A (2010) Crimean–Congo hemorrhagic fever: risk for emergence of new endemic foci in Europe? *Travel Medicine and Infectious Disease* 8, 139–143.
- Mangold AJ, Bargues MD and Mas-Coma S (1998) Mitochondrial 16S rDNA sequences and phylogenetic relationships of species of *Rhipicephalus* and other tick genera among Metastriata (Acari: Ixodidae). *Parasitology Research* 84, 478–484.
- Mauritsen T, Bader J, Becker T, Behrens J, Bittner M, Brokopf R, Brovkin V, Claussen M, Crueger T, Esch M, Fast I, Fiedler S, Fläschner D, Gayler V, Giorgetta M, Goll DS, Haak H, Hagemann S, Hedemann C, Hohenegger C, Ilyina T, Jahns T, Jimenéz-de-la-Cuesta D, Jungclaus J, Kleinen T, Kloster S, Kracher D, Kinne S, Kleberg D, Lasslop G, Kornblueh L, Marotzke J, Matei D, Meraner K, Mikolajewicz U, Modali K, Möbis B, Müller WA, Nabel JEMS, Nam CCW, Notz D, Nyawira SS, Paulsen H, Peters K, Pincus R, Pohlmann H, Pongratz J, Popp M, Raddatz TJ, Rast S, Redler R, Reick CH, Rohrschneider T, Schemann V, Schmidt H, Schnur R, Schulzweida U, Six KD, Stein L, Stemmler I, Stevens B, von Storch JS, Tian F, Voigt A, Vrese P, Wieners KH, Wilkenskjeld S, Winkler A and Roeckner E (2019) Developments in the MPI-M Earth system model version 1.2 (MPI-ESM1.2) and its response to increasing CO<sub>2</sub>. Journal of Advances in Modeling Earth Systems 11, 998–1038.
- McGinley L, Hansford KM, Cull B, Gillingham EL, Carter DP, Chamberlain JF, Hernandez-Triana LM, Phipps LP and Medlock JM (2021) First report of human exposure to *Hyalomma marginatum* in England: further evidence of a *Hyalomma* moulting event in north-western Europe? *Ticks and Tick-borne Diseases* **12**, 101541.
- Medlock JM and Leach SA (2015) Effect of climate change on vector-borne disease risk in the UK. *The Lancet Infectious Diseases* 15, 721–730.
- Merow C, Smith MJ and Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* 36, 1058–1069.
- Moo-Llanes DA, de Oca-Aguilar ACM, Romero-Salas D and Sánchez-Montes S (2021) Inferring the potential distribution of an emerging rickettsiosis in America: the case of *Rickettsia parkeri*. *Pathogens* (*Basel, Switzerland*) 10, 592.
- Naimi B (2017) Package 'usdm'. Uncertainty analysis for species distribution models. Available at https://cran.r-project.org/web/packages/usdm/
- Newbold T, Oppenheimer P, Etard A and Williams JJ (2020) Tropical and Mediterranean biodiversity is disproportionately sensitive to land-use and climate change. *Nature Ecology and Evolution* **4**, 1630–1638.
- Nuñez-Penichet C, Osorio-Olvera L, Gonzalez VH, Cobos ME, Jiménez L, DeRaad DA, Alkishe A, Contreras-Díaz RG, Nava-Bolaños A and Utsumi K (2021) Geographic potential of the world's largest hornet, *Vespa mandarinia* Smith (Hymenoptera: Vespidae), worldwide and particularly in North America. *PeerJ* 9, e10690.
- **Open Source Geospatial Foundation** (2022) GDAL, Geospatial Data Abstraction Software Library. Available at https://gdal.org/
- Ortega-Guzmán L, Rojas-Soto O, Santiago-Alarcon D, Huber-Sannwald E and Chapa-Vargas L (2022) Climate predictors and climate change projections for avian haemosporidian prevalence in Mexico. *Parasitology* 149, 1129–1144.
- Oshima N, Yukimoto S, Deushi M, Koshiro T, Kawai H, Tanaka TY and Yoshida K (2020) Global and Arctic effective radiative forcing of anthropogenic gases and aerosols in MRI-ESM2.0. *Progress in Earth and Planetary Science* 7, 1–21.

- Owens HL, Campbell LP, Dornak LL, Saupe EE, Barve N, Soberón J, Ingenloff K, Lira-Noriega A, Hensz CM, Myers CE and Peterson AT (2013) Constraints on interpretation of ecological niche models by limited environmental ranges on calibration areas. *Ecological Modelling* 263, 10–18.
- Perez G, Bastian S, Agoulon A, Bouju A, Durand A, Faille F, Lebert I, Rantier Y, Plantard O and Butet A (2016) Effect of landscape features on the relationship between *Ixodes ricinus* ticks and their small mammal hosts. *Parasites & Vectors* 9, 1–18.
- Peterson AT and Soberón J (2012) Species distribution modeling and ecological niche modeling: getting the concepts right. *Natureza a Conservacao* 10, 102–107.
- Phillips SJ and Schapire RE (2004) A Maximum Entropy Approach to Species Distribution Modeling. In Proceedings of the twenty-first international conference on Machine learning, p. 83. doi: 10.1145/ 1015330.1015412
- QGIS Geographic Information System (2022) QGIS Association. Available at https://www.qgis.org
- Raghavan RK, Heath ACG, Lawrence KE, Ganta RR, Peterson AT and Pomroy WE (2020) Predicting the potential distribution of *Amblyomma* americanum (Acari: Ixodidae) infestation in New Zealand, using maximum entropy-based ecological niche modelling. *Experimental and Applied* Acarology 80, 227–245.
- **R Core Team** (2022) *R: A Language and Environment for Statistical Computing* (4.22). R Foundation for Statistical Computing. Vienna, Austria. Available at https://www.r-project.org/
- Requena-García F, Cabrero-Sañudo F, Olmeda-García S, González J and Valcárcel F (2017) Influence of environmental temperature and humidity on questing ticks in central Spain. *Experimental and Applied Acarology* 71, 277–290.
- Rocklöv J and Dubrow R (2020) Climate change: an enduring challenge for vector-borne disease prevention and control. *Nature Immunology* 21, 479–483.
- Ros-García A, M'ghirbi Y, Bouattour A and Hurtado A (2011) First detection of *Babesia occultans* in *Hyalomma* ticks from Tunisia. *Parasitology* 138, 578–582.
- Sakai Y, Sudo M and Osakabe M (2012) Seasonal changes in the deleterious effects of solar ultraviolet-B radiation on eggs of the twospotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Applied Entomology and Zoology* **47**, 67–73.
- Sedaghat MM, Sarani M, Chinikar S, Telmadarraiy Z, Moghaddam AS, Azam K, Nowotny N, Fooks AR and Shahhosseini N (2017) Vector prevalence and detection of Crimean–Congo haemorrhagic fever virus in Golestan Province, Iran. *Journal of Vector Borne Diseases* 54, 353.
- Sellar AA, Jones CG, Mulcahy JP, Tang Y, Yool A, Wiltshire A, O'Connor FM, Stringer M, Hill R, Palmieri J, Woodward S, de Mora L, Kuhlbrodt T, Rumbold ST, Kelley DI, Ellis R, Johnson CE, Walton J, Abraham NL, Andrews MB, Andrews T, Archibald AT, Berthou S, Burke E, Blockley E, Carslaw K, Dalvi M, Edwards J, Folberth GA, Gedney N, Griffiths PT, Harper AB, Hendry MA, Hewitt AJ, Johnson B, Jones A, Jones CD, Keeble J, Liddicoat S, Morgenstern O, Parker RJ, Predoi V, Robertson E, Siahaan A, Smith RS, Swaminathan R, Woodhouse MT, Zeng G and Zerroukat M (2019) UKESM1: description and evaluation of the U.K. earth system model. Journal of Advances in Modeling Earth Systems 11, 4513–4558.
- Semenza JC and Suk JE (2018) Vector-borne diseases and climate change: a European perspective. FEMS Microbiology Letters 365, 1–9.
- Shemshad K, Rafinejad J, Kamali K, Piazak N, Sedaghat MM, Shemshad M, Biglarian A, Nourolahi F, Beigi EV and Enayati AA (2012) Species diversity and geographic distribution of hard ticks (Acari: Ixodoidea: Ixodidae) infesting domestic ruminants, in Qazvin Province, Iran. *Parasitology Research* 110, 373–380.
- Sillero N and Barbosa AM (2021) Common mistakes in ecological niche models. International Journal of Geographical Information Science 35, 213–226.
- Sillero N, Arenas-Castro S, Enriquez-Urzelai U, Vale CG, Sousa-Guedes D, Martínez-Freiría F, Real R and Barbosa AM (2021) Want to model a species niche? A step-by-step guideline on correlative ecological niche modelling. *Ecological Modelling* 456, 109671.
- Soberon J and Peterson AT (2005) Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics* 2, 1–10.
- Sofizadeh A, Telmadarraiy Z, Rahnama A, Gorganli-Davaji A and Hosseini-Chegeni A (2014) Hard tick species of livestock and their

bioecology in Golestan province, north of Iran. *Journal of Arthropod-borne Diseases* **8**, 108.

- Steven JP, Miroslav D and Schapire RE (2021) MaxEnt software for modeling species niches and distributions, version 3.4. 4. Available at https://biodiversityinformatics.amnh.org/open\_source/maxent/
- Sudo M and Osakabe M (2015) Joint effect of solar UVB and heat stress on the seasonal change of egg hatching success in the herbivorous false spider mite (Acari: Tenuipalpidae). *Environmental Entomology* 44, 1605–1613.
- Sultankulova KT, Shynybekova GO, Kozhabergenov NS, Mukhami NN, Chervyakova OV, Burashev YD, Zakarya KD, Nakhanov AK, Barakbayev KB and Orynbayev MB (2022) The prevalence and genetic variants of the CCHF virus circulating among ticks in the southern regions of Kazakhstan. Pathogens (Basel, Switzerland) 11, 841.
- Suzán G, Esponda F, Carrasco-Hernández R, Aguirre AA and Ostfeld RS (2012) Habitat fragmentation and infectious disease ecology. In Aguirre AA, Ostfeld R and Daszak P (eds). New Directions in Conservation Medicine: Applied Cases of Ecological Health. New York, USA: Oxford University Press, pp. 135–150. doi: 10.5860/choice.50-3886
- Telmadarraiy Z, Chinikar S, Vatandoost H, Faghihi F and Hosseini-Chegeni A (2015) Vectors of Crimean–Congo hemorrhagic fever virus in Iran. Journal of Arthropod-borne Diseases 9, 137.
- Turkes M, Turp MT, An N, Ozturk T and Kurnaz ML (2020) Impacts of climate change on precipitation climatology and variability in Turkey. In Harmancioglu NB and Altinbilek D (eds). Water Resources of Turkey. Cham, Switzerland: Springer, pp. 467–491. doi: 10.1007/978-3-030-11729-0\_14
- Ulbrich U, May W, Li L, Lionello P, Pinto JG and Somot S (2006) The Mediterranean climate change under global warming. In Lionello P, Malanotte-Rizzoli P and Boscolo R (eds). *Developments in Earth and*

*Environmental Sciences.* Amsterdam, Netherlands: Elsevier, pp. 399–415. doi: 10.1016/S1571-9197(06)80011-X

- Valcárcel F, González J, González MG, Sánchez M, Tercero JM, Elhachimi L, Carbonell JD and Sonia Olmeda A (2020) Comparative ecology of *Hyalomma lusitanicum* and *Hyalomma marginatum* Koch, 1844 (Acarina: Ixodidae). Insects 11, 1–17.
- Vatansever Z, Estrada-Pena A and Ergonul O (2007) Crimean–Congo hemorrhagic fever in Turkey. In Ergonul O and Whitehouse CA (eds). Crimean– Congo Hemorrhagic Fever: a Global Perspective. Dordrecht, Netherlands: Springer, pp. 59–74. doi: 10.1007/978-1-4020-6106-6\_6
- Vial L, Stachurski F, Leblond A, Huber K, Vourc'h G, René-Martellet M, Desjardins I, Balança G, Grosbois V, Pradier S, Gély M, Appelgren A and Estrada-Peña A (2016) Strong evidence for the presence of the tick *Hyalomma marginatum* Koch, 1844 in southern continental France. *Ticks* and *Tick-borne Diseases* 7, 1162–1167.
- Warren DL and Seifert SN (2011) Ecological niche modeling in MaxEnt: the importance of model complexity and the performance of model selection criteria. *Ecological Applications* **21**, 335–342.
- Williams HW, Cross DE, Crump HL, Drost CJ and Thomas CJ (2015) Climate suitability for European ticks: assessing species distribution models against null models and projection under AR5 climate. *Parasites and Vectors* 8, 1–15.
- Wu B, Li X, Liu J and Bao R (2022) Predicting the potential habitat for Ornithodoros tick species in China. Veterinary Parasitology 311, 109793.
- Zhao GP, Wang YX, Fan ZW, Ji Y, Liu MJ, Zhang WH, Li XL, Zhou SX, Li H, Liang S, Liu W, Yang Y and Fang LQ (2021) Mapping ticks and tickborne pathogens in China. *Nature Communications* 12, 1–13.