#### PERBANDINGAN JUMLAH TROMBOSIT PRIMIGRAVIDA DENGAN MULTIGRAVIDA PADA KEJADIAN PREEKLAMPSIA DI RUMAH SAKIT IBNU SINA GRESIK PERIODE TAHUN 2019 – 2021

### Anita Dahliana<sup>1\*</sup>, Ummy Maisarah Rasyidah<sup>1</sup>, Arya Rizky Fernanda<sup>1</sup>, Nita Damayanti<sup>2</sup>

<sup>1</sup>Faculty of Medicine, University of Surabaya (Ubaya), Surabaya, I. Raya Kalirungkut, Kali Rungkut, Surabaya, Jawa Timur 60293 <sup>2</sup>Faculty of Medicine, Alkhairaat University, Il Diponegoro No 39 Palu 94221, Sulawesi Tengah

<sup>2</sup>Faculty of Medicine, Alkhairaat University, Jl. Diponegoro No.39 Palu 94221, Sulawesi Tengah, Indonesia

\*Corresponding author: Telp: +6281216402066, email: anitadahliana@staff.ubaya.ac.id

## ABSTRAK

Latar belakang: Preeklampsia adalah suatu kondisi dimana ibu hamil dengan masa kehamilan lebih dari 20 minggu memiliki tanda utama hipertensi dan proteinuria. Penelitian ini bertujuan untuk menganalisis perbedaan jumlah trombosit antara primigravida dan multigravida pada kejadian preeklampsia. Metode: Penelitian ini merupakan penelitian cross-sectional. Pengumpulan data menggunakan probability sampling yaitu purposive sampling dengan 61 rekam medik ibu preeklampsia yang diambil sesuai kriteria inklusi di RS Ibnu Sina Gresik yang terdiri dari 14 primigravida dan 47 multigravida dari tahun 2019 sampai dengan 2021. Hasil: Pasien preeklampsia dengan jumlah trombosit terendah adalah 170 kelompok, terendah.000/L dan tertinggi 490.000/µL dengan rerata 282.066/µL. Pada primigravida jumlah trombosit 170.000/L dan tertinggi 342.000/L dengan rata-rata 277.857/µL, sedangkan pada kelompok multigravida memiliki jumlah trombosit terendah 173.000/µL dan tertinggi 490.000/µL dengan rata-rata 283.319/ L. Kedua kelompok dibandingkan dan hasilnya menunjukkan bahwa tidak ada perbedaan bermakna jumlah trombosit antara primigravida dan multigravida (p>0,05). Simpulan: Tidak terdapat perbedaan bermakna antara jumlah trombosit primigravida dan multigravida pada pasien preeklampsia.

Keywords: Gravida, Primigravida, Multigravida, Preeclampsia, Platelet.

### ABSTRACT

**Background**: Preeclampsia is a condition in which pregnant women with a gestation period of more than 20 weeks have the main signs of hypertension and proteinuria. This study aims to analyze the difference in the number of platelets between primigravida and multigravida in the incidence of preeclampsia. **Methods**: This research was a cross-sectional study. Data collection used probability sampling, namely purposive sampling with 61 medical records of preeclampsia mothers taken according to the inclusion criteria at the Ibnu Sina Hospital Gresik consisting of 14 primigravidas and 47 multigravidas from 2019 to 2021. **Results**: Preeclampsia patients with the lowest platelet count was 170 group, the lowest.000/L and the highest was 490.000/ $\mu$ L with a mean of 282.066/ $\mu$ L. In the primigravida platelet count was 170.000 / L and the highest was 342.000 / L with a mean of 277.857 /  $\mu$ L, meanwhile the multigravida group had the lowest platelet count of 173.000 /  $\mu$ L and the highest was 490.000 /  $\mu$ L with a mean of 283.319 / L. The two groups were compared and the results showed that there was no significant difference in platelet count between primigravida and multigravida (p>0.05). **Conclusion**: There is no

significant difference between primigravida and multigravida platelet count in preeclampsia patients.

Keywords: Gravida, Primigravida, Multigravida, Preeclampsia, Platelet.

# **INTRODUCTION**

Preeclampsia is one of the main problems that cause maternal and perinatal mortality and morbidity in the world.<sup>1</sup> The estimated prevalence of preeclampsia in the world is 4.6% of all pregnancies.<sup>2</sup> The prevalence of deaths caused by preeclampsia in Asia alone reaches 9%, including 70.000 maternal deaths and 500.000 fetal deaths in the world each year.<sup>3</sup> Preeclampsia is caused by various factors such as age, gravida status, history of high blood pressure, genetics and so on. Preeclampsia occurs due to endothelial dysfunction which in turn will lead to conditions of hypertension, proteinuria and decreased platelet count.<sup>4</sup>

Gravida status is a risk factor for preeclampsia. Gravida status is divided into two, namely primigravida or pregnant for the first time and multigravida, namely pregnant more than once. Gravida status can affect the number of human leukocyte antigen (HLA) in the placenta, which will cause placental ischemia and hypertension. In multigravida, gravida status is the safest for the incidence of preeclampsia.<sup>5</sup>

In pregnant women, the hemodilution effect and increased utilization of platelets in the maternal circulation can reduce the platelet count. Conditions where the number of platelets has decreased significantly can be thrombocytopenia. said to be Thrombocytopenia is defined as a platelet than 150.000 count of less μ/L. Thrombocytopenia is the second most common cause of blood disorders in pregnancy after anemia. Thrombocytopenia occurs with a prevalence of 7-10% of all pregnancies. A low platelet count can increase the risk of developing an epidural addition, new bleeding hematoma. In conditions can occur if the platelet count is below 20.000  $\mu/L$  and internal bleeding is

AGUSTUS 2023

below 10.000 µ/L.<sup>6</sup> Therefore, platelet count has a significant impact on primigravida and multigravida pregnancies.Incidence of thrombocytopenia in pregnancy occurs in 8-9% of all deliveries.<sup>7</sup> The presence of thrombocytopenia in pregnancy does not necessarily result in preeclampsia.<sup>8</sup> Thus, this study aims to analyse the differences in the number of platelets between primigravida and multigravida in the incidence of preeclampsia Gravida status is a risk factor for preeclampsia. Gravida status is divided into two, namely primigravida or pregnant for the first time and multigravida, namely pregnant more than once. Gravida status can affect the number of human leukocyte antigen (HLA) in the placenta, which will cause placental ischemia and hypertension. In multigravida, gravida status is the safest for the incidence of preeclampsia.5

# **METHODS**

This is a cross-sectional study conducted in Ibnu Sina Regional Hospital in Gresik East Java. The hospital is a 316-beds hospital with a multi-specialistic service, located in an urban setting, around one hour drive from Surabaya, the capital of the province.

We conducted a simple random sampling to select 61 medical records from the list of patients with preeclamsia admitted to the inpatient wards in Ibnu Sina Hospital in 2019-2021. The eligibility criteria were patients with the gestation age of 20 weeks or more and had obtained blood test for platelet count. Patients with Immune Thrombocytopenic Purpura (ITP), infection disorders, smoking, and platelet count < 100.000 /  $\mu$ L were excluded from the study. The sample size was determined based on the sample size calculation using the alpha of 5 and the power of 95. We collected these following data from the medical record: age, past medical history including pregnancy, blood pressure, the level of protein in the urine, and the platelet count.

Based on their pregnancy history, we categorized the patients into primigravida and multigravida. When patients underwent multiple blood tests, we recorded the newest data for the platelet count for the analysis. The data collecting process was carried out for 1 week.

As the data were normally distributed, we conducted an independent T-test to compare the platelet count between the groups of primipara and multipara.

We obtained the Ethical Clearance from the Ethical Committee of Universitas Surabaya (no 45a / KE / IV / 2022).

To ensure the confidentiality of the patients, all data that can refer to the identity of the patients had been omitted during the data collection, the analysis, and in the report and publication.

# RESULT

There are two groups of classification in this study, advanced maternal age - AMA (>35 Years old) and reproductive age - RA (20 - 34 years old).

Table 1. Age distribution from each group
of multigravida and primigravida

A go	Frecuency	Percentage
Age	(n)	(%)
Ν		
Reproductive		
Age (20-34	35	74,4
Years old)		
Advanced	12	
Maternal Age	12	25,6
(>35 Years old)		
Total	47	100
F	Primigravida	
Reproductive		
Age (20-34	13	92,8
Years old)		
Advanced		
Maternal Age	1	7,2
(>35 Years old)		
Total	14	100

Table 2 shows that there are 35 patients on the RA group while for AMA, there are only 12 patients in the multigravida group. The same thing occurred in the primigravida group where there are 13 RA patients and only 1 AMA patient. Most of the patients in this study are in the RA group, which consists of 48 patients while AMA only has 13 patients in total.

The normality test (Kolmogorov-Smirnov) is conducted in this study and the result is sig. 0,2 on both multigravida or primigravida which means the data is distributed normally (>Sig 0,05).

Table 2. Platelet Count distribution from	n
each group of multigravida and	
nrimigravida	

primgru	Iuu		
	Multigravida	Primigravida	p-
	(Mean ±	(Mean ±	value
	SD)	SD)	
Frequency	47	14	
Platelet	$283.319 \pm$	$277.857 \pm$	
Count	71.312,57	55.801,65	0.263
p-values	0,2	263	

Based on the data that were shown on the table 2. There is a slight difference of platelet count in multigravida and primigravida. Analysis of data comparing platelet count between primigravidas and multigravidas obtained using the independent T Test, resulted in sig.(2-tailed) 0.263, thus it can be concluded that there was no significant difference in the comparison of platelet count between primigravidas and multigravidas.

### DISCUSSION

From the gravida status, it shows that most of the patients were in the multigravida group with as many as 47 patients and a percentage of 77.1%, while the primigravida group only had 14 patients with a percentage of 22.9%. From the data available from 2019 to 2021 at the Ibnu Sina Hospital Gresik, the number of primigravida preeclampsia patients is less when compared to multigravida. This is in line with a study conducted at the Surabaya Muhammadiyah Hospital that states there are fewer primigravida patients than multigravida patients with a total of 43 primigravida patients with a percentage of 41% while multigravida patients are 62 patients with a percentage of 59%.<sup>10</sup>

The pathomechanism underlying gravida status is related to the immunologic theory. The theoretical basis for what occurs comes from the imperfect formation of human leukocyte antigen (HLA), which causes a rejection of placentation. The trophoblast implantation process into the maternal decidual tissue will also be disrupted, causing interference with spiral the arteries. Furthermore, the spiral arteries will narrow and be inelastic which can cause placental ischemia.<sup>11</sup> Stress occuring in pregnant women can cause an increase in corticotropicreleasing hormone (CRH) bv the hypothalamus, resulting in the increase of cortisol levels. Cortisol can increase the vascular concentration of angiotensin II, which in turn causes vasoconstriction of blood vessels and hypertension.<sup>12</sup>

Based on the results of this study, the incidence of preeclampsia in both the primigravida and multigravida groups have the same risk. This result is because the number of multigravidas was more than primigravidas, while in another study primigravidas were more likely to cause preeclampsia than multigravidas.<sup>13</sup>

Mean platelet count in the multigravida group was 283.319 / µL, consisting of platelet counts ranging from 173.000 / µL to 490.000  $/\mu$ L. While the average number of platelets in the primigravida group was 277.857/µL, ranging from  $170.000/\mu$ L to  $342.000/\mu$ L. The result of the Unpaired T-test showed a pvalue of 0.263, thus it can be concluded that there was no significant difference between the two groups because the p-value was more than 0.05. This study showed no significant platelet difference in count between primigravida and multigravida preeclampsia patients, although multigravida platelet levels were slightly higher than primigravidas.

Number of platelets decreased during pregnancy and will continue to decrease during pregnancy. Where, 9.9% have a platelet count lower than 150.000 /  $\mu$ L<sup>14</sup>

Likewise, a study stated there was no significant difference in the platelet count of preeclampsia patients and the results of the study concluded that the change in platelet count was not due to preeclampsia, but due to the pregnancy itself. These two studies prove that the decrease in platelet count is not affected by the incidence of preeclampsia, but changes that occur due to pregnancy itself.<sup>15</sup> There is also a study which compared the number of platelets in mild preeclampsia and severe preeclampsia and looked at the patient's gestational period. This study showed that platelet count in severe preeclampsia was much lower than in mild preeclampsia. Gestation period in preeclampsia patients also has an influence on the patient's platelet count. From this study, it was concluded that one of the factors that affect platelet count in preeclampsia patients the condition of severity of is the preeclampsia itself and the patient's gestation period, where it is found that early gestation period does not show significant changes, but the patient will experience changes in a longer gestation period.<sup>16</sup>

In this study, there was no significant in platelet count between difference primigravidas and multigravidas. Platelet count was not affected by gravida status because in this study the patient had preeclampsia. This is in line with a study which said that preeclampsia patients did experience changes in their platelet counts even though they had varying gravida status.<sup>17</sup> The pathomechanism underlying the platelet count in preeclampsia is imperfect vascular development of the placenta.<sup>18</sup> This development involves the trophoblast which proangiogenic will produce and antiangiogenic factors in preeclampsia.<sup>19</sup> Proangiogenic factors function to maintain blood vessels while antiangiogenic factors work the opposite, namely damaging blood vessels. Antiangiogenic factors consist of soluble Fms-like tyrosine kinase1 (sFlt-1) and soluble endglin (sEng). sFlt-1 will bind to vascular endothelial growth factor (VEGF) thereby inhibiting VEGF from binding to the endothelium causing endothelial dysfunction. sEng binds to TGF-β thereby inhibiting TGFβ to bind to endothelial receptors and causes a decrease in nitric oxide which functions as a vascular vasodilator. Increased antiangiogenic factors causes maternal endothelial dysfunction. This endothelial dysfunction leads to platelet activation and increased use of platelets, as well as thrombus formation.<sup>20</sup> From the explanation above, it can be concluded that gravida status does not affect the platelet count. The decrease in platelet in this study was caused count bv preeclampsia. In addition, there are other factors that can affect the platelet count such as the pregnancy itself, the severity of preeclampsia and the patient's gestational period.<sup>21</sup>

There are some limitation that may affect the result of this study. Currently, Gresik is the only city in which the research is being conducted which is classified as a small city, where many other factors affect the respondents. Only 61 medical record included in this study as responses, more participants in future studies should be looked for to improve research outcomes. Further research should notice other additional factors, such as antenatal check-up history and dietary intake, which can raise the risk of preeclampsia.

# CONCLUSION

To conclude this study, there is no significant difference between platelet count from the preeclampsia patients, whether they were primigravida or multigravida. Improving patient outcome should be the first priority, and medical staff should be expected to between understand the platelet count primigravidas and multigravidas. Trombositopenia can occurs in preeclampsia patient and should be given corticosteroid dexamethasone 40mg/d oral to the patient.

# REFEREENCE

1. Nakimuli A, Nakubulwa S, Kakaire O, Osinde MO, Mbalinda SN, Kakande N, et al. The burden of maternal morbidity and mortality attributable to hypertensive disorders in pregnancy: A prospective cohort study from Uganda. BMC Pregnancy Childbirth [Internet]. 2016;16(1):1–8. Available from: http://dx.doi.org/10.1186/s12884-016-1001-1

- Yang Y, Le Ray I, Zhu J, Zhang J, Hua J, Reilly M. Preeclampsia Prevalence, Risk Factors, and Pregnancy Outcomes in Sweden and China. JAMA Netw open. 2021 May;4(5):e218401.
- 3. Rana S, Lemoine E, Granger J, Karumanchi SA. Preeclampsia: Pathophysiology, Challenges, and Perspectives. Circ Res. 2019;124(7):1094–112.
- 4. Fox R, Kitt J, Leeson P, Aye CYL, Lewandowski AJ. Preeclampsia: Risk factors, diagnosis, management, and the cardiovascular impact on the offspring. J Clin Med. 2019;8(10):1–22.
- Triche EW, Harland KK, Field EH, Rubenstein LM, Saftlas AF. Maternalfetal HLA sharing and preeclampsia: Variation in effects by seminal fluid exposure in a case-control study of nulliparous women in Iowa. J Reprod Immunol [Internet]. 2014;101– 102(1):111–9. Available from: http://dx.doi.org/10.1016/j.jri.2013.06.00 4
- Tyas BD, Lestari P, Aldika Akbar MI. Maternal Perinatal Outcomes Related to Advanced Maternal Age in Preeclampsia Pregnant Women. J Fam Reprod Heal. 2019 Dec;13(4):191–200.
- Subtil SFC, Mendes JMB, Areia ALF de A, Moura JPAS. Update on Thrombocytopenia in Pregnancy. Rev Bras Ginecol e Obstet Rev da Fed Bras das Soc Ginecol e Obstet. 2020 Dec;42(12):834–40.
- Habas E, Rayani A, Ganterie R. Thrombocytopenia in hypertensive disease of pregnancy. J Obstet Gynecol India. 2013;63(2):96–100.
- 9. Ives CW, Sinkey R, Rajapreyar I, Tita ATN, Oparil S. Preeclampsia-Pathophysiology and Clinical Presentations: JACC State-of-the-Art Review. J Am Coll Cardiol. 2020 Oct;76(14):1690–702.

- Rafida M, Mochtar NM, Artiningtyas ND, Anas M. Relationship of Age, Body Mass Index, and Gravida in Pregnant Women With Preeclampsia in Muhammadiyah Hospital Surabaya. Proc 4th Int Conf Sustain Innov 2020–Health Sci Nurs (ICoSIHSN 2020). 2021;33(ICoSIHSN 2020):37–42.
- Miller EC, Wilczek A, Bello NA, Tom S, Wapner R, Suh Y. Pregnancy, preeclampsia and maternal aging: From epidemiology to functional genomics. Ageing Res Rev. 2022 Jan;73:101535.
- 12. Barbot M, Ceccato F, Scaroni C. The pathophysiology and treatment of hypertension in patients with Cushing's syndrome. Front Endocrinol (Lausanne). 2019;10(MAY):1–9.
- Arwan B, Sriyanti R. Relationship between Gravida Status , Age , BMI ( Body Mass Index ) and Preeclampsia. Andalas Obstet Gynecol J. 2020;4(1):13– 21.
- Reese JA, Peck JD, Deschamps DR, McIntosh JJ, Knudtson EJ, Terrell DR, et al. Platelet Counts during Pregnancy. N Engl J Med. 2018;379(1):32–43.
- Thalor N, Singh K, Pujani M, Chauhan V, Agarwal C, Ahuja R. A correlation between platelet indices and preeclampsia. Hematol Transfus cell Ther. 2019;41(2):129–33.
- Dadhich S, Agrawal S, Soni M, Choudhary R, Jain R, Sharma S, et al. Predictive value of platelet indices in development of preeclampsia. J SAFOG. 2012;4(1):17–21.
- 17. Sachan R, Patel ML, Vandana, Sachan P, Shyam R. Role of platelet count and mean platelet volume and red cell distribution width in the prediction of preeclampsia in early pregnancy. J Fam Med Prim care. 2021 Feb;10(2):838–43.
- Melchiorre K, Giorgione V, Thilaganathan B. The placenta and preeclampsia: villain or victim? Am J Obstet Gynecol. 2022 Feb;226(2S):S954–62.
- 19. Helmo FR, Lopes AMM, Carneiro ACDM, Campos CG, Silva PB, Dos Reis

Monteiro MLG, et al. Angiogenic and antiangiogenic factors in preeclampsia. Pathol Res Pract. 2018 Jan;214(1):7–14.

- 20. Zhou CC, Ahmad S, Mi T, Xia L, Abbasi S, Hewett PW, et al. Angiotensin II induces soluble fms-Like tyrosine kinase-1 release via calcineurin signaling pathway in pregnancy. Circ Res. 2007 Jan;100(1):88–95.
- Vázquez-Rodríguez JG, Sánchez-Brito LO. Severity of preeclampsia: data from a high specialty hospital in Mexico City. Rev Med Inst Mex Seguro Soc. 2020;58(4):444–9.



p-ISSN : 2657-179X, e-ISSN : 2656-7822

# Medika Alkhairaat: Jurnal Penelitian Kedokteran dan Kesehatan

# Vol. 5, No. 2, April 2023, pp 45-102

EFFECT OF GIVING SPINACH LEAF PUDDING (AMARANTHUS HYBRIDUS) PREVENTING CALCIUM DEFICIENCY AND IRON DEFICIENCY ANEMIA (Ummy Maisarah Rasyidah, Anita Dahliana, Winnie Nirmala Santosa)

FAKTOR-FAKTOR YANG MEMPENGARUHI STATUS GIZI BALITA DI DESA SARONDA KECAMATAN BAJO BARAT (Fitriana Ibrahim, Sumarni, Jumriana Ibriani, Nia Rahmadani)

GAGAL VENTRIKEL KANAN AKUT: PENEGAKAN DIAGNOSIS DAN MANAJEMEN TATALAKSANA (Sidhi Laksono Purwowiyoto, Indira Khairunnisa Effendi)

PENGARUH PEMBERIAN ASAM KLOROGENAT PADA MENCIT SWISS WEBSTER DENGAN MODEL UNILATERAL URETERAL OBSTRUCTION (Masita Muchtar, Nur Arfian, Junaedy Yunus)

GAMBARAN NYERI KEPALA PRIMER DAN INSOMNIA PADA SISWA-SISWI. SMA AL-AZHAR MANDIRI PALU

(Nur Kartika Sari, Ruslan Ramlan Ramli, Marina Musyawwirina Desiprimayanti RR)

PENGARUH PEMBERIAN ASAM KLOROGENAT TERHADAP EKSPRESI mRNA TLR-4, mRNA MCP-1, JUMLAH MAKROFAG M1 DAN M2 PADA GINIAL MENCIT SWISS WEBSTER DENGAN MODEL UNILATERAL URETERAL OBSTRUCTION (UUO) (Tiara Meirani Valeria Savista, Junaedy Yunus, Nur Arfian)

PERBANDINGAN JUMLAH TROMBOSIT PRIMIGRAVIDA DENGAN MULTIGRAVIDA PADA KEJADIAN PREEKLAMPSIA DI RUMAH SAKIT IBNU SINA GRESIK PERIODE TAHUN 2019 – 2021 (Anita Dabliana, Ummy Maisarah Basyidah, Arya Bizky Fernanda, Nita

(Anita Dahliana, Ummy Maisarah Rasyidah, Arya Rizky Fernanda, Nita Damayanti)

> PUBLISHED BY: FAKULTAS KEDOKTERAN UNIVERSITAS ALKHAIRAAT JL. DIPONEGORO NO. 39 PALU

> > Cp: 081334845084

MEDIKA	ALKHAI	RAAT 8			p-ISSN: 2657-179X e-ISSN: 2656-7822	
Jurnial	PENIELIT	'iani kej	DOKT	eran d	Published By: FK UNISA PALU	
Home	Submissions	Contact	Current	Archives	Announcements	
Publi	cation Ethics	Author Guide	elines (	Cover Issue	Editorial Team 👻	
		Abou	t the Journ	al 🔻		
				Search		

Home / Editorial Team

#### **Editor In Chief**

Dr. Misriyani, M.Sc (Medical Faculty, Alkhairaat University, Indonesia) (<u>Google Scholar</u>) (<u>SINTA</u>) (<u>Scopus</u>)

#### Vice Editor In Chief

dr. Andi Handriyati, M.Kes, Sp.A (Medical Faculty, Alkhairaat University, Indonesia)

#### **Editorial Board**

dr. H.A Mukramin Amran, Sp. Rad (Medical Faculty, Alkhairaat University, Indonesia) dr. Ruslan Ramlan Ramli, Sp. S (Medical Faculty, Alkhairaat University, Indonesia) drg. Lutfiah Sahabuddin, M.KM (Medical Faculty, Alkhairaat University, Indonesia) dr. Nur Faisah, M.Kes, Sp.S (Medical Faculty, Alkhairaat University, Indonesia) drg. Nita Damayanti, M.Kes (Medical Faculty, Alkhairaat University, Indonesia) drg. Nita Damayanti, M.Kes (Medical Faculty, Alkhairaat University, Indonesia) dr. Wijoyo Halim, M.Kes, Sp.S (Medical Faculty, Alkhairaat University, Indonesia) dr. Nur Meity, M.Med.Ed (Medical Faculty, Alkhairaat University, Indonesia) dr. Nayalisa, Diantamaela, M.Sc (Medical Faculty, Alkhairaat University, Indonesia) dr. Salmah Suciaty, M.Kes (Medical Faculty, Alkhairaat University, Indonesia)

# **Peer Reviewers**

Dr. dr. Ilhamjaya Patellongi, M.Kes (Department of Physiology, Hasanuddin University, Makassar, Indonesia) (<u>Google Scholar</u>) (<u>SINTA</u>)

I Made Wisnu Adhi Putra, M.Sc (Department of Biology, Dhyana Pura University, Bali, Indonesia) (<u>Scopus</u>)

Prof. Dr. Ahyar Ahmad, Ph.D. (Department of Chemistry, Hasanuddin University, Makassar, Indonesia) (<u>Scopus</u>)

dr. Rizalinda Sjahril, M.Sc, Ph.D. (Department of Microbiology, Medical Faculty, Hasanuddin University) (<u>Scopus</u>)

dr. Isra Wahid, Ph.D. (Department of Parasitology, Medical Faculty, Hasanuddin University) (<u>Scopus</u>)

Bohari, S.Gz., M.Kes (Department of Nutrition, Faculty of Public Health, Tadulako University, Indonesia) (<u>Scopus</u>)

Dr. Mudyawati, S.Si, M.Kes (Midwifery Study Program, Tahirah Al Baeti Midwifery Academy Bulukumba, Indonesia) (<u>Google Scholar</u>)

Make a Submission



# Call For Paper: Volume 5 Nomor 3 (2023)

2023-08-02



Read More >

# **Current Issue**

# Vol 5 No 2 (2023): Agustus



Cp: 081334845084

Published: 2023-08-02

# Articles

# The EFFECT OF GIVING SPINACH LEAF PUDDING (AMARANTHUS HYBRIDUS) PREVENTING CALCIUM DEFICIENCY AND IRON DEFICIENCY ANEMIA

Ummy Maisarah Rasyidah, Anita Dahliana, Winnie Nirmala Santosa 45-49

🖹 PDF

## FAKTOR-FAKTOR YANG MEMPENGARUHI STATUS GIZI BALITA DI DESA SARONDA KECAMATAN BAJO BARAT

Fitriana Ibrahim, Sumarni, Jumriana Ibriani, Nia Rahmadani 50-56

🔁 PDF

### GAGAL VENTRIKEL KANAN AKUT: PENEGAKAN DIAGNOSIS DAN MANAJEMEN TATALAKSANA

Sidhi Laksono, Indira Khairunnisa Effendi 57-70

🔁 PDF

# PENGARUH PEMBERIAN ASAM KLOROGENAT PADA MENCIT SWISS WEBSTER DENGAN MODEL UNILATERAL URETERAL OBSTRUCTION

Masita Muchtar, Nur Arfian, Junaedy Yunus

71-80

🖹 PDF

## GAMBARAN NYERI KEPALA PRIMER DAN INSOMNIA PADA SISWA-SISWI SMA AL-AZHAR MANDIRI PALU

Marina Musyawwirina Desiprimayanti, Nur Kartika Sari 81-87



PENGARUH PEMBERIAN ASAM KLOROGENAT TERHADAP EKSPRESI mRNA TLR-4, mRNA MCP-1, JUMLAH MAKROFAG M1 DAN M2 PADA GINJAL MENCIT SWISS WEBSTER DENGAN MODEL UNILATERAL URETERAL OBSTRUCTION (UUO)

Tiara Meirani Valeria Savista, Junaedy Yunus, Nur Arfian 88-96

🖾 PDF

# PERBANDINGAN JUMLAH TROMBOSIT PRIMIGRAVIDA DENGAN MULTIGRAVIDA PADA KEJADIAN PREEKLAMPSIA DI RUMAH SAKIT IBNU SINA GRESIK PERIODE TAHUN 2019 – 2021

Anita Dahliana, Ummy Maisarah Rasyidah, Arya Rizky Fernanda, Nita Damayanti 97-102



View All Issues >

Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan (<u>p-ISSN: 2657-179X</u> & <u>e-ISSN: 2656-7822</u>) is a scientific journal published by the medical faculty of the alkhairaat university. This journal specializes in the study of medicine and health. The managers invite scientists, scholars, professionals, and researchers in the disciplines of medicine and health to publish their research results after the selection of manuscripts, review of partners, and editing process. This journal is published every April, August, and December.

Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan is a peer-reviewed and open-access journal that focuses on promoting medical sciences generated from basic sciences, clinical, and



**community or public health research** to integrate research in all aspects of human health. This journal publishes **original articles, reviews**, and also interesting **case reports**.

The manuscript must be **original research**, written in **Bahasa (Indonesia) or English**, and not be simultaneously submitted to **another journal** or **conference**.

For the author interested in submitting the manuscript, kindly register yourself. The **author guidelines** can be viewed <u>here</u>, and the manuscript template can be <u>downloaded here</u>

\_\_\_\_\_

Journal Title	: Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan
Subjects	: Promoting medical sciences generated from basic sciences, clinical, and compublic health research to integrate research in all aspects of human health.
language	: English and Bahasa, The manuscript is written in Bahasa (Indonesia) or Engl
ISSN	: <u>p-ISSN: 2657-179X</u> l <u>e-ISSN: 2656-7822</u>
Frequency	: 3 issues per year (April, August, and December)
DOI	: https://doi.org/10.31970/ma
Editor-In-Chief	: Misriyani ( <u>Google Scholar)(SINTA</u> )( <u>Scopus</u> )

### Publisher

: Fakultas Kedokteran Universitas Alkhairaat

Citation Analysis : <u>Google Scholar</u>, <u>Garuda</u>

Jurnal Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan has been accredited by the Indonesian Ministry of Research, Technology, and Higher Education (RISTEKDIKTI) number: 225/E/KPT/2022 since 07 December 2022, valid thru April 2024. Available in Sinta with Status : Terakreditasi Peringkat 5.



**Make a Submission** 



### **MEDIKA ALKHAIRAAT : JURNAL PENELITIAN KEDOKTERAN DAN KESEHATAN**

**Q** FAKULTAS KEDOKTERAN UNIVERSITAS ALKHAIRAAT

#### ₱ P-ISSN : 2657179X <> E-ISSN : 26567822

0 Impact Factor						
fine 11 Google Citations						
Sinta 5 Current Acreditation						
	Google Scholar	<b>▶</b> <u>Garuda</u>	🚱 <u>Website</u>	S Editor URL		
History Accreditation						
2020	2021	2022		2023	2024	
<u>Garuda</u> <u>Google Scho</u>	lar					

GAMBARAN TINGKAT PENGETAHUAN, PENDIDIKAN DAN SOSIO- EKONOMI BUMIL KEK DI PUSKESMAS TALISE, PUSKESMAS SINGGANI DAN PUSKESMAS KAWATUNA 2021 Fakultas Kedokteran Universitas Alkhairaat Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 13-18 2023 DOI: 10.31970/ma.v5i1.112 OAccred : Sinta 5 IDENTIFIKASI FAKTOR RISIKO GIZI KURANG PADA ANAK BALITA Fakultas Kedokteran Universitas Alkhairaat Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 7-12 2023 POI: 10.31970/ma.v5i1.113 OAccred : Sinta 5

 HUBUNGAN ANTARA LAMA HIPERTENSI DENGAN FUNGSI KOGNITIF PADA PASIEN LANSIA DI RSUD SANJIWANI GIANYAR

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 33-39

 2023
 DOI: 10.31970/ma.v5i1.114
 O Accred : Sinta 5

 KORELASI PEMANFAATAN APLIKASI WHATSAPP SEBAGAI MEDIA KIE TENTANG KB TERHADAP MINAT KB PADA PUS DI BULUKUMBA

 Fakultas Kedokteran Universitas Alkhairaat

 <sup>\*</sup> Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 25-32

 <sup>\*</sup> DOI: 10.31970/ma.v5i1.115

 <sup>•</sup> Accred : Sinta 5

#### KEPUASAN PASIEN RAWAT INAP DI RUMAH SAKIT SIS ALJUFRI PALU TAHUN 2022

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 19-24

 <sup>1</sup> 2023

 <sup>1</sup> DOI: 10.31970/ma.v5i1.116

 <sup>O</sup> Accred : Sinta 5

#### PEMBERIAN ASI EKSKLUSIF TERHADAP KEJADIAN STUNTING PADA BALITA DI WILAYAH KERJA PUSKESMAS KAYAMANYA KECAMATAN POSO KOTA KABUPATEN POSO TAHUN 2022

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 1-6

 <sup>1</sup> 2023

 <sup>1</sup> DOI: 10.31970/ma.v5i1.117

 O\_Accred : Sinta 5

#### KARAKTERISTIK DEMOGRAFI PENDERITA HIPERTENSI DI WILAYAH KERJA PUSKESMAS KAWATUNA PALU TAHUN 2022

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 5 No 1 (2023): April 40-44

 2023
 DOI: 10.31970/ma.v5i1.118
 O Accred : Sinta 5

 GAMBARAN FAKTOR RISIKO TERJADINYA KATARAK DI RUMAH SAKIT UMUM ANUTAPURA PALU DAN KLINIK SPESIALIS MATA MITRA

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 4 No 1 (2022): April 19-24

 <sup>1</sup> 2022

 <sup>1</sup> DOI: 10.31970/ma.v4i1.92

 OAccred : Sinta 5

 KARAKTERISTIK PENGGUNA NARKOBA DI POLI JIWA RSU MADANI PALU PERIODE OKTOBER-DESEMBER TAHUN 2021

 Fakultas Kedokteran Universitas Alkhairaat

 Medika Alkhairaat

 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 4 No 1 (2022): April 1-7

 D 2022

 Modika Alkhairaat

 Occered : Sinta 5

#### HUBUNGAN ANTARA LAMA MENDERITA ULKUS DIABETIKUM DENGAN TINGKAT DEPRESI TAHUN 2021

 Fakultas Kedokteran Universitas Alkhairaat
 Medika Alkhairaat : Jurnal Penelitian Kedokteran dan Kesehatan Vol 4 No 1 (2022): April 8-13

 <sup>1</sup> 2022

 <sup>1</sup> DOI: 10.31970/ma.v4i1.90

 O Accred : Sinta 5

View more ...