

Clinical Implications of Hand Position and Lower Limb Length Measurement Method on Y-Balance Test Scores and Interpretation

Kim Hébert-Losier, PhD, PT

Faculty of Health, Sport and Human Performance, Adams Centre for High Performance, University of Waikato, Mount Maunganui, Tauranga, New Zealand

Context: The Lower Quarter Y-Balance Test (LQ-YBT) was developed to provide an effective and efficient screen for injury risk in sports. Earlier protocol recommendations for the LQ-YBT involved the athlete placing the hands on the hips and the clinician normalizing scores to lower limb length measured from the anterior-superior iliac spine to the lateral malleolus. The updated LQ-YBT protocol recommends the athlete's hands be free moving and the clinician measure lower limb length to the medial malleolus.

Objective: To investigate the effect of hand position and lower limb length measurement method on LQ-YBT scores and their interpretation.

Design: Cross-sectional study.

Setting: National Sports Institute of Malaysia.

Patients or Other Participants: A total of 46 volunteers, consisting of 23 men (age = 25.7 ± 4.6 years, height = 1.70 ± 0.05 m, mass = 69.3 ± 9.2 kg) and 23 women (age = 23.5 ± 2.5 years, height = 1.59 ± 0.07 m, mass = 55.7 ± 10.6 kg).

Intervention(s): Participants performed the LQ-YBT with hands on hips and hands free to move on both lower limbs.

Main Outcome Measure(s): In a single-legged stance, participants reached with the contralateral limb in each of the anterior, posteromedial, and posterolateral directions 3 times. Maximal reach distances in each direction were normalized to lower limb length measured from the anterior-superior iliac spine to the lateral and medial malleoli. Composite scores (average of the 3 normalized reach distances) and anterior-reach differenc-

es (in raw units) were extracted and used to identify participants at risk for injury (ie, anterior-reach difference ≥4 cm or composite score ≤94%). Data were analyzed using paired *t* tests, Fisher exact tests, and magnitude-based inferences (effect size [ES], ±90% confidence limits [CL]).

Results: I observed differences between hand positions in normalized anterior-reach distances that were trivial ($t_{91} = -2.075$, $P = .041$; ES = 0.12, 90% CL = ±0.10). In contrast, reach distances were greater when the hands moved freely for the normalized posteromedial ($t_{91} = -6.404$, $P < .001$; ES = 0.42, 90% CL = ±0.11), posterolateral ($t_{91} = -6.052$, $P < .001$; ES = 0.58, 90% CL = ±0.16), and composite scores ($t_{91} = -7.296$, $P < .001$; ES = 0.47, 90% CL = ±0.11). A similar proportion of the cohort was classified as at risk with the hands on the hips (35% [$n = 16$]) and the hands free to move (43% [$n = 20$]; $P = .52$). However, the participants classified as at risk with the hands on the hips were not all categorized as at risk with the hands free to move and vice versa. The lower limb length measurement method exerted trivial effects on LQ-YBT outcomes.

Conclusions: Hand position exerted nontrivial effects on LQ-YBT outcomes and interpretation, whereas lower limb length measurement method had trivial effects.

Key Words: balance, injury risk, musculoskeletal system, injury prevention

Key Points

- Scientists and clinicians currently use different protocols when administering the Lower Quarter Y-Balance Test (LQ-YBT).
- Hand position (on the hips versus free to move) had a nontrivial effect on the LQ-YBT scores and their interpretation.
- Lower limb length measurement method (anterior-superior iliac spine to the medial or lateral malleolus) had a trivial effect on LQ-YBT outcomes.
- When using the LQ-YBT, upper limb placement needs to be clearly documented for replication purposes.
- Researchers should investigate individual responses to changes in hand position to better understand the mechanistic contribution of the upper limbs to LQ-YBT performance.

The Lower Quarter Y-Balance Test (LQ-YBT) involves maintaining single-legged balance while reaching as far as possible with the contralateral limb in 3 directions.¹ It is a simplified version of the Star Excursion Balance Test, which involves 8 reach directions.¹ The LQ-YBT has demonstrated good-to-excellent intrarater and interrater reliability^{2–5}; stability over time⁴; and the ability to predict injury incidence in several athletic

populations, including high school basketball players,⁴ collegiate athletes,^{6–8} and professional soccer players.⁹ It is used not only to screen for injury risk in athletes but also to monitor rehabilitation progression and outcomes after injury or surgery,^{10,11} examine the effects of training interventions in uninjured populations,¹² and assess dynamic balance across the age spectrum.^{13,14}

The widespread application of the LQ-YBT has led to the development and use of a variety of protocols.⁵ When the test was first used as an injury-screening tool, Plisky et al⁴ illustrated participants performing the test with their hands placed on their hips (ie, the study protocol did not explicitly state the hand position, but the photograph showed an individual performing the test with hands on hips). A few years later, Plisky et al⁵ illustrated participants performing the LQ-YBT with their hands free to move. Consequently, the LQ-YBT research contains data from the test being performed with both the hands on the hips^{3,4,11,12} and the hands free to move.^{5,6,8,13} Here again, investigators^{3–5,8,12,13} have not explicitly stated hand position but have provided photographs of individuals performing the LQ-YBT. Given that upper limb movement can improve performance during balance,¹⁵ mobility,¹⁵ and lower limb strength tests,¹⁶ it is reasonable to assume that permitting upper limb movement during the LQ-YBT could influence outcomes and augment reach distance. Similarly, when the LQ-YBT was first used to define injury-risk cutoff scores, Plisky et al⁴ normalized reach distances to lower limb length measured from the anterior-superior iliac spine (ASIS) to the lateral malleolus, but later, Plisky et al⁵ recommended normalizing scores to measures taken from the ASIS to the medial malleolus. Whereas the latter method^{3,5,6,14,17} is more frequently used than the former,^{4,8} differences in lower limb length measurement methods could affect LQ-YBT scores and their interpretation.¹⁸

In the absence of age-, sex-, and population-specific injury-risk cutoff scores, the initial thresholds reported by Plisky et al⁴ are often cited and used for reference.^{6,7,10,19} In particular, high school basketball players who presented with an anterior-reach distance difference ≥ 4 cm or a composite reach score $\leq 94\%$ of lower limb length were more likely to sustain a lower limb injury.⁴ In later years, Butler et al⁷ determined that collegiate American football players with composite-reach scores of less than 89.6% were at greater risk of injury, whereas the anterior-reach distance difference could not predict injury incidence. Other than the population groups investigated, differences among study results might be due in part to differences in LQ-YBT protocols: Butler et al⁷ used the Y-Balance Test Kit and adhered to the more recent protocol that allows the hands to move freely, measures lower limb length to the medial malleolus, and permits the stance foot to lift from the contact surface.⁵

Therefore, the purpose of this study was to investigate the effect of hand position and lower limb length measurement method on LQ-YBT scores and their interpretation using the conventional injury-risk cutoff scores. I hypothesized that participants would be able to reach farther with their hands free to move, thereby reducing the number of participants classified as being at risk for injury. I also expected longer lower limb length values when measuring from the ASIS to the lateral malleolus, which would lead to lower normalized LQ-YBT scores.

METHODS

Design

The cross-sectional research design with repeated measures required participants to attend 1 experimental

session in the biomechanics laboratory of the National Sports Institute of Malaysia. I conducted half of the test sessions in the morning (9 AM to noon) and half in the afternoon (2 to 5 PM), thereby balancing the effect of time of day on dynamic postural control.²⁰ To minimize the potential influence of testing order and fatigue on LQ-YBT performance, hand position and test side were block randomized so an equal number of participants began with either their hands on their hips or their hands free to move and began with the right or left lower limb. The first lower limb length measure taken was also alternated between the lateral and medial malleoli. The independent variables of interest were hand position (on the hips or free to move) and lower limb length measurement method (lateral malleolus or medial malleolus). The dependent variables were LQ-YBT maximal reach distance in the anterior, posteromedial, and posterolateral directions normalized to lower limb length; composite reach score normalized to lower limb length; absolute anterior-reach distance difference; lower limb length; and the proportions of the cohort identified as at risk and not at risk for injury. All participants provided written informed consent, and the study was approved by the Institutional Review Board at the National Sports Institute of Malaysia.

Participants

A total of 46 participants (age range = 20–38 years), including 23 men (age = 25.7 ± 4.6 years, height = 1.70 ± 0.05 m, mass = 69.3 ± 9.2 kg, right-foot dominant = 20) and 23 women (age = 23.5 ± 2.5 years, height = 1.59 ± 0.07 m, mass = 55.7 ± 10.6 kg, right-foot dominant = 20), completed the study protocol. I defined *foot dominance* as the foot used to kick a ball. Inclusion criteria were good self-reported general health (ie, no known disease, infection, or illness) and no musculoskeletal injury, pathologic joint condition, or other medical condition within the 3 months before the study that could affect LQ-YBT performance. Based on the short-form International Physical Activity Questionnaire,²¹ I categorized 24, 17, and 5 participants as having high, moderate, and low physical activity levels, respectively. Participants were instructed to refrain from strenuous physical activity or resistance training on the day of testing.

Procedures

I recorded sex, age, height, body mass, foot dominance, injury history, and physical activity level. With participants standing in an upright position without shoes and with their weight evenly distributed between limbs,⁶ the length of each lower limb was measured from the most inferior aspect of the ASIS to the most distal aspect of the lateral and medial malleoli. The method for measuring lower limb length (lateral malleolus or medial malleolus) was randomized and performed by a single examiner (not the author) throughout the study. Intrarater reliability of lower limb length measures was not assessed, but researchers²² have demonstrated excellent intrarater reliability values (intra-class correlation coefficient = 0.985–0.990). Due to the novelty of the task, participants subsequently completed a 5-minute warm-up on a cycling ergometer (RevMaster, LeMond, CA) at a self-selected light intensity before the LQ-YBT. Before the experimental trials, participants were

familiarized with the LQ-YBT and performed 6 practice trials on each foot in each reach direction and with each hand position to minimize any potential learning effect.² The familiarization period was followed by a 2-minute rest period in quiet standing.

Metric cloth measuring tapes were affixed to the floor to reconstruct the Y-shaped reach directions. A 23-cm long, 12.5-cm wide, 15.5-cm deep cardboard box weighing 150 g was used as a reach indicator. While standing barefoot on 1 limb in the middle of the Y shape, participants reached as far as possible with their free limb 3 times in the anterior, posteromedial, and posterolateral directions sequentially, with the 2 posterior directions located 135° from the anterior direction. The 3 trials were completed with 1 limb and then the other limb in the anterior-reach direction before the posteromedial-reach direction and then the posterolateral-reach direction. Between trials, the reach foot was placed on the ground beside the stance foot. After all the trials for a given hand position were completed, participants rested for 2 minutes before performing the trials with the alternate hand position. Trials were disregarded and repeated when a participant lost balance, lifted or moved the stance foot from the floor, touched down with the reach foot, kicked the reach indicator, placed the reach foot on top of the indicator, did not return to the starting position in a controlled manner, or removed the hands from the hips during the hands-on-hips trials.⁴ Throughout the study, a single examiner (not the author) recorded the reach distances from all trials. Before data collection, the examiner completed a series of training sessions with a qualified physical therapist to promote standardization of testing procedures, which included performing supervised LQ-YBT assessments a minimum of 20 times. Intrarater reliability was not assessed; however, the LQ-YBT has demonstrated good to excellent intrarater reliability values (intraclass correlation coefficient = 0.85–0.91).⁵

The greatest reach distance in each direction for each limb and hand position was retained for analysis and subsequently normalized to lower limb length. For each limb and hand position, a composite reach score was also calculated by summing the greatest reach distances in each of the 3 directions and normalizing the value to 3 times that of the lower limb length:

$$\begin{aligned} & \text{Composite reach score (\%)} \\ & \frac{\text{Anterior reach} + \text{posteromedial reach} \\ & \quad + \text{posterolateral reach}}{3 \times \text{Lower limb length}} \times 100 \end{aligned}$$

Finally, the absolute difference between the right and left anterior reach distances was calculated. The number of participants considered at risk or not at risk for injury based on their anterior-reach distance differences (≥ 4 cm) and composite reach scores ($\leq 94\%$ of lower limb length) was computed following the initial thresholds identified by Plisky et al.⁴ All data were entered into Excel (version 2007; Microsoft Corp, Redmond, WA) for further analysis.

Statistical Analysis

Means and standard deviations were computed for all variables. The effect of hand position on normalized

anterior, posteromedial, posterolateral, and composite scores was investigated using paired *t* tests, and the α level was set at .05. Data were analyzed for practical meaningfulness using magnitude-based inferences. Magnitudes of the standardized effect sizes (ESs) were interpreted using thresholds of <0.2 (*trivial*), 0.2 (*small*), 0.6 (*moderate*), 1.2 (*large*), and 2.0 (*very large*).²³ The uncertainty of the ES was expressed using 90% confidence limits (CL) in a plus/minus form (ie, ES, \pm CL), and I qualitatively evaluated the chance that the true value of the ES was practically meaningful using the following standardized thresholds: $<0.5\%$ (*almost certainly not*), 1% to 5% (*very unlikely*), $>5\%$ to 25% (*unlikely*), $>25\%$ to 75% (*possibly*), $>75\%$ to 95% (*likely*), $>95\%$ to 99.5% (*very likely*), and $>99.5\%$ (*almost certainly*).²³

The effect of hand position on the proportion of participants classified as at risk or not at risk for injury was also investigated using Fisher exact tests from 2×2 tables of frequencies and analyzed for practical meaningfulness. Differences between the 2 hand positions in the percentage of the cohort classified as at risk were interpreted using thresholds of $<10\%$ (*trivial*), 10% (*small*), 30% (*moderate*), 50% (*large*), and 70% (*very large*).²³ The same statistical procedures were used to analyze the effect of lower limb length measurement method on lower limb lengths; normalized anterior, posteromedial, posterolateral, and composite scores; and the proportions of the participants classified as at risk and not at risk for injury. Consistent with the original protocol used to establish the at-risk thresholds,⁴ the lower limb length measured using the lateral malleolus and the hands-on-hips position was considered the reference condition when investigating the effects of hand position and lower limb length measurement method on LQ-YBT scores and their interpretation. Results from the alternate conditions are provided as supplemental material. All data processing and analyses were performed using Excel (version 2007; Microsoft Corp).

RESULTS

Hand Position

Hand position had a significant but likely (89.5% likelihood) trivial effect ($t_{91} = -2.075$, $P = .041$; ES = 0.12, 90% CL = ± 0.10) on the normalized anterior reach distance (Figure 1). In contrast, participants reached farther with the hands-free-to-move than the hands-on-hips method in the posteromedial ($t_{91} = -6.404$, $P < .001$; ES = 0.42, 90% CL = ± 0.11), posterolateral ($t_{91} = -6.052$, $P < .001$; ES = 0.58, 90% CL = ± 0.16), and composite directions ($t_{91} = -7.296$, $P < .001$; ES = 0.47, 90% CL = ± 0.11), with small and almost certain (99.9% likelihood) differences in ES measures.

Absolute anterior-reach distance with the hands on the hips (3.2 ± 2.7 cm) did not differ from the hands-free-to-move condition (3.8 ± 2.5 cm; $t_{45} = -1.326$, $P = .19$; ES = 0.24, 90% CL = ± 0.31). Similar proportions of the cohort were identified as *at risk* and *not at risk* for injury based on the anterior-reach distance difference when the LQ-YBT was performed with the hands on the hips ($n = 16$ [35%] and $n = 30$ [65%], respectively) compared with the hands free to move ($n = 20$ [43%] and $n = 26$ [57%], respectively).




Figure 1. The effect of hand position (on the hips versus free to move) on Lower Quarter Y-Balance Test scores normalized to lower limb length measured from the anterior-superior iliac spine to the lateral malleolus. Error bars represent standard deviations. ^a Indicates difference between conditions ($P < .05$).

The difference between the 2 hand positions in the percentage of the cohort classified as at risk was trivial (8.7%) and not significant ($P = .52$). However, only 7 of the 16 participants categorized as at risk with the hands on the hips were also categorized at risk with the hands free to move (Figure 2A). The proportion of participants identified as at risk based on the normalized composite score was also similar between the hands-on-hips (15%) and the hands-free-to-move conditions (9%; $P = .52$), with a trivial difference in the percentage (6.5%). Again, not all participants identified as at risk with the hands on the hips were identified as at risk with the hands free to move (Figure 2B). Whereas the results for the effect of hand position are for scores normalized to lower limb lengths

measured to the lateral malleolus, the results were almost identical for lengths measured to the medial malleolus (see Supplemental Figures 1 and 2, available online at <http://dx.doi.org/10.4085/1062-6050-52.8.02.S1> and <http://dx.doi.org/10.4085/1062-6050-52.8.02.S2>).

Lower Limb Length Measurement Method

The lower limb was longer when measured from the ASIS to the lateral malleolus than to the medial malleolus ($t_{91} = -5.423$, $P < .001$; Figure 3); however, this difference was almost certainly (99.9% likelihood) trivial ($ES = -0.09$, 90% $CL = \pm 0.04$). Similarly, whereas significant, the differences in the normalized anterior ($t_{91} = 5.497$, $P < .001$; $ES = 0.04$, 90% $CL = \pm 0.01$), posteromedial ($t_{91} =$




Figure 2. Venn diagram illustrating the numbers of participants classified as at risk and not at risk when performing the Lower Quarter Y-Balance Test with the hands on the hips and the hands free to move on the basis of the A, anterior-reach distance difference (≥ 4 cm) and, B, composite reach score ($\leq 94\%$) normalized to lower limb length measured from the anterior-superior iliac spine to the lateral malleolus. Diagrams are not precisely to scale.




Figure 3. The effect of lower limb length measurement method (anterior-superior iliac spine to the lateral malleolus versus anterior-superior iliac spine to the medial malleolus) on lower limb length and Lower Quarter Y-Balance Test scores normalized to lower limb length when performed with the hands on the hips. Error bars represent standard deviations. ^a Indicates difference between conditions ($P < .05$).

5.386, $P < .001$; ES = 0.05, 90% CL = ± 0.01), posterolateral ($t_{91} = 5.362$, $P < .001$; ES = 0.06, 90% CL = ± 0.02), and composite reach scores ($t_{91} = -4.093$, $P < .001$; ES = 0.06, 90% CL = ± 0.02) between lower limb length measurement methods were almost certainly (99.9% likelihood) trivial.

The proportions of the cohort identified as at risk and not at risk based on the normalized composite score were nearly identical ($P > .99$) when measured using the lateral malleolus ($n = 10$ [22%] and $n = 36$ [78%], respectively) and the medial malleolus ($n = 11$ [24%] and $n = 35$ [76%], respectively). Only 1 participant was categorized differently between the 2 measurement methods. Whereas the results pertaining to the effect of the lower limb length measurement method are for data from the LQ-YBT performed with the hands on the hips, the results were almost identical when the hands were free to move (see Supplemental Figure 3, available online at <http://dx.doi.org/10.4085/1062-6050-52.8.02.S3>).

DISCUSSION

The purpose of this study was to investigate the effects of hand position and method of measuring lower limb length on LQ-YBT scores and their interpretation using the original injury-risk cutoff thresholds established by Plisky et al.⁴ As anticipated, participants reached farther when their hands were free to move than when they were placed on their hips, with nontrivial differences detected for the normalized posteromedial, posterolateral, and composite reach scores. Whereas a similar number of individuals were identified as at risk for lower limb injury (ie, anterior-reach distance difference ≥ 4 cm or composite reach score $\leq 94\%$) with the hands on the hips and the hands free to move, only a subset of these individuals were categorized as at risk under both conditions, suggesting that different aspects of dynamic balance were involved. Measuring lower limb length from the ASIS to the lateral malleolus compared

with the medial malleolus affected most comparisons, but the method of measuring lower limb length had an almost certainly trivial effect on these comparisons (ie, lower limb length values and normalized anterior-, posteromedial-, posterolateral-, and composite-reach scores). Furthermore, the classification of participants based on the normalized composite scores was the same between lower limb length measurement methods, with the exception of 1 individual whose scores were on the borderline of the cutoff threshold. Hence, clinicians and scientists should consider that hand position during the LQ-YBT can significantly and non-trivially affect test scores and their interpretation and may elicit different neuromuscular-control strategies. In contrast, whether the lateral or medial malleolus is used to measure lower limb length for subsequent score normalization is a minimal concern.

When using the LQ-YBT in a preparticipation examination of 200 National Collegiate Athletic Association Division I collegiate athletes, Smith et al⁶ found that athletes with an anterior-reach distance difference of ≥ 4 cm had greater odds of injury; however, the normalized composite reach score did not predict injury incidence. In that study, athletes completed the task with their hands free to move, and the researchers measured lower limb length from the ASIS to the medial malleolus with the athletes standing and evenly distributing their weight between the lower limbs. Smith et al⁶ proposed assessing anterior-reach distance difference as part of a preparticipation screening to identify individuals at risk for injury and eliminating the posteromedial- and posterolateral-reach directions, thereby reducing the time needed for LQ-YBT assessment and allowing either more individuals or more tests to be included in a preparticipation examination of fixed duration. If a clinician is assessing only the anterior-reach distance difference, measuring lower limb length would become unnecessary because the metric represents the absolute difference (in centimeters) between the right and left anterior-reach distances (ie, no normalization). In this

study, the anterior-reach distance difference did not differ between the 2 hand positions examined ($P = .19$), suggesting less concern about upper body position when assessing this particular metric. However, more prospective studies are needed to verify whether the anterior-reach distance difference on the LQ-YBT is sufficient to identify individuals as at risk for lower limb injury.

In the current study, the anterior-reach distance difference identified a greater number (approximately 3 times more) of participants at potential risk for lower limb injury than did the composite score, which was likely due to the lack of sensitivity and specificity of the composite cutoff score used ($\leq 94\%$ of lower limb length) to identify individuals as at risk. The composite cutoff score should be based on sex, sport, and age.^{8,24} In the absence of population-specific cutoff scores or access to the Move2Perform injury-risk algorithm (monthly paid subscription; Move2Perform LLC, Evansville, IN), the threshold of 94% is often used as a clinical guideline or scientific reference,^{6,7,10,19} which was the reason for using this threshold in the current study. That aside, only 1 of the 7 participants identified as at risk in my study using the composite score was also identified as at risk using the anterior-reach distance difference when the hands were placed on the hips. In contrast, when the hands were free to move, 3 of 4 participants categorized as at risk using the composite score were also categorized as at risk using the anterior-reach distance difference. These results highlight that, particularly with the hands on the hips, anterior-reach distance difference and composite reach scores may identify individuals with distinct functional movement-impairment patterns.

Similarly, whereas the proportion of the cohort categorized as at risk for injury was similar between the 2 hand positions investigated, the individuals categorized as at risk were not necessarily the same. These findings suggest a shift in the neuromuscular-control strategies used to perform the LQ-YBT when the upper limbs are restricted compared with free moving, which is supported by empirical evidence of change in dynamic task performance when upper limb motion is restricted.^{16,25,26} For instance, limiting upper limb motion during running increases shoulder and pelvic rotations about the vertical axis²⁵ and can impede the recovery of gait stability measures after external perturbations.²⁶ In my study, restricting upper limb motion led to a change in the categorization of certain individuals from at risk to not at risk for injury (and vice versa). More in-depth investigations are required to further explain the individual responses to change in the relative contribution of the upper limb to LQ-YBT score observed in this study. Comparing LQ-YBT scores between restricted and free upper limb motion may provide a method for assessing the effectiveness of using the upper limb for dynamic balance, with implications for balance recovery after an unexpected disturbance in sports or the elderly population. For now, clinicians and scientists should consider that upper limb position during the LQ-YBT can influence the categorization of individuals within a cohort.

Authors^{10,11,13,14} of an increasing body of literature have provided data pertaining to the LQ-YBT in various population groups. Several factors influence LQ-YBT scores, including age,¹⁴ sex,^{17,19} and level²⁷ and type²⁸ of sport participation. In addition, variations in protocols and procedures affect LQ-YBT performance. For instance,

anterior-reach distance is greater and associated with less hip flexion at the point of maximal reach when executed from the ground without using a reach indicator than on the Y-Balance Test kit using a reach indicator.²⁹ This study provided additional insights into the LQ-YBT by highlighting that performance differences also exist between hands-on-hips and hands-free-to-move protocols, which are both frequently used. In future studies, researchers may seek to compare the kinematics of the LQ-YBT using the 2 hand positions to better understand the movement control of this task. Restricting upper limb motion during functional performance testing is believed to provide a more specific assessment of lower limb function,^{15,30} whereas permitting upper limb motion is believed to be more natural and functional.^{31,32} I cannot recommend using 1 protocol over another because the clinical aims need to be considered. Within health, research, and sport centers, consistently using 1 protocol is important to establish baseline scores, track changes over time, and determine population-specific injury-risk cutoff scores. When using published LQ-YBT data as a reference, following the protocols described is important, as variations could influence scores and the classification of individuals as at risk or not at risk. As such, placing the hands on the hips is advised if adhering to the cutoff scores established by Plisky et al.,⁴ whereas allowing the hands to move freely can be recommended if referring to cutoff scores provided by Butler et al.⁷ When reporting results from the LQ-YBT, clinicians and scientists alike should specifically describe the upper limb placement for replication purposes using both photographs and explicit writing to avoid misinterpretation of protocols. A direct comparison or agglomeration of results among studies involving the LQ-YBT without accounting for hand position is not advised; however, the concern is minimal regarding whether lower limb length is measured from the ASIS to the lateral or medial malleolus because of a trivial effect on LQ-YBT scores and almost no effect on the categorization of individuals.

CLINICAL RELEVANCE

The LQ-YBT is a convenient, reliable, and valid tool used to assess dynamic balance and predict the occurrence of lower limb injuries in athletes.^{5,6,9} Across studies and health, research, and sport centers, hand position and the method for measuring lower limb length have differed, with no previous knowledge about how such variations could influence test scores and their interpretation. This study provided evidence that hand position does significantly and nontrivially affect LQ-YBT scores and their interpretation, whereas the effect of lower limb length measurement method is trivial. Direct comparisons between, or inferences from, different studies or centers without considering hand position is not advised. More in-depth investigations into individual responses to a change in hand position on LQ-YBT performance are required to further comprehend the mechanistic contribution of upper limb motion to this dynamic task. When using the LQ-YBT, upper limb placement needs to be clearly documented for replication purposes.

ACKNOWLEDGMENTS

The data for this study were collected at the National Sports Institute of Malaysia. I thank Lilyana Nooryusra Binti Abdul

Gaffor and Amirah Binti Zahiran for their contributions to the data-collection process, as well as the volunteers for their participation in this study.

REFERENCES

1. Hertel J, Braham RA, Hale SA, Olmsted-Kramer LC. Simplifying the Star Excursion Balance Test: analyses of subjects with and without chronic ankle instability. *J Orthop Sports Phys Ther.* 2006;36(3):131–137.
2. Hertel J, Miller SJ, Denegar CR. Intratester and intertester reliability during the Star Excursion Balance Tests. *J Sport Rehabil.* 2000;9(2):104–116.
3. Shaffer SW, Teyhen DS, Lorenson CL, et al. Y-Balance Test: a reliability study involving multiple raters. *Mil Med.* 2013;178(11):1264–1270.
4. Plisky PJ, Rauh MJ, Kaminski TW, Underwood FB. Star Excursion Balance Test as a predictor of lower extremity injury in high school basketball players. *J Orthop Sports Phys Ther.* 2006;36(12):911–919.
5. Plisky PJ, Gorman PP, Butler RJ, Kiesel KB, Underwood FB, Elkins B. The reliability of an instrumented device for measuring components of the star excursion balance test. *N Am J Sports Phys Ther.* 2009;4(2):92–99.
6. Smith CA, Chimera NJ, Warren M. Association of Y Balance Test reach asymmetry and injury in Division I athletes. *Med Sci Sports Exerc.* 2015;47(1):136–141.
7. Butler RJ, Lehr ME, Fink ML, Kiesel KB, Plisky PJ. Dynamic balance performance and noncontact lower extremity injury in college football players: an initial study. *Sports Health.* 2013;5(5):417–422.
8. Lehr ME, Plisky PJ, Butler RJ, Fink ML, Kiesel KB, Underwood FB. Field-expedient screening and injury risk algorithm categories as predictors of noncontact lower extremity injury. *Scand J Med Sci Sports.* 2013;23(4):e225–e232.
9. Gonell AC, Romero JA, Soler LM. Relationship between the Y Balance Test scores and soft tissue injury incidence in a soccer team. *Int J Sports Phys Ther.* 2015;10(7):955–966.
10. Boyle MJ, Butler RJ, Queen RM. Functional movement competency and dynamic balance after anterior cruciate ligament reconstruction in adolescent patients. *J Pediatr Orthop.* 2016;36(1):36–41.
11. Clagg S, Paterno MV, Hewett TE, Schmitt LC. Performance on the modified star excursion balance test at the time of return to sport following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2015;45(6):444–452.
12. Cloak R, Nevill A, Wyon M. The acute effects of vibration training on balance and stability amongst soccer players. *Eur J Sport Sci.* 2016;16(1):20–26.
13. Faigenbaum AD, Myer GD, Fernandez IP, et al. Feasibility and reliability of dynamic postural control measures in children in first through fifth grades. *Int J Sports Phys Ther.* 2014;9(2):140–148.
14. Lee DK, Kang MH, Lee TS, Oh JS. Relationships among the Y Balance Test, Berg Balance Scale, and lower limb strength in middle-aged and older females. *Braz J Phys Ther.* 2015;19(3):227–234.
15. Markovic G, Dizdar D, Jukic I, Cardinale M. Reliability and factorial validity of squat and countermovement jump tests. *J Strength Cond Res.* 2004;18(3):551–555.
16. Milosevic M, McConville KM, Masani K. Arm movement improves performance in clinical balance and mobility tests. *Gait Posture.* 2011;33(3):507–509.
17. Alnahdi AH, Alderaa AA, Aldali AZ, Alsobayel H. Reference values for the Y Balance Test and the lower extremity functional scale in young healthy adults. *J Phys Ther Sci.* 2015;27(12):3917–3921.
18. Terry MA, Winell JJ, Green DW, et al. Measurement variance in limb length discrepancy: clinical and radiographic assessment of interobserver and intraobserver variability. *J Pediatr Orthop.* 2005;25(2):197–201.
19. Engquist KD, Smith CA, Chimera NJ, Warren M. Performance comparison of student-athletes and general college students on the Functional Movement Screen and the Y Balance Test. *J Strength Cond Res.* 2015;29(8):2296–2303.
20. Gribble PA, Tucker WS, White PA. Time-of-day influences on static and dynamic postural control. *J Athl Train.* 2007;42(1):35–41.
21. Craig CL, Marshall AL, Sjöström M, et al. International Physical Activity Questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc.* 2003;35(8):1381–1395.
22. Neelly K, Wallmann HW, Backus CJ. Validity of measuring leg length with a tape measure compared to a computed tomography scan. *Physiother Theory Pract.* 2013;29(6):487–492.
23. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3–13.
24. Cook G, Plisky P. *YBT*. Chatham, VA: Functional Movement Systems; 2015:1–30.
25. Arellano CJ, Kram R. The metabolic cost of human running: is swinging the arms worth it? *J Exp Biol.* 2014;217(pt 14):2456–2461.
26. Bruijn SM, Meijer OG, Beek PJ, van Dieën JH. The effects of arm swing on human gait stability. *J Exp Biol.* 2010;213(pt 23):3945–3952.
27. Bullock GS, Arnold TW, Plisky PJ, Butler RJ. Basketball players' dynamic performance across competition levels [published online February 5, 2016]. *J Strength Cond Res.* DOI: 10.1519/JSC.000000000001372.
28. Bressel E, Yonker JC, Kras J, Heath EM. Comparison of static and dynamic balance in female collegiate soccer, basketball, and gymnastics athletes. *J Athl Train.* 2007;42(1):42–46.
29. Fullam K, Caulfield B, Coughlan GF, Delahunty E. Kinematic analysis of selected reach directions of the Star Excursion Balance Test compared with the Y-Balance Test. *J Sport Rehabil.* 2014;23(1):27–35.
30. Hébert-Losier K, Beaven CM. The MARS for squat, countermovement, and standing long jump performance analyses: are measures reproducible? *J Strength Cond Res.* 2014;28(7):1849–1857.
31. Laffaye G, Wagner P, Tombleson T. Countermovement jump height: gender and sport-specific differences in the force-time variables. *J Strength Cond Res.* 2013;28(4):1096–1105.
32. Feltner ME, Frascchetti DJ, Crisp RJ. Upper extremity augmentation of lower extremity kinetics during countermovement vertical jumps. *J Sports Sci.* 1999;17(6):449–466.

SUPPLEMENTAL MATERIAL

Supplemental Figure 1.

Found at DOI: <http://dx.doi.org/10.4085/1062-6050-52.8.02.S1>

Supplemental Figure 2.

Found at DOI: <http://dx.doi.org/10.4085/1062-6050-52.8.02.S2>

Supplemental Figure 3.

Found at DOI: <http://dx.doi.org/10.4085/1062-6050-52.8.02.S3>

Address correspondence to Kim Hébert-Losier, PhD, PT, Faculty of Health, Sport and Human Performance, Adams Centre for High Performance, University of Waikato, 52 Miro Street, Mount Maunganui, Tauranga 3116, New Zealand. Address e-mail to kim.hebert-losier@waikato.ac.nz.

Online First