

Chemical and nutritional characteristics, and microbial degradation of rapeseed meal recalcitrant carbohydrates

Citation for published version (APA):

Long, C., Qi, X-L., & Venema, K. (2022). Chemical and nutritional characteristics, and microbial degradation of rapeseed meal recalcitrant carbohydrates: A review. *Frontiers in nutrition*, 9, 948302. <https://doi.org/10.3389/fnut.2022.948302>

Document status and date:

Published: 01/01/2022

DOI:

[10.3389/fnut.2022.948302](https://doi.org/10.3389/fnut.2022.948302)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



OPEN ACCESS

EDITED BY

Shabir Ahmad Mir,
Government College for Women,
Srinagar, India

REVIEWED BY

Fang Geng,
Chengdu University, China
Manzoor Ahmad Shah,
Government College for Women
Anantnag, India
Mudasir Bashir Mir,
Govt. Degree College Baramulla, India

*CORRESPONDENCE

Koen Venema
k.venema@maastrichtuniversity.nl

SPECIALTY SECTION

This article was submitted to
Food Chemistry,
a section of the journal
Frontiers in Nutrition

RECEIVED 19 May 2022

ACCEPTED 31 August 2022

PUBLISHED 28 September 2022

CITATION

Long C, Qi X-L and Venema K (2022)
Chemical and nutritional
characteristics, and microbial
degradation of rapeseed meal
recalcitrant carbohydrates: A review.
Front. Nutr. 9:948302.
doi: 10.3389/fnut.2022.948302

COPYRIGHT

© 2022 Long, Qi and Venema. This is
an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](#).
The use, distribution or reproduction
in other forums is permitted, provided
the original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

Chemical and nutritional characteristics, and microbial degradation of rapeseed meal recalcitrant carbohydrates: A review

Cheng Long^{1,2}, Xiao-Long Qi¹ and Koen Venema^{2*}

¹Animal Science and Technology College, Beijing University of Agriculture, Beijing, China, ²Faculty of Science and Engineering, Centre for Healthy Eating and Food Innovation, Maastricht University - Campus Venlo, Venlo, Netherlands

Approximately 35% of rapeseed meal (RSM) dry matter (DM) are carbohydrates, half of which are water-soluble carbohydrates. The cell wall of rapeseed meal contains arabinan, galactomannan, homogalacturonan, rhamnogalacturonan I, type II arabinogalactan, glucuronoxylan, XXGG-type and XXXG-type xyloglucan, and cellulose. Glycoside hydrolases including in the degradation of RSM carbohydrates are α -L-Arabinofuranosidases (EC 3.2.1.55), endo- α -1,5-L-arabinanases (EC 3.2.1.99), Endo-1,4- β -mannanase (EC 3.2.1.78), β -mannosidase (EC 3.2.1.25), α -galactosidase (EC 3.2.1.22), reducing-end-disaccharide-lyase (pectate disaccharide-lyase) (EC 4.2.2.9), (1 \rightarrow 4)-6-O-methyl- α -D-galacturonan lyase (pectin lyase) (EC 4.2.2.10), (1 \rightarrow 4)- α -D-galacturonan reducing-end-trisaccharide-lyase (pectate trisaccharide-lyase) (EC 4.2.2.22), α -1,4-D-galacturonan lyase (pectate lyase) (EC 4.2.2.2), (1 \rightarrow 4)- α -D-galacturonan glycanohydrolase (endo-polygalacturonase) (EC 3.2.1.15), Rhamnogalacturonan hydrolase, Rhamnogalacturonan lyase (EC 4.2.2.23), Exo- β -1,3-galactanase (EC 3.2.1.145), endo- β -1,6-galactanase (EC 3.2.1.164), Endo- β -1,4-glucanase (EC 3.2.1.4), α -xylosidase (EC 3.2.1.177), β -glucosidase (EC 3.2.1.21) endo- β -1,4-glucanase (EC 3.2.1.4), exo- β -1,4-glucanase (EC 3.2.1.91), and β -glucosidase (EC 3.2.1.21). In conclusion, this review summarizes the chemical and nutritional compositions of RSM, and the microbial degradation of RSM cell wall carbohydrates which are important to allow to develop strategies to improve recalcitrant RSM carbohydrate degradation by the gut microbiota, and eventually to improve animal feed digestibility, feed efficiency, and animal performance.

KEYWORDS

rapeseed meal, animal performance, cell wall polysaccharides, carbohydrates, gut microbial degradation

Introduction

The main ingredients of animal feeds are plant carbohydrates, which consist of more than 70% of dry matter (DM). Plant carbohydrates can be classified as non-structural carbohydrates (low molecular weight sugars, oligosaccharides, and storage polysaccharides) and structural polysaccharides (non-starch polysaccharides) (1). Non-starch polysaccharides (NSP) cannot be broken down by endogenous digestive enzymes, nor be absorbed in the upper gastrointestinal tract (GIT). However, they can be (partly) degraded by the gut microbiota in the large intestine, and absorbed by the host in the form of short-chain fatty acids (SCFA), in this manner contributing to additional energy extraction from the diet. In addition, NSP may affect the digestion of other nutrients by the means of physical hindrance or physiological changes in the gut, such as increased digesta viscosity (2). Thus, it is vital to understand the NSP composition and structure of feed ingredients to improve animal feed efficiency.

Rapeseed meal (RSM), a byproduct of rapeseed oil production, is not only a suitable protein source but also a potential energy source as animal feed. Non-structural carbohydrates make up ~3–5% of the DM. Meal production processes could affect fiber production from RSM. These include expeller pressed (rapeseed oil is physically extracted using heat), cold pressed (rapeseed oil is physically extracted without heat treatment) and solvent extraction (rapeseed oil is extracted from the meal by physical expeller extraction followed by solvent washing). Previous research showed that expeller-pressed meal had more crude fiber compared to the cold-pressed meal (3). Another research demonstrated that solvent-extraction meal had more neutral detergent fiber and acid detergent fiber compared to the cold-pressed meal (4). Non-starch polysaccharides constitute 20–40% of RSM (5, 6), which are represented by pectic polysaccharides (homogalacturonan, rhamnogalacturonan I, arabinan, galactomannan, and arabinogalactan), hemicelluloses (xyloglucan and glucuronoxylan), and cellulose (7). Reports show that the NSP can only be degraded for 3–6% in chickens (5, 8, 9), and around 58–68% in pigs. This is rather low compared to other NSP-rich feed ingredients, such as sugar beet pulp (~85% of NSP is degraded by pigs) (1). Knowledge about the relationship between the chemical structure of NSP and GIT degradation is scant.

Plant cell walls are composed of a primary and a secondary layer, which are both built from polysaccharides, lignin, and protein. Polysaccharides contribute the most to the cell wall composition. The primary cell wall of RSM contains pectin, cellulose, and xyloglucan. The cellulose microfibrils are interlinked with xyloglucan *via* hydrogen bonds forming a stiff network (10). Pectins are linked to each other and cross-linked with (Hemi)cellulose (11, 12). In the secondary cell wall of

RSM, the main carbohydrates are 4-*o*-methylglucuronoxylan, xyloglucan, and cellulose (13). This review aims to provide an overview of the carbohydrate composition of RSM, as well as the microbial degradation of RSM. Rapeseed derives from several species belonging to the genus *Brassicaceae*. In the current review, *B. napus* is discussed.

Nutritional characteristics of RSM carbohydrates

Carbohydrates in rapeseed meal (*B. napus*) are mainly pectins and (Hemi)celluloses and comprise 35–36% of the RSM dry matter (DM) (7, 14). The common feedstuff analysis shows commercial RSM contains 12.1% crude fiber and 34% nitrogen-free extract (NFE) (Table 1). Carbohydrates of RSM can be categorized into non-structural carbohydrates and structural carbohydrates: a portion of the NFE is non-structural carbohydrates, whereas the crude fiber and the remainder of the NFE are structural polysaccharides. The composition of the different categories of RSM carbohydrates is displayed in Table 1.

Non-structural carbohydrates

The non-structural carbohydrates in RSM are comprised of low molecular weight sugars, oligosaccharides, and storage polysaccharides (Figure 1).

Low molecular weight sugars

Low molecular weight sugars mainly are sucrose (2.3–2.9%), fructose (0.05–0.16%), and glucose (0.05–0.16%) (15).

Oligosaccharides

The primary oligosaccharides found in RSM are raffinose, stachyose, galactinol, and Myo-inositol. Of those, stachyose has the highest concentration (0.4–0.5%), followed by raffinose (0.05–0.16%), galactinol (0.1%), and Myo-inositol (0.1%) (6).

Storage polysaccharides

Only low concentrations of storage polysaccharides are present in RSM. The primary storage polysaccharide is starch. The starch content of the seed approaches 50% during early development, but starch almost completely disappears when energy stores are converted into oil as the seed matures. Rommi et al. (14) reported that starch concentrations found in the intact seed were the same as dehulled rapeseed press cakes (0.2% of DM), while the starch content of canola meal was up to 2.5% of DM (17). Starch granules are comprised of two

main macromolecules: amylose and amylopectin. Amylose is a linear polymer consisting of α -1,4-linked D-glucose units (GU) and includes 500–600 GU in its native form, which can be subdivided into 1–20 chains. Amylopectin is a large branched macromolecule (molecular weight 10^7 - 10^9 kDa), with ~5% of

its total linkages consisting of β -1,6-linked GU plus a large number of short α -1,4-linked linear GU chains (12–70 GU) (18).

TABLE 1 Carbohydrates composition of rapeseed meal (oil-free dry matter)^a.

Component, %	RSM
Crude fiber	12.1
Ether extract	3.8–4.1
Acid detergent fiber	17.3
Neutral detergent fiber	22.7
Lignin	2.6
Non-structural carbohydrates ^b	3.2
Structural carbohydrates ^c	29
Nitrogen free extract	34
Monosaccharides composition, mol %	
Glucose	40
Arabinose	17
Galactose	10
Xylose	9
Fructose	1
Mannose	2
Rhamnose	2
Uronic acid	20
Water-soluble carbohydrates	18.7
Monosaccharides and sucrose	16.7
Polysaccharides	2.0
Insoluble carbohydrates	15.8
Carbohydrate total	34.5–36

^aAdapted from references (7, 14–16).

^bFructose, glucose, galactose, sucrose, galactinol, raffinose and stachyose.

^cPectins, cellulose residue, amylose, arabinan, arabinogalactan, xyloglucan.

Structural polysaccharides

The largest part of the RSM carbohydrates is made up of structural polysaccharides (Figure 1). These include cellulose, hemicellulose (xyloglucan and xylan), and pectic polysaccharides (homogalacturonan, rhamnogalacturonan I, arabinan, and arabinogalactan).

Polysaccharides composition

RSM contains 34.5% NSP of its DM, and RSM is high in glucosyl (40 mol%), arabinosyl (19 mol%), and uronyl residues (18 mol%) (7, 19). Water-soluble and water-insoluble carbohydrates comprise 18.7 and 15.8% of the total carbohydrates content of RSM, respectively (Table 1). Water-soluble carbohydrates mainly contain glucosyl (64 mol%) and some galactosyl residues (17 mol%) (7), while water-insoluble carbohydrates are mainly glucosyl (32 mol%), arabinosyl (25 mol%), uronyl (18 mol%), and xylosyl residues (12%) (Table 2).

The detailed monosaccharide constituent compositions of the Water unextractable solids (WUS), which were sequentially extracted with (i) chelating agent (ChSS) to release calcium-bound pectins, (ii) dilute alkali (DASS) to release pectins tightly bound to hemicellulose, (iii) 4, and (iv) 6 molar alkali (4 MASS and 6 MASS) to release hemicelluloses, leaving cellulose that remains in the residue (RES), are presented in Table 2. The composition of the glycosidic residues indicates the presence of α -1,5-linked arabinan branched at O-2, galactomannan, homogalacturonan, rhamnogalacturonan I, type II arabinogalactan, glucuronoxylan, XXGG-type and XXXG-type xylo(X)glucan(G), and cellulose in RSM (7, 14, 20). Arabinan consists of a linear backbone of α -L-1,5-linked arabinose (Ara) units with α -1,2-linked or α -1,3-linked Ara units (Figure 2). Galactomannan has a backbone of β -1,4-linked




FIGURE 1 Schematic representation of carbohydrates presented in rapeseed meal.

TABLE 2 Molar sugar composition of RSM fractions^a.

	Molar composition (mol%)						
	Rha	Ara	Xyl	Man	Gal	Glc	UA
RSM	2	19	8	6	10	40	15
WSS	Tr ^b	7	1	7	17	64	5
WUS	1	25	12	4	8	32	18
CHSS	2	15	4	2	4	3	71
DASS	Tr	44	6	3	10	9	29
4MASS	Tr	17	20	15	13	30	5
6MASS	1	29	22	2	13	23	11
RES	4	10	6	4	6	40	20

^aAdapted from reference (7).

^bTrace amounts.

mannose units, substituted with α -1,6-linked galactose units (Figure 1). Homogalacturonan (HG) is the simplest form of pectin, consisting of a linear chain of α -1,4-linked galacturonic acid, and part of its carboxyl groups esterified with acetyl or methyl groups (Figure 2). Rhamnogalacturonan I (RG-I) consists of alternating rhamnose (Rha) and galacturonic acid (GalA) residues [-2)- α -Rha-(1,4)- α -D-GalA-(1,-), which is highly ramified with single terminal β -D-Gal and/or α -D-Ara at position O-4 or O-3 of the rhamnosyl residues where the α -D-GalA residues are often O-acetyl esterified at O-2 and/or O-3 (Figure 2) (21). Type II arabinogalactan (AG) is composed of a β -1,3-galactan backbone and β -1,6-galactan side chains (Figure 2). The side chains are variably decorated with L-arabinose. Glucuronoxylan (GX) has a linear backbone of β -(1-4)-linked D-xylosyl (Xyl) residues, which can be ramified with acetyl and arabinosyl residues, and some of the Xyl residues are decorated with a single α -D-glucuronic acid (GlcA) or 4-O-methyl-D-glucuronic acid (MeGlcA) residue at O2 (Figure 2) (22, 23). Both XXGG- and XXXG-type xyloglucan (XG) exist in RSM. XXGG (Figure 2) consists of a β -1,4-linked D-glucosyl (Glc) backbone carrying various side chains of D- β -1,2-Gal-D- α -1,6-Xyl and L- α -1,2-Ara-D- β -D- α -1,6-Xyl (24), whereas, XXXG (Figure 2) comprises a β -1,4-linked D-glucosyl backbone carrying continuous side chains of D- α -1,6-Xyl, D- β -1,2-Gal-D- α -1,6-Xyl, and L- α -1,2-Ara-D- β -1,2-Gal-D- α -1,6-Xyl in every 4 residues of D- β -1,4-Glc (24). Cellulose consists of long linear chains of β -1,4-linked D-Glc residues, with a degree of polymerization between 2,000 and 14,000 residues (Figure 2). The backbone of these polysaccharides can also be esterified with methyl-esters, ethyl-esters, and glycosyls (arabinosyl, galactosyl, mannosyl, glucosyl, xylosyl, or frucosyl).

Except for containing glucosyl residues, RES (Table 2) also has quite some arabinosyl and uronyl residues (10 and 20%, respectively), which indicates that some pectic polysaccharides are tightly associated with cellulose microfibrils. This suggests

that the cell wall polysaccharide matrix of rapeseed meal is strongly interlinked. The specific structures of dietary fibers of RSM are still not entirely understood. The main restriction may be the analysis method, but nowadays, the comprehensive microarray polymer profiling (CoMPP) technique is a powerful tool for probing cell wall structure studies (25). The profiles generated by CoMPP provide a global snapshot of cell-wall composition. It cannot only detect the amount of the particular polysaccharides but also their linkage type as discussed in the next section.

Glycosidic linkage type

A covalent link between one carbohydrate molecule and a second carbohydrate molecule is called a glycosidic bond. Glycosidic linkage is the type of bond between two adjacent glycosides in the chain of polysaccharides (26).


The proposed sugar linkage compositions of RSM fractions are shown in Table 3. The data can only be used in a qualitative manner instead of quantitative, due to the poor DMSO solubility of the fractions and the high amount of uronic acids (which are not detected with this method) present in some samples (27).

Microbial degradation of RSM cell wall carbohydrates

Low molecular weight sugars and starch can be 100% digested and/or absorbed, while (oligo- and poly-)saccharides are considered indigestible in the small intestine due to the lack of the necessary enzymes in monogastric animals (8). Indeed, mammalian genomes do not encode most of the enzymes needed to degrade the structural polysaccharides present in plant material. Instead, a complex mutual dependence has developed between the mammalian host and symbiotic gut microorganisms that do possess the ability to access the abundant source of energy in carbohydrates that are indigestible by the host. The gut microbiota has glycoside hydrolases (GH) that can degrade the oligo- and polysaccharides into small oligomers and monosaccharides which are subsequently taken up and fermented.

Glycoside hydrolases

Glycoside hydrolases (GHs) are a vast repertoire of cell wall-degrading enzymes that hydrolyze glycosidic bonds between two or more carbohydrate modules or sugar and a non-sugar moiety within carbohydrates or oligosaccharides (28, 29). GH families widely exist in prokaryotic, eukaryotic, and archaea species. A total of 173 GH families have been identified until now (accessed on May-2022, <http://www.cazy.org/Glycoside-Hydrolases.html>).



Carbohydrate binding modules (CBMs) are the non-catalytic part of cell-wall-degrading enzymes, and they are attached to the GH catalytic modules. Usually, CBMs have to recognize and bind to the specific polysaccharides first, before the GHs cleave the polysaccharides (30).

Degradation of arabinan

Arabinan can be hydrolyzed by α -L-Arabinofuranosidases (EC 3.2.1.55) and endo- α -1,5-L-arabinanases (EC 3.2.1.99), which are found in GH families 3, 43, 51, 54, and 62, and which release arabinosyl oligomers and L-arabinose (31). A previous study reported that *Bacillus subtilis* contained a series of arabinan-degrading genes, *abnA*, *abn2*, *abfA*, and *abf2*, which were induced by arabinose and arabinan, repressed by glucose, and subjected to temporal regulation (32). *AbnA* and *abn2* encode extracellular endo- α -1,5-L-arabinanases belonging to GH43, which hydrolyzes arabinan (branched) and linear α -1,5-L-arabinan and produces arabinose and arabino-oligosaccharides (33). These resulting products, are subsequently transported into the cell by different transport systems. Arabinose enters the cell mainly through the *araE* permease (34), and the uptake of arabinose oligomers occurs most likely *via araNPQ*, an ABC-type transporter (35). These products are further digested into the monosaccharide arabinose by two intracellular arabinofuranosidases, *abfA* and *abf2*, which

are α -L-arabinofuranosidases (EC 3.2.1.55) belonging to GH51. *AbfA* acts preferentially on (1 \rightarrow 5) arabinofuranosyl linkages, and in contrast, *abf2* is most active on (1 \rightarrow 2) and (1 \rightarrow 3) linkages (36). After this, L-arabinose is converted into D-xylose-5-phosphate, which is further catabolized through the pentose phosphate pathway. The induction mechanism of these genes is mediated through negative control by the key regulator of arabinose metabolism, *araR*. The transcriptional repression of the *abfA* and *abf2* genes is achieved by a tightly controlled mechanism but the regulation of *abnA* is more flexible.

The presence of α -L-arabinofuranosidases (Table 4) has also been determined in *Bifidobacterium longum* subsp. *longum* (37), *Bacteroides thetaiotaomicron* VPI-5482 (38), *Sulfolobus solfataricus* P2 (39), *Anoxybacillus kestanbolensis* AC26Sari (40), *Monoglobus pectinilyticus* (41), and *Roseburia faecis* M72/1 (42). Endo- α -1,5-L-arabinanases (Table 4) are present in *Paenibacillus polymyxa* (43), *Bacillus licheniformis* (44), *Caldicellulosiruptor saccharolyticus* (45), *Bacillus subtilis* (36), *Bacillus thermodenitrificans* (46), *Pseudomonas fluorescens* subsp. *cellulosa* (47), *Monoglobus pectinilyticus* (41), and *Roseburia faecis* M72/1 (42).

Degradation of galactomannan

Endo-1,4- β -mannanase (EC 3.2.1.78), β -mannosidase (EC 3.2.1.25) and α -galactosidase (EC 3.2.1.22) are involved

TABLE 3 Sugar linkage composition of RSM fractions (mol%)^a.

	WUS	DASS	4 MASS	6 MASS
t-Ara ^b	18	41	16	15
1,2-Ara	–	1	1	–
1,5-Ara	10	9	2	11
1,2,5-Ara	6	29	10	4
Total Ara	34	80	29	30
t-Xyl	9	3	7	12
1,2-Xyl	2	1	10	3
1,4-Xyl	3	1	3	7
Total Xyl	14	5	20	22
1,4,6-Man	–	–	3	1
Total Man	–	–	3	1
t-Fuc	2	1	2	5
1,2,4-Fuc	7	3	–	8
Total Fuc	9	4	2	13
t-Gal	3	–	4	6
1,2-Gal	–	–	5	7
1,3-Gal	–	–	7	–
1,3,6-Gal	–	7	4	–
Total Gal	3	7	20	13
1,4-Glc	25	5	17	6
1,4,6-Glc	16	–	11	15
Total Glc	41	5	28	21
T/B ^c	1.10	1.15	1.04	1.36

^aAdapted from reference (14).

^bt, terminal.

^cT/B, ratio terminally linked residues: branching points.

Ara, arabinose; Xyl, xylose; Man, mannose; Fuc, fucose; Gal, galactose; Glc, glucose; UA, uronic acid.

in the degradation of galactomannan into monosaccharides (48). β -mannanase degrades randomly within the main chain of galactomannans and produces shorter galactomanno-oligosaccharides that can be further hydrolyzed by β -mannosidase and α -galactosidase. β -mannosidase hydrolyses β -1,4-linked mannose residues from the non-reducing end of the galactomanno-oligosaccharides and α -galactosidase hydrolyses terminal α -1,6-linked galactose residues from galactomannans or the galactomanno-oligosaccharides. The gut microbes *Bacillus subtilis* YH12 (49), *Bacillus subtilis* TD7 (50), *Bacillus subtilis* Bs5 (51), *Bacillus licheniformis* DSM13 (52), *Sphingomonas* sp. JB13 (53), *Sphingobacterium* sp. GN25 (54), *Klebsiella oxytoca* KUB-CW2-3 (55), *Enterobacter* sp. strain N18 (56), *Flavobacterium* sp. (57), *Pseudomonas cellulosa* (47), *Monoglobus pectinilyticus* (41), and *Bacteroides ovatus* (58) are reported to have endo-1,4- β -mannanase (Table 4). While *Bifidobacterium* sp. (59), *Bifidobacterium longum* subsp. *longum* NCC2705 (60), *Bacteroides ovatus* (61, 62), *Cellvibrio mixtus* (63), *Bacteroides thetaiotaomicron* (64), *Pseudomonas cellulosa* (47), and *Kitasatospora* sp. (65) are reported to have

β -mannosidase (Table 4). Moreover, *Lactobacillus acidophilus* NCFM (66), *Lactobacillus crispatus* ST1 (67), *Lactobacillus brevis* (68), *Bifidobacterium animalis* subsp. *lactis* BI-04 (66), *Arthrobacter* sp. C2-2 (69), *Bacillus megaterium* (70), *Dictyoglomus thermophilum* (71), *Bacillus stearothermophilus* NCIM-5146 (72), *Bacillus stearothermophilus* NUB 3621 (73), *Bacteroides ovatus* 0038-1 (74), *?ifidobacterium bifidum* NCIMB41171 (75), *?ifidobacterium adolescentis* DSM20083 (76), *?ifidobacterium breve* 203 (77), *Clostridium stercorarium* (78), and *Monoglobus pectinilyticus* (41) are reported to have α -galactosidases (Table 4).

Aspergillus niger contains galactomannan-degradation genes, *aglA*, *aglB*, and *aglC* (encoding α -galactosidases), and *mndA* (encoding a β -mannosidase). *AglA* and *aglB* have been classified into GH27, while *aglC* has been classified into GH36, and *mndA* belongs to GH2. The metabolism mechanism of galactomannan is that α -galactosidase (encoded by *aglC*) and β -mannosidase (encoded by *mndA*) hydrolyze the galactomannan to the oligosaccharide Gal2Man5, where *mndA* cleaves single mannose units from the non-reducing end of the substrate until it reaches a galactose side-group (79), and afterward the non-reducing galactose group is hydrolyzed by α -galactosidase (*aglA*, *aglB*, or *aglC*). *AglB* and *mndA* play a major role in the degradation of galactomannan in *A. niger*. The expression of *aglA* is high on galactose and galactose-containing oligosaccharides, but is fully repressed in the presence of glucose (80). Little is known about genes, and especially their regulation, in members of the gut microbiota.

Degradation of homogalacturonan

Homogalacturonan (HG) can be cleaved by α -1,4-L-galacturonan reducing-end-disaccharide-lyase (pectate disaccharide-lyase) (EC 4.2.2.9), (1 \rightarrow 4)-6-O-methyl- α -D-galacturonan lyase (pectin lyase) (EC 4.2.2.10), (1 \rightarrow 4)- α -D-galacturonan reducing-end-trisaccharide-lyase (pectate trisaccharide-lyase) (EC 4.2.2.22), α -1,4-D-galacturonan lyase (pectate lyase) (EC 4.2.2.2), and (1 \rightarrow 4)- α -D-galacturonan glycanohydrolase (endo-polygalacturonase) (EC 3.2.1.15) (89, 124–128). Pectin lyase (EC 4.2.2.10) provides cleavage of α -1,4-linked D-galacturonan methyl ester to give oligosaccharides with 4-deoxy-6-O-methyl- α -D-galact-4-enuronosyl groups at their non-reducing ends, while α -1,4-D-galacturonan lyase (EC 4.2.2.2) cleaves α -1,4-linked D-galacturonan to give oligosaccharides with 4-deoxy- α -D-galact-4-enuronosyl groups at their non-reducing ends. Afterwards, pectate disaccharide-lyase (EC 4.2.2.9) hydrolyzes these oligosaccharides to (1,4- α -D-galacturonosyl)_{n-2} and 4-(4-deoxy- α -D-galact-4-enuronosyl)-D-galacturonate, and (1,4- α -D-galacturonosyl)_{n-2} will be cleaved by pectin lyase (EC 4.2.2.10 or EC 2.2.2) again until the disaccharide results. Polygalacturonase (EC 3.2.1.15) randomly hydrolyzes (1 \rightarrow 4)- α -D-galactosiduronic linkages

TABLE 4 Gut microbes containing the indicated plant cell wall degrading enzymes.

Microbial enzymes	Microorganism
α -L-arabinofuranosidases	<i>Bifidobacterium longum</i> subsp. <i>longum</i> (37), <i>Bacteroides thetaiotaomicron</i> VPI-5482 (38), <i>Sulfolobus solfataricus</i> P2 (39), <i>Anoxybacillus kestanbolensis</i> AC26Sari (40), <i>Monoglobus pectinilyticus</i> (41), and <i>Roseburia faecis</i> M72/1 (42)
endo- α -1,5-L-arabinanases	<i>Paenibacillus polymyxa</i> (43), <i>Bacillus licheniformis</i> (44), <i>Caldicellulosiruptor saccharolyticus</i> (45), <i>Bacillus. Subtilis</i> (36), <i>Bacillus. Thermodenitrificans</i> (46), <i>Pseudomonas fluorescens</i> subsp. <i>cellulosa</i> (47), <i>Monoglobus pectinilyticus</i> (41), and <i>Roseburia faecis</i> M72/1 (42)
endo-1,4- β -mannanase	<i>Bacillus subtilis</i> YH12 (49), <i>Bacillus subtilis</i> TD7 (50), <i>Bacillus subtilis</i> Bs5 (51), <i>Bacillus licheniformis</i> DSM13 (52), <i>Sphingomonas</i> sp. JB13 (53), <i>Sphingobacterium</i> sp. GN25 (54), <i>Klebsiella oxytoca</i> KUB-CW2-3 (55), <i>Enterobacter</i> sp. strain N18 (56), <i>Flavobacterium</i> sp. (57), <i>Pseudomonas cellulosa</i> (47), <i>Monoglobus pectinilyticus</i> (41), and <i>Bacteroides ovatus</i> (58)
β -mannosidase	<i>Bifidobacterium</i> sp. (59), <i>Bifidobacterium longum</i> subsp. <i>longum</i> NCC2705 (60), <i>Bacteroides ovatus</i> (61, 62), <i>Cellvibrio mixtus</i> (63), <i>Bacteroides thetaiotaomicron</i> (64), <i>Pseudomonas cellulose</i> (47), <i>Kitasatospora</i> sp. (65)
α -Galactosidases	<i>Lactobacillus acidophilus</i> NCFM (66), <i>Lactobacillus crispatus</i> ST1 (67), <i>Lactobacillus brevis</i> (68), <i>Bifidobacterium animalis</i> subsp. <i>lactis</i> Bl-04 (66), <i>Arthrobacter</i> sp. C2-2 (69), <i>Bacillus megaterium</i> (70), <i>Dictyoglomus thermophilum</i> (71), <i>Bacillus stearothermophilus</i> NCIM-5146 (72), <i>Bacillus stearothermophilus</i> NUB 3621 (73), <i>Bacteroides ovatus</i> 0038-1 (74), <i>?ifidobacterium bifidum</i> NCIMB41171 (75), <i>?ifidobacterium adolescentis</i> DSM20083 (76), <i>?ifidobacterium breve</i> 203 (77), <i>Clostridium stercorarium</i> (78), and <i>Monoglobus pectinilyticus</i> (41)
Pectate disaccharide-lyase	<i>Caldicellulosiruptor bescii</i> (81), <i>Bacteroides thetaiotaomicron</i> (82), <i>Bacteroides ovatus</i> (83), <i>Bacillus pumilus</i> BK2 (84), <i>Eubacterium eligens</i> , <i>Faecalibacterium prausnitzii</i> (85), and <i>Monoglobus pectinilyticus</i> (41)
pectate lyase	in <i>Caldicellulosiruptor bescii</i> (81), genus <i>Bacillus</i> : <i>Bacillus subtilis</i> , <i>B. licheniformis</i> , <i>B. cereus</i> , <i>B. circulans</i> , <i>B. pasteurii</i> , <i>B. amyloliquefaciens</i> , and <i>B. pumilus</i> (86, 87), <i>Paenibacillus</i> sp. (88), <i>Clostridium cellulovorans</i> (89), <i>Streptomyces thermocarboxyus</i> (90), <i>Bacteroides thetaiotaomicron</i> (82), <i>Bacteroides ovatus</i> (83), <i>Eubacterium eligens</i> , <i>Faecalibacterium prausnitzii</i> (85), and <i>Monoglobus pectinilyticus</i> (41)
endo-polygalacturonase	<i>Caldicellulosiruptor bescii</i> (81), <i>Bacteroides thetaiotaomicron</i> (82), <i>Bacteroides ovatus</i> (83), <i>Eubacterium eligens</i> , <i>Faecalibacterium prausnitzii</i> (85), <i>Monoglobus pectinilyticus</i> (41), and <i>Bifidobacterium longum</i> subsp. <i>longum</i> (91)
rhamnogalacturonan lyase and hydrolase	<i>Caldicellulosiruptor bescii</i> (81), <i>Bacteroides thetaiotaomicron</i> (82), <i>Bacteroides ovatus</i> (83), <i>Bacillus subtilis</i> (92), <i>Bacillus licheniformis</i> (93), <i>Cellvibrio japonicus</i> (94), <i>Clostridium cellulolyticum</i> (95), <i>Bacillus licheniformis</i> DSM13 (93), <i>Pseudomonas cellulose</i> (94), <i>Penicillium chrysogenum</i> (96), and <i>Monoglobus pectinilyticus</i> (41)
exo- β -1,3-galactanase	<i>Monoglobus pectinilyticus</i> (41), <i>Bifidobacterium longum</i> subsp. <i>longum</i> (91), <i>Clostridium thermocellum</i> (97), <i>Phanerochaete chrysosporium</i> (98), <i>Sphingomonas</i> sp. (99), <i>Bacteroides thetaiotaomicron</i> (100), <i>Bacteroides ovatus</i> (100), <i>Bacteroides caccae</i> (101), and <i>Bacteroides cellulolyticus</i> (101)
endo- β -1,6-galactanase	<i>Streptomyces avermitilis</i> NBRC14893 (102), <i>Bacteroides ovatus</i> (100), <i>Bacteroides caccae</i> (101), <i>Bacteroides cellulolyticus</i> (101), and <i>Bifidobacterium longum</i> subsp. <i>longum</i> (91)
exo- β -1,6-galactobiohydrolase	<i>Monoglobus pectinilyticus</i> (41), <i>Bacteroides thetaiotaomicron</i> (101), <i>Streptomyces avermitilis</i> (102), <i>Bacteroides ovatus</i> (100), <i>Bacteroides caccae</i> (101), <i>Bacteroides cellulolyticus</i> (101), and <i>Bifidobacterium longum</i> subsp. <i>longum</i> (91)
endo- β -1,4-xylanase	<i>Pseudomonas boreopolis</i> G22 (103), <i>Bacteroides ovatus</i> (82), <i>Monoglobus pectinilyticus</i> (41), <i>Bacteroides thetaiotaomicron</i> (100), <i>Bacteroides caccae</i> (101), <i>Bacteroides cellulolyticus</i> (101), <i>Clostridium thermocellum</i> (104), <i>Bacillus subtilis</i> (105), and <i>Streptomyces turgidiscabies</i> (106)
exo- β -1,4 xylanase	<i>Monoglobus pectinilyticus</i> (41), <i>Luteimicrobium xylanilyticum</i> (107), <i>Amycolatopsis mediterranei</i> (107), <i>Clostridium thermocellum</i> (104), <i>Bacillus subtilis</i> (105), and <i>Streptomyces turgidiscabies</i> (106)
endo and exo- β -1,4-glucanase	<i>Monoglobus pectinilyticus</i> (41), <i>Caldicellulosiruptor kronotskyensis</i> (108), <i>Roseburia</i> sp. (42), <i>Eubacterium rectale</i> group (42), <i>Ruminococcus champanellensis</i> (109), <i>Ruminococcus bromii</i> (110), <i>Ruminiclostridium cellulolyticum</i> (111), and <i>Phaeoacremonium minimum</i> (112)
α -D-xylosidase	<i>Sulfolobus solfataricus</i> P2 (39), <i>Talaromyces thermophilus</i> (113), <i>Cellvibrio japonicus</i> (114), <i>Bacteroides thetaiotaomicron</i> (82), <i>Bacteroides ovatus</i> (82), and <i>Monoglobus pectinilyticus</i> (41)
β -glucosidase	<i>Bifidobacterium adolescentis</i> (115), <i>Bacteroides ovatus</i> (116), <i>Listeria innocua</i> (117), <i>Streptomyces venezuelae</i> (118), <i>Pyrococcus furiosus</i> (119), <i>Cellvibrio japonicus</i> (114), <i>Caldicellulosiruptor saccharolyticus</i> (120), <i>Microbispora bispora</i> (121), <i>Thermoanaerobacter brockii</i> (122), <i>Thermobifida fusca</i> (103), <i>Pseudomonas</i> sp. (123), <i>Monoglobus pectinilyticus</i> (41), <i>Ruminococcus champanellensis</i> (109), and <i>Ruminococcus bromii</i> (110)

in pectate and other galacturonans. Pectate disaccharidase (Table 4) has been reported in *Caldicellulosiruptor bescii* (81), *Bacteroides thetaiotaomicron* (82), *Bacteroides ovatus* (83), *Bacillus pumilus* BK2 (84), *Eubacterium eligens*, *Faecalibacterium prausnitzii* (85), and *Monoglobus pectinilyticus* (41). Pectate lyase (Table 4) has been reported in *Caldicellulosiruptor bescii* (81), genus *Bacillus*: *Bacillus subtilis*, *B. licheniformis*, *B. cereus*, *B. circulans*, *B. pasteurii*, *B. amyloliquefaciens*, and *B. pumilus* (86, 87), *Paenibacillus* sp. (88), *Clostridium cellulovorans* (89), *Streptomyces thermocarboxydus* (90), *Bacteroides thetaiotaomicron* (82), *Bacteroides ovatus* (83), *Eubacterium eligens*, *Faecalibacterium prausnitzii* (85), and *Monoglobus pectinilyticus* (41). Endo-polygalacturonase (Table 4) has been reported in *Caldicellulosiruptor bescii* (81), *Bacteroides thetaiotaomicron* (82), *Bacteroides ovatus* (83), *Eubacterium eligens*, *Faecalibacterium prausnitzii* (85), *Monoglobus pectinilyticus* (41), and *Bifidobacterium longum* subsp. *longum* (91).

Degradation of rhamnogalacturonan I

Utilization of rhamnogalacturonan I (RG-I) by microbes is mediated by a series of enzymes, which is well-studied in *Bacillus subtilis* (92). Two main enzymes, a hydrolase, and a lyase are involved in the degradation of the RG-I backbone, whereas a few other enzymes are responsible for the breakdown of the RG-I side chains. Rhamnogalacturonan hydrolase (rhamnogalacturonan α -L-rhamnopyranohydrolase) cleaves α -1,2 linkages between GalA and Rha (129). Rhamnogalacturonan lyase [α -L-rhamnopyranosyl-(1 \rightarrow 4)- α -D-galactopyranosyluronate endolyase (EC 4.2.2.23)] cleaves the α -1,4 linkages of RG-I resulting in a double bond in the on-reducing GalA residue (130). Rhamnogalacturonan lyase and hydrolase (Table 4) have been reported in *Caldicellulosiruptor bescii* (81), *Bacteroides thetaiotaomicron* (82), *Bacteroides ovatus* (83), *Bacillus subtilis* (92), *Bacillus licheniformis* (93), *Cellvibrio japonicus* (94), *Clostridium cellulolyticum* (95), *Bacillus licheniformis* DSM13 (93), *Pseudomonas cellulose* (94), *Penicillium chrysogenum* (96), and *Monoglobus pectinilyticus* (41).

Degradation of type II arabinogalactan

Exo- β -1,3-galactanase (EC 3.2.1.145) cleaves the β -1,3-galactan backbone of type II arabinogalactan *via* bypassing the β -1,6-galactan side chains and releasing β -1,6-galactooligosaccharides and their derivatives (131). The β -1,6-galactan side chains are hydrolyzed to β -1,6-galactooligosaccharides of various degrees of polymerization (DP) by endo- β -1,6-galactanase (EC 3.2.1.164) (132, 133). On

the other hand, exo- β -1,6-galactobiohydrolase releases β -1,6-galactobiose (β -1,6-Gal2) from the non-reducing terminal end of β -1,6-galactooligosaccharides, and α -L-arabinofuranosidase (EC 3.2.1.55) releases arabinofuranose (Araf) from α -1,3-Araf-substituted β -1,6-galactooligosaccharides (134). Exo- β -1,3-galactanase (Table 4) has been reported in *Monoglobus pectinilyticus* (41), *Bifidobacterium longum* subsp. *longum* (91), *Clostridium thermocellum* (97), *Phanerochaete chrysosporium* (98), *Sphingomonas* sp. (99), *Bacteroides thetaiotaomicron* (100), *Bacteroides ovatus* (100), *Bacteroides caccae* (101), and *Bacteroides cellulosilyticus* (101). Endo- β -1,6-galactanase (Table 4) had been reported in *Streptomyces avermitilis* NBRC14893 (102), *Bacteroides ovatus* (100), *Bacteroides caccae* (101), *Bacteroides cellulosilyticus* (101), and *Bifidobacterium longum* subsp. *longum* (91). Exo- β -1,6-galactobiohydrolase (Table 4) has been reported in *Monoglobus pectinilyticus* (41), *Bacteroides thetaiotaomicron* (101), *Streptomyces avermitilis* (102), *Bacteroides ovatus* (100), *Bacteroides caccae* (101), *Bacteroides cellulosilyticus* (101), and *Bifidobacterium longum* subsp. *longum* (91).

Degradation of glucuronoxylan

Two enzymes, β -(1, 3–5)-D-xylan xylanohydrolase (endo- β -1,4 xylanase) (EC 3.2.1.8) and 1,4- β xylohydrolase (exo- β -1,4 xylanase) (EC 3.2.1.37), are involved in degrading the β -1,4 xylosyl linkages in unsubstituted domains along the xylan backbone of glucuronoxylan (GX) (135, 136). Glucuronoxylanase cleaves glucuronosyl moieties which are substituted as monomeric side chains on the xylan backbone (137). Endo- β -1,4-xylanase (Table 4) has been reported in *Pseudomonas boreopolis* G22 (103), *Bacteroides ovatus* (82), *Monoglobus pectinilyticus* (41), *Bacteroides thetaiotaomicron* (100), *Bacteroides caccae* (101), *Bacteroides cellulosilyticus* (101), *Clostridium thermocellum* (104), *Bacillus subtilis* (105), and *Streptomyces turgidiscabies* (106). Exo- β -1,4 xylanase (Table 4) has been reported in *Monoglobus pectinilyticus* (41), *Luteimicrobium xylanilyticum* (107), *Amycolatopsis mediterranei* (107), *Clostridium thermocellum* (104), *Bacillus subtilis* (105), and *Streptomyces turgidiscabies* (106).

Degradation of xyloglucan

A set of glucanases and glycosidases are involved in cleaving xyloglucan (XG) into monosaccharides by two-step degradation (138, 139). Endo- β -1,4-glucanase (EC 3.2.1.4) hydrolyzes XG into large fragments, which are further degraded into monosaccharides by α -xylosidase (EC 3.2.1.177) and β -glucosidase (EC 3.2.1.21) (140). Endo- β -1,4-glucanase (Table 4) has been reported in *Monoglobus pectinilyticus* (41), *Caldicellulosiruptor kronotskyensis* (108), *Roseburia* sp. (42),

Eubacterium rectale group (42), *Ruminococcus champanellensis* (109), *Ruminococcus bromii* (110), *Ruminiclostridium cellulolyticum* (111), and *Phaeoacremonium minimum* (112). Exo- β -1,4 xylanase (Table 4) has been reported in *Monoglobus pectinilyticus* (41), *Luteimicrobium xylanilyticum* (107), *Amycolatopsis mediterranei* (107), *Clostridium thermocellum* (104), *Bacillus subtilis* (105), *Streptomyces turgidiscabies* (106), *Ruminiclostridium cellulolyticum* (111), and *Phaeoacremonium minimum* (112). While α -D-xylosidase (Table 4) has been reported in *Sulfolobus solfataricus* P2 (39), *Talaromyces thermophilus* (113), *Cellvibrio japonicus* (114), *Bacteroides thetaiotaomicron* (82), *Bacteroides ovatus* (82), and *Monoglobus pectinilyticus* (41). Moreover, β -glucosidase (Table 4) has been reported in *Bifidobacterium adolescentis* (115), *Bacteroides ovatus* (116), *Listeria innocua* (117), *Streptomyces venezuelae* (118), *Pyrococcus furiosus* (119), *Cellvibrio japonicus* (114), *Caldicellulosiruptor saccharolyticus* (120), *Microbispora bispora* (121), *Thermoanaerobacter brockii* (122), *Thermobifida fusca* (103), *Pseudomonas* sp. (123), *Monoglobus pectinilyticus* (41), *Ruminococcus champanellensis* (109), and *Ruminococcus bromii* (110).

Degradation of cellulose

The enzymes of the cellulase system consist of endo- β -1,4-glucanase (EC 3.2.1.4), exo- β -1,4-glucanase (EC 3.2.1.91), and β -glucosidase (EC 3.2.1.21). Endo- β -1,4-glucanase and exo- β -1,4-glucanase cleave cellulose to cellodextrins and cellobiose, which are then degraded to glucose by β -glucosidase (141, 142). Endo and exo- β -1,4-glucanase (Table 4) have been reported in *Monoglobus pectinilyticus* (41), *Caldicellulosiruptor kronotskyensis* (108), *Roseburia* sp. (42), *Eubacterium rectale* group (42), *Ruminococcus champanellensis* (109), *Ruminococcus bromii* (110), *Ruminiclostridium cellulolyticum* (111), *Paenibacillus* sp. (143), and *Phaeoacremonium minimum* (112). The enzyme β -glucosidase (Table 4) has been reported in *Bifidobacterium adolescentis* (115), *Bacteroides ovatus* (116), *Listeria innocua* (117), *Streptomyces venezuelae* (118), *Pyrococcus furiosus* (119), *Cellvibrio japonicus* (114), *Caldicellulosiruptor saccharolyticus* (120), *Microbispora bispora* (121), *Thermoanaerobacter brockii* (122), *Thermobifida fusca* (103), *Pseudomonas* sp. (123), *Monoglobus pectinilyticus* (41), *Ruminococcus champanellensis* (109), and *Ruminococcus bromii* (110).

Processing to increase use of recalcitrant fibers

Our previous studies have already shown that physical processing technologies cannot significantly increase fiber degradability in monogastric animals (19, 144, 145). However,

the utilization of enzymes added to feed is a promising method to improve fiber fermentability.

Enzymatic and chemical modification

Supplementation of cell wall degrading enzymes to improve feed efficiency for pigs gets more and more attention from the feed industry nowadays. Supplementation of cell wall degrading enzymes can remove side-chains of polysaccharides in plant cell wall, which make them more accessible to bacterial enzymes (145). In addition, cell wall degrading enzymes are able to reduce the digesta viscosity, which might affect absorption of other nutrients, by cleaving the viscous polysaccharides (e.g., pectin). In the end, carbohydrases can depolymerize polysaccharides to oligosaccharides, which have potential prebiotic effects on the gut microbiota, leading to the health benefits for the host animal.

A previous study showed that pre-processing with sulfuric acid could increase utilization of carbohydrates from lignocellulosic biomass (146). Mild acid-treated rye (together with heat treatment) was reported to improve the release of arabinosyl residues in chickens (147). Alkaline pretreatment of rapeseed meal in the feed improved its fermentation in pigs (7). Therefore, chemical treatment in feed might be a promising method to improve fiber fermentation in pigs.

We have shown recently that RSM processed by enzymatic and chemical treatment led to modulation of the gut microbiota using a newly developed *in vitro* swine large intestinal model (SLIM) (148). In brief, our studies demonstrated that both enzymatic (cellulase or 2 different pectinases) and chemical (6N sodium hydroxide) pretreatment on RSM shifted its cell wall polysaccharide structure, subsequently altering microbial community composition and functional profile compared to untreated RSM, and eventually increased fiber degradability as evaluated by SCFA production in SLIM. Moreover, it was validated in pigs, by the mobile nylon bag technique, that cellulase and alkaline treatment on RSM improved the overall degradation of RSM (149–151).

Potential emerging technologies

High hydrostatic pressure, a non-thermal pasteurization technology, is applied to physical modification of chemical structure of compound (e.g., polysaccharides). High hydrostatic pressure could promote the water into interior of the matrix. As a result, high pressure (100–1,000 MPa) leads to the destruction of cell wall matrix (affecting non-covalent bonds, including hydrophobic interactions, hydrogen bonds, and van der Waals forces) and reduce substrate particle size. Research demonstrated that pectin (extracted from potato peel waste) could increase galacturonic acid content and decrease the esterification degree of pectin after high hydrostatic pressure

at 200 MPa for 5 min (152). Another study reported that high hydrostatic pressure could significantly reduce the molecular weight, degree of esterification and degree of acylation of sugar beet pectin at 250–500 MPa for 30 min (153). These reports have shown that high hydrostatic pressure could modify the polysaccharide structure, which might have a potential application value in improving fiber degradability of RSM in monogastric animals.

Cold plasma (CP) processing is a technology that causes the ionization of neutral gas (argon, helium, and nitrogen) and generating active species (free radicals, electrons, ions), which could induce oxidative damage in the cell wall compounds, such as polysaccharides. Therefore, CP could change structure and properties of polysaccharides. Study showed that CP could increase the viscosity and enhance the emulsion stabilization capacity of xanthan gum (154). Prasertsung et al. (155) reported that CP could degrade starches and cellulose into sugars and glucose (155). Therefore, CP might modify polysaccharides of RSM by inducing oxidative damage in its cell wall polysaccharides.

Conclusions and perspectives

Rapeseed meal is not only a promising protein ingredient but also a potentially energy source for non-ruminant animal diets. RSM contains a high amount of cell wall polysaccharides, even higher when compared to soybean meal commonly used in the feed industry. However, information on degradation and utilization of RSM carbohydrates upon feeding these diets is insufficient, and for this it is essential to understand the characterization and quantification of RSM carbohydrates. For the sake of improving the degradation of RSM carbohydrates, it is necessary to understand the mode of actions of RSM polysaccharide degradation by microbes, discover factors that limit their utilization and absorption in the GIT, and develop strategies to improve efficiency and productivity. The use of exogenous enzymes, carbohydrases, becomes a good solution to improve RSM carbohydrate degradation, and eventually to improve animal digestibility, feed efficiency, and animal performance.

The current review summarized the polysaccharide types of RSM, gut microbiota which could degrade the polysaccharides, and glycoside hydrolases which used by the microorganism. The information offered above could be used to develop novel

engineering microbes and enzymic preparations to improve RSM utilization in the future studies. One possible way is to deep mine the genomes of these microbes for encoded the carbohydrases by bioinformatic tools. What is more, these enzymes can be overexpressed in an expression host and subsequently used to process RSM to investigate its degradability in livestock.

Author contributions

CL prepared the original draft and edited the manuscript. X-LQ and KV critically reviewed the manuscript. All authors listed have approved the final manuscript for publication.

Funding

This research has been partly made possible with the support of the Dutch Province of Limburg, with a grant to the Centre for Healthy Eating and Food Innovation (HEFI) of Maastricht University—Campus Venlo to KV. This study was also supported by the Young Teachers' Research and Innovation Ability Improvement Program of Beijing University of Agriculture (QJKC2022026), Technology Innovation spark action support program of Beijing University of Agriculture, and 2022 Beijing Municipal Education Commission Classification Development Project.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Pustjens AM, De Vries S, Bakuwel M, Gruppen H, Gerrits WJ, Kabel MA. Unfermented recalcitrant polysaccharide structures from rapeseed (*Brassica napus*) meal in pigs. *Ind Crops Prod.* (2014) 58:271–9. doi: 10.1016/j.indcrop.2014.04.036
2. Kiarie E, Romero LF, Nyachoti CM. The role of added feed enzymes in promoting gut health in swine and poultry. *Nutr Res Rev.* (2013) 26:71–88. doi: 10.1017/S0954422413000048

3. Leming R, Lember A. Chemical composition of expeller-extracted and cold-pressed rapeseed cake. *Agrarartheadus*. (2005) 16:96–109.
4. Agyekum AK, Woyengo TA. Nutritive value of expeller/cold-pressed canola meal and pre-pressed solvent-extracted carinata meal for broiler chicken. *Poult Sci*. (2022) 101:101528. doi: 10.1016/j.psj.2021.101528
5. Slominski BA, Campbell LD. Non-starch polysaccharides of canola meal: quantification, digestibility in poultry and potential benefit of dietary enzyme supplementation. *J Sci Food Agric*. (1990) 53:175–84. doi: 10.1002/jsfa.2740530205
6. Simbaya J, Slominski BA, Rakow G, Campbell LD, Downey RK, Bell JM. Quality characteristics of yellow-seeded Brassica seed meals: protein, carbohydrate, and dietary fiber components. *J Agric Food Chem*. (1995) 43:2062–6. doi: 10.1021/jf00056a020
7. Pustjens AM, Schols HA, Kabel MA, Gruppen H. Characterisation of cell wall polysaccharides from rapeseed (*Brassica napus*) meal. *Carbohydr Polym*. (2013) 98:1650–6. doi: 10.1016/j.carbpol.2013.07.059
8. Slominski BA. Hydrolysis of galactooligosaccharides by commercial preparations of α -galactosidase and β -fructofuranosidase: potential for use as dietary additives. *J Sci Food Agric*. (1994) 65:323–30. doi: 10.1002/jsfa.2740650310
9. Meng X, Slominski B. Nutritive values of corn, soybean meal, canola meal, and peas for broiler chickens as affected by a multienzyme preparation of cell wall degrading enzymes. *Poult Sci*. (2005) 84:1242–51. doi: 10.1093/ps/84.8.1242
10. Carpita NC, Gibeault DM. Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *Plant J*. (1993) 3:1–30. doi: 10.1111/j.1365-313X.1993.tb00007.x
11. Seymour GB, Knox JP. *Pectins and Their Manipulation*. Oxford: Blackwell Publishing (2002).
12. Zykwincka A, Thibault JF, Ralet MC. Competitive binding of pectin and xyloglucan with primary cell wall cellulose. *Carbohydr Polym*. (2008) 74:957–61. doi: 10.1016/j.carbpol.2008.05.004
13. Pustjens AM. Fate of rapeseed meal polysaccharides during digestion in pigs and poultry: effect of processing and enzyme addition. In: *Fiber Fermentation in Pigs And Poultry Sense and Nonsense of Its Manipulation*. Wageningen: Wageningen University and Research (2013). p. 64.
14. Rommi K, Hakala TK, Holopainen U, Nordlund E, Poutanen K, Lantto R. Effect of enzyme-aided cell wall disintegration on protein extractability from intact and dehulled rapeseed (*Brassica rapa* L. and *Brassica napus* L.) press cakes. *J Agric Food Chem*. (2014) 62:7989–97. doi: 10.1021/jf501802e
15. Bell J. Nutrients and toxicants in rapeseed meal: a review. *J Anim Sci*. (1984) 58:996–1010. doi: 10.2527/jas1984.584996x
16. Nair J, Penner GB, Yu P, Lardner H, McAllister T, Damiran D, et al. Evaluation of canola meal derived from *Brassica juncea* and *Brassica napus* seed as an energy source for feedlot steers. *Can J Anim Sci*. (2015) 95:599–607. doi: 10.4141/cjas-2015-055
17. Slominski B, Campbell L. The carbohydrate content of yellow-seeded canola. In: *Proceedings of the 8th International Rapeseed Congress*. Saskatoon (1991). p. 9–11.
18. Gallant D, Bouchet B, Buleon A, Perez S. Physical characteristics of starch granules and susceptibility to enzymatic degradation. *Eur J Clin Nutr*. (1992) 46:S3–16.
19. De Vries S, Pustjens A, Kabel M, Kwakkel R, Gerrits W. Effects of processing technologies and pectolytic enzymes on degradability of nonstarch polysaccharides from rapeseed meal in broilers. *Poult Sci*. (2014) 93:589–98. doi: 10.3382/ps.2013-03476
20. Ghosh P, Ghosal P, Thakur S, Lerouge P, Loutelier-Bourhis C, Driouch A, et al. Cell wall polysaccharides of Brassica campestris seed cake: isolation and structural features. *Carbohydr Polym*. (2004) 57:7–13. doi: 10.1016/j.carbpol.2004.01.012
21. Colquhoun IJ, De Ruiter GA, Schols HA, Voragen AG. Identification by NMR spectroscopy of oligosaccharides obtained by treatment of the hairy regions of apple pectin with rhamnogalacturonase. *Carbohydr Res*. (1990) 206:131–44. doi: 10.1016/0008-6215(90)84012-J
22. Ebringerová A, Heinze T. Xylan and xylan derivatives-biopolymers with valuable properties. 1. Naturally occurring xyans structures, isolation procedures and properties. *Macromol Rapid Commun*. (2000) 21:542–56. doi: 10.1002/1521-3927(20000601)21:9<542::AID-MARC542>3.0.CO;2-7
23. Peña MJ, Zhong R, Zhou G-K, Richardson EA, O'Neill MA, Darvill AG, et al. Arabidopsis irregular xylem8 and irregular xylem9: implications for the complexity of glucuronoxylan biosynthesis. *Plant Cell*. (2007) 19:549–63. doi: 10.1105/tpc.106.049320
24. Larsbrink J, Rogers TE, Hemsworth GR, Mckee LS, Tauzin AS, Spadiut O, et al. A discrete genetic locus confers xyloglucan metabolism in select human gut Bacteroidetes. *Nature*. (2014) 506:498–502. doi: 10.1038/nature12907
25. Kračun SK, Fangel JU, Rydahl MG, Pedersen HL, Vidal-Melgosa S, Willats WGT. Carbohydrate microarray technology applied to high-throughput mapping of plant cell wall glycans using comprehensive microarray polymer profiling (CoMPP). High-throughput glycomics and glycoproteomics. *Methods Protoc*. (2017) 147–65. doi: 10.1007/978-1-4939-6493-2_12
26. Forsee WT, Springfield JD, Schutzbach J. Effect of phospholipids on alpha-1, 2-mannosidase activity. *J Biol Chem*. (1982) 257:9963–7. doi: 10.1016/S0021-9258(18)33971-1
27. Hilz H, Bakx EJ, Schols HA, Voragen AG. Cell wall polysaccharides in black currants and bilberries-characterisation in berries, juice, and press cake. *Carbohydr Polym*. (2005) 59:477–88. doi: 10.1016/j.carbpol.2004.11.002
28. Flint HJ, Bayer EA. Plant cell wall breakdown by anaerobic microorganisms from the mammalian digestive tract. *Ann N Y Acad Sci*. (2008) 1125:280–8. doi: 10.1196/annals.1419.022
29. Van Den Brink J, De Vries RP. Fungal enzyme sets for plant polysaccharide degradation. *Appl Microbiol Biotechnol*. (2011) 91:1477. doi: 10.1007/s00253-011-3473-2
30. Crouch LI, Labourel A, Walton PH, Davies GJ, Gilbert HJ. The contribution of non-catalytic carbohydrate binding modules to the activity of lytic polysaccharide monoxygenases. *J Biol Chem*. (2016) 291:7439–49. doi: 10.1074/jbc.M115.702365
31. Cartmell A, Mckee LS, Pena MJ, Larsbrink J, Brumer H, Kaneko S, et al. The structure and function of an arabinan-specific alpha-1,2-arabinofuranosidase identified from screening the activities of bacterial GH43 glycoside hydrolases. *J Biol Chem*. (2011) 286:15483–95. doi: 10.1074/jbc.M110.215962
32. Raposo MP, Inacio JM, Mota LJ, De Sa-Nogueira I. Transcriptional regulation of genes encoding arabinan-degrading enzymes in *Bacillus subtilis*. *J Bacteriol Res*. (2004) 186:1287–96. doi: 10.1128/JB.186.5.1287-1296.2004
33. Leal TF, De Sa-Nogueira I. Purification, characterization and functional analysis of an endo-arabinanase (AbnA) from *Bacillus subtilis*. *FEMS Microbiol Lett*. (2004) 241:41–8. doi: 10.1016/j.femsle.2004.10.003
34. Sá-Nogueira I, Ramos SS. Cloning, functional analysis, and transcriptional regulation of the *Bacillus subtilis* araE gene involved in L-arabinose utilization. *J Bacteriol Res*. (1997) 179:7705–11. doi: 10.1128/jb.179.24.7705-7711.1997
35. Sá-Nogueira I, Nogueira TV, Soares S, De Lencastre H. The *Bacillus subtilis* L-arabinose (ara) operon: nucleotide sequence, genetic organization and expression. *Microbiology*. (1997) 143:957–69. doi: 10.1099/00221287-143-3-957
36. Inacio JM, De Sa-Nogueira I. Characterization of abn2 (yxiA), encoding a *Bacillus subtilis* GH43 arabinanase, Abn2, and its role in arabinopolysaccharide degradation. *J Bacteriol*. (2008) 190:4272–80. doi: 10.1128/JB.00162-08
37. Komeno M, Hayamizu H, Fujita K, Ashida H. Two novel α -L-Arabinofuranosidases from *Bifidobacterium longum* subsp. *longum* belonging to glycoside hydrolase family 43 cooperatively degrade arabinan. *Appl Environ Microbiol*. (2019) 85:e02582–18. doi: 10.1128/AEM.02582-18
38. Mahowald MA, Rey FE, Seedorf H, Turnbaugh PJ, Fulton RS, Wollam A, et al. Characterizing a model human gut microbiota composed of members of its two dominant bacterial phyla. *Proc Natl Acad Sci USA*. (2009) 106:5859–64. doi: 10.1073/pnas.0901529106
39. Morana A, Paris O, Maurelli L, Rossi M, Cannio R. Gene cloning and expression in *Escherichia coli* of a bi-functional β -D-xylosidase/ α -L-arabinosidase from *Sulfolobus solfataricus* involved in xylan degradation. *Extremophiles*. (2007) 11:123–32. doi: 10.1007/s00792-006-0020-7
40. Canakci S, Kacagan M, Inan K, Belduz AO, Saha BC. Cloning, purification, and characterization of a thermostable α -L-arabinofuranosidase from *Anoxybacillus kestanbolensis* AC26Sari. *Appl Microbiol Biotechnol*. (2008) 81:61. doi: 10.1007/s00253-008-1584-1
41. Kim CC, Healey GR, Kelly WJ, Patchett ML, Jordens Z, Tannock GW, et al. Genomic insights from *Monoglobus pectinilyticus*: a pectin-degrading specialist bacterium in the human colon. *ISME J*. (2019) 13:1437–56. doi: 10.1038/s41396-019-0363-6
42. Sheridan PO, Martin JC, Lawley TD, Browne HB, Harris HM, Bernalier-Donadille A, et al. Polysaccharide utilization loci and nutritional specialization in a dominant group of butyrate-producing human colonic Firmicutes. *Microbiol Genom*. (2016) 2:e000043. doi: 10.1099/mgen.0.000043
43. Wang S, Yang Y, Zhang J, Sun J, Matsukawa S, Xie J, et al. Characterization of abnZ2 (yxiA1) and abnZ3 (yxiA3) in *Paenibacillus polymyxa*, encoding two novel endo-1,5- α -L-arabinanases. *Bioresour Bioprocess*. (2014) 1:14. doi: 10.1186/s40643-014-0014-8

44. Seo E-S, Lim Y-R, Kim Y-S, Park C-S, Oh D-K. Characterization of a recombinant endo-1, 5- α -L-arabinanase from the isolated bacterium *Bacillus licheniformis*. *Biotechnol Bioprocess Eng.* (2010) 15:590–4. doi: 10.1007/s12257-009-3138-5
45. Hong M-R, Park C-S, Oh D-K. Characterization of a thermostable endo-1, 5- α -L-arabinanase from *Caldicellulosiruptor saccharolyticus*. *Biotechnol Lett.* (2009) 31:1439. doi: 10.1007/s10529-009-0019-0
46. Takao M, Akiyama K, Sakai T. Purification and characterization of thermostable endo-1, 5- α -L-arabinase from a strain of *Bacillus thermodenitrificans*. *Appl Environ Microbiol.* (2002) 68:1639–46. doi: 10.1128/AEM.68.4.1639-1646.2002
47. Hogg D, Woo E-J, Bolam DN, Mckie VA, Gilbert HJ, Pickersgill RW. Crystal structure of mannanase 26A from *Pseudomonas cellulosa* and analysis of residues involved in substrate binding. *J Biol Chem.* (2001) 276:31186–92. doi: 10.1074/jbc.M010290200
48. Ademark P, De Vries RP, Häggglund P, Stålbbrand H, Visser J. Cloning and characterization of *Aspergillus niger* genes encoding an α -galactosidase and a β -mannosidase involved in galactomannan degradation. *Eur J Biochem.* (2001) 268:2982–90. doi: 10.1046/j.1432-1327.2001.02188.x
49. Liu H-X, Gong J-S, Li H, Lu Z-M, Li H, Qian J-Y, et al. Biochemical characterization and cloning of an endo-1, 4- β -mannanase from *Bacillus subtilis* YH12 with unusually broad substrate profile. *Process Biochem.* (2015) 50:712–21. doi: 10.1016/j.procbio.2015.02.011
50. Li C-Y, Liu F-F, Ye J, Liu J-F, Yang S-Z, Zhang H-Z, et al. A low-temperature active endo- β -1, 4-mannanase from *Bacillus subtilis* TD7 and its gene expression in *Escherichia coli*. *Appl Environ Biotechnol.* (2018) 3:17–25. doi: 10.26789/AEB.2018.02.004
51. Huang JL, Bao LX, Zou HY, Che SG, Wang GX. High-level production of a cold-active B-mannanase from *Bacillus subtilis* Bs5 and its molecular cloning and expression. *Mol Genet Microbiol Virol.* (2012) 27:147–53. doi: 10.3103/S0891416812040039
52. Songsiririthigul C, Buranabanyat B, Haltrich D, Yamabhai M. Efficient recombinant expression and secretion of a thermostable GH26 mannan endo-1, 4- β -mannosidase from *Bacillus licheniformis* in *Escherichia coli*. *Microb Cell Factories.* (2010) 9:20. doi: 10.1186/1475-2859-9-20
53. Zhou J, Zhang R, Gao Y, Li J, Tang X, Mu Y, et al. Novel low-temperature-active, salt-tolerant and proteases-resistant endo-1, 4- β -mannanase from a new *Sphingomonas* strain. *J Biosci Bioeng.* (2012) 113:568–74. doi: 10.1016/j.jbiosc.2011.12.011
54. Zhang R, Zhou J, Gao Y, Guan Y, Li J, Tang X, et al. Molecular and biochemical characterizations of a new low-temperature active mannanase. *Folia Microbiol.* (2015) 60:483–92. doi: 10.1007/s12223-015-0391-1
55. Pongsapipatana N, Damrongteerapap P, Chantorn S, Sintuprapa W, Keawsoompong S, Nitisinprasert S. Molecular cloning of kman coding for mannanase from *Klebsiella oxytoca* KUB-CW2-3 and its hybrid mannanase characters. *Enzyme Microb Technol.* (2016) 89:39–51. doi: 10.1016/j.enzmictec.2016.03.005
56. You J, Liu J-F, Yang S-Z, Mu B-Z. Low-temperature-active and salt-tolerant β -mannanase from a newly isolated *Enterobacter* sp. strain N18. *J Biosci Bioeng.* (2016) 121:140–6. doi: 10.1016/j.jbiosc.2015.06.001
57. Zakaria MM, Ashiuchi M, Yamamoto S, Yagi T. Optimization for β -mannanase production of a psychrophilic bacterium, *Flavobacterium* sp. *Biosci Biotechnol Biochem.* (1998) 62:655–60. doi: 10.1271/bbb.62.655
58. Reddy SK, Bågenholm V, Pudlo NA, Bouraoui H, Koropatkin NM, Martens EC, et al. A β -mannan utilization locus in *Bacteroides ovatus* involves a GH36 α -galactosidase active on galactomannans. *FEBS Lett.* (2016) 590:2106–18. doi: 10.1002/1873-3468.12250
59. Cordeiro RL, Pirolla RaS, Persinoti GF, Gozzo FC, De Giuseppe PO, Murakami MT. N-glycan utilization by bifidobacterium gut symbionts involves a specialist β -mannosidase. *J Mol Biol.* (2019) 431:732–47. doi: 10.1016/j.jmb.2018.12.017
60. Sonnenburg JL, Chen CT, Gordon JI. Genomic and metabolic studies of the impact of probiotics on a model gut symbiont and host. *PLoS Biol.* (2006) 4:e413. doi: 10.1371/journal.pbio.0040413
61. Krishnaswamyreddy S. *galactmannan Degradation by Fungi and Gut Bacteria: Structural Enzymology and Fine-Tuned Substrate Specificity*. Lund: Lund University (2016).
62. Bågenholm V, Reddy SK, Bouraoui H, Morrill J, Kulcinskaja E, Bahr CM, et al. Galactomannan catabolism conferred by a polysaccharide utilization locus of *Bacteroides ovatus* enzyme synergy and crystal structure of a β -mannanase *J Biol Chem.* (2017) 292:229–43. doi: 10.1074/jbc.M116.746438
63. Dias FM, Vincent F, Pell G, Prates JA, Centeno MS, Tailford LE, et al. Insights into the molecular determinants of substrate specificity in glycoside hydrolase family 5 revealed by the crystal structure and kinetics of *Cellvibrio mixtus* mannosidase 5A. *J Biol Chem.* (2004) 279:25517–26. doi: 10.1074/jbc.M401647200
64. Tailford LE, Money VA, Smith NL, Dumon C, Davies GJ, Gilbert HJ. Mannose foraging by *Bacteroides thetaiotaomicron* structure and specificity of the β -mannosidase, BtMan2A. *J Biol Chem.* (2007) 282:11291–9. doi: 10.1074/jbc.M610964200
65. Rahmani N, Kashiwagi N, Lee J, Niimi-Nakamura S, Matsumoto H, Kahar P, et al. Mannan endo-1,4- β -mannosidase from *Kitasatospora* sp. isolated in Indonesia and its potential for production of manno oligosaccharides from mannan polymers. *AMB Express.* (2017) 7:100. doi: 10.1186/s13568-017-0401-6
66. Hachem MA, Fredslund F, Andersen JM, Jonsgaard Larsen R, Majumder A, Ejby M, et al. Raffinose family oligosaccharide utilisation by probiotic bacteria: insight into substrate recognition, molecular architecture and diversity of GH36 α -galactosidases. *Biotatal Biotransformat.* (2012) 30:316–25. doi: 10.3109/10242422.2012.674717
67. Ojala T, Kuparinen V, Koskinen JP, Alatalo E, Holm L, Auvinen P, et al. Genome sequence of *Lactobacillus crispatus* ST1. *J Bacteriol Res.* (2010) 192:3547–8. doi: 10.1128/JB.00399-10
68. Comfort DA, Bobrov KS, Ivanen DR, Shabalin KA, Harris JM, Kulminkskaya AA, et al. Cold-active β -galactosidase from *Arthrobastrum* sp. C2-2 α -galactosidase (Tm GalA) confirms the mechanistic commonality of clan GH-D glycoside hydrolases. *Biochemistry.* (2007) 46:3319–30. doi: 10.1021/bi061521n
69. Skalova T, Dohnálek J, Spiwok V, Lipovová P, Vondráčková E, Petroková H, et al. Cold-active β -galactosidase from *Arthrobastrum* sp. C2-2 forms compact 660 kDa hexamers: crystal structure at 19 Å resolution. *J Mol Biol.* (2005) 353:282–94. doi: 10.1016/j.jmb.2005.08.028
70. Huang Y, Zhang H, Ben P, Duan Y, Lu M, Li Z, et al. Characterization of a novel GH36 α -galactosidase from *Bacillus megaterium* and its application in degradation of raffinose family oligosaccharides. *Int J Biol Macromol.* (2018) 108:98–104. doi: 10.1016/j.ijbiomac.2017.11.154
71. Schröder C, Janzer V-A, Schirrmacher G, Claren J, Antranikian G. Characterization of two novel heat-active α -galactosidases from thermophilic bacteria. *Extremophiles.* (2017) 21:85–94. doi: 10.1007/s00792-016-0885-z
72. Gote M, Khan M, Gokhale D, Bastawde K, Khire J. Purification, characterization and substrate specificity of thermostable α -galactosidase from *Bacillus stearothermophilus* (NCIM-5146). *Process Biochem.* (2006) 41:1311–7. doi: 10.1016/j.procbio.2006.01.003
73. Lisowska B. *Genomic Analysis and Metabolic Modelling of Geobacillus thermoglucosidarius NCIMB 11955*. Bath: University of Bath (2016).
74. Gherardini F, Babcock M, Salyers AA. Purification and characterization of two alpha-galactosidases associated with catabolism of guar gum and other alpha-galactosides by *Bacteroides ovatus*. *J Bacteriol Res.* (1985) 161:500–6. doi: 10.1128/jb.161.2.500-506.1985
75. Goulas T, Goulas A, Tzortzis G, Gibson GR. A novel α -galactosidase from *Bifidobacterium bifidum* with transgalactosylating properties: gene molecular cloning and heterologous expression. *Appl Microbiol Biotechnol.* (2009) 82:471–7. doi: 10.1007/s00253-008-1750-5
76. Van Laere K, Hartemink R, Beldman G, Pitson S, Dijkema C, Schols H, et al. Transglycosidase activity of *Bifidobacterium adolescentis* DSM 20083 α -galactosidase. *Appl Microbiol Biotechnol.* (1999) 52:681–8. doi: 10.1007/s002530051579
77. Zhao H, Lu L, Xiao M, Wang Q, Lu Y, Liu C, et al. Cloning and characterization of a novel α -galactosidase from *Bifidobacterium breve* 203 capable of synthesizing Gal- α -1, 4 linkage. *FEMS Microbiol Lett*285. (2008) 278–83. doi: 10.1111/j.1574-6968.2008.01246.x
78. Kimura T, Sakka K, Ohmiya K. Cloning, sequencing, and expression of the gene encoding the *Clostridium stercorarium* α -Galactosidase Aga36A in *Escherichia coli*. *Biosci Biotechnol Biochem.* (2003) 67:2160–6. doi: 10.1271/bbb.67.2160
79. Ademark P, Lundqvist J, Häggglund P, Tenkanen M, Torto N, Tjerneld F, et al. Hydrolytic properties of a β -mannosidase purified from *Aspergillus niger*. *J Biotechnol.* (1999) 75:281–9. doi: 10.1016/S0168-1656(99)00172-8
80. De Vries RP, Van Den Broeck HC, Dekkers E, Manzanares P, De Graaff LH, Visser J. Differential expression of three α -galactosidase genes and a single β -galactosidase gene from *Aspergillus niger*. *Appl Environ Microbiol.* (1999) 65:2453–60. doi: 10.1128/AEM.65.6.2453-2460.1999
81. Hamouda HI, Ali N, Su H, Feng J, Lu M, Li F-L. Exploration of two pectate lyases from *Caldicellulosiruptor bescii* reveals that the CBM66 module has a crucial role in pectic biomass degradation. *Appl Environ Microbiol.* (2020) 86:e00787–20. doi: 10.1128/AEM.00787-20
82. Martens EC, Lowe EC, Chiang H, Pudlo NA, Wu M, McNulty NP, et al. Recognition and degradation of plant cell wall polysaccharides by two human gut symbionts. *PLoS Biol.* (2011) 9:e1001221. doi: 10.1371/journal.pbio.1001221

83. Luis AS, Martens EC. Interrogating gut bacterial genomes for discovery of novel carbohydrate degrading enzymes. *Curr Opin Chem Biol.* (2018) 47:126–33. doi: 10.1016/j.cbpa.2018.09.012
84. Klug-Santner BG, Schnitzhofer W, Vršanská M, Weber J, Agrawal PB, Nierstrasz VA, et al. Purification and characterization of a new bioscouring pectate lyase from *Bacillus pumilus* BK2. *J Biotechnol.* (2006) 121:390–401. doi: 10.1016/j.jbiotec.2005.07.019
85. Lopez-Siles M, Khan TM, Duncan SH, Harmsen HJ, Garcia-Gil LJ, Flint HJ. Cultured representatives of two major phylogroups of human colonic *Faecalibacterium prausnitzii* can utilize pectin, uronic acids, and host-derived substrates for growth. *Appl Environ Microbiol.* (2012) 78:420–8. doi: 10.1128/AEM.06858-11
86. Hugouvieux-Cotte-Pattat N, Condemine G, Shevchik VE. Bacterial pectate lyases, structural and functional diversity. *Environ Microbiol Rep.* (2014) 6:427–40. doi: 10.1111/1758-2229.12166
87. Bekli S, Aktas B, Gencer D, Aslim B. Biochemical and molecular characterizations of a novel pH and temperature-stable pectate lyase from *Bacillus amyloliquefaciens* S6 for industrial application. *Mol Biotechnol.* (2019) 61:681–93. doi: 10.1007/s12033-019-00194-2
88. Yaniv O, Jindou S, Frolow F, Lamed R, Bayer EA. A simple method for determining specificity of carbohydrate-binding modules for purified and crude insoluble polysaccharide substrates. In: *Biomass Conversion*. Totowa, NJ: Humana Press (2012). p. 101–7.
89. Tamaru Y, Doi RH. Pectate lyase A, an enzymatic subunit of the *Clostridium cellulovorans* cellulosome. *PNAS.* (2001) 98:4125–9. doi: 10.1073/pnas.071045598
90. Tonouchi A, Hara Y, Umehara R, Sanuki T, Fukusawa T, Miyairi K. Cloning of the gene encoding an endo-acting pectate lyase from *Streptomyces thermocarboxydus*. *Biosci Biotechnol Biochem.* (2010) 74, 433–6. doi: 10.1271/bbb.90693
91. Fujita K, Sakaguchi T, Sakamoto A, Shimokawa M, Kitahara K. *Bifidobacterium longum* subsp. *longum* exo- β -1, 3-galactanase, an enzyme for the degradation of type II arabinogalactan. *Appl Environ Microbiol.* (2014) 80:4577–84. doi: 10.1128/AEM.00802-14
92. Ochiai A, Itoh T, Kawamata A, Hashimoto W, Murata K. Plant cell wall degradation by saprophytic *Bacillus subtilis* strains: gene clusters responsible for rhamnogalacturonan depolymerization. *Appl Environ Microbiol.* (2007) 73:3803–13. doi: 10.1128/AEM.00147-07
93. Silva IR, Larsen DM, Meyer AS, Mikkelsen JD. Identification, expression, and characterization of a novel bacterial RGI Lyase enzyme for the production of bio-functional fibers. *Enzyme Microb Technol.* (2011) 49:160–6. doi: 10.1016/j.enzmictec.2011.04.015
94. Mckie VA, Vincken J-P, Voragen AG, Van Den Broek LA, Stimson E, Gilbert HJ. A new family of rhamnogalacturonan lyases contains an enzyme that binds to cellulose. *Biochem J.* (2001) 355:167–77. doi: 10.1042/bj3550167
95. Pagès S, Valette O, Abdou L, Bélaïch A, Bélaïch J-P. A rhamnogalacturonan lyase in the *Clostridium cellulolyticum* cellulosome. *J Bacteriol Res.* (2003) 185:4727–33. doi: 10.1128/JB.185.16.4727-4733.2003
96. Kunishige Y, Iwai M, Nakazawa M, Ueda M, Tada T, Nishimura S, et al. Crystal structure of exo-rhamnogalacturonan lyase from *Penicillium chrysogenum* as a member of polysaccharide lyase family 26. *FEBS Lett.* (2018) 592:1378–88. doi: 10.1002/1873-3468.13034
97. Ichinose H, Kuno A, Kotake T, Yoshida M, Sakka K, Hirabayashi J, et al. Characterization of an exo- β -1, 3-galactanase from *Clostridium thermocellum*. *Appl Environ Microbiol.* (2006) 72:3515–23. doi: 10.1128/AEM.72.5.3515-3523.2006
98. Ichinose H, Yoshida M, Kotake T, Kuno A, Igarashi K, Tsumuraya Y, et al. An exo- β -1, 3-galactanase having a novel β -1, 3-galactan-binding module from *Phanerochaete chrysosporium*. *J Biol Chem.* (2005) 280:25820–9. doi: 10.1074/jbc.M501024200
99. Sakamoto T, Tanaka H, Nishimura Y, Ishimaru M, Kasai N. Characterization of an exo- β -1, 3-d-galactanase from *Sphingomonas* sp. 24T and its application to structural analysis of larch wood arabinogalactan. *Appl Microbiol Biotechnol.* (2011) 90:1701–10. doi: 10.1007/s00253-011-3219-1
100. Fujita K, Sasaki Y, Kitahara K. Degradation of plant arabinogalactan proteins by intestinal bacteria: characteristics and functions of the enzymes involved. *Appl Microbiol Biotechnol.* (2019) 103:7451–7. doi: 10.1007/s00253-019-10049-0
101. Cartmell A, Muñoz-Muñoz J, Briggs JA, Ndeh DA, Lowe EC, Baslé A, et al. A surface endogalactanase in *Bacteroides thetaiotaomicron* confers keystone status for arabinogalactan degradation. *Nat Microbiol.* (2018) 3:1314–26. doi: 10.1038/s41564-018-0258-8
102. Ichinose H, Kotake T, Tsumuraya Y, Kaneko S. Characterization of an endo- β -1, 6-galactanase from *Streptomyces avermitilis* NBRC14893. *Appl Environ Microbiol.* (2008) 74:2379–83. doi: 10.1128/AEM.01733-07
103. Chen S, Wilson DB. Proteomic and transcriptomic analysis of extracellular proteins and mRNA levels in *Thermobifida fusca* grown on cellobiose and glucose. *J Bacteriol Res.* (2007) 189:6260–5. doi: 10.1128/JB.00584-07
104. Sharma K, Fontes CM, Najmudin S, Goyal A. Molecular organization and protein stability of the *Clostridium thermocellum* glucuronoxylan endo- β -1, 4-xylanase of family 30 glycoside hydrolase in solution. *J Struct Biol.* (2019) 206:335–44. doi: 10.1016/j.jsb.2019.04.005
105. St. John FJ, Rice JD, Preston JF. Characterization of XynC from *Bacillus subtilis* subsp. *subtilis* strain 168 and analysis of its role in depolymerization of glucuronoxylan. *J Bacteriol Res.* (2006) 188:8617–26. doi: 10.1128/JB.01283-06
106. Maehara T, Yagi H, Sato T, Ohnishi-Kameyama M, Fujimoto Z, Kamino K, et al. GH30 glucuronoxylan-specific xylanase from *Streptomyces turgidiscabies* C56. *Appl Environ Microbiol.* (2018) 84:e01850–17. doi: 10.1128/AEM.01850-17
107. Lee SH, Lee MJ, Cho H-Y, Lee JS, Rhee YH, Shin D-H, et al. Genetic and functional characterization of a novel GH10 endo- β -1, 4-xylanase with a ricin-type β -trefoil domain-like domain from *Luteimicrobium xylanilyticum* HY-24. *Int J Biol Macromol.* (2018) 106:620–8. doi: 10.1016/j.ijbiomac.2017.08.063
108. Conway JM, Pierce WS, Le JH, Harper GW, Wright JH, Tucker AL, et al. Multidomain, surface layer-associated glycoside hydrolases contribute to plant polysaccharide degradation by *Caldicellulosiruptor* species. *J Biol Chem.* (2016) 291:6732–47. doi: 10.1074/jbc.M115.707810
109. Morais S, David YB, Bensoussan L, Duncan SH, Koropatkin NM, Martens EC, et al. Enzymatic profiling of cellulosomal enzymes from the human gut bacterium, *Ruminococcus champanellensis*, reveals a fine-tuned system for cohesin-dockerin recognition. *Environ Microbiol.* (2016) 18:542–56. doi: 10.1111/1462-2920.13047
110. Ze X, David YB, Laverde-Gomez JA, Dassa B, Sheridan PO, Duncan SH, et al. Unique organization of extracellular amylases into amyloosomes in the resistant starch-utilizing human colonic Firmicutes bacterium *Ruminococcus bromii*. *MBio.* (2015) 6:e01058–15. doi: 10.1128/mBio.01058-15
111. Ravachol J, De Philip P, Borne R, Mansuelle P, Maté MJ, Perret S, et al. Mechanisms involved in xyloglucan catabolism by the cellulosome-producing bacterium *Ruminoclostridium cellulolyticum*. *Sci Rep.* (2016) 6:1–17. doi: 10.1038/srep22770
112. Matsuzawa T, Kameyama A, Yaoi K. A novel isoprimeverose-producing enzyme from *Phaeoacremonium minimum* is active with low concentrations of xyloglucan oligosaccharides. *FEBS Open Bio.* (2019) 9:92–100. doi: 10.1002/2211-5463.12549
113. Guerfali M, Gargouri A, Belghith H. Catalytic properties of *Talaromyces thermophilus* α -1-arabinofuranosidase and its synergistic action with immobilized endo- β -1, 4-xylanase. *J Mol Catal.* (2011) 68:192–9. doi: 10.1016/j.molcatb.2010.11.003
114. Larsbrink J, Izumi A, Ibatullin FM, Nakhai A, Gilbert HJ, Davies GJ, et al. Structural enzymatic characterization of a glycoside hydrolase family 31 α -xylosidase from *Cellvibrio japonicus* involved in xyloglucan saccharification. *Biochem J.* (2011) 436:567–80. doi: 10.1042/BJ20110299
115. Florindo RN, Souza VP, Manzine LR, Camilo CM, Marana SR, Polikarpov I, et al. Structural and biochemical characterization of a GH3 β -glucosidase from the probiotic bacteria *Bifidobacterium adolescentis*. *Biochimie.* (2018) 148:107–15. doi: 10.1016/j.biochi.2018.03.007
116. Hemsworth GR, Thompson AJ, Stepper J, Sobala LF, Coyle T, Larsbrink J, et al. Structural dissection of a complex *Bacteroides ovatus* gene locus conferring xyloglucan metabolism in the human gut. *Open Bio.* (2016) 6:160142. doi: 10.1098/rsob.160142
117. Nakajima M, Yoshida R, Miyana A, Abe K, Takahashi Y, Sugimoto N, et al. Functional and structural analysis of a β -glucosidase involved in β -1, 2-glucan metabolism in *Listeria innocua*. *PLoS ONE.* (2016) 11:e0148870. doi: 10.1371/journal.pone.0148870
118. Zmudka MW, Thoden JB, Holden HM. The structure of DesR from *Streptomyces venezuelae*, a β -glucosidase involved in macrolide activation. *Protein Sci.* (2013) 22:883–92. doi: 10.1002/pro.2204
119. Cota J, Corrêa TL, Damasio AR, Diogo JA, Hoffmam ZB, Garcia W, et al. Comparative analysis of three hyperthermophilic GH1 and GH3 family members with industrial potential. *N Biotechnol.* (2015) 32:13–20. doi: 10.1016/j.nbt.2014.07.009
120. Hong M-R, Kim Y-S, Park C-S, Lee J-K, Kim Y-S, Oh D-K. Characterization of a recombinant β -glucosidase from the thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *J Biosci Bioeng.* (2009) 108:36–40. doi: 10.1016/j.jbiosc.2009.02.014
121. Wright RM, Yablonsky MD, Shalita ZP, Goyal A, Eveleigh D. Cloning, characterization, and nucleotide sequence of a gene encoding *Microbispora bispora* BglB, a thermostable beta-glucosidase expressed in *Escherichia coli*.

- Appl Environ Microbiol.* (1992) 58:3455–65. doi: 10.1128/aem.58.11.3455-3465.1992
122. Breves R, Bronnenmeier K, Wild N, Lottspeich F, Staudenbauer WL, Hofemeister J. Genes encoding two different beta-glucosidases of *Thermoanaerobacter brockii* are clustered in a common operon. *Appl Environ Microbiol.* (1997) 63:3902–10. doi: 10.1128/aem.63.10.3902-3910.1997
123. Mckie VA, Black GW, Millward-Sadler SJ, Hazlewood GP, Laurie JI, Gilbert HJ. Arabinanase A from *Pseudomonas fluorescens* subsp. *cellulosa* exhibits both an endo- and an exo-mode of action. *Biochem J.* (1997) 323:547–55. doi: 10.1042/bj3230547
124. Nagel CW, Vaughn RH. The degradation of oligogalacturonides by the polygalacturonase of *Bacillus polymyxa*. *Arch Biochem Biophys.* (1961) 94:328–32. doi: 10.1016/0003-9861(61)90047-9
125. Edstrom R, Phaff H. Purification and certain properties of pectin trans-eliminase from *Aspergillus fonscaeus*. *J Biol Chem.* (1963) 239:2403–8. doi: 10.1016/S0021-9258(18)93866-4
126. Nasuno S, Starr M. Polygalacturonic acid trans-eliminase of *Xanthomonas campestris*. *Biochem J.* (1967) 104:178. doi: 10.1042/bj1040178
127. Mayans O, Scott M, Connerton I, Gravesen T, Benen J, Visser J, et al. Two crystal structures of pectin lyase A from *Aspergillus* reveal a pH driven conformational change and striking divergence in the substrate-binding clefts of pectin and pectate lyases. *Structure.* (1997) 5:677–89. doi: 10.1016/S0969-2126(97)00222-0
128. Shevchik VE, Kester HC, Benen JA, Visser J, Robert-Baudouy J, Hugouvieux-Cotte-Pattat N. Characterization of the exopolygalacturonate lyase PelX of *Erwinia chrysanthemi* 3937. *J Bacteriol Res.* (1999) 181:1652–63. doi: 10.1128/JB.181.5.1652-1663.1999
129. Mutter M, Beldman G, Schols HA, Voragen A. Rhamnogalacturonan [alpha]-L-rhamnopyranohydrolase (a novel enzyme specific for the terminal nonreducing rhamnosyl unit in rhamnogalacturonan regions of pectin). *Plant Physiol.* (1994) 106:241–50. doi: 10.1104/pp.106.1.241
130. Mutter M, Colquhoun IJ, Schols HA, Beldman G, Voragen A. Rhamnogalacturonase B from *Aspergillus aculeatus* is a rhamnogalacturonan [alpha]-L-rhamnopyranosyl-(1->4)-[alpha]-D-galactopyranosyluronide lyase. *Plant Physiol.* (1996) 110:73–7. doi: 10.1104/pp.110.1.73
131. Fujita K, Takashi Y, Obuchi E, Kitahara K, Suganuma T. Characterization of a novel beta-L-Arabinofuranosidase in *Bifidobacterium longum* functional elucidation of a DUF1680 protein family member. *J Biol Chem.* (2014) 289:5240–9. doi: 10.1074/jbc.M113.528711
132. St John FJ, González JM, Pozharski E. Consolidation of glycosyl hydrolase family 30: a dual domain 4/7 hydrolase family consisting of two structurally distinct groups. *FEBS Lett.* (2010) 584:4435–41. doi: 10.1016/j.febslet.2010.09.051
133. Aspeborg H, Coutinho PM, Wang Y, Brumer H, Henrissat B. Evolution, substrate specificity and subfamily classification of glycoside hydrolase family 5 (GH5). *BMC Evol Biol.* (2012) 12:186. doi: 10.1186/1471-2148-12-186
134. Fujita K, Sakamoto A, Kaneko S, Kotake T, Tsumuraya Y, Kitahara K. Degradative enzymes for type II arabinogalactan side chains in *Bifidobacterium longum* subsp. *longum*. *Appl Microbiol Biotechnol.* (2019) 103:1299–310. doi: 10.1007/s00253-018-9566-4
135. Takenishi S, Tsujisaka Y. On the modes of action of three xylanases produced by a strain of *Aspergillus niger* van Tieghem. *Agric Biol Chem.* (1975) 39:2315–23. doi: 10.1271/bbb1961.39.2315
136. Comtat J, Joseleau J-P. Mode of action of a xylanase and its significance for the structural investigation of the branched L-arabino-D-glucurono-D-xylan from redwood (*Sequoia sempervirens*). *Carbohydr Res.* (1981) 95:101–12. doi: 10.1016/S0008-6215(00)85299-4
137. Nishitani K, Nevins D. Glucuronoxylan xylanohydrolase. A unique xylanase with the requirement for appendant glucuronosyl units. *J Biol Chem.* (1991) 266:6539–43. doi: 10.1016/S0021-9258(18)38151-1
138. Koyama T, Hayashi T, Kato Y, Matsuda K. Degradation of xyloglucan by wall-bound enzymes from soybean tissue I. Occurrence of xyloglucan-degrading enzymes in soybean cell wall. *Plant Cell Physiol.* (1981) 22:1191–8.
139. Koyama T, Hayashi T, Kato Y, Matsuda K. Degradation of xyloglucan by wall-bound enzymes from soybean tissue II: degradation of the fragment heptasaccharide from xyloglucan and the characteristic action pattern of the alpha-D-xylosidase in the enzyme system. *Plant Cell Physiol.* (1983) 24:155–62. doi: 10.1093/pcp/24.2.155
140. Hayashi T. Xyloglucans in the primary cell wall. *Annu Rev Plant Biol.* (1989) 40:139–68. doi: 10.1146/annurev.pp.40.060189.001035
141. Béguin P, Aubert J-P. The biological degradation of cellulose. *FEMS Microbiol Rev.* (1994) 13:25–58. doi: 10.1111/j.1574-6976.1994.tb00033.x
142. Teeri TT. Crystalline cellulose degradation: new insight into the function of cellobiohydrolases. *Trends Biotechnol.* (1997) 15:160–7. doi: 10.1016/S0167-7799(97)01032-9
143. Adlakhia N, Sawant S, Anil A, Lali A, Yazdani SS. Specific fusion of beta-1, 4-endoglucanase and beta-1, 4-glucosidase enhances cellulolytic activity and helps in channeling of intermediates. *Appl Environ Microbiol.* (2012) 78:7447–54. doi: 10.1128/AEM.01386-12
144. De Vries S, Pustjens A, Schols H, Hendriks W, Gerrits W. Improving digestive utilization of fiber-rich feedstuffs in pigs and poultry by processing and enzyme technologies: a review. *Anim Feed Sci Technol.* (2012) 178:123–38. doi: 10.1016/j.anifeeds.2012.10.004
145. De Vries S, Pustjens AM, Kabel MA, Salazar-Villanea S, Hendriks WH, Gerrits WJ. Processing technologies and cell wall degrading enzymes to improve nutritional value of dried distillers grain with solubles for animal feed: an *in vitro* digestion study. *J Agric Food Chem.* (2013) 61:8821–8. doi: 10.1021/jf4019855
146. Liu Z-S, Wu X-L, Kida K, Tang Y-Q. Corn stover saccharification with concentrated sulfuric acid: effects of saccharification conditions on sugar recovery and by-product generation. *Bioresour Technol.* (2012) 119:224–33. doi: 10.1016/j.biortech.2012.05.107
147. Teitge D, Campbell G, Classen H, Thacker P. Heat pretreatment as a means of improving the response to dietary pentosanase in chicks fed rye. *Can J Anim Sci.* (1991) 71:507–13. doi: 10.4141/cjas91-060
148. Long C, De Vries S, Venema K. Polysaccharide source altered ecological network, functional profile, and short-chain fatty acid production in a porcine gut microbiota. *Benef Microbes.* (2020) 11:591–610. doi: 10.3920/BM2020.0006
149. Long C, De Vries S, Venema K. Differently pre-treated rapeseed meals affect *in vitro* swine gut microbiota composition. *Front Microbiol.* (2020) 11:570985. doi: 10.3389/fmicb.2020.570985
150. Long C, Rosch C, De Vries S, Schols H, Venema K. Cellulase and alkaline treatment improve intestinal microbial degradation of recalcitrant fibers of rapeseed meal in pigs. *J Agric Food Chem.* (2020) 68:11011–25. doi: 10.1021/acs.jafc.0c03618
151. Long C, Venema K. Pretreatment of rapeseed meal increases its recalcitrant fiber fermentation and alters the microbial community in an *in vitro* model of swine large intestine. *Front Microbiol.* (2020) 11:588264. doi: 10.3389/fmicb.2020.588264
152. Xie F, Zhang W, Lan X, Gong S, Wu J, Wang Z. Effects of high hydrostatic pressure and high pressure homogenization processing on characteristics of potato peel waste pectin. *Carbohydr Polym.* (2018) 196:474–82. doi: 10.1016/j.carbpol.2018.05.061
153. Peng X-Y, Mu T-H, Zhang M, Sun H-N, Chen J-W, Yu M. Effects of pH and high hydrostatic pressure on the structural and rheological properties of sugar beet pectin. *Food Hydrocoll.* (2016) 60:161–9. doi: 10.1016/j.foodhyd.2016.03.025
154. Bulbul V, Bhushette PR, Zambare RS, Deshmukh R, Annature US. Effect of cold plasma treatment on *Xanthan gum* properties. *Polym Test.* (2019) 79:106056. doi: 10.1016/j.polymertesting.2019.106056
155. Prasertsung I, Aroonraj K, Kamwilaisak K, Saito N, Damrongsakkul S. Production of reducing sugar from cassava starch waste (CSW) using solution plasma process (SPP). *Carbohydr Polym.* (2019) 205:472–9. doi: 10.1016/j.carbpol.2018.10.090