

Assessing balance in people with bilateral vestibulopathy using the Mini-Balance Evaluation Systems Test (Mini-BESTest): feasibility and comparison with healthy control data

Citation for published version (APA):

Zhu, M., van Stiphout, L., Karabulut, M., Pérez Fornos, A., Guinand, N., Meijer, K., van de Berg, R., & McCrum, C. (2023). Assessing balance in people with bilateral vestibulopathy using the Mini-Balance Evaluation Systems Test (Mini-BESTest): feasibility and comparison with healthy control data. *Journal of Neurology*, 270(9), 4423-4433. Advance online publication. <https://doi.org/10.1007/s00415-023-11795-y>

Document status and date:

Published: 01/09/2023

DOI:

[10.1007/s00415-023-11795-y](https://doi.org/10.1007/s00415-023-11795-y)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

CC BY

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 02 Oct. 2023



Assessing balance in people with bilateral vestibulopathy using the Mini-Balance Evaluation Systems Test (Mini-BESTest): feasibility and comparison with healthy control data

Meichan Zhu^{1,2,3} · Lisa van Stiphout¹ · Mustafa Karabulut¹ · Angélica Pérez Fornos⁴ · Nils Guinand⁴ · Kenneth Meijer² · Raymond van de Berg¹ · Christopher McCrum²

Received: 20 April 2023 / Revised: 23 May 2023 / Accepted: 24 May 2023
© The Author(s) 2023

Abstract

Objectives Bilateral vestibulopathy (BVP) leads to unsteadiness when walking, which worsens in darkness or on uneven ground, as well as falls. Since simple balance tests struggle to distinguish between BVP and healthy participants, we aimed (1) to test if the Mini-BESTest is feasible in BVP, (2) how people with BVP perform on the Mini-BESTest and (3) to compare these scores with healthy reference data.

Methods Fifty participants with BVP completed the Mini-BESTest. 12-month falls incidence was obtained by questionnaire. To compare the overall and sub-scores between our participants with BVP and those of healthy participants from the literature ($n = 327$; obtained via PubMed searches), Mann–Whitney U tests were used. Sub scores within the BVP group were also compared. Spearman correlations were used to investigate the relationships between Mini-BESTest score and age.

Results No floor or ceiling effects were observed. Participants with BVP had significantly lower Mini-BESTest total scores than the healthy group. Anticipatory, reactive postural control and sensory orientation sub scores of the Mini-BESTest were significantly lower in BVP, while dynamic gait sub scores were not significantly different. A stronger negative correlation between age and Mini-BESTest total score was found in BVP than in the healthy group. Scores did not differ between patients with different falls history.

Conclusion The Mini-BESTest is feasible in BVP. Our results confirm the commonly reported balance deficits in BVP. The stronger negative association between age and balance in BVP might reflect the age-related decline in the remaining sensory systems with which people with BVP compensate.

Keywords Balance · Mini-Balance Evaluation Systems Test · Bilateral vestibulopathy

Introduction

Bilateral vestibulopathy (BVP) is characterized by bilateral hypofunction of the vestibular organ or nerves which is diagnosed using the criteria reported by the Barany Society

✉ Meichan Zhu
z.meichan@maastrichtuniversity.nl

✉ Christopher McCrum
chris.mccrum@maastrichtuniversity.nl

¹ Division of Balance Disorders, Department of Otorhinolaryngology and Head and Neck Surgery, School for Mental Health and Neuroscience, Maastricht University Medical Center, Maastricht, The Netherlands

² Department of Nutrition and Movement Sciences, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, Maastricht, The Netherlands

³ Department of Otorhinolaryngology, Guangzhou Twelfth People's Hospital (Guangzhou Otolaryngology-Head and Neck Surgery Hospital), No. 1 Tianqiang Road, Tianhe District, Guangzhou 510620, Guangdong, China

⁴ Service of Otorhinolaryngology and Head and Neck Surgery, Department of Clinical Neurosciences, Geneva University Hospitals, Geneva, Switzerland

[1]. The symptoms of BVP include oscillopsia and postural instability. Importantly, BVP has many different aetiologies with many different disease presentations [2]. About 30–40% of people with BVP suffer from oscillopsia during head motion, due to reduced or absent vestibular-ocular reflex (VOR) [3]. In addition, people with BVP have a higher risk of falls [4], increased gait variability [5, 6] and often report difficulty and instability while walking in dimly lit environments and on uneven ground [7–9]. Consequently, avoidance of falls and reduction in mobility may occur and negatively impact societal participation, which may be linked with the increase in depression and reduced quality of life in people with vestibular disorders [10–15].

Deficient vestibular function may be defined using video head impulse testing (vHITs), caloric testing and torsion swing test and evaluation of dynamic visual acuity (DVA), cervical and ocular vestibular-evoked myogenic potentials for otolith function and Romberg testing for balance [1, 16, 17]. However, McCrum et al. [6] found no clear correlation between various gait variability parameters and the results of the caloric, vHIT and DVA tests, indicating that specific, objective assessment of balance and gait may be required in BVP to gain a better picture of a patient's deficits.

Regarding the increasing falls risk in BVP, the data on exactly how falls occur is limited [4, 8, 9, 12, 18, 19]. People with BVP compensate with a combination of the remaining proprioceptive and visual function. As such, their balance might not be sufficiently examined by tasks which use zero or only a single sensory perturbation [20]. Tools such as the Berg Balance Scale (BBS), the Timed "Up & Go" Test (TUG) and the Romberg test, which is recommended by the Barany Society [1], may therefore, not be suitable [20]. Herssens et al. [20] reported that few studies have concentrated on the balance abilities of people with BVP during walking. This is critical since evaluation of the systems underlying balance deficits and falls in BVP is needed to identify and intervene on those deficits [20]. As Horak et al. [21] outlined, balance (or postural control) is a complex skill based on the interaction of dynamic sensorimotor processes, including active alignment of the head and body (with respect to gravity, support surfaces, the visual surround and internal references), integration and weighting of sensory information from somatosensory, vestibular and visual systems, with each of these broad factors being weighted according to the current task, goals and environment. For this reason, Herssens et al. [20] recommended that further research on the Balance Evaluation Systems Test [22] in BVP is needed. The refined Mini-Balance Evaluation Systems Test (Mini-BESTest) assesses multiple dynamic balance tasks related to four subcomponents of balance (anticipatory postural adjustments, reactive postural control, sensory orientation, and dynamic gait) [23]. However, its

feasibility and ability to assess the specific balance deficits reported in people with BVP has yet to be evaluated.

To address the current gap in research and practice, this study investigated balance function in people with BVP using the Mini-BESTest. We aimed to determine whether the Mini-BESTest can be used with people with BVP (in terms of completion, floor and ceiling effects) and to explore how people with BVP perform in terms of their overall score and their subcomponent scores and if these reveal consistent balance deficits within this group (are specific subcomponents of balance more severely affected in BVP?) and in relation to healthy reference data from the literature (in terms of total score and if subscores focused on sensory function show larger differences). We expected that people with BVP would record lower overall scores than healthy participants and that subcomponents more clearly associated with sensory function would reveal larger deficits.

Methods

Participants

As part of a large prospective study, participants were recruited from Maastricht University Medical Center (MUMC+) and other tertiary referral centres in the Netherlands. Within the MUMC+, all patients diagnosed with BVP at the outpatient clinic of the Department of Otorhinolaryngology were asked to participate in the study in the period from June 2021 to June 2022. Ethical approval was granted by the azM/UM Medical Ethical Committee (METC: NL72200.068.19). Before participating in the study, each participant provided written informed consent. In the current manuscript, we report and analyse the Dutch version Mini-BESTest results from the larger project. Since this was a secondary analysis of the larger trial, no a priori sample size calculation was performed for the current analysis. Sensitivity power analyses are reported in the data analysis section below.

Fifty-three people previously diagnosed with BVP participated in this research. Recruited participants were formerly diagnosed with BVP according to Barany Society BVP diagnostic criteria [1]. On the testing day of the larger project, the diagnosis procedure was repeated, at which point two people were excluded because they no longer had BVP according to the criteria, leaving 51 participants. In the current study, 50 people with BVP are included because one participant did not perform the Mini-BESTest. The exclusion criteria were: a history of other neurological disorders; age under 18 and unable to discontinue vestibulo-suppressive medication. In addition to the procedures described below, participants provided information on their falls incidence in

the previous 12 months by completing a questionnaire based on the recommendations of Lamb et al. [24] and Lord et al. [25] translated into Dutch (materials available at McCrum [26]), that led with the question “In the past year, have you had any fall including a slip or trip in which you lost your balance and landed on the floor or ground or lower level?”.

Mini-BESTest Assessment

The Mini-BESTest was used in this study to evaluate the participants’ balance [23]. The Mini-BESTest evaluates balance tasks associated with various components of balance, namely anticipatory, reactive postural control, sensory orientation and dynamic gait. It comprises of 14 items and each item is scored on a 3-level ordinal scale from 0 (severe = unable to perform), 1 (moderate) and 2 (normal performance) with a maximum score of 28. The Mini-BESTest items are shown in Table 1. The Mini-BESTest was administered and rated by a trained member of the research team. We followed the full standardised equipment, testing and assessment procedures, which can be found at <https://www.bestest.us/> with additional information in the scientific publication [23]. The test was performed as the first assessment in the morning of the larger project’s testing day to ensure mental or physical fatigue would not affect the results.

Healthy reference data

To compare our results from participants with BVP to representative healthy control data, 21 articles from the literature were identified (see the preprint mentioned in the Acknowledgements for full details). From these, we obtained complete individual data for Mini-BESTest total and sub scores from 190 healthy participants in four articles [27–30] and individual Mini-BESTest total scores from an additional 137 healthy participants in four articles [31–34]. These data were combined to form healthy control reference data groups for

our analyses (see below). The available group-level data in the remaining 11 articles [35–45] were used for additional visual comparison.

Data analysis

Due to the current study conducting secondary analyses of a larger project, no a priori sample size calculation was performed. To indicate the statistical power of the current analyses, we performed sensitivity power analyses for the current sample sizes and tests with $\alpha = 0.05$ and $\beta = 0.2$ using G*Power (Version 3.1.9.4) [46] which revealed effect sizes of 0.42 for Wilcoxon signed rank tests ($N = 49$), 0.44 ($N = 50$ vs. $n = 327$) and 0.46 ($N = 50$ vs. $N = 190$) for Mann–Whitney tests.

To assess Mini-BESTest sub score differences, a Friedman test within the BVP group was performed with sub score as the repeated measures factor, with post hoc Bonferroni-adjusted Wilcoxon signed rank tests for pairwise comparisons (note that to account for the different number of items in the sub scores, the sub scores were divided by the number of items before these analyses). To compare the total scores and sub scores between our participants with BVP and the obtained individual data of healthy control participants from the literature, Mann–Whitney U tests were used. The statistical analyses were conducted in Jamovi version 2.2.5 [47]. In addition to the statistical tests, the BVP data were visually compared to the summary values from all other studies obtained from the literature using GraphPad Prism 9 (GraphPad Software, San Diego, California USA, <http://www.graphpad.com>). All quantitative variables are expressed as mean \pm SD, unless stated otherwise. Effect sizes for non-parametric tests were calculated as Cohen’s d according to Fritz et al. [48].

Table 1 Sections and items of the Mini-BESTest [23]

Section	Item
Anticipatory	1. Sit to stand
	2. Rise to toes
	3. Stand on one leg
Reactive postural control	4. Compensatory stepping correction—forward
	5. Compensatory stepping correction—backward
	6. Compensatory stepping correction—lateral
Sensory orientation	7. Stance (feet together); eyes open, firm surface
	8. Stance (feet together); eyes closed, foam surface
	9. Incline—eyes closed
Dynamic gait	10. Change in gait speed
	11. Walk with head turns—horizontal
	12. Walk with pivot turns
	13. Step over obstacles
	14. Timed Up and Go with dual task

Results

Participant characteristics

We collected 49 total scores, 49 reactive postural control sub scores and 50 of all other sub scores from our participants with BVP. For one participant, the reactive postural control part of the Mini-BESTest could not be completed for practical reasons and therefore, only the other three sub scores were included in the analyses for this participant. Regarding the healthy control reference data, 327 healthy participants from eight articles [27–34] formed the control group for the Mini-BESTest total score and 190 healthy participants from four articles [27–30] formed the control group for the Mini-BESTest sub scores. Note that two of the contacted authors provided data for more participants than were reported in the identified study [33, 34]. For studies evaluating participants twice or using two assessors to score the performance, we always used the data of the first trial or first assessor reported. One participant in the dataset of Nakhostin-Ansari et al. [30] was excluded from the current analysis due to age (16 years old). Healthy participants were selected from the large public dataset of Santos, Duarte [28] based on the health and clinical information provided to form a younger adult group and middle to older aged adult group. The summary data obtained from the remaining 11 articles [35–45] found with the PubMed search were collated for visual comparison. Demographic data are illustrated in Table 2. The mean ages and sex distributions between the BVP and healthy groups were not statistically significantly different (Table 2). Significant group differences were found in height, body weight and body mass index (BMI) between two groups (Table 2).

Mini-BESTest feasibility

Regarding the Mini-BESTest completion and scores, one participant with BVP scored the maximum (28) and one participant with BVP scored the minimum score (0). We, therefore, confirm that in this sample, no floor or ceiling effects were observed. Other than the reasons for exclusion

mentioned above, no complications or concerning situation arose during the testing.

Mini-BESTest total scores

A Mann–Whitney U test revealed that Mini-BESTest total scores were significantly lower than those observed in the healthy control reference data (Fig. 1; $U(N_{\text{BVP}}=49, N_{\text{Healthy}}=327)=4564.00, p<0.001, d=0.52$) and this effect size exceeded what our sensitivity power analysis determined we could detect. Participants with BVP scored 20.78 ± 5.41 while the healthy controls scored 24.24 ± 2.72 . Mean and standard deviation values from the current study and the identified studies in the literature are shown in Fig. 2, which seem to agree with our statistical analysis, in that the majority of studies on healthy adults report higher mean Mini-BESTest total scores than our BVP group.

Mini-BESTest sub scores

Within the BVP group, the Friedman test revealed a significant sub score effect ($\chi^2_{\text{F}}(3)=29.86, p<0.001$) and post hoc Wilcoxon signed rank tests revealed significant differences




Fig. 1 Mini-BESTest total scores in our BVP and healthy control groups. The boxplots indicate the medians, interquartile range and minimum and maximum values, with the cross indicating the mean values. Maximum score possible indicated by the dotted line

Table 2 Demographic data of the participants with BVP and Healthy Control Reference Groups

	BVP	N	Healthy control	N	Statistical comparison
Age, years	57.56 ± 12.39	50	60.07 ± 17.40	305*	<i>t</i> test $p=0.328$
Male, <i>n</i> (%)	28 (56.0)	50	165 (54.1)	305*	Chi square $\chi^2=0.06, p=0.802$
Height, cm	172.46 ± 8.92	50	164.79 ± 9.20	221*	<i>t</i> test $p<0.001$
Body weight, kg	85.44 ± 21.47	50	66.19 ± 12.90	221*	<i>t</i> test $p<0.001$
BMI	28.55 ± 6.26	50	24.30 ± 3.85	221*	<i>t</i> test $p<0.001$

Mean \pm standard deviation unless otherwise indicated

*Demographic data was not obtained from all studies ($n=327$)




Fig. 2 Mini-BESTest total scores (circles: mean \pm standard deviation; squares: medians) for the current participants with BVP and healthy reference values found in the literature (left to right is young-

est to oldest mean age). Values are shown here as originally reported, meaning that different rounding conventions have been used

between the relative dynamic gait sub score and all other sub scores ($p < 0.001$). No other sub score comparisons were statistically significant (anticipatory vs. reactive postural control: $p = 0.11$; anticipatory vs. sensory orientation: $p = 0.22$; reactive postural control vs. sensory orientation: $p = 0.38$). Mann–Whitney U tests revealed that, compared to the healthy control reference data (Fig. 3), the scores of the participants with BVP were significantly lower for anticipatory, reactive postural control and sensory orientation sub scores [$U(N_{\text{BVP}} = 50, N_{\text{Healthy}} = 190) = 2364.50, p < 0.001,$

$d = 0.75$; $U(N_{\text{BVP}} = 50, N_{\text{Healthy}} = 190) = 3737, p = 0.028, d = 0.3$; and $U(N_{\text{BVP}} = 50, N_{\text{Healthy}} = 190) = 1223.50, p < 0.001, d = 1.22$, respectively], while the dynamic gait sub scores were not statistically different [$U(N_{\text{BVP}} = 50, N_{\text{Healthy}} = 190) = 4374.50, p = 0.367, d = 0.11$]. Participants with BVP scored (mean \pm SD) $4.34 \pm 1.12, 4.00 \pm 1.79, 4.22 \pm 1.96$ and 8.22 ± 2.11 and the healthy controls scored $5.27 \pm 0.93, 4.66 \pm 1.25, 5.66 \pm 0.68$ and 8.43 ± 1.10 for anticipatory, reactive postural control, sensory orientation and dynamic gait sub scores, respectively. Note that the




Fig. 3 Mini-BESTest sub scores in our BVP and healthy control groups. The boxplots indicate the medians, interquartile range and minimum and maximum values, with the points indicating the mean values. Maximum score possible indicated by the dotted lines

reactive postural control score, while statistically significant, did not exceed the effect size for which our analysis was powered and should be interpreted cautiously. In this analysis, the sensory orientation sub score showed the largest effect, and this is also supported by visual comparison with the literature values in Fig. 4, which does support our expectation that sub scores more directly related to sensory function would show larger deficits.

Mini-BESTest total scores and falls history

In the 49 participants with BVP with Mini-BESTest total scores, 25 did not report falling (Mini-BESTest Mean \pm SD, Median: 21.4 ± 5.8 , 23) and 24 did report falling (20.1 ± 5.01 , 21), 18 with two or more falls and six with a single fall. A Mann–Whitney U tests did not find a statistically significant difference in the Mini-BESTest total score between these groups [$U(N_{\text{nofalls}} = 25, N_{\text{falls}} = 24) = 242, p = 0.249, d = 0.235$].

Exploratory analyses on the age–balance association

In our data from the participants with BVP, we observed a potential age effect on the Mini-BESTest scores. To further explore this, we conducted Spearman correlations between participant age and Mini-BESTest total scores for the BVP and healthy control data separately. For the participants with BVP, a significant negative correlation between Mini-BESTest total score and age was found [Spearman's correlation coefficient = -0.67 (95% CI -0.74 to -0.35), $p < 0.001$], and for the healthy control group a significant negative correlation was also found [Spearman's correlation coefficient = -0.32 (95% CI -0.32 to -0.11), $p < 0.001$]. Scatter plots of these data can be seen in Fig. 5. This difference in

age effect is also visible when visualizing the Mini-BESTest total scores separated into young (20–39 years), middle-aged (40–59 years) and older (60+ years) adult age groups (Fig. 6) and when comparing sub score data across these age groups (Table 3).

Discussion

We aimed to determine whether the Mini-BESTest can be used with people with BVP and to explore how people with BVP perform in terms of their overall score and their subcomponent scores and if these reveal consistent balance deficits within this group and in relation to healthy reference data. Our findings supported our hypothesis that BVP would perform significantly worse than healthy controls on the Mini-BESTest total score. Regarding the sub scores, our expectation that subcomponents with more sensory disturbance would reveal larger deficits was partly supported, since the sensory orientation sub score was significantly lower in BVP compared to the healthy control data and the effect size of this difference was the largest difference of the sub score comparisons. However, when compared to the other sub scores within the BVP group, the sensory orientation was only significantly lower than the dynamic gait sub score.

Regarding the application and use of the Mini-BESTest in the BVP population, 49 of our 50 participants had complete scores. The reactive postural control part of the test could not be completed for one participant for practical reasons unrelated to the patient's ability. Floor and ceiling effects were not observed, as only one patient (2.04%) scored 0 and only one patient (2.04%) scored the maximum score which is similar to previous studies on mixed balance disorders [49], Parkinson's disease [50] and chronic stroke [51]. We experienced no other issues related to feasibility or safety when performing the test. As a result, we conclude that the Mini-BESTest can be used in the BVP population, as has been shown with other patient populations such as people with balance disorders [49], Parkinson's disease [50, 52], chronic stroke [51] and subacute stroke [53]. While feasible, there is currently no data on the reliability of the Mini-BESTest in the BVP population, which future work should address.

The Mini-BESTest scores in our participants with BVP (mean \pm SD: 20.78 ± 5.41 , median: 22, interquartile range: 19.0–25.0) are similar to those in some previous reports in people with chronic stroke [51] ($n = 106$, median: 19.0, interquartile range: 14.0–22.0) and Parkinson's disease [50, 52] (King et al. [50]: $n = 97$, median: 23, interquartile range: 20.0–24.0, mean \pm SD: 21.8 ± 3.6 ; Leddy et al. [52]: $n = 80$, Mean \pm SD: 20.2 ± 7.0). Our participants with BVP were significantly taller and heavier, with a higher BMI than the healthy control group which may in part be due to the




Fig. 4 Mini-BESTest sub scores (circles: mean ± standard deviation; squares: medians) for the current participants with BVP and healthy reference values found in the literature (left to right is young-

est to oldest mean age). Values are shown here as originally reported, meaning that different rounding conventions have been used

Netherlands being a taller than average population, combined with the negative effects on daily life physical activity due to BVP [15]. We cannot clearly determine the potential role these differences might have in the Mini-BESTest performance with the current data. However, the age-Mini-BESTest total score association results did not differ much when we reanalysed the data with a partial Spearman

correlation with BMI as a control variable (BVP: $-0.62, p < 0.001$; control: $-0.27, p < 0.001$), suggesting that this difference is not a major contributing factor. Finally, we did not find significant differences in Mini-BESTest total scores between participants with BVP with and without falls in the previous 12 months, despite previous reports that the Mini-BESTest can significantly distinguish between people with

Fig. 5 Scatter plots indicating the relationship between age and Mini-BESTest total score in BVP and healthy control groups





Fig. 6 Mini-BESTest total score across young (20–39 years), middle-aged (40–59 years) and older (60+ years) adult age groups in our participants with BVP and healthy control data from the literature. The boxplots indicate the medians, interquartile range and minimum and maximum values, with the crosses indicating the mean values

and without falls history in other populations [42, 51, 52, 54]. The percentage of participants with BVP with a recent history of falls (48%) was slightly higher than the literature average of 42% reported in a previous review on falls in BVP [4]. While not the main purpose of this study, we also performed area under the receiver operator characteristic curve and sensitivity and specificity analyses for Mini-BESTest total and sub scores and these results can be found in the supplement (Online Resource 1).

We observed a significant association between age and Mini-BESTest total score in both BVP and healthy groups. One of the included studies with healthy control data has previously reported an age-balance relationship [38]. The association was much stronger in the BVP group which we suggest may be related to less possibility for sensory compensation and re-weighting with increasing age. Aside from the vestibular

system, proprioception and vision are important for balance control and all show a gradual age-related decline [25, 55–57] and previous experimental studies have demonstrated more substantial balance performance decline when multiple sensory systems are perturbed [20, 58–61]. These findings imply that close attention might need to be paid to proprioceptive and visual system health in people with BVP to identify and help patients with less possibility for sensory compensation. Sensory substitution devices may also have a role to play here [62].

The current study has some limitations that should be kept in mind. Since participants in this study were volunteering to participate in a full day of assessments as part of the larger study, this may have influenced who chose to participate. We did not record physical activity, sport or physical therapy or rehabilitation history, so prior experience with balance tasks and training may have varied across the participants. One important consideration is that, while sub scores of the Mini-BESTest have been analysed here, these sub scores are not necessarily independent and significant correlation between the sub scores and tasks may exist [63]. While we have reported sub score-specific differences, these should be interpreted with caution and should not necessarily be used for clinical decision making. One participant behaved nervously and did not appear to perform the tasks correctly for psychosomatic reasons (note that this was not the participant that scored 0). Since we had no a priori criteria for how to deal with this situation, we included this participant in the analysis, but also repeated the analysis excluding the participant to examine if this individual score would have an effect on the overall results (this did not lead to any changes in the statistical significance outcomes of the tests, the direction of effects or the conclusions).

In conclusion, our findings demonstrate that the Mini-BESTest can feasibly be used in people with BVP. The Mini-BESTest reveals significantly reduced balance performance in people with BVP compared to healthy control reference data. Within both BVP and healthy groups, a significant negative association between balance performance and age

Table 3 Age group Mini-BESTest total and sub scores (mean \pm SD) for the obtained data in BVP and healthy control groups

Variables	Age group	BVP	N	Control	N
Total score (maximum: 28)	(20–39)	26.25 \pm 1.50	4	24.28 \pm 2.21	50
	(40–59)	22.52 \pm 4.21	23	26.26 \pm 1.43	42
	(60+)	17.95 \pm 5.53	22	23.69 \pm 2.74	213
Anticipatory (maximum 6)	(20–39)	5.50 \pm 0.58	4	5.37 \pm 0.80	46
	(40–59)	4.71 \pm 0.86	24	5.85 \pm 0.49	20
	(60+)	3.73 \pm 1.12	22	5.14 \pm 0.99	124
Reactive postural control (maximum 6)	(20–39)	5.50 \pm 0.58	4	4.52 \pm 1.11	46
	(40–59)	4.52 \pm 1.70	23	5.40 \pm 0.94	20
	(60+)	3.18 \pm 1.68	22	4.59 \pm 1.31	124
Sensory orientation (maximum 6)	(20–39)	5.75 \pm 0.50	4	5.50 \pm 0.69	46
	(40–59)	4.38 \pm 0.82	24	6.00 \pm 0.00	20
	(60+)	3.77 \pm 1.07	22	5.67 \pm 0.72	124
Dynamic gait (maximum 10)	(20–39)	9.50 \pm 0.58	4	8.65 \pm 1.08	46
	(40–59)	8.88 \pm 1.51	24	9.05 \pm 0.69	20
	(60+)	7.27 \pm 2.47	22	8.24 \pm 1.12	124

was found, with the association being twice as high in BVP, indicating that reduced capacity for sensory compensation with increasing age might need to be considered in the assessment and care of people with BVP.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00415-023-11795-y>.

Acknowledgements We thank the authors of the following articles for responding to, and fulfilling our requests for data: Akizuki et al. [27], Dominguez-Olivan et al. [29], Giardini et al. [33], Harro, Garascia [34], Kim et al. [32], Naghdi et al. [31], Nakhostin-Ansari et al. [30]. We also thank Ilaria Arcolin for helpful advice on Mini-BESTest sub score use and interpretation (see limitations). A preprint version of this manuscript is available at OSF Preprints [64].

Author contributions MZ: data curation, formal analysis, funding acquisition, investigation, project administration, visualization, writing—original draft, and writing—review and editing. LvS: conceptualization, data curation, funding acquisition, investigation, project administration, and writing—review and editing. MK: data curation, investigation, and writing—review and editing. APF: conceptualization and writing—review and editing. NG: conceptualization and writing—review and editing. KM: conceptualization, resources, supervision, and writing—review and editing. RvdB: conceptualization, funding acquisition, resources, supervision, and writing—review and editing. CM: conceptualization, data curation, formal analysis, methodology, project administration, supervision, visualization, writing—original draft, and writing—review and editing.

Funding Meichan Zhu was funded by the China Scholarship Council. Lisa van Stiphout was supported by funding from MED-EL (Innsbruck, Austria). Mustafa Karabulut was granted by the Ministry of National Education in the Republic of Türkiye.

Data and code availability Anonymised data of the participants with BVP (some demographic data excluded) and the R code for the statistical analysis performed in Jamovi can be found at: <http://www.doi.org/10.17605/OSF.IO/TFKN3> The healthy control data obtained from the literature are not publicly shared since the authors of this article are not the owners of that data, but they are available on request.

Declarations

Conflicts of interest The authors declare that there is no conflicts of interest relevant to the content of this article.

Ethical standard statement Ethical approval for the study was granted by the azM/UM Medical Ethical Committee (METC: NL72200.068.19) and was conducted in accordance with the Declaration of Helsinki and good clinical practice guidelines. Before participating in the study, each participant provided written informed consent.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Strupp M, Kim JS, Murofushi T, Straumann D, Jen JC, Rosengren SM, Della Santina CC, Kingma H (2017) Bilateral vestibulopathy: diagnostic criteria consensus document of the Classification Committee of the Barany Society. *J Vestib Res* 27(4):177–189. <https://doi.org/10.3233/VES-170619>
2. Lucieer F, Vonk P, Guinand N, Stokroos R, Kingma H, van de Berg R (2016) Bilateral vestibular hypofunction: insights in etiologies, clinical subtypes, and diagnostics. *Front Neurol* 7:26. <https://doi.org/10.3389/fneur.2016.00026>
3. Batuecas-Caletrio A, Trinidad-Ruiz G, Rey-Martinez J, Matíño-Soler E, Martín Sanz E, Pérez Fernández N (2020) Oscillopsia in bilateral vestibular hypofunction: not only gain but saccades too. *Ear Hear* 41(2):323–329. <https://doi.org/10.1097/aud.0000000000000760>

4. Herssens N, How D, van de Berg R, McCrum C (2022) Falls among people with bilateral vestibulopathy: a review of causes, incidence, injuries, and methods. *JAMA Otolaryngol Head Neck Surg* 148(2):187–192. <https://doi.org/10.1001/jamaoto.2021.3673>
5. Schniepp R, Wuehr M, Neuhaeuser M, Kamenova M, Dimitriadis K, Klopstock T, Strupp M, Brandt T, Jahn K (2012) Locomotion speed determines gait variability in cerebellar ataxia and vestibular failure. *Mov Disord* 27(1):125–131. <https://doi.org/10.1002/mds.23978>
6. McCrum C, Lucieer F, van de Berg R, Willems P, Perez Fornos A, Guinand N, Karamanidis K, Kingma H, Meijer K (2019) The walking speed-dependency of gait variability in bilateral vestibulopathy and its association with clinical tests of vestibular function. *Sci Rep* 9(1):18392. <https://doi.org/10.1038/s41598-019-54605-0>
7. Lee SU, Kim HJ, Kim JS (2020) Bilateral vestibular dysfunction. *Semin Neurol* 40(1):40–48. <https://doi.org/10.1055/s-0039-3402066>
8. Schniepp R, Schlick C, Schenkel F, Pradhan C, Jahn K, Brandt T, Wuehr M (2017) Clinical and neurophysiological risk factors for falls in patients with bilateral vestibulopathy. *J Neurol* 264(2):277–283. <https://doi.org/10.1007/s00415-016-8342-6>
9. Swanenburg J, Zurbrugg A, Straumann D, Hegemann SCA, Palla A, de Bruin ED (2017) A pilot study investigating the association between chronic bilateral vestibulopathy and components of a clinical functional assessment tool. *Physiother Theory Pract* 33(6):454–461. <https://doi.org/10.1080/09593985.2017.1323362>
10. Alghwiri A, Alghadir A, Whitney SL (2013) The vestibular activities and participation measure and vestibular disorders. *J Vestib Res* 23(6):305–312. <https://doi.org/10.3233/ves-130474>
11. Molnár A, Maihoub S, Mavrogeni P, Tamás L, Szirmai Á (2022) Depression scores and quality of life of vertiginous patients, suffering from different vestibular disorders. *Eur Arch Otorhinolaryngol* 279(11):5173–5179. <https://doi.org/10.1007/s00405-022-07366-y>
12. Herdman SJ, Blatt P, Schubert MC, Tusa RJ (2000) Falls in patients with vestibular deficits. *Am J Otol* 21(6):847–851
13. MacDowell SG, Wellons R, Bissell A, Knecht L, Naquin C, Karpinski A (2018) The impact of symptoms of anxiety and depression on subjective and objective outcome measures in individuals with vestibular disorders. *J Vestib Res* 27(5–6):295–303. <https://doi.org/10.3233/ves-170627>
14. Schlick C, Schniepp R, Loidl V, Wuehr M, Hesselbarth K, Jahn K (2016) Falls and fear of falling in vertigo and balance disorders: A controlled cross-sectional study. *J Vestib Res* 25(5–6):241–251. <https://doi.org/10.3233/VES-150564>
15. Lucieer FMP, Van Hecke R, van Stiphout L, Duijn S, Perez-Fornos A, Guinand N, Van Rompaey V, Kingma H, Joore M, van de Berg R (2020) Bilateral vestibulopathy: beyond imbalance and oscillopsia. *J Neurol* 267(Suppl 1):241–255. <https://doi.org/10.1007/s00415-020-10243-5>
16. Starkov D, Strupp M, Pleshkov M, Kingma H, van de Berg R (2021) Diagnosing vestibular hypofunction: an update. *J Neurol* 268(1):377–385. <https://doi.org/10.1007/s00415-020-10139-4>
17. van de Berg R, Rosengren S, Kingma H (2018) Laboratory examinations for the vestibular system. *Curr Opin Neurol* 31(1):111–116. <https://doi.org/10.1097/wco.0000000000000526>
18. Schniepp R, Huppert A, Decker J, Schenkel F, Schlick C, Rasoul A, Dieterich M, Brandt T, Jahn K, Wuehr M (2021) Fall prediction in neurological gait disorders: differential contributions from clinical assessment, gait analysis, and daily-life mobility monitoring. *J Neurol*. <https://doi.org/10.1007/s00415-021-10504-x>
19. Dobbels B, Lucieer F, Mertens G, Gilles A, Moyaert J, van de Heyning P, Guinand N, Perez Fornos A, Herssens N, Halle-mans A, Vereeck L, Vanderveken O, Van Rompaey V, van de Berg R (2020) Prospective cohort study on the predictors of fall risk in 119 patients with bilateral vestibulopathy. *PLoS ONE* 15(3):e0228768. <https://doi.org/10.1371/journal.pone.0228768>
20. Herssens N, Verbecque E, McCrum C, Meijer K, van de Berg R, Saeys W, Vereeck L, Van Rompaey V, Halle-mans A (2020) A systematic review on balance performance in patients with bilateral vestibulopathy. *Phys Ther* 100(9):1582–1594. <https://doi.org/10.1093/ptj/pzaa083>
21. Horak FB (2006) Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls? *Age Ageing* 35(Suppl 2):ii7–ii11. <https://doi.org/10.1093/ageing/af077>
22. Horak FB, Wrisley DM, Frank J (2009) The Balance Evaluation Systems Test (BESTest) to differentiate balance deficits. *Phys Ther* 89(5):484–498. <https://doi.org/10.2522/ptj.20080071>
23. Franchignoni F, Horak F, Godi M, Nardone A, Giordano A (2010) Using psychometric techniques to improve the Balance Evaluation Systems Test: the mini-BESTest. *J Rehabil Med* 42(4):323–331. <https://doi.org/10.2340/16501977-0537>
24. Lamb SE, Jorstad-Stein EC, Hauer K, Becker C (2005) Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 53(9):1618–1622. <https://doi.org/10.1111/j.1532-5415.2005.53455.x>
25. Lord SR, Sherrington C, Menz HB, Close JCT (2011) Falls in older people: risk factors and strategies for prevention, 2nd edn. Cambridge University Press, New York
26. McCrum C (2020) Falls history questionnaire material in English, German and Dutch. <https://osf.io/hmjef/>. Accessed 1 June 2021
27. Akizuki K, Echizenya Y, Kaneno T, Yabuki J, Ohashi Y (2018) Dynamic balance assessment using an unstable board in community-dwelling elderly people. *J Phys Ther Sci* 30(8):1086–1091. <https://doi.org/10.1589/jpts.30.1086>
28. Santos DA, Duarte M (2016) A public data set of human balance evaluations. *PeerJ* 4:e2648. <https://doi.org/10.7717/peerj.2648>
29. Dominguez-Olivan P, Gasch-Gallen A, Aguas-Garcia E, Bengoetxea A (2020) Validity and reliability testing of the Spanish version of the BESTest and mini-BESTest in healthy community-dwelling elderly. *BMC Geriatr* 20(1):444. <https://doi.org/10.1186/s12877-020-01724-3>
30. Nakhostin-Ansari A, Nakhostin Ansari N, Mellat-Ardakani M, Nematizad M, Naghdi S, Babaki M, Farhangian M, Habibi AH, Tafakhori A, Hasson S (2020) Reliability and validity of Persian versions of Mini-BESTest and Brief-BESTest in persons with Parkinson's disease. *Physiother Theory Pract*. <https://doi.org/10.1080/09593985.2020.1822967>
31. Naghdi S, Nakhostin Ansari N, Forogh B, Khalifelloo M, Honar-pisheh R, Nakhostin-Ansari A (2020) Reliability and validity of the Persian version of the mini-balance evaluation systems test in patients with stroke. *Neurol Ther* 9(2):567–574. <https://doi.org/10.1007/s40120-020-00207-2>
32. Kim A, Darakjian N, Finley JM (2017) Walking in fully immersive virtual environments: an evaluation of potential adverse effects in older adults and individuals with Parkinson's disease. *J Neuroeng Rehabil* 14(1):16. <https://doi.org/10.1186/s12984-017-0225-2>
33. Giardini M, Arcolin I, Guglielmetti S, Godi M, Capelli A, Corna S (2021) Balance performance in patients with post-acute COVID-19 compared to patients with an acute exacerbation of chronic obstructive pulmonary disease and healthy subjects. *Int J Rehabil Res*. <https://doi.org/10.1097/mrr.0000000000000510>
34. Harro CC, Garascia C (2019) Reliability and validity of computerized force platform measures of balance function in healthy older adults. *J Geriatr Phys Ther* 42(3):E57–e66. <https://doi.org/10.1519/jpt.0000000000000175>
35. Ohtsu H, Yoshida S, Minamisawa T, Katagiri N, Yamaguchi T, Takahashi T, Yomogida SI, Kanzaki H (2020) Does the balance

- strategy during walking in elderly persons show an association with fall risk assessment? *J Biomech* 103:109657. <https://doi.org/10.1016/j.jbiomech.2020.109657>
36. McIntosh EI, Zettl JL, Vallis LA (2017) Stepping responses in young and older adults following a perturbation to the support surface during gait. *J Mot Behav* 49(3):288–298. <https://doi.org/10.1080/00222895.2016.1204262>
 37. Morishita S, Hirabayashi R, Tsubaki A, Aoki O, Fu JB, Onishi H, Tsuji T (2019) Assessment of the Mini-Balance Evaluation Systems Test, Timed Up and Go test, and body sway test between cancer survivors and healthy participants. *Clin Biomech* 69:28–33. <https://doi.org/10.1016/j.clinbiomech.2019.07.002>
 38. O'Hoski S, Winship B, Herridge L, Agha T, Brooks D, Beauchamp MK, Sibley KM (2014) Increasing the clinical utility of the BESTest, mini-BESTest, and brief-BESTest: normative values in Canadian adults who are healthy and aged 50 years or older. *Phys Ther* 94(3):334–342. <https://doi.org/10.2522/ptj.20130104>
 39. Vervoort G, Bengevoord A, Strouwen C, Bekkers EM, Heremans E, Vandenberghe W, Nieuwboer A (2016) Progression of postural control and gait deficits in Parkinson's disease and freezing of gait: a longitudinal study. *Parkinson Relat Disord* 28:73–79. <https://doi.org/10.1016/j.parkreldis.2016.04.029>
 40. Shanahan J, Coman L, Ryan F, Saunders J, O'Sullivan K, Ni Bhriain O, Clifford AM (2016) To dance or not to dance? A comparison of balance, physical fitness and quality of life in older Irish set dancers and age-matched controls. *Public Health* 141:56–62. <https://doi.org/10.1016/j.puhe.2016.07.015>
 41. Bonora G, Mancini M, Carpinella I, Chiari L, Ferrarin M, Nutt JG, Horak FB (2017) Investigation of anticipatory postural adjustments during one-leg stance using inertial sensors: evidence from subjects with Parkinsonism. *Front Neurol* 8:361. <https://doi.org/10.3389/fneur.2017.00361>
 42. Yingyongyudha A, Saengsirisuwan V, Panichaporn W, Boonsinsukh R (2016) The Mini-Balance Evaluation Systems Test (Mini-BESTest) demonstrates higher accuracy in identifying older adult participants with history of falls than do the BESTest, Berg Balance Scale, or Timed Up and Go Test. *J Geriatr Phys Ther* 39(2):64–70. <https://doi.org/10.1519/jpt.0000000000000050>
 43. Löfgren N, Benka Wallén M, Sorjonen K, Conradsson D, Franzén E (2017) Investigating the Mini-BESTest's construct validity in elderly with Parkinson's disease. *Acta Neurol Scand* 135(6):614–621. <https://doi.org/10.1111/ane.12640>
 44. Kelley KK, Aaron D, Hynds K, Machado E, Wolff M (2014) The effects of a therapeutic yoga program on postural control, mobility, and gait speed in community-dwelling older adults. *J Altern Complement Med* 20(12):949–954. <https://doi.org/10.1089/acm.2014.0156>
 45. Bao T, Carender WJ, Kinnaird C, Barone VJ, Peethambaran G, Whitney SL, Kabeto M, Seidler RD, Sienko KH (2018) Effects of long-term balance training with vibrotactile sensory augmentation among community-dwelling healthy older adults: a randomized preliminary study. *J Neuroeng Rehabil* 15(1):5. <https://doi.org/10.1186/s12984-017-0339-6>
 46. Faul F, Erdfelder E, Lang AG, Buchner A (2007) G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 39(2):175–191. <https://doi.org/10.3758/bf03193146>
 47. The jamovi project. jamovi (Version 2.3) [Computer Software] (2022). <https://www.jamovi.org>
 48. Fritz CO, Morris PE, Richler JJ (2012) Effect size estimates: current use, calculations, and interpretation. *J Exp Psychol Gen* 141:2–18. <https://doi.org/10.1037/a0024338>
 49. Godi M, Franchignoni F, Caligari M, Giordano A, Turcato AM, Nardone A (2013) Comparison of reliability, validity, and responsiveness of the mini-BESTest and Berg Balance Scale in patients with balance disorders. *Phys Ther* 93(2):158–167. <https://doi.org/10.2522/ptj.20120171>
 50. King LA, Priest KC, Salarian A, Pierce D, Horak FB (2012) Comparing the Mini-BESTest with the Berg Balance Scale to evaluate balance disorders in Parkinson's Disease. *Parkinsons Dis* 2012:375419. <https://doi.org/10.1155/2012/375419>
 51. Tsang CS, Liao LR, Chung RC, Pang MY (2013) Psychometric properties of the Mini-Balance Evaluation Systems Test (Mini-BESTest) in community-dwelling individuals with chronic stroke. *Phys Ther* 93(8):1102–1115. <https://doi.org/10.2522/ptj.20120454>
 52. Leddy AL, Crouner BE, Earhart GM (2011) Utility of the Mini-BESTest, BESTest, and BESTest sections for balance assessments in individuals with Parkinson disease. *J Neurol Phys Ther* 35(2):90–97. <https://doi.org/10.1097/NPT.0b013e31821a620c>
 53. Oyama C, Otaka Y, Onitsuka K, Takagi H, Tan E, Otaka E (2018) Reliability and validity of the Japanese version of the Mini-balance evaluation systems test in patients with subacute stroke. *Prog Rehabil Med* 3:20180015. <https://doi.org/10.2490/prm.20180015>
 54. Schlenstedt C, Brombacher S, Hartwigsen G, Weisser B, Möller B, Deuschl G (2016) Comparison of the Fullerton advanced balance scale, mini-BESTest, and Berg Balance Scale to predict falls in Parkinson disease. *Phys Ther* 96(4):494–501. <https://doi.org/10.2522/ptj.20150249>
 55. Whitney S, Morris LO (2006) Multisensory impairment in older adults: evaluation and intervention. In: Calhoun KH, Eibling DE, Wax MK, Kost KM (eds) *Geriatric otolaryngology*. Taylor & Francis, New York, pp 109–123
 56. Agrawal Y, Van de Berg R, Wuyts F, Walther L, Magnusson M, Oh E, Sharpe M, Strupp M (2019) Presbyvestibulopathy: diagnostic criteria. *J Vestib Res* 29:161–170. <https://doi.org/10.3233/ves-190672>
 57. Paraskevoudi N, Balci F, Vataki A (2018) "Walking" through the sensory, cognitive, and temporal degradations of healthy aging. *Ann N Y Acad Sci*. <https://doi.org/10.1111/nyas.13734>
 58. Eysel-Gosepath K, McCrum C, Epro G, Bruggemann GP, Karanidis K (2016) Visual and proprioceptive contributions to postural control of upright stance in unilateral vestibulopathy. *Somatosen Mot Res* 33(2):72–78. <https://doi.org/10.1080/08990220.2016.1178635>
 59. Baloh RW, Jacobson KM, Beykirch K, Honrubia V (1998) Static and dynamic posturography in patients with vestibular and cerebellar lesions. *Arch Neurol* 55(5):649–654. <https://doi.org/10.1001/archneur.55.5.649>
 60. Herdman SJ, Sandusky AL, Hain TC, Zee DS, Tusa RJ (1994) Characteristics of postural stability in patients with aminoglycoside toxicity. *J Vestib Res* 4(1):71–80
 61. Dozza M, Wall C 3rd, Peterka RJ, Chiari L, Horak FB (2007) Effects of practicing tandem gait with and without vibrotactile biofeedback in subjects with unilateral vestibular loss. *J Vestib Res* 17(4):195–204
 62. Kingma H, Felipe L, Gerards MC, Gerits P, Guinand N, Perez-Fornos A, Demkin V, van de Berg R (2019) Vibrotactile feedback improves balance and mobility in patients with severe bilateral vestibular loss. *J Neurol* 266(Suppl 1):19–26. <https://doi.org/10.1007/s00415-018-9133-z>
 63. Godi M, Arcolin I, Leavy B, Giardini M, Corna S, Franzén E (2021) Insights into the Mini-BESTest Scoring System: comparison of 6 different structural models. *Phys Ther*. <https://doi.org/10.1093/ptj/pzab180>
 64. Zhu M, van Stiphout L, Karabulut M, Fornos AP, Guinand N, Meijer K, van de Berg R, McCrum C (2023) Assessing balance in people with bilateral vestibulopathy using the mini-balance evaluation systems test (Mini-BESTest). *OSF Preprints*. <https://doi.org/10.31219/osf.io/nwa2y>