



# EEG Signature of Out-of-Body Experiences Induced by Virtual Reality: A Novel Methodological Approach

Charlotte Martial<sup>1,2</sup>, H el ena Cassol<sup>1</sup>, Mel Slater<sup>3,4</sup>, Pierre Bourdin<sup>3,4</sup>, Armand Mensen<sup>5</sup>, Oliva Ramon<sup>3,4</sup>, Steven Laureys<sup>1,2,6\*</sup>, and Pablo N u nez<sup>1\*</sup>

## Abstract

■ Out-of-body experiences (OBEs) are subjective experiences of seeing one's own body and the environment from a location outside the physical body. They can arise spontaneously or in specific conditions, such as during the intake of dissociative drug. Given its unpredictable occurrence, one way to empirically study it is to induce subjective experiences resembling an OBE using technology such as virtual reality. We employed a complex multisensory method of virtual embodiment in a virtual reality scenario with 7 healthy participants to induce virtual OBE-like experiences. Participants performed 2 conditions in a randomly determined order. For both conditions, the participant's viewpoint was lifted out of the virtual body toward the ceiling of the virtual room, and real body movements were (visuo-tactile ON

condition) or were not (visuo-tactile OFF condition) translated into movements on the virtual body below—the latter aiming to maintain a feeling of connection with the virtual body. A continuous 128-electrode EEG was recorded. Participants reported subjective experiences of floating in the air and of feeling high up in the virtual room at a strong intensity, but a weak to moderate feeling of being “out of their body” in both conditions. The EEG analysis revealed that this subjective experience was associated with a power shift that manifested in an increase of delta and a decrease of alpha relative power. A reduction of theta complexity and an increase of beta-2 connectivity were also found. This supports the growing body of evidence revealing a prominent role of delta activity during particular conscious states. ■

## INTRODUCTION

It is now known that humans can experience a disturbed sense of self in varying situations and conditions, sometimes amounting to an out-of-body experience (OBE), which refers to the subjective experience of seeing one's body and the environment from a location outside the physical body (Blanke, Landis, Spinelli, & Seeck, 2004). OBEs are often reported in near-death experiences (NDEs; Martial, Cassol, Laureys, & Gosseries, 2020; Martial, Simon, et al., 2020), but also in many other types of contexts, such as sensory deprivation (Blackmore, 1982), drug intake (Timmermann et al., 2018), and sleep paralysis (Herrero, Gallo, Gasca-Rol n, Gleiser, & Forcato, 2023). Their incidence is estimated around 10–25% in the normal population (Alvarado, 2000; Irwin, 1985; Blackmore, 1982). Considering the unpredictable occurrence of OBEs, induction methods may be used to empirically study the phenomenon in laboratory settings. Recently, several researchers have successfully induced subjective experiences resembling OBE using non-invasive techniques such as hypnosis (e.g., Facco et al., 2019; Martial

et al., 2019; Palmieri et al., 2014) or virtual reality (VR; e.g., van Heugten-van der Kloet, Cosgrave, van Rheede, & Hicks, 2018; Bourdin, Barberia, Oliva, & Slater, 2017; Ehrsson, 2007). A few researchers also had the opportunity to study OBE artificially induced by direct electrical cortical stimulation in epileptic patients (e.g., Blanke, Ortigue, Landis, & Seeck, 2002). Current theories assume that OBEs may be a distorted body representation elicited by illusory perceptions arising because of a failure to integrate complex somatosensory and vestibular information (Blanke et al., 2002). In parallel, some psychological theories have also been suggested, such as the one proposing that the OBE is triggered by changes in perception of the physical body boundaries and dissociative processes associated with disruption of somatic inputs (Irwin, 2000; Blackmore, 1984).

Considering that a recurrent feature of the OBE is a convincing impression of reality with all the qualities of three-dimensional perception (Blanke & Mohr, 2005; Blanke et al., 2004; Blackmore, 1987, 1982), VR is a good candidate for studying OBEs in laboratory settings. The particularity and originality of VR is that this method allows the deliberate creation of rich visual experiences that may not match bodily signals and perceptions, which turns out to be essential in the study of the phenomenon of OBE (Tseng & Juan, 2013). Using VR may thus allow to disentangle various components of self-consciousness such

<sup>1</sup>University of Li ge, Belgium, <sup>2</sup>University Hospital of Li ge, Belgium, <sup>3</sup>University of Barcelona, Spain, <sup>4</sup>Institute of Neurosciences of the University of Barcelona, Spain, <sup>5</sup>University of Bern, Switzerland, <sup>6</sup>University Laval, Qu bec, Canada

\*Equally contributed.

as bodily self-location and the sense of ownership and help for the development of a neurophysiological model of OBE and, more broadly, of the sense of selfhood.

In laboratory settings, the VR experience can be accompanied by neurophysiological techniques such as EEG, a technique that enables the detection of fast synchronized firing of cortical neurons (Núñez et al., 2019). By recording the EEG activity during the experiment, the dynamics of the ongoing oscillations can be detected and characterized by means of a variety of spectral and nonlinear methods (O'Neill et al., 2018; Ruiz-Gómez et al., 2018), as well as measures of statistical interdependence between brain regions (functional connectivity; Friston, 2011). Spectral methods are based on representations of the power spectrum of a signal and enable the detection of shifts in the EEG power distribution over the different conditions in the experiment. The continuous wavelet transform (CWT) is a time–frequency representation of a time series that is well-suited to the analysis of nonstationary signals such as EEG compared with traditional Fourier analysis (Rioul & Vetterli, 1991). In particular, the relative power (RP) derived from the CWT has been shown to be affected by disorders of consciousness, where higher power in the theta, alpha, and beta-1 bands has been linked to higher levels of consciousness (Piarulli et al., 2016). The Lempel-Ziv complexity (LZC) is a nonlinear measure that reflects the compressibility of a signal and can be used to measure the local diversity of the EEG over time and has been shown to decrease during general anesthesia (Schartner et al., 2015). Thus, EEG-derived measures may provide useful tools to characterize altered states of consciousness such as OBE.

So far, no study has ever used a 128-system EEG to measure changes in brain activity associated to an OBE experimentally induced by VR. In the present preliminary study, we used a protocol adapted from Bourdin et al. (2017) to induce OBEs using a method of virtual embodiment in a virtual environment in healthy participants. We wanted to induce a “full body ownership illusion” (Kilteni, Groten, & Slater, 2012) through a multisensory virtual environment, including a wide field-of-view head-tracked stereo head-mounted display, real-time motion capture, a virtual reflection of their body in a virtual mirror, a virtual shadow, and synchronous vibrotactile stimulation on the person’s real body. These experimental conditions may permit to induce a dissociative state with a drift of the subjective experience of the body to a position outside one’s usual bodily borders. More specifically, participants performed two distinct experimental conditions in a randomly determined order. For both conditions, the participant’s viewpoint was lifted out of the virtual body toward the ceiling of the virtual room, and real body movements were (visuo-tactile ON [VT] condition) or were not (visuo-tactile OFF [noVT] condition) translated into movements on the virtual body below. The latter condition aimed to maintain a feeling of connection with the virtual body, as shown in Bourdin et al. (2017). In their study, Bourdin et al.

(2017) found that there was a shift in self-location up and out of the body in the noVT condition but less in the VT one. We expect to replicate these results, as well as to study associated electrical brain activity using a 128-system EEG. We investigated the spectral changes of the EEG during this potential subjective experience drift by means of CWT-based RP and complexity using the LZC, and we also applied the phase-locking value (PLV) to quantify the phase synchrony between electrodes.

## METHODS

### Participants


Seven healthy participants (three women; mean age =  $34 \pm 7$  years) participated in the study. They were recruited via word of mouth and announcements at the Faculty of Psychology of the University of Barcelona. Participants had to be aged at least 18 years and not suffer from epilepsy. Participants were compensated for participating in the study. Because the present study was intended to be a proof-of-concept, we did not perform a power analysis.

The present study was approved by the ethical committee of the Faculty of Psychology of the University of Barcelona in accordance with the tenets of the Declaration of Helsinki and its later amendments, and all procedures were followed.

### VR Apparatus

The virtual environment was the one used in Bourdin et al. (2017). The environment was designed and rendered using Unity3D (Unity Technologies). It consisted of a virtual room with some virtual standard furniture (see Figure 1; please also see <https://www.youtube.com/watch?v=u2Dc6BKqIP0>). This was displayed in an Oculus head-mounted-display (Oculus VR) at 75 Hz using a 1080-px OLED panel, split to 75 Hz 960 × 1080 per eye rendering. We used an Intersense Wand joystick connected to an Intersense IS-9006 hybrid inertial/acoustic 6-DOF tracking system (Intersense) with data streamed over a VRPN network (Taylor et al., 2001) to the computer running the application. The virtual body used was a 3-D model of a female or male character available from Rocketbox Studios according to the participant’s gender. Participant movements were tracked by a Natural Point’s Optitrack infrared system (NaturalPoint, Inc.), including 12 cameras, tracking three triplets of reflective markers on each foot, wrist, and forearm. We used the Natural Point’s Tracking Tools software (NaturalPoint, Inc.) with the VRPN protocol (Taylor et al., 2001) to calculate in real time the positions and orientations of each marker and to stream to the VE. Tactile stimulations were administered using four vibrotactile devices (vibrated for 100 msec at a frequency of 150 Hz) on both wrists and ankles of the

**Figure 1.** The virtual environment and setup. (A) One of the participants wearing the equipment, including the head-mounted display and the EEG. (B) The participant can see their virtual body from a first person perspective, including the wand. (C) Part of the virtual environment from a first person perspective, including the mirror and the virtual balls. Images (B) and (C) were taken from Bourdin, Barberia, Oliva, and Slater (2017).



individual and controlled by an Arduino (Arduino SRL.) board. See Bourdin et al. (2017) for more details.

### Procedure

After being instructed about the experimental procedure, participants were invited to sign the informed consent. They were asked to what extent they had experience with VR (using a Likert scale as follows: 1 = “novice” to 7 = “expert”). They were then invited to wear a black tracking suit and black socks. For each participant, we scaled the virtual body based on the measures of their limbs. Participants were seated in a chair with their arms on their thighs and their legs lying stretched on a low table. We attached vibrators to their wrists and ankles, and several tracking markers were collocated on their limbs. We set up the EEG cap (gHamp recorded 128 active EEG channels). Afterward, a resting-state recording was performed for 4 min with eyes open. We then set up the head-mounted display (on top of the EEG cap). During the whole VR experience, we tried to ensure as much as possible that the head-mounted display did not disturb the EEG signal. Once the full setup was completed,


instructions were given thanks to a recorded voice. The experimental session started with the instruction of orally describing the virtual environment, permitting the participant to explore and become familiar with the virtual environment. When looking down at themselves in the virtual environment, participants could see a virtual body (with black clothes) from a first-person perspective in the same posture as their real body, moving congruently and synchronously with their real movements. They could also see this virtual body in a virtual mirror placed in front (see Figure 1). This was followed by the three trials of a Mental Ball-Dropping task in which participants were invited to estimate the time (by clicking a button of the wand) an imaginary apple would take to fall from their hand to the floor (Lenggenhager, Mouthon, & Blanke, 2009). This was used as an estimate of the sense of self location. After the three Mental Ball-Dropping task trials, two tasks were performed to induce body ownership through synchronous visuomotor and visuotactile stimulation. First, the voice asked them to look toward their legs and follow with their feet the path that virtually appeared on the low table, as described in Kokkinara and Slater (2014). Second, participants felt virtual balls

hitting the virtual body through corresponding vibrotactile stimulations. After all these familiarization tasks, participants randomly performed two conditions: the noVT condition and the VT condition (see Figure 2). Each condition was divided into two phases: an initial embodiment phase (*in-the-body* phase) that was the same for both conditions, and the out-of-body phase (*out-the-body* phase), which was different for the two conditions. In the latter, for both conditions, the viewpoint of the participant was lifted out of the virtual body toward the ceiling of the virtual room, and just behind the body, so that the body could be seen below. The noVT condition consisted of the balls going up with the invisible body (i.e., the elevated viewpoint) and striking the space around the participant's visual center of perception. In this condition, real body movements were not translated into movements on the virtual body below; the virtual body below remained stationary. By contrast, the VT condition consisted of the balls striking the virtual body that participants could see below but they still felt the vibrations. In this condition, real body movements were translated into movements on the virtual body below. After the *in-the-body* phase of each of the two conditions, participants were asked

to what extent they felt as if the body they were seeing was their own body (using a Likert scale as followed: 1 = "I did not feel that at all" to 7 = "I felt it in its maximum intensity") and to what extent they felt as if the body they were seeing belonged to someone else (using a Likert scale as followed: 1 = "I did not feel that at all" to 7 = "I felt it in its maximum intensity"). After the *out-the-body* phase of the two conditions, participants were asked these two same questions, and four additional questions (see Table 1 for details). The *obe* and *otherbodyobe* questions of this set of questions (see Table 1 for details) were used to specifically assess potential subjective experience resembling OBE. The experimental session ended with a black screen. In total, the experiment lasted between 60 and 90 min, including about 40 min in the VR.

After removing trackers and vibrators with the minimum interaction possible, participants were invited to answer two Visual Analog Scales asking them to what extent they felt absorbed in the tasks (ranging from 0 to 10) and to what extent they felt dissociated in the tasks (ranging from 0 to 10; subjective ratings from Vanhaudenhuyse et al., 2019). Finally, they were debriefed about the experiment.

**Figure 2.** The OBE and VTO conditions. (A) The VT condition in which the balls hit the virtual body that participants could see below but they still felt the vibrations. (B) The noVT condition in which the balls went up with the invisible body (i.e., the elevated viewpoint) and hit the space around the participant's visual center of perception. Images were taken from Bourdin et al. (2017).



**Table 1.** Participants' Responses to the Body Ownership and Out-of-body Questions for Both Conditions (noVT and VT)

Variable Name	Questions on A Likert Scale (1 = "I Did Not Feel That at All" to 7 = "I Felt It to the Maximum Intensity")	noVT	VT	z Statistic	p	Effect Size	Bourdin et al. (2017)'s	Bourdin et al. (2017)'s
							Median Scores	Median Scores
							noVT	VT
Immediately after the <i>in-the-body</i> phase: Please indicate to what extent you felt each of the sensations that I will indicate now.								
<i>mybody</i>	I felt as if the body I was seeing was my own body.	5 (2–6)	4 (3–6)	4	.850	–0.200	6 (5–6)	6 (5–6)
<i>otherbody</i>	I felt as if the body I was seeing belonged to someone else.	2 (1–4)	1 (1–3)	5.5	1	0.100	3 (1.5–5)	2.5 (1.5–3.5)
Immediately after the <i>out-of-body</i> phase: Please indicate to what extent you felt each of the sensations that I will indicate now. When you answer these questions, please refer to your experience when you were watching the room from above.								
<i>mybodyobe</i>	I felt as if the body I was seeing was my own body.	3 (1–3)	4 (2–6)	5	.281	–0.524	2 (2–5.5)	4 (2–5)
<i>otherbodyobe</i>	I felt as if the body I was seeing belonged to someone else.	4 (3–5)	2 (1–3)	9	.197	0.800	6 (3.5–7)	4.5 (2–6)
<i>floatingobe</i>	I felt as if I was floating in air.	7 (6–7)	5 (4–6)	10	.098	1	7 (7–7)	6 (6–7)
<i>elevatedobe</i>	I felt as if I was in an elevated position in the room.	7 (6–7)	6 (5–7)	4.5	.586	0.500	7 (6–7)	7 (7–7)
<i>connectionobe</i>	I felt a connection with the body, as if I was looking down at myself.	1 (1–2)	5 (3–6)	0	.054	–1	2.5 (2–6)	5 (4–7)
<i>obe</i>	I felt out of my body.	2 (1–4)	3 (2–5)	2	.170	–0.733	6 (4.5–7)	6 (4–7)

The latest two columns report median scores reported by the participants from Bourdin et al. (2017).

## EEG Analyses

### Preprocessing

The EEG recordings were first segmented into three sequences corresponding to the resting state (4 min), noVT (32 sec), and VT (32 sec) conditions. Each segment was filtered between 1 and 70 Hz by means of zero-phase finite impulse response filter, as well as with a 50-Hz finite impulse response notch filter to remove powerline artifacts. Both filters were designed by means of a Hamming window, with a filter order of 2000. Afterward, an independent component analysis (ICA) was carried out on each of the sequences to discard ICA components with clear artifacts, such as heartbeats and eye blinking, after visual inspection. Subsequently, each sequence was segmented into 1-sec epochs and the epochs in which artifacts remained after the ICA were discarded. One-second epochs were used as a tradeoff between epoch length and the total number of discarded segments. All subsequent analyses were conducted on five frequency bands: delta ( $\delta$ , 1–4 Hz), theta ( $\theta$ , 4–8 Hz), alpha ( $\alpha$ , 8–13 Hz), beta-1 ( $\beta_1$ , 13–19 Hz), and beta-2 ( $\beta_2$ , 19–30 Hz). The gamma ( $\gamma$ , 30–70 Hz) band was not included because of the presence of high-frequency muscle activity present in parts of the recordings, which were unavoidable because of the length of the experiment and the nature of VR headsets. Nonetheless, the EEGs were rigorously visually inspected after filtering to ensure these artifacts were successfully removed.

### CWT

The CWT is a time–frequency representation of a time series that provides a good compromise between the time and the frequency resolutions compared with the short-term Fourier transform (Tallon-Baudry, Bertrand, Delpuech, & Pernier, 1996). The CWT provides control over the frequency and time resolution in which neuro-electric components can be detected (Samar, Bopardikar, Rao, & Swartz, 1999). To adequately model the EEG signal, a biologically plausible fit was provided with the Morlet wavelet acting as the “mother wavelet” (Núñez et al., 2017; Roach & Mathalon, 2008). Two of its parameters (center frequency and bandwidth) need to be adjusted and were set to 1 to obtain a good balance between the time ( $\Delta t$ ) and frequency resolutions ( $\Delta f$ ) at low frequencies (Núñez et al., 2017; Bachiller et al., 2015). In addition, the Heisenberg box was set with a width of 2 times  $\Delta t$  and  $\Delta f$  as a tradeoff between frequency and time resolutions (Núñez et al., 2017; Bachiller et al., 2015). Finally, the cone of influence was taken into account to overcome the issues of edge effects introduced by discontinuities in the epochs (Torrence & Compo, 1998).

### RP

To analyze the spectral content of each sequence, the RP was computed from the CWT. The RP characterizes the

contribution to the power spectrum of each frequency band and is computed as the sum of the contribution of each spectral component in each band to the total power (Núñez et al., 2017). The RP was computed as follows:

$$PSD(f) = \sum_{n=1}^{n=N} CWT_n(f, n\Delta t), \quad (1)$$

$$RP_{band} = \sum_{f=f_{b1}}^{f_{b2}} PSD(f), \quad (2)$$

where  $CWT_n$  is the normalized CWT scalogram (absolute value of the CWT squared) to the sum of the absolute values in the 1- to 30-Hz range to ensure the RP was a relative measure,  $n$  is the number of temporal samples in the CWT of a single epoch, and  $f_{b1}$  and  $f_{b2}$  represent the lower and upper frequency limits of each band.

### LZC

The LZC is a measure of complexity and uncertainty, which is related to the number of distinct patterns in a symbolic sequence and the rate in which they appear, with higher LZC values indicating a higher level of complexity in the data (Schartner et al., 2015; Abásolo, Hornero, Gómez, García, & López, 2006). In EEG data, the grand-average LZC over channels can be considered a measure of differentiation, as it measures the diversity of patterns across time and space (Schartner et al., 2015). The LZC algorithm can only be computed on strings composed of a limited number of symbols; therefore, an EEG time series needs to be transformed into a symbolic sequence before its computation. To achieve this, a coarse graining of the EEG is performed by means of a thresholding procedure as follows (Abásolo et al., 2006):

$$s(t) = \begin{cases} 0 & \text{if } x(t) < T \\ 1 & \text{if } x(t) > T \end{cases} \quad (3)$$

where the threshold  $T$  is the median of the time series  $x(t)$ . After conversion into a binary sequence, the LZC algorithm is applied. A full description of the LZC implementation used in this study can be found in Abásolo et al. (2006). The LZC was normalized by dividing it with the upper bound of the complexity, which has been proven to be  $n/\log_\alpha(n)$ , where  $\alpha$  is the number of different symbols in the set and  $n$  is the length of the sequence in samples (Lempel & Ziv, 1976) and was applied across individual frequency bands (Kim, Lee, & Vlisides, 2022; Tanabe et al., 2022).

### Connectivity Estimation

In the present study, we chose a phase-based connectivity measure, the PLV (Lachaux, Rodriguez, Martinerie, & Varela, 1999), which quantifies the phase synchrony between two signals in a specific frequency band by means

of estimates of the instantaneous frequency (Colclough et al., 2016). The PLV looks for latencies in which the phase difference between the two signals are stable across time, which is known as “phase locking.” For resting-state recordings, the PLV is computed over the epochs in which the recording is divided (in the present study, 1-sec epochs were used, following the artifact rejection step of the preprocessing). The PLV can be computed as follows (Bruña, Maestú, & Pereda, 2018):

$$PLV_{X,Y} = \frac{1}{T} \left| \sum_{t=1}^T e^{-i(\phi_X(t) - \phi_Y(t))} \right|,$$

where  $\phi$  is the instantaneous phase of the signals  $X$  and  $Y$  and  $T$  is the length of each epoch in samples.

To eliminate spurious correlations because of volume conduction and field spread, the time series were orthogonalized before the computation of the connectivity. For each epoch, each channel was orthogonalized with respect to a linear projection of the other by means of a pairwise linear regression, and the results of the PLV after both regressions were averaged (O’Neill et al., 2018; Brookes, Woolrich, & Barnes, 2012).

### Statistical Analyses

Demographic data and scores are expressed as mean and standard deviation ( $SD$ ) or as medians with interquartile range (Q1–Q3) for asymmetric distribution. We compared the scores obtained after the VT and noVT condition using paired-samples Wilcoxon signed-ranks tests ( $p < .05$ ).

Regarding the EEG data, an exploratory analysis was performed to check the distribution of the RP and PLV values from all three conditions by means of the Shapiro–Wilk test (normality) and Levene test (homoscedasticity). Both conditions were not met by the data, and therefore non-parametric tests were conducted to evaluate the statistical

differences between the conditions. Friedman tests were performed to determine interactions across the three conditions (resting state, VT, and noVT) on the five frequency bands under study. If statistically significant differences were found, pairwise Wilcoxon signed-ranks tests were performed to evaluate differences between specific conditions. A false discovery rate (FDR) correction was completed to control for the number of comparisons with a significance level set to  $\alpha = .05$  (Benjamini & Hochberg, 1995). Both the statistical analyses and the signal processing were achieved using MATLAB® software (Version R2019a, Mathworks).


## RESULTS

### Participants

Participants reported having a little to no level of prior experience with VR (median = 2.28; interquartile range [IQR] = 1–2.5).

Regarding the *in-the-body* phase, participants reported higher responses to *mybody* question than to *otherbody* question for both conditions, suggesting an overall body ownership for both conditions (see Table 1 for details).

Regarding the *out-the-body* phase of the noVT condition, participants reported particularly high responses to *floatingobe* and *elevatedobe* questions. In the noVT condition, they tended to disown the virtual body (*mybodyobe* and *connectionobe* question), whereas in the VT condition, participants tended to report connection with the virtual body that they saw below (*connectionobe* question). For both conditions, participants reported they had felt “out of their body” at a moderate intensity (*obe* question). No significant differences between the scores were observed between the two conditions, although the *connectionobe* question was closed to the threshold value (see Table 1 for details).




**Figure 3.** Distribution plots depicting the RP values averaged over all electrodes for all three conditions under study. Statistically significant between-conditions differences are indicated with blue rectangles ( $p < .05$ , Friedman test, FDR corrected for multiple comparisons), whereas statistically significant post hoc differences between pairs of conditions in bands that showed significant group interactions are indicated with red brackets ( $p < .05$ , Wilcoxon signed-ranks test).

Participants reported a high score of absorption (median = 7.34; IQR = 7.7–8.65) and a moderate score of dissociation (median = 5.58; IQR = 5–7.05).

### EEG Results


The RP values averaged over all electrodes are displayed in Figure 3. Statistically significant differences were found in the delta ( $p = .018$ ,  $\chi^2(2) = 8$ , Friedman test;  $p = .030$  after FDR correction), alpha ( $p = .005$ ,  $\chi^2(2) = 10.57$ , Friedman test;  $p = .025$  after FDR correction), and beta-1 bands ( $p = .018$ ,  $\chi^2(2) = 8$ , Friedman test;  $p = .030$  after FDR correction). In the delta band, the statistical tests showed significant differences between the resting state and the noVT condition ( $p = .015$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .046$  after FDR correction). In the alpha band, the statistical tests revealed significant

differences between the resting state and both the VT ( $p = .015$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .023$  after FDR correction) and noVT conditions ( $p = .015$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .023$  after FDR correction). In the beta-1 band, significant differences between the resting state and noVT condition ( $p = .015$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .046$  after FDR correction) were found. Figure 4 shows topographic maps of the RP values and the statistical differences between conditions ( $p < 0.05$ , Wilcoxon signed-ranks test). In the delta band, the noVT condition displayed a significant increase in power in the fronto-central area compared with the resting-state condition that was more localized to the central region in the VT condition. In the theta band, there were, for the most part, no clear differences between conditions, except for some very localized areas in the central regions between the noVT and VT conditions, whereas the



**Figure 4.** Topographic maps of (A) the RP values and (B) the statistical comparisons between conditions for each electrode location and frequency band. (A) Each column corresponds to a specific condition of the VR experiment, whereas each row corresponds to a frequency band under study. (B) Each column represents statistically significant between-conditions differences ( $p < .05$ , Wilcoxon signed-ranks test), whereas each row corresponds to a frequency band under study. RS = resting state; VT = visuo-tactile ON; noVT = visuo-tactile OFF.





**Figure 5.** Distribution plots depicting the Lempel-Ziv complexity (LZC) values averaged over all electrodes for all three conditions under study in the theta band. Statistically significant post hoc differences between pairs of conditions are indicated with red brackets ( $p < .05$ , Wilcoxon signed-ranks test). Only the theta band is shown for ease of visualization.

alpha band displayed a clear decrease in centro-parietal RP during both conditions. Interestingly, in the beta-1 band, the noVT condition showed a localized decrease in RP in the frontal and central areas that was not as apparent in the VT condition.


The LZC values averaged over all electrodes in the theta band are shown in Figure 5. Only the theta band is shown for ease of visualization, as it was the only band with statistically significant differences between conditions. As previously mentioned, statistically significant differences were found in the theta band ( $p = .028$ ,  $\chi^2(2) = 7.14$ , Friedman

test;  $p = .140$  after FDR correction). In the said band, the statistical tests indicated that there were significant differences only between the resting state and noVT condition ( $p = .015$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .046$  after FDR correction).

The PLV values averaged over all electrode pairs are displayed in Figure 6. Statistically significant differences were again found in the alpha ( $p = .021$ ,  $\chi^2(2) = 7.71$ , Friedman test;  $p = .105$  after FDR correction) and beta-2 bands ( $p = .049$ ,  $\chi^2(2) = 6$ , Friedman test;  $p = .124$  after FDR correction). In the alpha band, the statistical tests revealed significant differences between the resting state and the VT ( $p = .031$ ,  $W = 28$ , Wilcoxon signed-ranks test;  $p = .046$  after FDR correction) and between the VT and noVT condition ( $p = .015$ ,  $W = 26$ , Wilcoxon signed-ranks test;  $p = .070$  after FDR correction). In the beta-2 band, the statistical tests showed significant differences between the resting state and the noVT condition ( $p = .046$ ,  $W = 2$ , Wilcoxon signed-ranks test;  $p = .109$  after FDR correction).

## DISCUSSION

At a phenomenological level, our VR protocol induced a feeling of subjective body ownership over the virtual body during the embodied (*in-the-body*) phase for both VT and noVT conditions in a sample of healthy participants who reported having a rather low level of prior experience with VR. Our participants' median scores for the two conditions were nonetheless slightly lower than the median scores found in Bourdin et al.'s (2017) study. Next, in the *out-the-body* phase of the noVT condition, although participants reported particularly high responses to the questions asking if they felt high up in the virtual room and floating in the air, they also reported they had felt



**Figure 6.** Distribution plots depicting the grand-average PLV values averaged over all electrode pairs for all three conditions under study. Statistically significant between-conditions differences are indicated with blue rectangles ( $p < .05$ , Friedman test), whereas statistically significant post hoc differences between pairs of conditions in bands that showed significant group interactions are indicated with red brackets ( $p < .05$ , Wilcoxon signed-ranks test).

“out of their body” (*obe* and *otherbodyobe* question) at a weak to moderate intensity. Nevertheless, participants tended to report a disownership of the previous owned body, although this feeling seems weaker than what participants from Bourdin et al.’s (2017) study reported. The scores did not significantly differ between the two conditions for all questions, although the *connectionobe* question tended to be significant. Indeed, as expected, in the *out-the-body* phase of the VT condition, participants tended to report a maintained feeling of connection with the virtual body, which was much less reported in the noVT condition. Moreover, our participants reported high scores of absorption and a moderate score of dissociation for the whole VR experiment session. Therefore, at a phenomenological level, the present VR protocol combining the first-person perspective, visuomotor and visuotactile synchronous stimulation, seemed to induce a feeling of subjective body ownership over a virtual body during the *in-the-body* phase and a strong sensation of floating in the air but a weak to moderate feeling of being “out of their body” in the *out-the-body* phase for both conditions.

In terms of EEG activity changes, the VT and noVT conditions did not differ much, except for PLV values. Nonetheless, it is worth mentioning that for the grand-average RP values, significant differences were found only between the resting state and noVT conditions in the delta and beta-1 bands, and for the LZC in the theta band. The altered alpha and beta-1 connectivity patterns showed an increase of high-frequency connectivity associated to the noVT condition (but not the VT condition), whereas grand-average alpha connectivity was lower in the VT condition compared with both the eyes-open resting state and noVT condition. EEG beta phase connectivity has been shown to increase during transitions between the baseline to unresponsive, and unresponsive to baseline states in propofol-induced sedation compared with consciousness and unconsciousness (Lee et al., 2017), thereby perhaps indicating that the OBE brain activity could behave similarly to a state of transition between states of consciousness.

Regarding the noVT condition specifically, the EEG RP analysis demonstrated a decrease in parietal alpha band RP, a localized decrease in beta-1 band RP in the frontal and central areas, as well as a significant increase in delta band power in the fronto-central area. Interestingly, this is consistent with the EEG pattern found by Timmermann et al. (2019) who studied immersion into a N,N-Dimethyltryptamine (DMT)-induced state: The DMT experience was associated with decreases in total spectral power in alpha and beta bands paralleled by increases in spontaneous signal diversity and the emergence of delta (and theta) oscillations during peak effects of the hallucinatory experience. Importantly, DMT is known to induce OBE associated with vivid visual imagery and somatic effects (Lawrence, Carhart-Harris, Griffiths, & Timmermann, 2022; Timmermann et al., 2018), just like

what we intended to study in the present work. Interestingly, a constantly growing body of evidence from research has recently revealed a prominent role of delta activity during conscious states (see Frohlich, Toker, & Monti, 2021, for a discussion), whereas a historically rooted consensus was that the delta rhythm is an indicator of unconsciousness (or highly diminished consciousness), such as in anesthesia, slow wave sleep, and coma.

Alpha power was shown to be related to self-identification and self-location during a VR experiment inducing multisensory conflicts, that is, in which participants were shown a life-sized, back facing virtual body or a cubic control object while synchronously or asynchronously stroking the back of the participant and the character or cubic control object (Lenggenhager, Halje, & Blanke, 2011). EEG alpha power significantly decreased in the sensorimotor areas when participants were asynchronously stroked compared with the visual baseline (Lenggenhager et al., 2011). The authors suggest that alpha oscillations in sensorimotor cortex and medial pFC may capture self-location (Lenggenhager et al., 2011). More generally, alpha band power over central areas has been repeatedly associated with body perception and sensorimotor processing (see Pineda, 2005, for a review). The pattern of decreases in alpha power observed in the present study might be related to modifications of self-location induced by the two VR conditions. In this study, alpha is the band that shows the most RP values decreases, mostly in central areas. This alpha decrease is nonetheless not consistent with what was observed in a previous study that invited NDE experiencers to recall the OBE they experienced during their NDE using hypnosis (Martial et al., 2019). This may be because of the fact that the latter was based on the recall of an autobiographical memory, thus probably reflecting working memory load.

The significant decrease of theta band LZC in the noVT condition compared with the resting-state condition could be linked to an artificially induced state of altered consciousness. Another complexity-based index, the perturbational complexity index is one of the most reliable indicators of the level of consciousness and is based on the complexity of electrocortical responses after TMS (Casali et al., 2013). In addition, complexity of brain activity by itself without TMS has also been shown to be reduced in anesthesia (Schartner et al., 2015). The results here suggest that complexity might be partially reduced, pointing toward the experience of OBE-like sensations of floating in the air and feeling high up followed patterns similar to those of altered levels of consciousness.

Interestingly, Bourdin et al. (2017) showed that a virtual OBE results in a reduction of death anxiety. The authors hypothesized that it may be because of an implicit learning that consciousness may be separated from the physical body. Extrapolating from this study, one can assume that OBEs artificially induced by VR may be used as a therapeutic tool in end-of-life patients because the authors showed

that their VR paradigm can have psychological effect, in relation to death anxiety.

Some limitations of this study should be acknowledged. First, the EEG recordings were obtained during long sessions where a restart of the protocol was not feasible, and thus, artifacts related to muscle activity had to be removed or dealt with during the preprocessing stage, which prevented the inclusion of the gamma band in the analyses. Second, it is worth noting that the low level of experience could indicate that the observed effects of the experiment were increased by the lack of expectations because of previous experience with VR and the novelty of the OBE-like experience was enhanced. Third, the presence of the EEG cap, which may have caused a feeling of discomfort or strengthened proprioception and perception from the real body, may have influenced the relatively weak to moderate perception of being out of their body reported by the participants. Fourth, with the notable exception of the RP, the results for the most part did not survive an FDR correction, which led to a more cautious approach to the interpretation of results. This was, however, to be expected given the small sample size. Nonetheless, the present results are encouraging of a follow-up study with a larger sample. Although the phenomenology reported by our participants did not significantly differ between the two conditions, and the noVT tended to induce a stronger feeling of dissociation with the virtual body, which they were seeing below from the ceiling of the virtual room, one can hypothesize that significant differences between both conditions would be observed in a larger sample such as in Bourdin et al. (2017). It would also be interesting to add further questions regarding the subjective experience of body ownership and bodily borders to detail their phenomenological experience. Future empirical studies are needed to better understand OBEs and other disturbed self-perceptions. However, this study represents a step in exposing the neurophysiological underpinnings of a virtually induced OBE.

In conclusion, at a phenomenological level, the present VR protocol seemed to induce a feeling of subjective body ownership over a virtual body during the *in-the-body* phase for both conditions. In the *out-the-body* phase, we succeeded in inducing subjective experiences of floating in the air and of feeling high up in the virtual room without any adverse effects in a small sample of healthy participants, but the feeling of being “out of their body” reported by our participants was relatively weak. Although some improvements can be done in the methodology, this study provides a proof-of-concept methodology for studying the phenomenon of OBEs and presents promising EEG results.

Reprint requests should be sent to Charlotte Martial, Coma Science Group, GIGA-Consciousness, University of Liège, Liège, Belgium Avenue de l’hôpital, 11 4000 Liège, Belgium, or via e-mail: [cmartial@uliege.be](mailto:cmartial@uliege.be).

## Data Availability Statement

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

## Author Contributions

C. M. and S. L. secured the funding. C. M., H. C., P. B., A. M., and O. R. collaborated on data acquisition. All authors contributed to conceptualization and study design. C. M. and P. N. did the analyses and wrote the original draft. All authors contributed to reviewing the manuscript in detail and approved the final manuscript.

## Funding Information

The study was further supported by the University and University Hospital of Liège, the BIAL Foundation (<https://dx.doi.org/10.13039/501100005032>), the Belgian National Funds for Scientific Research (FRS-FNRS), the European Union’s Horizon 2020 Framework Programme for Research and Innovation under the Specific Grant Agreement No. 945539 (Human Brain Project SGA3), the FNRS (<https://dx.doi.org/10.13039/501100002661>), PDR project (T.0134.21), the ERA-Net FLAG-ERA JTC2021 project ModelDXConsciousness (Human Brain Project Partnering Project), the fund Generet, the King Baudouin Foundation, the Télévie Foundation, the European Space Agency (ESA) and the Belgian Federal Science Policy Office (BELSPO) in the framework of the PRODEX Programme, the Public Utility Foundation “Université Européenne du Travail,” “Fondazione Europea di Ricerca Biomedica,” the Mind Science Foundation, the European Commission, the Fondation Leon Fredericq, the Mind-Care foundation, the DOCMA project (EU-H2020-MSCA-RISE-778234), the National Natural Science Foundation of China (Joint Research Project 81471100), and the European Foundation of Biomedical Research FERB Onlus. The work at UB was supported by the Immortality Project at UC Riverside, a John Templeton Foundation project and the Catalan Government (2014-SGR855).

## Diversity in Citation Practices

Retrospective analysis of the citations in every article published in this journal from 2010 to 2021 reveals a persistent pattern of gender imbalance: Although the proportions of authorship teams (categorized by estimated gender identification of first author/last author) publishing in the *Journal of Cognitive Neuroscience (JoCN)* during this period were  $M(\text{an})/M = .407$ ,  $W(\text{oman})/M = .32$ ,  $M/W = .115$ , and  $W/W = .159$ , the comparable proportions for the articles that these authorship teams cited were  $M/M = .549$ ,  $W/M = .257$ ,  $M/W = .109$ , and  $W/W = .085$  (Postle and Fulvio, *JoCN*, 34:1, pp. 1–3). Consequently, *JoCN*

encourages all authors to consider gender balance explicitly when selecting which articles to cite and gives them the opportunity to report their article's gender citation balance.

## REFERENCES

- Abásolo, D., Hornero, R., Gómez, C., García, M., & López, M. (2006). Analysis of EEG background activity in Alzheimer's disease patients with Lempel-Ziv complexity and central tendency measure. *Medical Engineering and Physics*, *28*, 315–322. <https://doi.org/10.1016/j.medengphy.2005.07.004>, PubMed: 16122963
- Alvarado, C. (2000). Out of body experiences. In E. Cardena, S. J. Lynn, & S. Krippner (Eds.), *Varieties of anomalous experience: Examining the scientific evidence* (pp. 183–218). Washington DC: American Psychological Association. <https://doi.org/10.1037/10371-006>
- Bachiller, A., Poza, J., Gómez, C., Molina, V., Suazo, V., & Hornero, R. (2015). A comparative study of event-related coupling patterns during an auditory oddball task in schizophrenia. *Journal of Neural Engineering*, *12*, 016007. <https://doi.org/10.1088/1741-2560/12/1/016007>, PubMed: 25474418
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, *57*, 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Blackmore, S. J. (1982). *Beyond the body: An investigation of out-of-the-body experiences*. London: Heinemann.
- Blackmore, S. J. (1984). A psychological theory of the out-of-body experience. *Journal of Parapsychology*, *48*, 201–218.
- Blackmore, S. J. (1987). Where am I? Perspectives in imagery and the out-of-body experience. *Journal of Mental Imagery*, *11*, 53–66.
- Blanke, O., Landis, T., Spinelli, L., & Seeck, M. (2004). Out-of-body experience and autoscopia of neurological origin. *Brain*, *127*, 243–258. <https://doi.org/10.1093/brain/awh040>, PubMed: 16019077
- Blanke, O., & Mohr, C. (2005). Out-of-body experience, heautoscopy, and autoscopic hallucination of neurological origin. Implications for mechanisms of corporeal awareness and self-consciousness. *Brain Research Reviews*, *50*, 184–199. <https://doi.org/10.1016/j.brainresrev.2005.05.008>, PubMed: 16019077
- Blanke, O., Ortigue, S., Landis, T., & Seeck, M. (2002). Stimulating illusory own-body perceptions. *Nature*, *419*, 269–270. <https://doi.org/10.1038/419269a>, PubMed: 12239558
- Bourdin, P., Barberia, I., Oliva, R., & Slater, M. (2017). A virtual out-of-body experience reduces fear of death. *PLoS One*, *12*, e0169343. <https://doi.org/10.1371/journal.pone.0169343>, PubMed: 28068368
- Brookes, M. J., Woolrich, M. W., & Barnes, G. R. (2012). Measuring functional connectivity in MEG: A multivariate approach insensitive to linear source leakage. *Neuroimage*, *63*, 910–920. <https://doi.org/10.1016/j.neuroimage.2012.03.048>, PubMed: 22484306
- Bruña, R., Maestú, F., & Pereda, E. (2018). Phase locking value revisited: Teaching new tricks to an old dog. *Journal of Neural Engineering*, *15*, 056011. <https://doi.org/10.1088/1741-2552/aacfe4>, PubMed: 29952757
- Casali, A. G., Gosseries, O., Rosanova, M., Boly, M., Sarasso, S., Casali, K. R., et al. (2013). A theoretically based index of consciousness independent of sensory processing and behavior. *Science Translational Medicine*, *5*, 198ra105. <https://doi.org/10.1126/scitranslmed.3006294>, PubMed: 23946194
- Colclough, G. L., Woolrich, M. W., Tewarie, P. K., Brookes, M. J., Quinn, A. J., & Smith, S. M. (2016). How reliable are MEG resting-state connectivity metrics? *Neuroimage*, *138*, 284–293. <https://doi.org/10.1016/j.neuroimage.2016.05.070>, PubMed: 27262239
- Ehrsson, H. H. (2007). The experimental induction of out-of-body experiences. *Science*, *317*, 1048. <https://doi.org/10.1126/science.1142175>, PubMed: 17717177
- Facco, E., Casiglia, E., Al Khafaji, B. E., Finatti, F., Duma, G. M., Mento, G., et al. (2019). The neurophenomenology of out-of-body experiences induced by hypnotic suggestions. *International Journal of Clinical and Experimental Hypnosis*, *67*, 39–68. <https://doi.org/10.1080/00207144.2019.1553762>, PubMed: 30702402
- Friston, K. J. (2011). Functional and effective connectivity: A review. *Brain Connectivity*, *1*, 13–36. <https://doi.org/10.1089/brain.2011.0008>, PubMed: 22432952
- Frohlich, J., Toker, D., & Monti, M. M. (2021). Consciousness among delta waves: A paradox? *Brain*, *144*, 2257–2277. <https://doi.org/10.1093/brain/awab095>, PubMed: 33693596
- Herrero, N. L., Gallo, F. T., Gasca-Rolín, M., Gleiser, P. M., & Forcato, C. (2023). Spontaneous and induced out-of-body experiences during sleep paralysis: Emotions, “aura” recognition, and clinical implications. *Journal of Sleep Research*, *32*, e13703. <https://doi.org/10.1111/jsr.13703>, PubMed: 36053735
- Irwin, H. J. (1985). *Flight of mind: A psychological study of the out-of-body experience*. Metuchen, New Jersey: Scarecrow Press.
- Irwin, H. J. (2000). The disembodied self: An empirical study of dissociation and the out-of-body experience. *Journal of Parapsychology*, *64*, 261–277.
- Kiltner, K., Grotten, R., & Slater, M. (2012). The sense of embodiment in virtual reality. *Presence: Teleoperators and Virtual Environments*, *21*, 373–387. [https://doi.org/10.1162/PRES\\_a\\_00124](https://doi.org/10.1162/PRES_a_00124)
- Kim, H., Lee, U., & Vlisides, P. E. (2022). Delirium and cortical complexity. *Journals of Gerontology: Series A*, *77*, 2219–2220. <https://doi.org/10.1093/gerona/glac163>, PubMed: 35943887
- Kokkinara, E., & Slater, M. (2014). Measuring the effects through time of the influence of visuomotor and visuotactile synchronous stimulation on a virtual body ownership illusion. *Perception*, *43*, 43–58. <https://doi.org/10.1068/p7545>, PubMed: 24689131
- Lachaux, J. P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Human Brain Mapping*, *8*, 194–208. [https://doi.org/10.1002/\(sici\)1097-0193\(1999\)8:4<194::aid-hbm4>3.0.co;2-c](https://doi.org/10.1002/(sici)1097-0193(1999)8:4<194::aid-hbm4>3.0.co;2-c)
- Lawrence, D. W., Carhart-Harris, R., Griffiths, R., & Timmermann, C. (2022). Phenomenology and content of the inhaled N, N-dimethyltryptamine (N, N-DMT) experience. *Scientific Reports*, *12*, 8562. <https://doi.org/10.1038/s41598-022-11999-8>, PubMed: 35610230
- Lee, M., Sanders, R. D., Yeom, S. K., Won, D. O., Seo, K. S., Kim, H. J., et al. (2017). Network properties in transitions of consciousness during Propofol-induced sedation. *Scientific Reports*, *7*, 1–13. <https://doi.org/10.1038/s41598-017-15082-5>, PubMed: 29196672
- Lempel, A., & Ziv, J. (1976). On the complexity of finite sequences. *IEEE Transactions on Information Theory*, *22*, 75–81. <https://doi.org/10.1109/TIT.1976.1055501>
- Lenggenhager, B., Halje, P., & Blanke, O. (2011). Alpha band oscillations correlate with illusory self-location induced by virtual reality. *European Journal of Neuroscience*, *33*, 1935–1943. <https://doi.org/10.1111/j.1460-9568.2011.07647.x>, PubMed: 21395867

- Lenggenhager, B., Mouthon, M., & Blanke, O. (2009). Spatial aspects of bodily self-consciousness. *Consciousness and Cognition*, *18*, 110–117. <https://doi.org/10.1016/j.concog.2008.11.003>, PubMed: 19109039
- Martial, C., Cassol, H., Laureys, S., & Gosseries, O. (2020). Near-death experience as a probe to explore (disconnected) consciousness. *Trends in Cognitive Sciences*, *24*, 173–183. <https://doi.org/10.1016/j.tics.2019.12.010>, PubMed: 31982302
- Martial, C., Mensen, A., Charland-Verville, V., Vanhaudenhuyse, A., Rentmeister, D., Ali Bahri, M., et al. (2019). Neurophenomenology of near-death experience memory in hypnotic recall: A within-subject EEG study. *Scientific Reports*, *9*, 14047. <https://doi.org/10.1038/s41598-019-50601-6>, PubMed: 31575924
- Martial, C., Simon, J., Puttaert, N., Gosseries, O., Charland-Verville, V., Nyssen, A.-S., et al. (2020). The near-death experience content (NDE-C): Development and psychometric validation. *Consciousness and Cognition*, *86*, 103049. <https://doi.org/10.1016/j.concog.2020.103049>, PubMed: 33227590
- Núñez, P., Poza, J., Bachiller, A., Gomez-Pilar, J., Lubeiro, A., Molina, V., et al. (2017). Exploring non-stationarity patterns in schizophrenia: Neural reorganization abnormalities in the alpha band. *Journal of Neural Engineering*, *14*, 046001. <https://doi.org/10.1088/1741-2552/aa6e05>, PubMed: 28424430
- Núñez, P., Poza, J., Gómez, C., Rodríguez-González, V., Hillebrand, A., Tola-Arribas, M. A., et al. (2019). Characterizing the fluctuations of dynamic resting-state electrophysiological functional connectivity: Reduced neuronal coupling variability in mild cognitive impairment and dementia due to Alzheimer's disease. *Journal of Neural Engineering*, *16*, 056030. <https://doi.org/10.1088/1741-2552/ab234b>, PubMed: 31112938
- O'Neill, G. C., Tewarie, P., Vidaurre, D., Liuzzi, L., Woolrich, M. W., & Brookes, M. J. (2018). Dynamics of large-scale electrophysiological networks: A technical review. *Neuroimage*, *180*, 559–576. <https://doi.org/10.1016/j.neuroimage.2017.10.003>, PubMed: 28988134
- Palmieri, A., Calvo, V., Kleinbub, J. R., Meconi, F., Marangoni, M., Barilaro, P., et al. (2014). "Reality" of near-death-experience memories: Evidence from a psychodynamic and electrophysiological integrated study. *Frontiers in Human Neuroscience*, *8*, 429. <https://doi.org/10.3389/fnhum.2014.00429>, PubMed: 24994974
- Piarulli, A., Bergamasco, M., Thibaut, A., Cologan, V., Gosseries, O., & Laureys, S. (2016). EEG ultradian rhythmicity differences in disorders of consciousness during wakefulness. *Journal of Neurology*, *263*, 1746–1760. <https://doi.org/10.1007/s00415-016-8196-y>, PubMed: 27294259
- Pineda, J. A. (2005). The functional significance of mu rhythms: Translating "seeing" and "hearing" into "doing". *Brain Research Reviews*, *50*, 57–68. <https://doi.org/10.1016/j.brainresrev.2005.04.005>, PubMed: 15925412
- Rioul, O., & Vetterli, M. (1991). Wavelets and signal processing. *IEEE Signal Processing Magazine*, *8*, 14–38. <https://doi.org/10.1109/79.91217>
- Roach, B. J., & Mathalon, D. H. (2008). Event-related EEG time–frequency analysis: An overview of measures and an analysis of early gamma band phase locking in schizophrenia. *Schizophrenia Bulletin*, *34*, 907–926. <https://doi.org/10.1093/schbul/sbn093>, PubMed: 18684772
- Ruiz-Gómez, S., Gómez, C., Poza, J., Gutiérrez-Tobal, G., Tola-Arribas, M., Cano, M., et al. (2018). Automated multiclass classification of spontaneous EEG activity in Alzheimer's disease and mild cognitive impairment. *Entropy*, *20*, 35. <https://doi.org/10.3390/e20010035>, PubMed: 33265122
- Samar, V. J., Bopardikar, A., Rao, R., & Swartz, K. (1999). Wavelet analysis of neuroelectric waveforms: A conceptual tutorial. *Brain and Language*, *66*, 7–60. <https://doi.org/10.1006/brln.1998.2024>, PubMed: 10080864
- Schartner, M., Seth, A., Noirhomme, Q., Boly, M., Bruno, M. A., Laureys, S., et al. (2015). Complexity of multi-dimensional spontaneous EEG decreases during propofol induced general anaesthesia. *PLoS One*, *10*, e0133532. <https://doi.org/10.1371/journal.pone.0133532>, PubMed: 26252378
- Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1996). Stimulus specificity of phase-locked and non-phase-locked 40 Hz visual responses in human. *Journal of Neuroscience*, *16*, 4240–4249. <https://doi.org/10.1523/JNEUROSCI.16-13-04240.1996>, PubMed: 8753885
- Tanabe, S., Parker, M., Lennertz, R., Pearce, R. A., Banks, M. I., & Sanders, R. D. (2022). Reduced electroencephalogram complexity in postoperative delirium. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, *77*, 502–506. <https://doi.org/10.1093/geron/ab352>, PubMed: 34958346
- Taylor, R.M., Hudson, T.C., Seeger, A., Weber, H., Juliano, J., & Helsler, A.T. (2001). VRPN: A device-independent, network-transparent VR peripheral system. In *Proceedings of the ACM symposium on virtual reality software and technology* (pp. 55–61). New York: ACM Press. <https://doi.org/10.1145/505008.505019>
- Timmermann, C., Roseman, L., Schartner, M., Milliere, R., Williams, L. T. J., Erritzoe, D., et al. (2019). Neural correlates of the DMT experience assessed with multivariate EEG. *Scientific Reports*, *9*, 16324. <https://doi.org/10.1038/s41598-019-51974-4>, PubMed: 31745107
- Timmermann, C., Roseman, L., Willimans, L., Erritzoe, D., Martial, C., Cassol, H., et al. (2018). DMT models the near-death experience. *Frontiers in Psychology*, *9*, 1424. <https://doi.org/10.3389/fpsyg.2018.01424>, PubMed: 30174629
- Torrence, C., & Compo, G. P. (1998). A practical guide to wavelet analysis. *Bulletin of the American Meteorological Society*, *79*, 61–78. [https://doi.org/10.1175/1520-0477\(1998\)079<0061:APGTWA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2)
- Tseng, P., & Juan, C. H. (2013). Virtual reality in the neuroscience of multisensory integration and consciousness of bodily self. *Journal of Neuroscience and Neuroengineering*, *2*, 387–392. <https://doi.org/10.1166/jnsne.2013.1063>
- Vanhaudenhuyse, A., Ledoux, D., Gosseries, O., Demertzi, A., Laureys, S., & Faymonville, M.-F. (2019). Can subjective ratings of absorption, dissociation, and time perception during "neutral hypnosis" predict hypnotizability?: An exploratory study. *International Journal of Clinical and Experimental Hypnosis*, *67*, 1–11. <https://doi.org/10.1080/00207144.2019.1553765>, PubMed: 30702397
- van Heugten-van der Kloet, D., Cosgrave, J., van Rheede, J., & Hicks, S. (2018). Out-of-body experience in virtual reality induces acute dissociation. *Psychology of Consciousness: Theory, Research, and Practice*, *5*, 346–357. <https://doi.org/10.1037/cns0000172>