

1 **Effects of 2-year physical activity and dietary intervention on adrenarchal and** 2 **pubertal development: the PANIC study**

3 Jani Liimatta^{1,2,3}, Christa E. Flück^{2,3}, Aino Mäntyselkä^{1,4}, Merja R. Häkkinen^{5,6}, Seppo Auriola⁵,
4 Raimo Voutilainen^{1,4}, Jarmo Jääskeläinen^{1,4}, Timo A. Lakka^{7,8,9}

5 ¹ Kuopio Pediatric Research Unit (KuPRU), University of Eastern Finland, Kuopio, Finland

6 ² Department of BioMedical Research (DBMR), University of Bern, Bern, Switzerland

7 ³ Pediatric Endocrinology, Diabetology, and Metabolism, Inselspital, Bern University Hospital, Bern,
8 Switzerland

9 ⁴ Department of Pediatrics, Kuopio University Hospital, Kuopio, Finland

10 ⁵ School of Pharmacy, University of Eastern Finland, Kuopio, Finland

11 ⁶ Department of Health Security, Finnish Institute for Health and Welfare (THL), Kuopio, Finland

12 ⁷ Institute of Biomedicine, School of Medicine, University of Eastern Finland, Kuopio, Finland

13 ⁸ Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, Kuopio, Finland

14 ⁹ Foundation for Research in Health Exercise and Nutrition, Kuopio Research Institute of Exercise
15 Medicine, Kuopio, Finland

16 **Correspondence and reprint requests:**

17 MD, PhD, Jani Liimatta (ORCID ID: 0000-0001-6076-1512)

18 Kuopio Pediatric Research Unit (KuPRU)

19 Department of Pediatrics, Kuopio University Hospital

20 P.O. box 100, FI-70029 Kuopio, Finland

21 Email: jani.liimatta@pshyvinvointialue.fi

1 **Clinical Trials registration:** The PANIC study has been registered at www.clinicaltrials.gov (No.

2 NCT01803776).

3 **Disclosure summary:** The authors have nothing to disclose.

4 **Funding:**

5 Open access funding provided by University of Eastern Finland (UEF) including Kuopio University
6 Hospital. The PANIC study has been supported financially by grants from Ministry of Education and
7 Culture of Finland, Academy of Finland, Ministry of Social Affairs and Health of Finland, Research
8 Committee of the Kuopio University Hospital Catchment Area (State Research Funding), Finnish
9 Innovation Fund Sitra, Social Insurance Institution of Finland, Finnish Cultural Foundation, Foundation
10 for Pediatric Research, Diabetes Research Foundation in Finland, Finnish Foundation for Cardiovascular
11 Research, Juho Vainio Foundation, Paavo Nurmi Foundation, Yrjö Jahnsson Foundation, and the city of
12 Kuopio.

13 This work was also supported by the Sigrid Jusélius Foundation (Helsinki, Finland; Research Fellowship to
14 J.L.), the Foundation for Pediatric Research (Helsinki, Finland; Research Fellowship to J.L.).

15 The funding sources have not been involved in: 1) design and conduct of the study; 2) collection,
16 management, analysis, and interpretation of data; 3) review or approval of the manuscript; or 4)
17 decision to submit the manuscript for publication.

18 **Key words:** Adrenarche, Puberty, DHEAS, Lifestyle Intervention, Physical Activity, Diet

19

20

21

1 Abstract

2 **Context:** Childhood overweight has been linked to earlier development of adrenarche and puberty, but
3 it remains unknown if lifestyle interventions influence sexual maturation in general populations.

4 **Objective:** To investigate if a 2-year lifestyle intervention influences circulating androgen concentrations
5 and sexual maturation in a general population of children.

6 **Design and participants:** A 2-year intervention study in which 421 prepubertal and mostly normal-
7 weight 6-9-year-old children were allocated either to a lifestyle intervention group (119 girls, 132 boys)
8 or a control group (84 girls, 86 boys).

9 **Intervention:** A 2-year physical activity and dietary intervention.

10 **Main outcome measures:** Serum dehydroepiandrosterone, dehydroepiandrosterone sulfate,
11 androstenedione, and testosterone concentrations, and clinical adrenarchal and pubertal signs.

12 **Results:** The intervention and control groups had no differences in body size and composition, clinical
13 signs of androgen action, and serum androgens at baseline. The intervention attenuated the increase of
14 dehydroepiandrosterone ($p = 0.032$), dehydroepiandrosterone sulfate ($p = 0.001$), androstenedione ($p =$
15 0.003), and testosterone ($p = 0.007$) and delayed pubarche ($p = 0.038$) in boys but it only attenuated the
16 increase of dehydroepiandrosterone ($p = 0.013$) and dehydroepiandrosterone sulfate ($p = 0.003$) in girls.
17 These effects of lifestyle intervention on androgens and the development of pubarche were
18 independent of changes in body size and composition but the effects of intervention on androgens were
19 partly explained by changes in fasting serum insulin.

20 **Conclusions:** A combined physical activity and dietary intervention attenuates the increase of serum
21 androgen concentrations and sexual maturation in a general population of prepubertal and mostly
22 normal-weight children, independently of changes in body size and composition.

1 **1. Introduction**

2 Sexual maturation from childhood to adulthood includes two main events, adrenarche and puberty
3 (1,2). Adrenarche refers to the maturation of the adrenocortical zona reticularis (ZR) leading to
4 increased production of adrenal androgens, dehydroepiandrosterone (DHEA), dehydroepiandrosterone
5 sulfate (DHEAS), androstenedione (A4), and 11-hydroxy-A4 (11OHA4) (1). Increased exposure to adrenal
6 androgens in adrenarche eventually leads to clinical signs, which include adult-type body odor, oily hair
7 and skin, microcomedonal acne, and the development of axillary and pubic hair. Adrenocorticotrophic
8 hormone (ACTH) stimulation from the hypothalamic-pituitary-adrenal (HPA) axis is necessary for ZR
9 development and the onset of adrenarche, but adrenal androgen production is not similarly dependent
10 and regulated by HPA axis as puberty is by pulsatile secretion of gonadotropins from the hypothalamic-
11 pituitary-gonadal (HPG) axis (1). The activation of the HPG axis and the onset of puberty are usually seen
12 a few years after adrenarche with testicular enlargement in boys and breast development in girls being
13 their first clinical signs (2).

14 Although adrenarche and puberty are independent events with different regulatory mechanisms, they
15 seem to be linked as pubertal development is advanced in girls with premature timing of adrenarche (3)
16 and in girls with higher circulating DHEAS concentrations in mid-childhood (4). Also, early timing of
17 adrenarche and puberty are associated with adiposity at the age of these events (5, 6) and weight gain
18 in early childhood (7-9). Secular trends of the increased prevalence of childhood obesity and earlier
19 timing of puberty have been shown to share similar patterns, and early weight gain leading to increased
20 fat mass and childhood obesity has been suggested to be one factor behind a trend of earlier sexual
21 maturation, especially in girls (6,10). Other possible factors behind earlier timing of sexual maturation
22 may be insulin and insulin-like growth factors (IGFs) as it is known that children with premature
23 adrenarche have signs of insulin resistance together with increased circulating IGF-1 concentrations (5,7)
24 and that there is a physiological decrease in insulin sensitivity during puberty (11).

1 Physical activity and dietary interventions have been found to decrease adiposity and improve glucose
2 tolerance in overweight and obese children (12). Given that adipose tissue is somehow regulating the
3 timing of maturation as suggested by the association between early timing of puberty and adiposity
4 (6,10), it could be hypothesized that lifestyle interventions affecting body fat content and insulin
5 sensitivity also influence adrenarchal and pubertal development. Supporting this hypothesis, earlier
6 studies have found that weight loss decreased circulating A4 and testosterone levels in obese children
7 (13) and that lifestyle modifications had beneficial effects on hormonal status in adolescent girls with
8 the polycystic ovary syndrome (PCOS) (14), a condition characterized by hyperandrogenism. Exercise has
9 also been found to improve insulin sensitivity in adult women with PCOS (15). Little is known, however,
10 whether lifestyle interventions influence adrenarchal and pubertal development in general populations
11 of children with mostly normal weight.

12 We have reported earlier that a 2-year combined physical activity and dietary intervention attenuated
13 the increase of insulin resistance in a general population of mostly normal weight primary school
14 children participating in the Physical Activity and Nutrition in Children (PANIC) study (16). It is not well
15 known how lifestyle interventions affect sexual maturation in general populations of mostly normal
16 weight children, and we therefore report here the effects of the same 2-year lifestyle intervention on
17 serum androgen concentrations and on adrenarchal and pubertal development in these children.

18

19 **2. Methods**

20 **2.1. Participants and design**

21 We investigated a cohort of children who had participated in the Finnish PANIC study. The PANIC study
22 is a non-randomized controlled trial on the effects of a combined physical activity and dietary
23 intervention on cardiometabolic risk factors and other health outcomes in a general population of

1 children from the city of Kuopio, Finland (17,18). The city of Kuopio has approximately 120 000
2 inhabitants and includes a central urban area and a few surrounding rural villages. The study protocol
3 has been approved by the Research Ethics Committee of the Hospital District of Northern Savo
4 (statement 69/2006), and the study has been carried out in accordance with the principles of the
5 Declaration of Helsinki. The parents or caregivers of children provided their written consent, and the
6 children gave their assent to participation. The PANIC study has been registered at
7 www.clinicaltrials.gov (No. NCT01803776).

8 A flow chart of this study is presented in Figure 1. A total of 736 children, aged 6-9 years, who started
9 the first grade in 16 primary schools of the city of Kuopio between 2007-2009, were invited to
10 participate. Altogether, 512 (70%) children accepted the invitation and participated in the baseline
11 examination between October 2007 and December 2009. The participants did not differ in sex, age,
12 height standard deviation score (SDS), or body mass index (BMI) SDS from all children who started the
13 first grade in the city of Kuopio between 2007-2009. A total of eight children were excluded at baseline
14 because of either disabilities that could hamper participation in the intervention study or withdrawal of
15 the families. Thus, 504 children participated in the 2-year intervention study.

16 Children from nine schools were allocated to an intervention group and children from other seven
17 schools to a control group to avoid contamination in the control group by local or national health
18 promotion programs that could have been initiated in the study region over the 2-year intervention
19 study. The intervention and control groups were proportionally matched according to the location of the
20 schools (urban vs. rural) to minimize sociodemographic differences between the groups. More children
21 were included in the intervention group as a larger number of dropouts were expected in this group.
22 The children, their parents, or caregivers, or people carrying out the examination visits or doing the
23 measurements were not blinded to the group assignment. A total of 66 children were lost over two
24 years because of moving elsewhere, lack of time or motivation, or unknown reasons.

1 We excluded 19 children from the present analyses because of suspected early signs of pubertal
2 development or missing pubertal data at baseline, and altogether 251 children (119 girls, 131 boys) from
3 the intervention group and 170 children (84 girls, 86 boys) from the control group completed the 2-year
4 follow-up period and were included in the present analyses (Figure 1). Data on biomarkers at baseline
5 and 2-year follow-up examinations were available for 241 children in the intervention group and for 158
6 children in the control group. Data on clinical adrenarchal and pubertal signs were available for 240
7 children in the intervention group and for 164 children in the control group. The partly incomplete data
8 on biomarkers were due either to missing blood samples or to hemolysis interfering with the analyses.

9 **2.2. Physical activity and dietary intervention**

10 The main goals for the individualized and family-based physical activity and dietary intervention were to
11 1) decrease the consumption of significant sources of saturated fat and particularly high-fat dairy and
12 meat products, 2) increase the consumption of significant sources of unsaturated fat and particularly
13 high-fat vegetable oil-based margarines, vegetable oils, and fish, 3) increase the consumption of
14 vegetables, fruits, and berries, 4) increase the consumption of significant sources of fiber and
15 particularly whole grain products, 5) decrease the consumption of significant sources of sugar and
16 particularly sugar-sweetened beverages and dairy products, and candies, 6) decrease the consumption of
17 significant sources of salt and the use of salt in cooking, 7) increase total physical activity by emphasizing
18 its diversity, 8) decrease total and particularly screen-based sedentary behavior, and 9) avoid excessive
19 energy intake.

20 The 2-year intervention consisted of six intervention visits that occurred 0.5, 1.5, 3, 6, 12, and 18 months
21 after baseline examinations. Each intervention visit included 30–45 min of physical activity counselling
22 and 30–45 min of dietary counselling for the children and their parents or caregivers. The children and
23 their parents or caregivers received individualized advice from a specialist in exercise medicine and a

1 clinical nutritionist on how to increase physical activity, decrease sedentary time, and improve diet
2 among the children in everyday conditions. Each visit had a specific topic of discussion (physical activity,
3 sedentary time, diet) in accordance with the goals of the intervention and included practical tasks on
4 these topics for the children. The children and their parents or caregivers also received fact sheets on
5 physical activity, sedentary time, and diet as well as verbal and written information on opportunities for
6 exercising in the city of Kuopio. Some material support was also given for physical activity, such as
7 exercise equipment and allowance for playing indoor sports. Of the children in the intervention group
8 who attended the baseline examination, 87% participated in all six visits, 92% in at least five visits, and
9 96% in at least four visits. The children in the intervention group, particularly those who did not attend
10 organized sports or exercise, were also encouraged to participate in after-school exercise clubs
11 organized at the nine schools by trained exercise instructors of the PANIC study. The children in the
12 control group were not allowed to attend these exercise clubs to avoid a non-intentional intervention in
13 the control group. Altogether, 83% of the children in the intervention group participated in at least one
14 of the after-school exercise clubs, and 41% attended these exercise clubs at least once a month.

15 In the control group, the children and their parents or caregivers received general verbal and written
16 advice on health-improving physical activity and diet only at baseline with no further lifestyle
17 counselling.

18 **2.3. Assessment of body size and composition, physical activity, and dietary factors**

19 Anthropometric measurements were performed in the morning after a 12-hour fast. Height was
20 measured thrice using a calibrated wall-mounted stadiometer to an accuracy of 0.1 cm with the children
21 standing in the Frankfurt plane, and the average of the two closest values was used in the analyses.
22 Weight was measured twice using a weight scale integrated into the InBody 720® bioelectrical
23 impedance device (Biospace, Seoul, South Korea) to an accuracy of 0.1 kg with the children having

1 emptied the bladder and wearing light underwear, and the mean of the two values was used in the
2 analyses. Body lean and fat masses were measured using the same InBody 720® bioelectrical impedance
3 device. BMI was calculated as body weight (kg) divided by body height (m) squared. Baseline growth
4 velocity (cm/year) was calculated from height values between the age of five years and the age at
5 baseline, and 2-year growth velocity from height values between the age at baseline and 2-year follow-
6 up examinations. Age- and sex- standardized height SDS and BMI SDS were calculated using the Finnish
7 national references (19). Physical activity was assessed using the PANIC Physical Activity Questionnaire
8 and dietary factors using 4-day food records (17).

9 **2.4. Assessment of adrenarchal and pubertal status**

10 A trained research physician assessed pubertal status according to the Tanner staging method (20,21).
11 For girls, breast development (Tanner B) was assessed by inspection and palpation and scored as 1-5.
12 For boys, testicular development (Tanner G) was assessed with orchidometer and scored as 1-5. For girls
13 and boys, development of pubic hair (Tanner P) was assessed with inspection and scored as 1-5.
14 Pubertal onset was defined as breast development (Tanner B) ≥ 2 for girls and testicular volume ≥ 4 ml
15 (Tanner G ≥ 2) for boys.

16 A trained research physician assessed also adrenarchal signs that included adult-type body odor,
17 greasiness of the hair and skin, acne, and development of axillary or pubic hair. Adult-type body odor
18 and greasiness of the hair and skin were assessed also by asking the parents. Clinical adrenarche was
19 defined if one or more of these clinical signs were present.

20 **2.5. Collection of birth data**

21 Birth data were collected retrospectively from the national Medical Birth Register and the local Kuopio
22 University Hospital register. Preterm birth was defined as if a child was born before 37.0 weeks of

1 gestational age. Birth weight SDS and birth length SDS were calculated using Finnish national references
2 (22). Being born small for gestational age was defined as birth weight and/or birth length ≤ -2 SDS.

3

4 **2.6. Biochemical analyses**

5 All blood samples were collected in the morning after a 12-hour fast, and the serum samples were
6 stored at -80°C until used for biochemical analyses.

7 Serum DHEAS concentrations were measured using an enzyme-linked immunosorbent assay (ELISA) kit
8 (cat # 1950, [RRID:AB_2819763](https://pubmed.ncbi.nlm.nih.gov/2819763/), Alpha Diagnostic International, San Antonio, TX, USA). The intra-assay
9 and inter-assay coefficients of variation (CV) for this assay were 7.5-11.5% and 7.0-11.0%, respectively.
10 The detection limit of the DHEAS immunoassay was $0.014\ \mu\text{mol/l}$. Biochemical adrenarche was defined
11 as serum DHEAS concentration $\geq 1\ \mu\text{mol/l}$ (37 ng/dL).

12 Serum concentrations of other androgens than DHEAS were measured using liquid chromatography –
13 tandem mass spectrometry (LC-MS/MS), as described previously (23). At the time of measuring serum
14 androgen concentrations at baseline and 2-year follow-up examination by LC-MS/MS, our in-house
15 method did not include androgens in the so-called 11-oxygenated androgen pathway (24). Therefore,
16 only androgens belonging to the classic pathway, including DHEA, DHEAS, A4, and testosterone, are
17 used in the present study.

18 Absolute changes in serum androgen levels over two years (Δ) were calculated by the following formula:
19 [steroid concentration at 2-year follow-up examination – steroid concentration at baseline]. Relative
20 changes (%) in serum androgen levels were calculated with the following formula: [(androgen
21 concentration at 2-year follow-up examination – androgen concentration at baseline examination) /
22 androgen concentration at baseline examination) x 100].

1 Serum IGF-1 concentrations were measured using an ELISA kit (cat # E20, [RRID:AB 2813791](#),
2 Mediagnost, Reutlingen, Germany). The intra-assay and inter-assay CVs for this assay were 5.1-6.6% and
3 7.7-9.2%, respectively. Serum luteinizing hormone (LH) concentrations were measured using an
4 electrochemiluminescence immunoassay (Cat # 11732234, [RRID:AB 2800498](#), Roche diagnostics GmbH,
5 Mannheim, Germany) with an inter- and intra-assay CVs as 1.6-1.9 and 1.4%, respectively. Biochemical
6 evidence for pubertal onset at 2-year follow-up examination was defined as serum LH concentration of
7 at least 0.3 U/l (25).

8 Serum insulin concentrations were analyzed using an ECLIA kit with the sandwich principle (Cat #
9 12017547, [RRID:AB 2756877](#), Roche Diagnostics, Mannheim, Germany). The intra-assay and inter-assay
10 CVs for this assay were 1.3–3.5% and 1.6–4.4%, respectively. A hexokinase method was used to analyze
11 plasma glucose (Roche Diagnostics). The intra-assay and inter-assay CVs for this method were 0.7–0.9 %
12 and 1.5–1.8 %, respectively. Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) was
13 calculated using the following formula: $[(\text{fasting insulin (mU/l)} \times \text{fasting glucose (mmol/l)}) / 22.5]$ (26).

14 **2.7. Statistical analyses**

15 All statistical analyses were performed using the IBM SPSS statistics software, Version 25.0 (IBM Corp.,
16 Armonk, NY, USA). A p-value less than 0.05 was used to indicate statistical significance. Distributions of
17 the variables were analyzed by the Shapiro-Wilk test and visual observation of the histograms. The
18 Mann-Whitney U-test and the Student's t-test were used to compare differences in continuous variables
19 between groups. For categorical variables, the Pearson's χ^2 test was used. Intervention effects on the
20 changes of serum androgen concentrations during the follow-up period (Δ steroid concentration) were
21 analyzed using linear regression models that were tested to meet assumptions of normality, linearity,
22 homoscedasticity, and absence of multicollinearity. In the case of continuous variables with skewed
23 distributions, logarithmic transformation was used before regression analyses.

1 **3. Results**

2 **3.1. Characteristics of children at baseline**

3 Children in the control group had slightly lower birth length SDS at baseline (Table 1) and slightly higher
4 serum IGF-1 concentration at baseline (Table 2) than children in the physical activity and dietary
5 intervention group. These differences between the groups reached statistical significance in boys but
6 not in girls. Serum androgen concentrations were comparable between the groups at baseline (Table 2).

7 **3.2. Intervention effects on physical activity and dietary factors**

8 Total and unsupervised physical activity increased in the intervention group but decreased in the control
9 group over two years (total physical activity, +7 vs -5 min/d, $p = 0.003$; unsupervised physical activity, +6
10 vs -9 min/d, $p < 0.001$). There were no differences in the changes of organized exercise between the
11 groups (+4 vs +3 min/d, $p = 0.444$). The effects of intervention on dietary factors over two years have
12 been published comprehensively elsewhere (17). In brief, the intervention improved diet quality, for
13 example increased the consumption of vegetables, fruits, and berries and replaced the consumption of
14 foods high in saturated fat with the consumption of foods high in unsaturated fat. There was no
15 difference in the change of protein intake between the groups.

16 **3.3. Intervention effects on height, growth velocity, and body composition**

17 There were no differences in measures of body size or composition between the intervention and
18 control groups at baseline or 2-year follow-up examinations (Table 1). The intervention had no effects
19 on height SDS, growth velocity, BMI SDS, or body fat percentage over two years in the whole study
20 population or in girls or in boys separately (Figure 2 and Table 1).

21 **3.4. Intervention effects on clinical signs of adrenarche and puberty**

22 The prevalence of pubarche at 2-year follow-up examination was lower among boys in the intervention
23 group than among boys in the control group (Table 1). Otherwise, we did not find any statistically

1 significant differences between the groups in the prevalence of clinical adrenarchal or pubertal signs.
2 When pubertal onset was defined by Tanner B \geq 2 and/or serum LH concentration \geq 0.3 U/l, girls in the
3 intervention group tended to have a lower prevalence of puberty at 2-year follow-up examination than
4 girls in the control group, but this difference did not reach the level of statistical significance ($p = 0.057$).

5 **3.5. Intervention effects on serum androgens and other biomarkers**

6 Serum DHEA concentration at 2-year follow-up examination was lower among children in the
7 intervention group than among children in the control group (Table 2). No differences in serum DHEAS,
8 A4, or testosterone concentrations were found between the groups at the 2-year follow-up
9 examination.

10 Serum DHEA, DHEAS, A4, and testosterone concentrations increased less over two years among children
11 in the intervention group than among children in the control group (Figure 3). In boys, the levels of all
12 androgens increased less in the intervention group than in the control group, but in girls this was the
13 case only for DHEA and DHEAS (Figure 3).

14 The intervention attenuated the increase of serum DHEA concentration over two years after adjustment
15 for gender, gestational age, birth length SDS, age, serum IGF-1 concentration, BMI SDS, and presence of
16 adrenarchal signs at baseline, and presence of pubertal signs at 2-year follow-up (Table 3; model 1). The
17 intervention also attenuated the increase of serum testosterone concentration over two years in boys
18 but not in girls after these adjustments. Further adjustment for the change of fasting serum insulin
19 concentration over two years had no statistically significant effect on these results (Table 3; model 2).
20 However, the change of HOMA-IR correlated positively with and the changes of serum DHEA, A4, and
21 testosterone concentrations over two years in girls ($r = 0.212$ and $p < 0.001$ for DHEA; $r = 0.199$ and $p =$
22 0.010 for A4; $r = 0.153$ and $p = 0.049$ for testosterone) but not in boys.

23

1 4. Discussion

2 This 2-year non-randomized controlled trial showed that the physical activity and dietary intervention
3 attenuated the increase of serum DHEAS, DHEA, A4, and testosterone concentrations in a general
4 population of initially prepubertal and mostly normal-weight children. The intervention attenuated the
5 increase of all these serum androgens in boys but only the increase of serum DHEA and DHEAS in girls.
6 The intervention also decreased the development of pubarche in boys but not in girls. The effects of
7 physical activity and dietary intervention on serum androgens and the development of pubarche were
8 independent of changes in body size and composition, but the intervention effects were partly mediated
9 by changes in fasting serum insulin.

10 Studies that have reported the effects of physical activity or dietary interventions on circulating
11 concentrations of androgens or other steroid hormones in children are scarce, and to the best of our
12 knowledge, there are no such studies in general populations of initially prepubertal and mostly normal-
13 weight children. In German obese prepubertal 8-year-old children who had received physical activity,
14 dietary, and other behavioral interventions, weight loss was associated with decreases in circulating
15 concentrations of glucocorticoids, A4, and testosterone but not DHEAS. (13). Children in our study
16 differed from those in the German study by being mostly normal weight and showing no intervention
17 effects on measures of body size or composition. Consistent with the results of the German study,
18 however, we found that the intervention did not decrease but attenuated the gradual increase of serum
19 androgen concentrations between the ages of 6 and 10 years. The same German group also reported
20 that successful weight loss was associated with increased steroid sulfation capacity in obese children
21 (27). In our study, DHEAS was the only measured sulfated steroid, and we did not detect major changes
22 in the absolute serum DHEAS concentration after intervention, although its relative gradual increase was
23 smaller in the intervention group than in the control group between the ages of 6 to 11 years.

1 A meta-analysis showed that physical activity, dietary, and other behavioral interventions improved the
2 free androgen index and clinical manifestations, including the Ferriman-Gallwey score of hairiness and
3 menstrual irregularities, in adolescent girls with PCOS (14), which is a condition characterized by
4 hyperandrogenism. Although hyperandrogenic adolescent girls are metabolically different from the
5 mostly normal-weight children participating in our study, the results of the meta-analysis also suggest
6 that lifestyle interventions can modulate androgen production and metabolism in children. In another
7 meta-analysis among PCOS women, lifestyle modifications were found to be associated with reduced
8 fasting blood glucose and serum insulin levels (28). Insulin is an important biomarker of sexual
9 maturation, and premature adrenarche and normal central puberty are known to be associated with
10 decreased insulin sensitivity (5,11). We have previously shown that the 2-year combined physical
11 activity and dietary intervention attenuated the increase of insulin resistance in the present population
12 of initially prepubertal and mostly normal-weight children (16). In the present study, we found that the
13 same intervention attenuated the increase of serum androgen concentrations and the change in HOMA-
14 IR was positively correlated with the changes in DHEA, A4, and testosterone concentrations during the
15 follow-up in girls. These results together suggest that physical activity and dietary interventions may
16 slow the development of adrenarche, especially in girls, and that this may be partly explained by
17 enhanced insulin sensitivity.

18 Previous studies in children have taught us that the nutritional and weight status are drivers for the
19 timing and intensity of adrenarche and puberty. For example, children with premature adrenarche are
20 more likely to be overweight or obese than those with later timing of adrenarche (5), and childhood
21 obesity is associated with earlier pubertal onset (6,10). These two developmental events seem to be
22 linked as children with premature adrenarche have more advanced pubertal development at the age of
23 12 years than those with normal timing of adrenarche (3). We and other research groups have also
24 provided evidence that some dietary factors, such as increased protein intake, are associated with

1 circulating adrenal androgen concentrations in children (29, 30). Our finding that the lifestyle
2 intervention attenuated the increase of serum androgens and decreased the prevalence of pubarche in
3 boys at the age of adrenarche suggests that changes in environmental factors, such as physical activity
4 and dietary factors, may have effects on the biological system regulating the timing and strength of the
5 sexual development in children. Our lifestyle intervention mainly affected serum androgens but not the
6 clinical signs of androgen action, except delaying pubarche in boys. The reason for this may be that the
7 duration of intervention of two years may have been too short given that approximately fifth of the
8 initially prepubertal and mostly normal-weight children had already entered adrenarche by baseline and
9 less than fourth of the children entered puberty over the next two years. It is also well known that the
10 onset of adrenarche and puberty is seen later in boys than in girls, which was also found in our study
11 cohort. Therefore, another explanation for observing the effect of our lifestyle intervention only on
12 pubarche in boys but not on other clinical signs of androgen action in either sex is that the intervention
13 started when many girls had already reached adrenarche.

14 The strengths of our study include the relatively large general population of initially prepubertal and
15 mostly normal-weight children examined, the long-term and controlled lifestyle intervention conducted,
16 the sensitive LC-MS/MS method used to measure most serum androgens, and the comprehensive
17 assessments of clinical androgenic and pubertal signs. Serum DHEAS was not measured by the LC-
18 MS/MS method, but it was well measurable by an immunoassay as it is abundant in the circulation
19 already in mid-childhood. Body size and composition in our cohort were comparable to those of the
20 national reference population (19) making it possible to generalize the results to other children of the
21 same age in Finland. We emphasized the individual needs of the families and parental involvement in
22 our intervention, both of which have been observed to improve adherence of families to lifestyle
23 interventions (31). Thus, only about 15% of the children in the intervention group dropped out during
24 the 2-year follow-up, and almost 90% of the children and their parents or caregivers participated in all

1 physical activity and dietary counselling sessions, indicating that the intervention was well accepted by
2 the participants.

3 A limitation of the study is that we did not randomly allocate the participants to the intervention and
4 control group, but instead allocated the children from nine schools to the intervention group and the
5 children from seven schools to the control group to avoid contamination in the control group by local or
6 national health promotion programs that could have been initiated in the study region during the
7 follow-up period. This type of allocation of the children to the study groups also enabled us to organize
8 after-school exercise clubs as part of the intervention at the nine-school premises and thus avoid a non-
9 intentional intervention in the control group. We matched the intervention and control groups
10 according to the location of the schools so that children from urban and rural areas were included in
11 both groups to minimize sociodemographic differences between the groups. There were only minor
12 differences in baseline characteristics between the intervention and control groups, suggesting fair
13 success in avoiding selection bias. Our individualized and family-based lifestyle intervention consisted
14 mainly of physical activity counselling for the children and their parents or caregivers, whereas providing
15 the children with supervised exercise played a much smaller role in our intervention. This kind of
16 physical activity intervention could be seen as a limitation of our study. However, we found that total
17 and unsupervised physical activity increased in the intervention group but decreased in the control
18 group, showing that this kind of intervention was able to increase physical activity among these children.
19 It is possible in an unblinded study design, which we used, that knowing the study group has an
20 influence on participants when answering the questionnaire or on investigators when performing the
21 assessments. Another limitation is the lack of data on circulating sex hormone-binding globulin
22 concentrations that the lifestyle intervention could have altered and thereby affected the bioavailability
23 of circulating androgens (32). One may question the reliability of the clinical assessment of adult-type
24 body odor and greasiness of hair and skin. Although our experience indicates that these signs of

1 androgen action are well detectable in clinical examinations and that parents usually recognize the
2 onset of these signs, we do agree that they are weak indicators of clinical adrenarche. We also agree
3 that defining biochemical adrenarche as a serum DHEAS concentration of $\geq 1 \mu\text{mol/l}$, which is commonly
4 used for this purpose, has its limitations because serum DHEAS concentrations are sexually dimorphic
5 and do not necessarily correlate straightforwardly with clinical signs of androgen action (33, 34).

6 To conclude, we found that the 2-year combined physical activity and dietary intervention in mid-
7 childhood attenuated the increase of serum androgen concentrations in a general population of initially
8 prepubertal and mostly normal-weight children, and that the intervention delayed pubarche in boys.
9 These intervention effects on serum androgens were independent of changes in body size and
10 composition but may have been mediated through altered insulin sensitivity, especially in girls. Future
11 studies should consider the limitations of our study and confirm its results, preferably with supervised
12 exercise interventions starting even at younger ages. Our findings highlight the importance of targeting
13 physical activity and dietary interventions at children with hyperandrogenic conditions, including those
14 with a normal body weight. Our findings also emphasize that a healthy diet and a physically active
15 lifestyle in childhood might prevent conditions like premature adrenarche, which, at least in some cases,
16 may precede later unfavorable health outcomes (35,36).

1 **Acknowledgements:**

2 We are grateful to all children and their parents and caregivers who have participated in the PANIC
3 study. We are also indebted to all members of the PANIC research team for their invaluable contribution
4 in the acquisition of the data throughout the study. We also want to thank Biocenter Finland and
5 Biocenter Kuopio for supporting their core LC-MS laboratory facility.

6 **Contribution statement:**

7 TAL and JJ designed the study. TAL and AM conducted the study. Steroid profiling with the LC-MS/MS
8 method was performed by MRH, SA, JJ, and RV. JL planned and performed the statistical analysis. JL,
9 CEF, JJ, and TAL interpreted the results. JL drafted the manuscript. All authors critically revised the
10 manuscript for its intellectual content and approved the final version of the manuscript. TAL is the
11 principal investigator of the PANIC study.

12 **Data availability:**

13 Information about the PANIC study and the data used in the present paper are available
14 at www.panicstudy.fi/en/etusivu. The data are not publicly available due to research ethical reasons and
15 because the owner of the data is the University of Eastern Finland and not the research group. However,
16 the corresponding author can provide further information on the PANIC study and the PANIC data on a
17 reasonable request.

19 **References**

- 20 1. Rosenfield RL. Normal and premature adrenarche. *Endocr Rev.* 2021;42(6):783-814.
- 21 2. Livadas S, Chrousos GP. Control of the onset of puberty. *Curr Opin Pediatr.* 2016;28(4):551-8.

- 1 3. Liimatta J, Utriainen P, Voutilainen R, Jääskeläinen J. Girls with a history of premature
2 adrenarche have advanced growth and pubertal development at the age of 12 years. *Front*
3 *Endocrinol (Lausanne)*. 2017;8:291.
- 4 4. Santos-Silva R, Fontoura M, Severo M, Santos AC. Dehydroepiandrosterone sulphate levels at 7
5 years old are positively associated with more advanced pubertal development between 10 and
6 13 years old in girls. *Clin Endocrinol (Oxf)*. 2022;97(6):747-54.
- 7 5. Utriainen P, Jääskeläinen J, Romppanen J, Voutilainen R. Childhood metabolic syndrome and its
8 components in premature adrenarche. *J Clin Endocrinol Metab*. 2007;92(11):4282-5.
- 9 6. Akslaegre L, Juul A, Olsen LW, Sørensen TIA. Age at puberty and the emerging obesity epidemic.
10 *PLoS One*. 2009;4(12):e8450.
- 11 7. Utriainen P, Voutilainen R, Jääskeläinen J. Girls with premature adrenarche have accelerated
12 early childhood growth. *J Pediatr*. 2009;154(6):882-7.
- 13 8. Dunger BD, Ahmed ML, Ong KK. Early and late weight gain and the timing of puberty. *Mol Cell*
14 *Endocrinol*. 2006;254-255:140-5.
- 15 9. Ong K, Potau N, Petry CJ, Jones R, Ness AR, Honour JW, de Zegher F, Ibáñez L, Dunger DB.
16 Opposing influences of prenatal and postnatal weight gain on adrenarche in normal boys and
17 girls. *J Clin Endocrinol Metab*. 2004;89(6):2647-51.
- 18 10. Li W, Liu Q, Deng X, Chen Y, Liu S, Story M. Association between obesity and pubertal timing: a
19 systematic review and meta-analysis. *Int J Environ Res Public Health*. 2017;14(10):1266.
- 20 11. Goran MI, Gower BA. Longitudinal study on pubertal insulin resistance. *Diabetes*.
21 2001;50(11):2444-50.
- 22 12. Ho M, Garnett SP, Baur LA, Burrows T, Stewart L, Neve M, Collins C. Impact of dietary and
23 exercise interventions on weight change and metabolic outcomes in obese children and

- 1 adolescents: a systematic review and meta-analysis of randomized trials. *JAMA pediatr.*
2 2013;167(8):759-68.
- 3 13. Reinehr T, Kulle A, Wolters B, Lass N, Welzel M, Riepe F, Holterhus PM. Steroid hormone profiles
4 in prepubertal obese children before and after weight loss. *J Clin Endocrinol Metab.*
5 2013;98(6):E1022-30.
- 6 14. Abdollahian S, Tehrani FR, Amiri M, Ghodsi D, Yarandi RB, Jafari M, Majd HA, Nahidi F. Effect of
7 lifestyle modifications in anthropometric, clinical, and biochemical parameters in adolescent
8 girls with polycystic ovary syndrome: a systematic review and meta-analysis. *BMC Endocr*
9 *Disord.* 2020;20(1):71.
- 10 15. Shele G, Genkil J, Speelman D. A systematic review of the effects of exercise on hormones in
11 women with polycystic ovary syndrome. *J Funct Morphol Kinesiol.* 2020;5(2):35.
- 12 16. Lakka T, Lintu N, Väistö J, Viitasalo A, Sallinen T, Haapala EA, Tompuri TT, Soininen S, Karjalainen
13 P, Schnurr TM, Mikkonen S, Atalay M, Kilpeläinen TO, Laitinen T, Laaksonen DE, Savonen K,
14 Brage S, Schwab U, Jääskeläinen J, Lindi V, Eloranta AM. A 2 year physical activity and dietary
15 intervention attenuates the increase in insulin resistance in a general population of children: the
16 PANIC study. *Diabetologia.* 2020;63(11):2270-81.
- 17 17. Viitasalo A, Eloranta AM, Lintu N, Väistö J, Venäläinen T, Kiiskinen S, Karjalainen P, Pe Itola J,
18 Lampinen EK, Haapala EA, Paananen J, Schwab U, Lindi V, Lakka TA. The effects of a 2-year
19 individualized and family-based lifestyle intervention on physical activity, sedentary behavior
20 and diet in children. *Prev Med.* 2016;87:81–8.
- 21 18. Venäläinen TM, Viitasalo AM, Schwab US, Eloranta AM, Haapala EA, Jalkanen HP, de Mello VD,
22 Laaksonen DE, Lindi VI, Ågren JJ, Lakka TA. Effect of a 2-y dietary and physical activity
23 intervention on plasma fatty acid composition and estimated desaturase and elongase activities

- 1 in children: the Physical Activity and Nutrition in Children Study. *Am J Clin*
2 *Nutr.* 2016;104(4):964–72.
- 3 19. Saari A, Sankilampi U, Hannila ML, Kiviniemi V, Kesseli K, Dunkel L. New Finnish growth
4 references for children and adolescents aged 0 to 20 years: length/height-for-age, weight-for-
5 length/height, and body mass index-for-age. *Ann Med.* 2011;43(3):235-48.
- 6 20. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis*
7 *Child.* 1969;44(235):291–303.
- 8 21. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. *Arch Dis*
9 *Child.* 1970;45(239):13–23.
- 10 22. Sankilampi U, Hannila ML, Saari A, Gissler M, Dunkel L. New population-based reference for
11 birth weight, length, and head circumference in singletons and twins from 23 to 43 gestation
12 weeks. *Ann Med.* 2013;45(5-6):446-54.
- 13 23. Häkkinen MR, Heinosalo T, Saarinen N, Linnanen T, Voutilainen R, Lakka T, Jääskeläinen J,
14 Poutanen M, Auriola S. Analysis by LC-MS/MS of endogenous steroids from human serum,
15 plasma, endometrium and endometriotic tissue. *J Pharm Biomed Anal.* 2018;152:165-72.
- 16 24. Rege J, Turcu AF, Kasa-Vubu JZ, Lerario AM, Auchus GC, Auchus RJ, Smith JM, White PC, Rainey
17 WE. 11-Ketotestosterone is the dominant circulating bioactive androgen during normal and
18 premature adrenarche. *J Clin Endocrinol Metab.* 2018;103(12):4589-98.
- 19 25. Harrington J, Palmert MR, Hamilton J. Use of local data to enhance uptake of published
20 recommendations: an example from the diagnostic evaluation of precocious puberty. *Arch Dis*
21 *Child.* 2014;99(1):15-20.
- 22 26. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model
23 assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin
24 concentrations in man. *Diabetologia.* 1985;28(7):412-9.

- 1 27. Reinehr T, Sánchez-Guijo A, Lass N, Wudy SA. Higher steroid sulfation is linked to successful
2 weight loss in obese children. *Endocr Connect*. 2018;7(10):1020-30.
- 3 28. Domecq JP, Prutsky G, Mullan RJ, Hazem A, Sundaresh V, Elamin MB, Phung OJ, Wang A, Hoeger
4 K, Pasquali R, Erwin P, Bodde A, Montori VM, Murad MH. Lifestyle modification programs in
5 polycystic ovary syndrome: systematic review and meta-analysis. *J Clin Endocrinol Metab*.
6 2013;98(12):4655-63.
- 7 29. Shi L, Wudy SA, Buyken AE, Hartmann MF, Remer T. Body fat and animal protein intakes are
8 associated with adrenal androgen secretion in children. *Am J Clin Nutr*. 2009;90(5):1321-8.
- 9 30. Mäntyselkä A, Jääskeläinen J, Eloranta AM, Väistö J, Voutilainen R, Ong K, Brage S, Lakka TA,
10 Lindi V. Associations of lifestyle factors with serum dehydroepiandrosterone sulphate and
11 insulin-like growth factor-1 concentration in prepubertal children. *Clin Endocrinol (Oxf)*.
12 2018;88(2):234-42.
- 13 31. Showel NN, Fawole O, Segal J, Wilson RF, Cheskin LJ, Bleich SN, Wu Y, Lau B, Wang Y. A
14 systematic review of home-based childhood obesity prevention studies. *Pediatrics*.
15 2013;132(1):193-200.
- 16 32. Aroda VR, Christophi CA, Edelstein SL, Perreault L, Kim C, Golden SH, Horton E, Mather KJ.
17 Circulating sex hormone binding globulin levels are modified with intensive lifestyle
18 intervention, but their changes did not independently predict diabetes risk in the Diabetes
19 Prevention Program. *BMJ Open Diabetes Res Care*. 2020;8(2):e001841.
- 20 33. Mäntyselkä A, Jääskeläinen J, Lindi V, Viitasalo A, Tompuri T, Voutilainen R, Lakka TA. The
21 presentation of adrenarche is sexually dimorphic and modified by body adiposity. *J Clin
22 Endocrinol Metab*. 2014;99(10):3889-94.
- 23 34. Utriainen P, Voutilainen R, Jääskeläinen J. Continuum of phenotypes and sympathoadrenal
24 function in premature adrenarche. *Eur J Endocrinol*. 2009;160(4):657-65.

1 35. Liimatta J, Utriainen P, Laitinen T, Voutilainen R, Jääskeläinen J. Cardiometabolic risk profile
2 among young adult females with a history of premature adrenarche. *J Endocr Soc.*
3 2019;3(10):1771-83.

4 36. Tennilä J, Jääskeläinen J, Utriainen P, Voutilainen R, Häkkinen M, Auriola S, Morin-Papunen L,
5 Liimatta J. PCOS features and steroid profiles among young adult women with a history of
6 premature adrenarche. *J Clin Endocrinol Metab.* 2021;106(9):e3335-45.

7
8 **Figure 1.** Flow chart of the study.

9 **Figure 2.** Absolute change (Δ) in height SDS (panel A), growth velocity (GV, panel B), body mass index
10 SDS (BMI, panel C), and body fat percentage (panel D) over 2-years among children in the combined
11 physical and dietary intervention group and among children in the control group. The mean age at
12 baseline was 7.6 years. Medians and interquartile ranges (IQRs) in boxes, ranges (IQR*1.5) in whiskers,
13 and outliers (circles) are provided in boxplot charts. No statistically significant differences were found
14 between the groups in any of these parameters (Mann-Whitney U test, p-values > 0.140), also when
15 analyzing girls and boys separately.

16 **Figure 3.** Relative unadjusted changes (%) in serum dehydroepiandrosterone sulfate (DHEAS; panels A-
17 C), dehydroepiandrosterone (DHEA; D-F), androstenedione (A4; G-I), and testosterone (J-L)
18 concentrations among children in the combined physical activity and dietary intervention group and
19 among children in the control group over two years. The mean age at baseline was 7.6 years. Medians
20 and interquartile ranges (IQRs) in boxes, ranges (IQR*1.5) in whiskers, and outliers (circles) are provided
21 in boxplot charts. Differences between the groups were analyzed using the Mann-Whitney U test, and p-
22 values were used to indicate statistical significance of the differences.

Table 1. Clinical data at baseline and 2-year follow-up examinations among children in the 2-year combined physical activity and dietary intervention group and among children in the control group.

	All children			Girls			Boys		
	Intervention (n = 251)	Control (n = 170)	P	Intervention (n = 119)	Control (n = 84)	P	Intervention (n = 132)	Control (n = 86)	P
At birth									
Gestational age, wk	39.9 (1.5)	39.8 (1.7)	0.520	39.7 (1.6)	39.7 (1.8)	0.953	39.9 (1.3)	39.8 (1.5)	0.374
Birth length SDS	-0.07 (1.03)	-0.36 (0.87)	0.003	-0.03 (1.05)	-0.28 (0.89)	0.084	-0.10 (1.02)	-0.43 (0.85)	0.014
Birth weight SDS	-0.02 (1.08)	-0.19 (0.92)	0.090	0.06 (1.10)	-0.18 (0.99)	0.123	-0.09 (1.05)	-0.21 (0.85)	0.390
Small for gestational age, % ^a	5	7	0.406	3	9	0.206	7	6	0.818
Preterm, % ^b	4	5	0.633	7	5	0.764	2	6	0.115
At baseline examination									
Age, y	7.6 (0.3)	7.6 (0.4)	0.733	7.6 (0.3)	7.5 (0.4)	0.095	7.6 (0.3)	7.7 (0.4)	0.269
Height SDS	0.17 (0.95)	0.09 (1.03)	0.388	0.10 (0.87)	0.09 (1.10)	0.937	0.24 (1.02)	0.09 (0.95)	0.274
Growth velocity, cm/y ^c	6.73 (0.89)	6.71 (0.71)	0.719	6.59 (0.64)	6.69 (0.66)	0.285	6.86 (1.06)	6.72 (0.75)	0.323
Body mass index SDS	-0.20 (1.04)	-0.24 (1.14)	0.736	-0.22 (1.02)	-0.37 (1.07)	0.333	-0.18 (1.07)	-0.11 (1.19)	0.651
Waist circumference, cm	56.4 (5.3)	56.6 (6.2)	0.742	55.5 (4.8)	55.3 (6.0)	0.711	57.2 (5.8)	57.9 (6.2)	0.383
Body fat percentage	19.5 (7.9)	19.6 (8.0)	0.958	21.9 (6.9)	20.9 (7.6)	0.337	17.3 (8.1)	18.2 (8.2)	0.447
Lean mass percentage	76.6 (8.4)	76.6 (8.5)	0.988	73.9 (7.3)	75.1 (8.0)	0.304	79.0 (8.6)	78.0 (8.7)	0.443
VAT-to-body fat mass, %	1.54 (1.25)	1.63 (1.30)	0.496	0.88 (0.65)	0.97 (0.74)	0.337	2.15 (1.36)	2.29 (1.40)	0.468
Clinical adrenarache, % ^d	16	18	0.502	24	23	0.816	8	14	0.171
Biochemical adrenarache, % ^e	19	18	0.869	20	17	0.611	17	19	0.794
Clinical and/or biochemical adrenarache, % ^f	31	31	0.989	36	32	0.570	26	29	0.585
Pubarache, % ^g	0	1	0.568	1	2	0.571	0	0	0.999
At 2-year follow-up examination									
Age, y	9.7 (0.4)	9.7 (0.5)	0.838	9.8 (0.4)	9.7 (0.5)	0.150	9.8 (0.4)	9.8 (0.5)	0.277
Height SDS	0.10 (0.95)	0.04 (1.04)	0.538	0.03 (0.86)	0.02 (1.10)	0.924	0.17 (1.02)	0.07 (0.99)	0.470
Growth velocity, cm/y ^h	5.43 (0.68)	5.48 (0.71)	0.514	5.56 (0.75)	5.55 (0.74)	0.970	5.32 (0.60)	5.40 (0.67)	0.341
Body mass index SDS	-0.17 (1.04)	-0.13 (1.08)	0.687	-0.19 (1.00)	-0.23 (1.02)	0.795	-0.15 (1.07)	-0.03 (1.14)	0.427
Waist circumference, cm	61.0 (6.9)	61.2 (7.7)	0.369	59.6 (6.1)	59.5 (7.5)	0.867	62.2 (7.4)	63.0 (7.6)	0.480
Body fat percentage	24.2 (9.6)	24.4 (9.7)	0.893	26.6 (8.7)	25.1 (8.8)	0.261	22.0 (10.0)	23.7 (10.6)	0.264
Lean mass percentage	72.7 (9.4)	72.6 (9.6)	0.904	70.4 (8.5)	71.8 (8.6)	0.252	74.9 (9.8)	73.3 (10.4)	0.263
VAT-to-body fat mass, %	1.39 (0.99)	1.35 (0.94)	0.679	0.90 (0.61)	0.98 (0.68)	0.420	1.85 (1.07)	1.71 (1.01)	0.346
Clinical adrenarache, % ^c	63	61	0.704	73	63	0.174	54	58	0.506
Biochemical adrenarache, % ^d	29	33	0.363	26	30	0.585	31	36	0.433
Clinical and/or biochemical adrenarache, % ^e	67	72	0.267	73	73	0.955	62	72	0.166
Pubarache, % ^f	8	10	0.318	15	15	0.989	1	6	0.038
Puberty, only clinical signs, % ^g	20	21	0.939	27	35	0.259	14	6	0.107
Puberty, clinical signs and/or biochemical evidence, % ⁱ	24	29	0.300	28	41	0.057	21	17	0.494

Continuous variables are expressed as means (standard deviations) and categorical variables as percentages. Differences between the intervention and control groups were analyzed using the student t-test for continuous variables and the χ^2 test for categorical variables, and p-values < 0.05 indicating statistically significant differences between the groups are bolded.

^a Birth length and/or birth weight ≤ -2 SDS; ^b Gestational age < 37.0 weeks; ^c Growth velocity between the age of five years and baseline examination; ^d At least one of the following signs: adult-type body odor, greasiness of hair and skin, comedones/acne, and axillary/pubertic hair; ^e serum DHEAS concentration $\geq 1 \mu\text{mol/L}$; ^f At least one clinical sign of adrenarche and/or serum DHEAS concentration $\geq 1 \mu\text{mol/L}$; ^g Tanner P ≥ 2 ; ^h growth velocity between baseline and follow-up examination; ⁱ Tanner B for girls or G for boys ≥ 2 and/or serum luteinizing hormone ≥ 0.3 U/L. SDS, standard deviation score; VAT, visceral adipose tissue mass.

Table 2. Biochemical data at baseline and 2-year follow-up examinations among children in the 2-year combined physical activity and dietary intervention group and among children in the control group.

	All children			Girls			Boys		
	Intervention (n = 250)	Control (n = 170)	p	Intervention (n = 119)	Control (n = 84)	p	Intervention (n = 131)	Control (n = 86)	p
At baseline examination (6-9 y)									
DHEA, nmol/L	1.70 (0.86-2.91)	1.33 (0.81-2.57)	0.313	1.92 (1.02-3.34)	1.52 (0.84-2.72)	0.382	1.40 (0.73-2.49)	1.24 (0.67-2.07)	0.44
DHEAS, $\mu\text{mol/L}$	0.59 (0.33-0.84)	0.56 (0.31-0.86)	0.505	0.57 (0.35-0.84)	0.49 (0.31-0.85)	0.457	0.60 (0.32-0.84)	0.58 (0.32-0.88)	0.84
A4, nmol/L	0.74 (0.47-1.07)	0.67 (0.40-0.99)	0.062	0.82 (0.47-1.21)	0.74 (0.44-1.10)	0.306	0.63 (0.46-0.98)	0.62 (0.35-0.87)	0.08
Testosterone, pmol/L	122 (83-168)	117 (76-178)	0.689	133 (92-194)	124 (92-190)	0.712	115 (79-154)	107 (71-174)	0.78
Insulin, U/L	4.2 (2.7-5.7)	4.2 (2.9-6.0)	0.466	4.7 (3.4-6.0)	4.3 (3.3-6.1)	0.736	3.9 (2.5-5.5)	4.0 (2.6-6.0)	0.23
HOMA-IR	0.91 (0.55-1.27)	0.91 (0.64-1.34)	0.525	1.00 (0.68-1.32)	0.92 (0.70-1.29)	0.630	0.85 (0.53-1.19)	0.89 (0.55-1.36)	0.23
IGF-1, nmol/L	21.7 (17.5-25.9)	22.9 (18.8-28.3)	0.013	22.5 (18.5-28.1)	24.1 (19.4-30.1)	0.264	20.2 (16.2-24.5)	22.1 (18.4-27.4)	0.01
At 2-year follow-up examination (9-11 y)									
DHEA, nmol/l	3.06 (1.71-4.75)	3.99 (2.13-6.49)	0.003	3.47 (1.93-5.23)	4.22 (2.70-6.76)	0.033	2.78 (1.57-4.42)	3.27 (2.00-6.38)	0.04
DHEAS, $\mu\text{mol/l}$	0.69 (0.38-1.05)	0.67 (0.42-1.19)	0.449	0.67 (0.36-1.02)	0.63 (0.36-1.02)	0.819	0.74 (0.40-1.10)	0.74 (0.42-1.22)	0.37
A4, nmol/l	1.20 (0.79-1.70)	1.20 (0.80-1.84)	0.613	1.36 (0.92-1.98)	1.33 (0.92-2.06)	0.991	1.07 (0.73-1.52)	1.07 (0.70-1.70)	0.63
Testosterone, pmol/l	202 (135-291)	206 (148-313)	0.383	250 (156-347)	215 (154-328)	0.540	185 (123-241)	197 (127-277)	0.12
Insulin, U/L	4.9 (3.5-7.2)	5.9 (3.9-7.9)	0.028	5.4 (4.0-8.2)	6.3 (4.1-8.2)	0.167	4.6 (3.1-6.6)	5.1 (3.5-7.4)	0.12
HOMA-IR	1.10 (0.77-1.61)	1.27 (0.84-1.76)	0.034	1.13 (0.88-1.81)	1.38 (0.87-1.84)	0.208	1.03 (0.68-1.50)	1.16 (0.76-1.71)	0.11
IGF-1, nmol/L	29.8 (24.1-35.8)	29.7 (24.3-35.6)	0.942	33.2 (26.8-40.3)	32.1 (26.8-40.3)	0.586	26.2 (22.9-33.5)	28.2 (22.9-32.1)	0.82

The values are medians (25th-75th percentile range). HOMA-IR was calculated using the following formula: [(insulin (mU/L) x glucose (mmol/L)) / 22.5]. Differences between the intervention and control groups were analyzed using the Mann-Whitney U test, and p-values < 0.05 indicating statistically significant differences between the groups are bolded. To have values in conventional units, multiply DHEA by 0.2884 (ng/mL), DHEAS by 36.85 ($\mu\text{g/dL}$), A4 by 0.2864 (ng/mL), testosterone by 0.2885 (pg/mL), and IGF-1 by 7.65 (ng/mL).

A4, androstenedione; DHEA, dehydroepiandrosterone; DHEAS, dehydroepiandrosterone sulfate; HOMA-IR, homeostatic model assessment for insulin resistance; IGF-1, insulin-like growth factor 1.

Table 3. Effects of the 2-year combined physical activity and dietary intervention on changes (Δ) in serum dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEAS), androstenedione (A4), and testosterone concentrations.

	Outcome (dependent) variable							
	Δ DHEA (nmol/l)		Δ DHEAS (μ mol/l)		Δ A4 (nmol/l)		Δ Testosterone (pmol/l)	
	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Model 1:								
All children (n = 421)	-1.23 (-1.85 to -0.62)	< 0.001	-0.05 (-0.14 to 0.04)	0.292	-0.12 (-0.27 to 0.03)	0.111	-121 (-237 to -5.27)	0.041
Girls only (n = 203)	-1.27 (-2.11 to -0.43)	0.003	-0.07 (-0.18 to 0.05)	0.244	-0.02 (-0.26 to 0.21)	0.855	22.9 (-9.24 to 53.1)	0.167
Boys only (n = 218)	-1.31 (-2.24 to -0.37)	0.007	-0.05 (-0.19 to 0.09)	0.503	-0.19 (-0.38 to 0.01)	0.063	-253 (-476 to -29.5)	0.027
Model 2:								
All children (n = 421)	-1.14 (-1.77 to -0.52)	< 0.001	-0.07 (-0.16 to 0.03)	0.169	-0.11 (-0.26 to 0.04)	0.156	-124 (-243 to -4.36)	0.042
Girls only (n = 203)	-1.20 (-2.07 to -0.34)	0.007	-0.07 (-0.19 to 0.04)	0.215	-0.04 (-0.24 to 0.25)	0.972	24.0 (-8.35 to 56.4)	0.145
Boys only (n = 218)	-1.15 (-2.09 to -0.22)	0.016	-0.05 (-0.20 to 0.09)	0.467	-0.19 (-0.39 to 0.01)	0.058	-256 (-485 to -28.0)	0.028

Values are from linear regression models adjusted for gestational age, birth length SDS, age, serum IGF-1 concentration, BMI SDS, and presence of adrenarchal signs at baseline, and presence of pubertal signs at 2-year follow-up (Model 1) and additionally for the change of fasting serum insulin concentration over two years (Model 2). The mean age at baseline was 7.6 years. P-values < 0.05 indicating statistically significant differences between the groups are bolded.




Figure 1
259x162 mm (x DPI)




Figure 2
259x163 mm (x DPI)




Figure 3
245x196 mm (x DPI)