



University of HUDDERSFIELD

University of Huddersfield Repository

Redmond, A.C, Hain, J and Bird, H.A

Distribution and effect of joint pain in people with hypermobility syndrome: pilot results By Redmond, A.C., Hain,J (married name Tranter)Bird, H.A.

Original Citation

Redmond, A.C, Hain, J and Bird, H.A (2006) Distribution and effect of joint pain in people with hypermobility syndrome: pilot results By Redmond, A.C., Hain,J (married name Tranter)Bird, H.A. *Annals of the Rheumatic Diseases*, 65 (SII). p. 241. ISSN 0003-4967

This version is available at <http://eprints.hud.ac.uk/5793/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Poster Presentations (THU0417)

Session: Back pain, mechanical musculoskeletal problems, local soft tissue disorders

Thursday, 22 June 2006

DISTRIBUTION AND EFFECT OF JOINT PAIN IN PEOPLE WITH HYPERMOBILITY SYNDROME: PILOT RESULTS

A.C. Redmond, J. Hain, H.A. Bird

Academic Unit of Musculoskeletal Disease, University of Leeds, Leeds, United Kingdom

Background: Joint hypermobility syndrome (HMS) is thought to be associated with increased pain and reduced health status and quality of life. Good quantitative data are rare however, and no studies exist comparing HMS patients with directly matched controls.

Objectives: The aim of the current study was to pilot a large national prevalence study of the effects of HMS on joint pain and health status.

Methods: Thirty-nine people with a diagnosis of hypermobility completed survey forms covering general health status (EuroQoL EQ5D, Manchester Foot Pain and Disability Questionnaire), foot pain (Visual Analogue Scale) and distribution of symptoms. Age and gender matched controls completed a modified version of the same survey forms.

Results: The sample comprised 39 hypermobiles (36F, 3M) and 37 controls (28F, 9M). The mean ages were 38 years (HMS) and 36 years (controls). Within the HMS group 16 (41%) knew their HMS type to be benign familial, three reported a diagnosis of EDS type 1, and five EDS type 3. Fifteen did not know.

People with HMS had significantly worse general health measured by EQ-5D (65pts vs 90**), much greater foot impairment (MFPDQ scores 16/33 vs 0/33**) and greater foot pain VAS scores (34mm vs 0.4mm**)(where ** represents $P < 0.001$).

Respondents provided data on Likert-type scales reporting frequency of symptoms in various joints and limb segments. The measures employed five point scales labelled with ranks "never", "seldom", "sometimes", "much of the time", and "constantly".

With the exception of headache, people with HMS reported significantly higher rates of symptoms in all axial and upper limb segments or joints. The most commonly affected joints were the shoulders, hands and lower back (see Table).

Frequency and location of symptoms

N (%) of HMS group	reporting	Median score HMS	Median score control	symptoms
[pic]	'sometimes'	group (+ descriptor)	group (+ descriptor)	Neck 27 (69%) 2 (sometimes)
1 (seldom)**	Shoulders 30 (77%)	3 (much of the time)	0 (never)	** Elbows 18 (46%) 1

(seldom) 0 (never) ** Hands 31 (79%) 3 (much of the time) 0 (never) ** Lower back 29 (74%) 3 (much of the time) 1 (seldom) ** Hips 30 (77%) 3 (much of the time) 0 (never) ** Knees 34 (87%) 3 (much of the time) 0 (never) ** Ankles 29 (74%) 3 (much of the time) 0 (never) ** Feet 28 (72%) 3 (much of the time) 0 (never) **

[pic]

|**Significantly different, adjusted P<0.001 |
|In the lower limb the hips, knees, ankles and feet all returned |
|medians of three indicating impairment "much of the time". All |
|responses were significantly worse than those of the control group. |
|Conclusion: Nearly 9/10 people with HMS report joint symptoms, most |
|commonly affecting the knees, hands, shoulders, wrists and hips. |
|Symptoms associated with HMS were significantly more frequent than in |
|the control group and were associated with increased pain and |
|significantly diminished health status in people with HMS. |
|[pic] |