



# Sarcoidosis mimicking nodal manifestations of marginal zone lymphoma

Johanna S. Enke<sup>1</sup> · Nic G. Reitsam<sup>2</sup> · Tina Schaller<sup>2</sup> · Rainer Claus<sup>2,3</sup> · Constantin Lapa<sup>1</sup> · Alexander Dierks<sup>1</sup>

Received: 19 December 2022 / Accepted: 21 January 2023  
© The Author(s) 2023

A 54-year-old man with newly diagnosed, histologically confirmed intraorbital marginal zone lymphoma (MZL) was referred for initial staging of disease. Whole-body <sup>18</sup>F-fluorodesoxyglucose (<sup>18</sup>F)FDG PET/CT was performed and showed only moderate uptake of the primary manifestation (A, red arrow) and mediastinal lymph nodes (A).

As [<sup>18</sup>F]FDG has limited sensitivity in the staging of MZL [1], an additional C-X-C motif chemokine receptor 4 (CXCR4)-directed PET/CT scan using [<sup>68</sup>Ga]Ga-PentixaFor ([<sup>68</sup>Ga]Ga-CPCR4.2) was performed, given that CXCR4 is overexpressed by most B- and T-cell neoplasms [2, 3]. Chemokine receptor-directed imaging demonstrated high tracer uptake of the intraorbital MZL-manifestation (B, red arrow) as well as multiple lymph nodes of the neck and thorax, the latter being rated as possible nodal MZL manifestations (B). Transbronchial fine-needle aspiration of a paratracheal lymph node (red star; [<sup>18</sup>F]FDG, SUV<sub>max</sub> = 4.77; [<sup>68</sup>Ga]Ga-CPCR4.2, SUV<sub>max</sub> = 7.18)

revealed no signs of lymphoma infiltration but characteristic epithelioid cell granulomas with pronounced CXCR4-expression in the surrounding rim of activated lymphocytes (C), consistent with the diagnosis of sarcoidosis, a multisystem inflammatory disorder of enormous heterogeneity in clinical presentation [4].

Since CXCR4 is abundantly involved in immune cell activation and several inflammatory processes, and especially expressed by macrophages and T-lymphocytes [5], intense CXCR4-expression in sarcoid lesions is biologically reasonable. To our knowledge, this is one of the first reports on CXCR4 visualization in sarcoidosis by means of PET/CT. While sarcoidosis (as other inflammatory conditions) might represent a pitfall in oncologic imaging using CXCR4-directed PET tracers, non-invasive detection of receptor expression could also benefit the diagnostic workup of sarcoidosis, especially in cardiac- or neurosarcoidosis, and should be further evaluated.

---


This article is part of the Topical Collection on Image of the month.

✉ Constantin Lapa  
Constantin.Lapa@uk-augsburg.de

<sup>1</sup> Nuclear Medicine, Faculty of Medicine, University of Augsburg, Augsburg, Germany

<sup>2</sup> Pathology, Faculty of Medicine, University of Augsburg, Augsburg, Germany

<sup>3</sup> Internal Medicine, Faculty of Medicine, University of Augsburg, Augsburg, Germany



**Funding** Open Access funding enabled and organized by Projekt DEAL.

**Data Availability** The data that support the findings of this study are available from the corresponding author, [CL], upon reasonable request.

## Declarations

**Ethical approval** Informed consent from the patient for publication of this case study was obtained.

**Conflict of interest** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Duell J, Krummenast F, Schirbel A, Klassen P, Samnick S, Rauert-Wunderlich H, et al. Improved primary staging of marginal-zone lymphoma by addition of CXCR4-directed PET/CT. *J Nucl Med Off Publ Soc Nucl Med.* 2021;62(10):1415–21.
2. Griffith JW, Sokol CL, Luster AD. Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu Rev Immunol.* 2014;32:659–702.
3. Stollberg S, Kämmerer D, Neubauer E, Schulz S, Simonitsch-Klupp I, Kiesewetter B, et al. Differential somatostatin and CXCR4 chemokine receptor expression in MALT-type lymphoma of gastric and extragastric origin. *J Cancer Res Clin Oncol.* 2016;142(11):2239–47.
4. Chen ES, Moller DR. Sarcoidosis—scientific progress and clinical challenges. *Nat Rev Rheumatol.* 2011;7(8):457–67.
5. Salvatore P, Pagliarulo C, Colicchio R, Napoli C. CXCR4-CXCL12-dependent inflammatory network and endothelial progenitors. *Curr Med Chem.* 2010;17(27):3019–29.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.