



TITLE:

A Note on Canonical Transforms Representing $SL(2, \mathbb{R})$, a Two-Fold Covering of $SL(2, \mathbb{R})$, in $L^2(\mathbb{R})$

AUTHOR(S):

IWAI, Toshihiro; REW, See-Gew

CITATION:

IWAI, Toshihiro ...[et al]. A Note on Canonical Transforms Representing $SL(2, \mathbb{R})$, a Two-Fold Covering of $SL(2, \mathbb{R})$, in $L^2(\mathbb{R})$. *Memoirs of the Faculty of Engineering, Kyoto University* 1985, 47(4): 262-268

ISSUE DATE:

1985-10-31

URL:

<http://hdl.handle.net/2433/281305>

RIGHT:

A Note on Canonical Transforms Representing $SL_2(2, \mathbf{R})$, a Two-Fold Covering of $SL(2, \mathbf{R})$, in $L^2(\mathbf{R})$

By

Toshihiro IWAI and See-Gew REW

(Received June 29, 1985)

Abstract

Canonical transforms representing unitarily $SL_2(2, \mathbf{R})$, a two-fold covering of $SL(2, \mathbf{R})$, are explicitly constructed in the form of integral transforms in $L^2(\mathbf{R})$ with the main stress laid on the composition of canonical transforms.

1. Introduction

In his book¹⁾, K. B. Wolf treated in detail canonical transformations associated with $SL(2, \mathbf{R})$ in the form of integral transforms. His method, due to M. Moshinsky and C. Quesne²⁾, is a quantization of the symplectic action of $SL(2, \mathbf{R})$ on \mathbf{R}^2 , the phase space for a classical system of one degree of freedom. In the case of several degrees of freedom, the integral transforms are discussed in reference 3).

It was D. Shale⁴⁾ who observed that this quantization determines a double-valued unitary representation of the symplectic group. N. R. Wallach showed in his book⁵⁾ that this quantization determines a unitary representation of the metaplectic group, a two-fold covering of the symplectic group, which becomes known as the Segal-Shale-Weil representation⁶⁾.

The two-fold covering causes a sign problem in the integral kernel. Though the problem has been solved in an abstract manner, it seems to require further investigation in an explicit manner. The present article gives integral transforms for $SL_2(2, \mathbf{R})$ with the phase factors assigned strictly, and shows that the integral transforms actually constitute a unitary representation of $SL_2(2, \mathbf{R})$, by composing the integral transforms in an explicit form.

2. Canonical Transforms

In this section, we review quantum mechanical canonical transforms associated with $SL(2, \mathbf{R})$. Assume that a unitary operator U on $L^2(\mathbf{R})$ induces a linear trans-

formation of x and $y = -id/dx$;

$$U \begin{bmatrix} x \\ y \end{bmatrix} U^{-1} = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}, \quad ad - bc = 1. \quad (2.1)$$

This transformation corresponds to the canonical transformation on \mathbf{R}^2 by $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for the classical system. It is known¹⁾ that U is expressed in the integral transform with the integral kernel, for $b \neq 0$,

$$K(x, \xi) = \frac{\alpha}{(2\pi|b|)^{1/2}} \exp \left[\frac{i}{2b} (dx^2 - 2x\xi + a\xi^2) \right] \quad (2.2)$$

where α is a complex constant with $|\alpha|=1$. To determine α , one has to pass on to the two-fold covering $SL_2(2, \mathbf{R})$ of $SL(2, \mathbf{R})$. We will discuss this in the next section.

On supposing that α is determined in some way or other, we calculate compositions of the integral transforms. Let

$$M_{21} = M_2 M_1, \quad M_k = \begin{bmatrix} a_k & b_k \\ c_k & d_k \end{bmatrix}, \quad k=1, 2, 21. \quad (2.3)$$

By U_k , $k=1, 2, 21$, we mean the unitary operator determined by (2.1) with coefficient matrices M_k^{-1} . For the unitary operators U_k , the integral kernels and the phase factors are denoted by K_k and α_k , respectively. Performing an iterated integral of the kernels yields

$$U_2 U_1 = \begin{cases} \frac{\alpha_1 \alpha_2}{\alpha_{21}} e^{i\pi/4} U_{21} & \text{for } \frac{b_{21}}{b_1 b_2} > 0, \\ \frac{\alpha_1 \alpha_2}{\alpha_{21}} e^{-i\pi/4} U_{21} & \text{for } \frac{b_{21}}{b_1 b_2} < 0. \end{cases} \quad (2.4)$$

Our purpose is to show that for suitably chosen phase factors α_k , Eq. (2.4) becomes exactly $U_2 U_1 = U_{21}$, which means that U is a unitary representation of $SL_2(2, \mathbf{R})$.

3. One-Parameter Subgroups

To realize how the two-fold covering $SL_2(a, \mathbf{R})$ emerges in the integral transforms, we consider one-parameter groups of unitary transformations corresponding to an Iwasawa decomposition of $SL(2, \mathbf{R})$;

$$SL(2, \mathbf{R}) \simeq KAN,$$

or

$$\begin{aligned} \begin{bmatrix} a & b \\ c & d \end{bmatrix} &= \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} e^\tau & 0 \\ 0 & e^{-\tau} \end{bmatrix} \begin{bmatrix} 1 & \zeta \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} e^\tau \cos \theta & e^\tau \zeta \cos \theta + e^{-\tau} \sin \theta \\ -e^\tau \sin \theta & -e^\tau \zeta \sin \theta + e^{-\tau} \cos \theta \end{bmatrix}. \end{aligned} \quad (3.1)$$

The parameters (θ, τ, ζ) are given by

$$e^{i\theta} = \frac{a-ic}{(a^2+c^2)^{1/2}}, \quad e^\tau = (a^2+c^2)^{1/2}, \quad \zeta = \frac{ab+cd}{a^2+c^2}. \quad (3.2)$$

The one-parameter subgroup $K(\theta)$ is associated with the harmonic oscillator, $A(\tau)$ with the dilatation, and $N(\zeta)$ with a free particle. The $K(\theta)$, diffeomorphic with S^1 , causes the problem of covering, but $A(\tau)$ and $N(\zeta)$ do not. Let $e^{-i\theta H}$, $e^{-i\tau D}$ and $e^{-i\zeta F}$ denote the one-parameter groups of unitary operators corresponding to $K(\theta)$, $A(\tau)$, and $N(\zeta)$, respectively, where H , D , and F , are given, as is well known, by

$$\begin{aligned} H &= \frac{1}{2} \left(-\frac{d^2}{dx^2} + x^2 \right), \\ D &= \frac{1}{i} \left(x \frac{d}{dx} + 1/2 \right), \\ F &= -\frac{1}{2} \frac{d^2}{dx^2}. \end{aligned} \quad (3.3)$$

Then $e^{-i\theta H}$ should be a representation of a two-fold covering of $K(\theta)$, while $e^{-i\tau D}$ and $e^{-i\zeta F}$ are representation of $A(\tau)$ and $N(\zeta)$, respectively.

To fix an idea of covering, we look into $e^{-i\theta H}$. The integral kernel of $e^{-i\theta H}$ is given from (2.2) as

$$K_\theta(x, \xi) = \frac{\alpha(\theta)}{(2\pi|\sin\theta|)^{1/2}} \exp \left[\frac{i}{2\sin\theta} ((x^2 + \xi^2) \cos\theta - 2x\xi) \right]. \quad (3.4)$$

The phase factor $\alpha(\theta)$ should be determined under the condition that U_θ should be a one-parameter group of unitary operators with the initial value $U_0 = id$. Put another way, $\alpha(\theta)$ must be determined in each interval $n\pi < \theta < (n+1)\pi$, $n=0, \pm 1, \pm 2, \dots$, so that U_θ may be continuous at $n\pi$. As a consequence we obtain

$$\alpha(\theta) = e^{-i\pi/4} \beta(\theta), \quad \alpha(\theta + 4n\pi) = \alpha(\theta),$$

with

$$\beta(\theta) = \begin{cases} -1 & \text{for } -2\pi < \theta < -\pi, \\ i & \text{for } -\pi < \theta < 0, \\ 1 & \text{for } 0 < \theta < \pi, \\ -i & \text{for } \pi < \theta < 2\pi \end{cases} \quad (3.5)$$

and

$$K_\theta(x, \xi) \rightarrow \begin{cases} -\delta(x-\xi) & \text{as } \theta \rightarrow -2\pi, \\ i\delta(x+\xi) & \text{as } \theta \rightarrow -\pi, \\ \delta(x-\xi) & \text{as } \theta \rightarrow 0, \\ -i\delta(x+\xi) & \text{as } \theta \rightarrow \pi. \end{cases} \quad (3.6)$$

We notice here that $\alpha(\theta)$ is periodic in θ with period 4π , while $K(\theta)$ has period 2π . For the U_θ , we can prove that $U_s U_t = U_{s+t}$ by using (2.4).

The operators $e^{-i\tau D}$ and $e^{-i\zeta F}$ cause no problem of covering, and are known to have respective integral kernels

$$K_\tau(x, \xi) = e^{-\tau/2} \delta(e^{-\tau}x - \xi) \tag{3.7}$$

and

$$K_\zeta(x, \xi) = \frac{\alpha(\zeta)}{(2\pi|\zeta|)^{1/2}} \exp\left[\frac{i}{2\zeta}(x-\xi)^2\right] \tag{3.8}$$

with $\alpha(\zeta) = e^{-i\pi(s\theta\pi\zeta)/4}$.

In view of the Iwasawa decomposition (3.1), we infer that compositions $e^{-i\zeta F} e^{-i\tau D} e^{-i\theta H}$ will provide a unitary representation of $SL_2(2, \mathbf{R})$, where we consider (θ, τ, ζ) as the parameters of $SL_2(2, \mathbf{R})$ with θ ranging over the interval of length 4π .

4. A Representation of $SL_2(2, \mathbf{R})$

We are going to determine the factor α in the expression (2.2) for the parameters (θ, τ, ζ) of $SL_2(2, \mathbf{R})$.

Let

$$\alpha = e^{-i\pi/4}\beta. \tag{4.1}$$

For $e^{-i\theta H}$, β takes values $\pm 1, \pm i$, and for $e^{-i\zeta F}$, $\beta = 1$ or i , depending on the sign of b .

Let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ be a matrix assigned by the parameters (3.1). For the M , there are two elements in $SL_2(2, \mathbf{R})$ assigned by the parameters (θ, τ, ζ) and $(\theta \pm 2\pi, \tau, \zeta)$, the sign \pm depending on the value of θ . To determine β , we consider the sign of

$$b = \zeta e^\tau \cos \theta + e^{-\tau} \sin \theta \tag{4.2}$$

for τ and ζ fixed arbitrarily. On setting

$$\tan \chi_0 = -\zeta e^{2\tau} \quad \text{with } |\chi_0| < \pi/2, \tag{4.3}$$

the sign of b is alternating, as θ goes into intervals $\chi_0 + n\pi < \theta < \chi_0 + (n+1)\pi$ in such a manner that, for example, $b > 0$ for $\chi_0 < \theta < \chi_0 + \pi$. In view of the values of β for $e^{-i\theta H}$, we define β to take the values such that

$$\beta = \begin{cases} -1 & \text{for } \chi_0 - 2\pi < \theta < \chi_0 - \pi, \\ i & \text{for } \chi_0 - \pi < \theta < \chi_0, \\ 1 & \text{for } \chi_0 < \theta < \chi_0 + \pi, \\ -i & \text{for } \chi_0 + \pi < \theta < \chi_0 + 2\pi, \end{cases} \tag{4.4}$$

If $\tau = \zeta = 0$, then $\chi_0 = 0$, so that Def. (4.4) reduces to (3.5). We turn to the case of $\theta = \tau = 0$, whence $\tan \chi_0 = -\zeta$ with $|\chi_0| < \pi/2$. If $\zeta > 0$, then $\chi_0 < 0$, so that $\chi_0 < 0 < \chi_0 + \pi$. Hence, from (4.4) we have $\beta = 1$. If $\zeta < 0$, then $\chi_0 > 0$, so that $\chi_0 - \pi < 0 < \chi_0$. Hence, $\beta = 1$. Thus, Def. (4.4) covers the values of β for $e^{-i\zeta F}$. We wish to show that the integral transforms, whose kernels (2.2) have the factors $\alpha = e^{-i\pi/4}\beta$ with β defined by (4.4), indeed give a unitary representation of $SL_2(2, \mathbf{R})$. Let

$M_k, k=1, 2, 21,$ be matrices in $SL(2, \mathbf{R})$ with parameters $(\theta_k, \tau_k, \zeta_k)$ such that $M_{21}=M_2M_1$. Then from (3.1) and (3.2) applied for $M_{21}=M_2M_1$, we have

$$\theta_{21} = \theta_2 + \arg [e^{\tau_1 + \tau_2} (\cos \theta_1 - \zeta_2 \sin \theta_1) + ie^{\tau_1 - \tau_2} \sin \theta_1]. \tag{4.5}$$

When the argument and the equality in Eq. (4.5) are taken modulo 2π , Eq. (4.5) gives the parameter θ_{21} for M_{21} . However, if we take Eq. (4.5) modulo 4π , the θ_{21} should be considered as a parameter of $SL_2(2, \mathbf{R})$. Moreover, Eq. (4.5) modulo 4π gives the multiplication law for $SL_2(2, \mathbf{R})$ together with similar equations obtained from the second and third equations in Eq. (3.2). See also reference 7) for parameters of the universal covering group of $SL(2, \mathbf{R})$.

We are now looking into Eq. (4.5). Let $Z(\theta_1)$ denote the quantity in (4.5) (enclosed by the square brackets), the values of $\tau_1, \tau_2,$ and ζ_2 being fixed. Let $\cot \kappa_1 = \zeta_2$ with $|\kappa_1| < \pi/2$ for ζ_2 given. Then for $\zeta_2 > 0,$ $\arg Z(\theta_1)$ ranges from -2π to $2\pi,$ when θ_1 goes from -2π to $2\pi,$ with $\arg Z(\kappa_1 + n\pi) = n\pi + \pi/2$ and $\arg Z(n\pi) = n\pi, n = -2, -1, 0, 1.$ For $\zeta_2 < 0,$ one has the same range of $\arg Z(\theta_1)$ as that for $\zeta_2 > 0,$ with the difference $\arg Z(\kappa_1 + n\pi) = (n-1)\pi + \pi/2$ and $\arg Z((n-1)\pi) = (n-1)\pi, n = -1, 0, 1, 2.$ Thus we know from Eq. (4.5) how θ_{21} depends on θ_1 and $\theta_2.$

We are now in a position to know the sign of b_{21}/b_1b_2 in the (θ_1, θ_2) -plane, and thereby are able to show by virtue of Eq. (2.4) that the integral transforms under consideration give a representation of $SL_2(2, \mathbf{R}).$ Let

$$\tan \chi_k = -\zeta_k e^{2\tau_k}, \quad k=1, 2, 21, \tag{4.6}$$

where $|\chi_k| < \pi/2$ for $k=1, 2.$ For $k=21,$ χ_k will be soon determined uniquely. For $(\chi_k, \tau_k, \zeta_k), b_k$ vanish, $k=1, 2, 21.$ Since one has $b_{21}=0$ when $b_1=b_2=0,$ Eq. (4.5) gives

$$\chi_{21} = \chi_2 + \arg Z(\chi_1), \text{ mod. } 4\pi, \tag{4.7}$$

by which χ_{21} is determined uniquely. To identify the sign of $b_{21}/b_1b_2,$ we draw orthogonal lines $\theta_1 = \chi_1 + n\pi$ and $\theta_2 = \chi_2 + m\pi$ in the (θ_1, θ_2) -plane, n and m being integers. We next draw the curves defined by $\theta_{21} = \chi_{21} + k\pi, k$ being integers, which pass the intersection points of the already drawn orthogonal lines, because at those points $b_1 = b_2 = b_{21} = 0.$ For example, the curve $\theta_{21} = \chi_{21}$ passes the points $(\chi_1 + l\pi, \chi_2 - l\pi), l$ being integers, and so on. Thus, the (θ_1, θ_2) -plane is broken up into curved triangles. The sign of b_{21}/b_1b_2 can now be definitely settled in each curved triangle.

We are ready to describe the composition of the integral transforms. Let $U_k, k=1, 2, 21,$ be the integral transforms with certain values of $\beta,$ which are determined by (4.4) with χ_k replaced for $\chi_0.$ The operators $\pm U_k$ are in two-to-one correspondence with $M_k.$ Now we can prove $U_{21} = U_2U_1$ by the use of (2.4). For example, in the region in which $\chi_1 - \pi < \theta_1 < \chi_1, \chi_2 - \pi < \theta_2 < \chi_2,$ and $\chi_{21} - 2\pi < \theta_{21} < \chi_{21}$

$-\pi$, b_{21}/b_1b_2 is positive. Moreover, from (4.1) and (4.4) with χ_k replaced for χ_0 , $\alpha_1 = \alpha_2 = ie^{-i\pi/4}$ and $\alpha_{21} = -e^{-i\pi/4}$. Then we have from (2.4)

$$U_1 U_2 = \frac{\alpha_1 \alpha_2}{\alpha_{21}} e^{i\pi/4} U_{21} = U_{21}. \tag{4.8}$$

In other regions the same reasoning holds. Thus we have proved $U_2 U_1 = U_{21}$ for $b_1 \neq 0$, $b_2 \neq 0$.

We are in the final stage of studying canonical transforms. So far we have defined integral transforms for $\theta \neq \chi_0 + n\pi$, n being integers. The last task is to define the canonical transforms for $\theta = \chi_0 + n\pi$. A definition can be made by finding the limits of the integral kernels as θ tends to $\chi_0 + n\pi$. We take the integral kernel (2.2) again to consider its limit as $b \rightarrow 0$. Since $ad - bc = 1$, we may understand that $a \neq 0$ when b is near zero. Hence the integral kernel can be written, by replacing d with $(1 + bc)/a$, in the form

$$K(x, \xi) = \frac{\alpha}{(2\pi|b|)^{1/2}} \exp\left(\frac{ic}{2a}x^2\right) \exp\left(\frac{ia}{2b}\left(\xi - \frac{x}{a}\right)^2\right) \tag{4.9}$$

It follows from (4.9) that, as $b \rightarrow 0$,

$$\int K(x, \xi) d\xi \rightarrow \frac{\alpha_0}{|\alpha_0|^{1/2}} e^{\pm i\pi/4} \exp\left(\frac{ic_0}{2a_0}x^2\right), \tag{4.10}$$

where the + sign or - sign is taken, according as $a/b > 0$ or $a/b < 0$, and the subscript 0 indicates the limit as $b \rightarrow 0$.

We are going to the limit of the kernel (4.9) as $\theta \rightarrow \chi_0$. The condition $|\chi_0| < \pi/2$ means that $a_0 = e^\tau \cos \chi_0 > 0$, so that $a/b > 0$ for $\theta \rightarrow \chi_0 + 0$, and $a/b < 0$ for $\theta \rightarrow \chi_0 - 0$. Thus, from (4.1) and (4.4), α tends to $\alpha_0 = e^{-i\pi/4}$ or $\alpha_0 = ie^{-i\pi/4}$ according as $\theta \rightarrow \chi_0 + 0$ or $\theta \rightarrow \chi_0 - 0$. Hence, the limit (4.10) has a unique value as $\theta \rightarrow \chi_0 \pm 0$, so that it follows from (4.9) that, as $\theta \rightarrow \chi_0$,

$$K(x, \xi) \rightarrow \frac{\exp\left(-\frac{i}{2}x^2 \tan \chi_0\right)}{|e^\tau \cos \chi_0|^{1/2}} \delta\left(\frac{x}{e^\tau \cos \chi_0} - \xi\right) \tag{4.11}$$

When $\tau = \zeta = 0$, one has $\chi_0 = 0$, so that Expression (4.11) reduces to $K(x, \xi) \rightarrow \delta(x - \xi)$.

In the same manner we obtain the limits of $K(x, \xi)$ as $\theta \rightarrow \chi_0 + n\pi$, n being integers;

$$K(x, \xi) \rightarrow \frac{i^k \exp\left(-\frac{i}{2}x^2 \tan \chi_0\right)}{|e^\tau \cos \chi_0|^{1/2}} \delta\left(\frac{(-1)^k x}{e^\tau \cos \chi_0} - \xi\right) \tag{4.12}$$

as $\theta \rightarrow \chi_0 - k\pi$, $k = -2, -1, 0, 1$. Expression (4.12) reduces to (3.6) when $\tau = \zeta = 0$.

When $\tau = 0$, one has $\chi_0 = 0$. Then, Expression (4.12) reads

$$K(x, \xi) \rightarrow i^k e^{-\tau/2} \delta((-1)^k e^{-\tau} - \xi). \tag{4.13}$$

If $\theta \rightarrow 0$, Expression (4.13) with $k = 0$ reduces to (3.7).

Thus, we have defined the unitary operators U for all parameters of $SL_2(2, \mathbf{R})$.

The composition of operators $U_2 U_1 = U_{2,1}$ is true for all parameters of $SL_2(2, \mathbf{R})$ because of the continuity of U .

In conclusion, we point out that the canonical transformations we have constructed have two invariant closed subspaces, which are the space of even functions and the space of odd functions.

References

- 1) Wolf, K.B., *Integral Transforms in Science and Engineering*, Plenum Press, New York, 1979.
- 2) Moshinsky, M. and Quesne, C., *J. Math. Phys.* **12**, 1772 (1971).
- 3) Guillemin, V. and Sternberg, S., *Symplectic Techniques in Physics*, Cambridge University Press, Cambridge, 1984.
- 4) Shale, D., *Trans. Amer. Math. Soc.* **103**, 149 (1962).
- 5) Wallach, N.R., *Symplectic Geometry and Fourier Analysis*, Math Sci Press, Brookline, Mass., 1977.
- 6) Lion, G. and Vergne, M., *The Weil Representation, Maslov Index and Theta Series*, Progress in Math. 6, Birkhauser, Boston, 1980.
- 7) Bargmann, V., in *Analytic Methods in Mathematical Physics*, ed. by Gilbert, R.P. and Newton, R.G., Gordon and Breach, New York, 1970.