




Priority areas for investment in more sustainable and climate-resilient livestock systems

Received: 16 June 2022

Accepted: 26 May 2023

Published online: 29 June 2023

 Check for updates


Camila Bonilla-Cedrez¹, Peter Steward², Todd S. Rosenstock³, Philip Thornton ⁴, Jacobo Arango⁵, Martin Kropff⁶ & Julian Ramirez-Villegas ^{5,7,8} 

Livestock production supports economic growth, jobs and nutrition, but contributes to and is vulnerable to climate change. A transition is thus needed for livestock systems to become more sustainable and climate resilient, with clear positive effects on the Sustainable Development Goals. It is unclear, however, where the global community should invest to support this change. We identified priority geographies for livestock system investments in 132 low- and middle-income countries (LMICs), at mid- and low latitudes. Our results show that adaptation and mitigation goals are inextricably linked for the vast majority of these countries. An equal weighting of adaptation and mitigation indicators suggests that the top five investment priorities are India, Brazil, China, Pakistan and Sudan. Across LMICs, these act as critical control points for the livestock sector's interactions with the climate system, land and livelihoods.

Livestock production supports society and generates nearly 40% of global agricultural gross domestic product (GDP). About 1.3 billion people¹, including almost 930 million poor Africans and South Asians², depend on it for their livelihoods. Many rely on livestock as the primary source of revenue, and keeping livestock can also act as insurance, offering some protection when other income streams fail¹. An important economic asset, livestock symbolize wealth and status across the Global South³. Through the provision of draught power and manure and the recycling of agricultural byproducts, livestock underpin crop production and the food system^{3–5}. Animal-sourced food also provides nutrient-dense diets that contribute to cognitive development, growth and well-being⁶. Thus, livestock positively affect economics, health and cultural development, all of which are pillars of the Sustainable Development Goals (SDGs). Among others, the livestock sector has substantial potential to contribute to SDG 8.4 (on decoupling economic growth from environmental degradation), SDG 12.1 (on sustainable consumption and production patterns), SDG 13.1

(on strengthening resilience and adaptive capacity), SDG 13.2 (on the integration of climate change measures in national policies) and SDG 13.b (on promoting mechanisms for raising the capacity for effective climate change-related planning and management).

Climate change seriously threatens livestock productivity and those who depend on it. More frequent extreme weather events, irregular precipitation and rising temperatures decrease yields and product quality, increase pest and disease outbreaks, increase mortality, cause price shocks and disrupt supply, with cascading effects on producers and consumers^{7,8}. For example, without adaptation, by 2100, the impact of heat stress on cattle alone will probably reach 4–10% of the 2005 production value⁹. Global estimates can, however, mask large regional differences. Reduction in milk and meat production in African and Asian countries may exceed 50 or even 70% under high-emission scenarios (Shared Socioeconomic Pathway 585) by 2100⁹. Despite uncertainties in the impact projections⁹, the risk is substantial^{10,11}. An analysis of 113 countries in Asia, Africa and Latin America suggested

¹Animal Production Systems Group, Wageningen University & Research, Wageningen, the Netherlands. ²International Center for Tropical Agriculture, Nairobi, Kenya. ³Biodiversity International, Montpellier, France. ⁴Clim-Eat, Netherlands Food Partnership, Utrecht, the Netherlands. ⁵International Center for Tropical Agriculture, Palmira, Colombia. ⁶Resilient Agrifood Systems, CGIAR System Organization, Montpellier, France. ⁷Plant Production Systems Group, Wageningen University & Research, Wageningen, the Netherlands. ⁸Biodiversity International, Rome, Italy.  e-mail: j.r.villegas@cgiar.org

BOX 1

Key concepts and terms on climate change adaptation and mitigation

Adaptation: Actions that decrease the vulnerability of farming systems to climate change (for example, farmers implementing water management systems to adapt to short- and long-term rainfall variations).

Adaptive capacity: Skills and capacity, such as knowledge, finance and technical capacity, that permit farmers or institutions to adjust their behaviour to mitigate the impact of climate hazards or capture opportunities.

Exposure: The degree to which a system is subject to a climate hazard.

Hazard: A climate event or process that can harm farmers, such as loss in productivity—a function of exposure and sensitivity.

Impact: Hazards' effect on farming systems and farmers, such as loss in productivity—a function of exposure and sensitivity.

Mitigation: Actions that decrease the rate of climate change by limiting or preventing GHG emissions and by enhancing activities that remove them from the atmosphere (for example, decreasing herd size to decrease their proportion of national and global GHG emissions).

Risk: The combination of exposure and vulnerability to hazards (the potential for adverse consequences).

that US\$994 billion per year in livestock value—around 55% of the total value of production (VOP) for five commodities—is exposed to various climate hazards, especially rainfall variability (US\$198 billion) and heat stress (US\$130 billion)¹². Adaptation of livestock production systems is imperative to maintain and enhance the benefits they provide.

These benefits, however, must be considered alongside the negative impacts of livestock on the climate system. Feed production, enteric emissions, manure management, grazing and land-use change release methane, nitrous oxide and carbon dioxide into the atmosphere. Livestock production accounts for about 5.8% of global annual greenhouse gas (GHG) emissions and about 31.5% of the food systems' contribution^{13,14}. The dominant livestock emission source varies by region. Herd size and enteric emissions drive climate impacts, for example, in rangeland systems¹⁵, while land-use change has an influence in mixed grazing systems. The expansion of livestock production systems also threatens tropical forests¹⁶. Given the impacts, actions to decrease agricultural emissions need to target livestock production systems.

The livestock sector affects 10 of 17 SDGs, but not all in positive ways¹⁷, meaning that the meeting of global goals will require transitioning to climate-resilient and low-emissions livestock production systems¹⁸. Yet, while nearly 100 countries prioritize livestock in their Nationally Determined Contributions (NDCs), public and private investors hesitate to target the livestock sector due to perceived risks and environmental concerns. Critical questions remain largely unanswered on where and in what to invest. In this study, we quantify countries and

livestock production systems that leverage the livestock agri-food systems in low- and middle-income countries (LMICs) to guide future adaptation and mitigation investments (Box 1). We detail proven actions for accelerating the transition to low-emission, climate-resilient systems. We discuss their scalability, enabling factors and constraints.

Results

Exposure to climate hazards and emissions by geographies

We found that substantial livestock production value and rural population are exposed to climate hazards (Table 1 and Fig. 1). Across the 132 countries in this analysis, US\$660 billion in VOP, 1.04 billion people, 470 million tropical livestock units (TLUs) and 671 million hectares of pasture are exposed to climate hazards (Table 1 and Supplementary Table 1). The most prevalent hazard combinations by percentage of total area were rainfall variability plus heat stress plus drought (33%), heat stress (28%) and rainfall variability plus heat stress (12%) (Fig. 1 and Supplementary Fig. 1). The levels of importance of the various climate hazards differ among countries and livestock production systems (Fig. 1 and Supplementary Figs. 1 and 2). Livestock systems in India, Nigeria and Sudan are the most exposed, with India's exposure exceeding that of other countries at least fivefold (Fig. 1). Within the study area, rainfed arid and humid regions where mixed farming (crops and livestock) systems are prevalent had the highest VOPs (US\$405 billion), rural populations (661 million) and TLUs (295 million) exposed to climate hazards (Table 1 and Supplementary Fig. 2). Regions such as the Horn of Africa, which predominantly includes rainfed arid rangelands, and West Africa (with both humid and arid conditions) have 14% of their VOP, 28% of their rural population, 39% of their pasture area and 32% of their TLUs exposed to climate hazards.

GHG emissions from livestock are substantial across the study region. Across the 132 countries included in this analysis, emissions were estimated to be 2,995 megatons (CO₂e) (Table 1, Fig. 1 and Supplementary Table 2). Seventy-one percent (2,132 megatons CO₂e) were directly from livestock production, including enteric emissions, manure and feeds, while 29% (863 megatons CO₂e) related to deforestation for pastures and soy production (Table 1 and Supplementary Table 2). Mixed rainfed systems in humid and arid lands produce the most carbon dioxide equivalents (CO₂e) by far, with ruminants representing 66 and 94% of their total emissions (Supplementary Fig. 3). Emissions from Brazil and India far exceed those of the other countries studied, but these emissions result from vastly different dynamics. In Brazil, 64% of total emissions are due to soybean and pasture-driven forest loss. In India, 99% of the emissions are from ruminants, predominantly from bovine milk production associated with mixed systems (Supplementary Fig. 4).

Priorities for adaptation and mitigation

Our analysis shows that no country or system with appreciable livestock production value and population has zero exposure to climate hazards or zero emissions (Fig. 1). Therefore, using this analysis to obtain a set of geographic priorities requires the identification of target outcomes, their relative importance and the implied trade-offs across scales. An equal weighting of adaptation and mitigation indicators (see Methods) suggests that the top five investment priorities are India, Brazil, China, Pakistan and Sudan (Figs. 1 and 2 and Supplementary Fig. 5). Across LMICs, these act as critical control points for the livestock sector's interactions with the climate system, land and livelihoods. These five countries combined account for 46% of the total VOP, 35% of the total rural population exposed to climate hazards and 51% of emissions. However, differential weighting indicators change the outcomes of investment prioritization. For example, a sole focus on mitigation prioritizes investments in Brazil, China, India, Pakistan and Bangladesh (Fig. 2 and Supplementary Fig. 7). In contrast, focusing only on adaptation prioritizes India, Sudan, South Africa, Nigeria and Chad (Fig. 2 and Supplementary Fig. 8).

Table 1 | Totals of GHG emissions and exposure to climate hazards per livestock system within the study area

Livestock production system	GHG emissions			Climate hazard exposure			
	Direct emissions (Mt CO ₂ e)	Indirect emissions (Mt CO ₂ e)	Total emissions (Mt CO ₂ e)	VOP (billion US\$)	Rural population (million)	Pasture area (million ha)	TLUs (million)
Mixed irrigated							
MIA	263	1	264	115.7	167.9	5.7	70.9
MIH	103	3	106	57.9	31.7	0.7	30.8
MIY	1	0	1	0	0	0	0
MIT	58	0	58	18.8	10.1	4.0	3.7
Mixed rainfed							
MRA	379	20	399	132.4	285.6	150.6	132.8
MRH	440	200	640	99.1	175.9	28.0	60.8
MRY	0	0	0	0	0	0	0.1
MRT	148	6	154	31.1	37.4	10.1	22.1
Rangelands							
LGA	250	22	272	50.1	108.7	326.7	71.4
LGH	99	56	155	21.6	46.2	49.4	14.7
LGY	4	0	4	0.1	0.2	3.4	0.3
LGT	77	4	81	24.0	11.4	38.8	8.8
Other	310	551	862	110	167	54	54
Total	2,132.3	863.2	2,995.6	660.7	1,042.0	671.7	470.5

The category other is as reported by Robinson et al.⁴¹ for systems of varying type (for example, root crop based, root crop mixed, forest based and tree based), with some component of livestock. LGA, rangelands arid; LGH, rangelands humid; LGT, rangelands temperate; LGY, rangelands hyperarid; MIA, mixed irrigated arid; MIH, mixed irrigated humid; MIT, mixed irrigated temperate; MIY, mixed irrigated hyperarid; MRA, mixed rainfed arid; MRH, mixed rainfed humid; MRT, mixed rainfed temperate; MRY, mixed rainfed hyperarid.




Fig. 1 | Adaptation and mitigation potential priorities. Adaptation priorities include areas where TLUs (L), pasture area (P), rural population (R) and VOP (V) are exposed to climate hazards. Mitigation priorities include reducing livestock direct and deforestation-linked GHG emissions (E). The bottom-left (top-right) corner of the colour scale indicates low (high) adaptation and mitigation potential priorities. Shades of blue show increasing mitigation potential, moving from light to dark blue, over the x axis of the colour scale, whereas shades of green show increasing adaptation potential, moving from light to dark green, over the y axis

of the colour scale. The climate hazards include climate variability (RF), heat stress (THI), drought (D) and flooding (F). The following ID codes for the various indicators, hazards and emissions categories reference datasets used in the analysis (see Supplementary Table 3): TLUs (001), VOP (002), pasture area (003), rural population (004), climate hazards (006–009), GHG emissions (011–017), bovine emissions (011 and 012), forest loss to pasture emissions (013–016), forest loss to soy emissions (013–015 and 017), pig emissions (011 and 012) and shoat emissions (011 and 012).




Fig. 2 | Priority countries for adaptation and mitigation. Priority countries for adaptation are shown in green (left) and those for mitigation are shown in blue (right). Larger shapes and fonts and more intense colours within each panel represent higher priorities. DRC, Democratic Republic of the Congo.



Fig. 3 | Adaptation and mitigation options available for livestock systems in LMICs, constraints to their adoption in the selected countries and the relative importance of each constraint for the adoption of each option in general. Darker colours represent a higher constraint for the adoption of a particular option in each country, as determined by the quantile in which

the country sits with respect to the global median. Indicators to represent each constraint include accessibility (research and development expenditure (percentage of GDP)), cost (GDP per capita), knowledge (literacy rate in the total adult population), labour (employment in agriculture) and land tenure (Rule of Law). Data from ref. 57.

For most geographies, it is virtually impossible to disentangle the importance and relationship between adaptation and mitigation. More specifically, adaptation and mitigation emerge as joint priorities most clearly in India, in which 32% of the rural population and 41% of the VOP in the study area are exposed to climate hazards, and which produces 13% of the emissions (Fig. 1). Across most of Sub-Saharan Africa, adaptation measures tend to greatly outweigh those of mitigation options (Fig. 1). The converse is true for countries in Latin America, where mitigation concerns are generally greater than adaptation concerns.

Climate actions and their adoption constraints

Adaptation and mitigation options already exist that would allow the livestock sector to meet its economic, environmental and climate mitigation and resilience goals (Fig. 3). Among other factors, low investment, lack of education and cultural, institutional and political barriers have led to generally low adoption rates for these actions.

Constraints to the adoption vary by region, country and options (Fig. 3). For instance, for the selected countries, cost is a barrier to adopting most options, and the relative importance of cost varies between 20 and 50%. In countries such as Argentina, Brazil and China, the main barrier to adopting a technology is the limited labour force due to the lower rural population and fewer people employed in agriculture. This barrier differs in Ethiopia, Nigeria and Sudan, where accessibility and knowledge are the main adoption constraints.

Discussion

From a global perspective, the weighting of adaptation and mitigation indicators highlights the geographies with the greatest problems, providing a regional focus. At local scales, however, these indicators emerge from farmers' decisions on system management and, as such, are interconnected; any choice of adaptive or mitigating priorities needs understanding of the trade-offs that impact other goals.

Adopting a livestock practice or technology nearly always involves trade-offs between adaptation and mitigation outcomes¹⁹. Thus, to avoid unintended consequences, actions need to be aligned with local demands and goals. The consideration that adaptation and mitigation need to be addressed jointly is especially important in areas where population growth and/or dietary change are most prominent²⁰. Yet, a lack of alignment at the policy level potentially hinders this objective. Roughly 50% of the NDCs that mention livestock note just one of the priorities²¹, and only 28 out of 184 countries' NDCs include soil-related targets²². This omission suggests that, as a global community, we are creating institutions and narratives that disincentivize or preclude action on adaptation, mitigation or both.

Our assessment of adaptation and mitigation options suggests that careful consideration of context specificities is needed for the scaling of options due to regional systems dynamics and adoption constraints (Fig. 3). In livestock systems in general, to maximize the benefits of improving and intensifying diets, reductions in herd size are necessary^{15–17}. However, reductions in animal numbers may be limited to: (1) humid and temperate mixed systems where feeding grain concentrates are a plausible option²³; (2) places where an alternative source of both feed and food protein are available^{15,23,24}; and (3) where livestock are not enmeshed in cultural identity^{25,26}. Manure management is another viable strategy for decreasing emissions, but its applicability is limited for open grazing systems. In the study region, 18% of cattle, 27% of sheep and 43% of goats roam in extensive grazing systems, where most excretion happens in the field; collection occurs after the manure is dry, if at all^{15,27}. In East Asian systems with a high concentration of animals, emissions are associated with unregulated manure disposal, and animals produce more manure than can be recycled in the agricultural area²⁸. In grasslands and agricultural land, 47% of the total potential mitigation arises from soil organic carbon protection and sequestration²⁹, as well as the restoration of degraded rangelands²³. The potential of restoration and improved land management actions to increase carbon storage and/or avoid emissions across global forests, wetlands, grasslands and agricultural lands is 23.8 GtCO₂e yr⁻¹ (ref. 30). Silvopastoral systems are a viable land management strategy for adaptation and mitigation in Latin America, where livestock is a major driver of deforestation and 11% of people are exposed to heat stress. Silvopastoral systems could increase productivity^{31,32}, decrease GHG emissions³³, improve carbon storage potential³² and aid adaptation to heat stress³⁴. However, the scalability of silvopastoral systems has been limited by several factors: lack of knowledge, high initial investment requirements and the extended periods before returns on investments are seen.

Climate risk reduction should be essential in transforming livestock systems; according to our analysis, 69% of the VOP is exposed to high levels of climate variability. Climate services and access to credit and insurance are proven climate risk management actions that have demonstrated potential and scalability for crop-based systems^{35,36}. In Senegal, where virtually all livestock systems are exposed to high or extreme climate variability, climate services increase farmer income by 10–25%³⁷. In the Horn of Africa, where pastoral drylands are regularly affected by drought and its interannual variability (Supplementary Figs. 1 and 2), the Predictive Livestock Early Warning System³⁸ seeks to improve livestock herd management, migration patterns and livestock health through providing forecasts on water and pasture availability. Likewise, the index-based livestock insurance approach has sold around 90,000 insurance policies in Kenya and Ethiopia, positively impacting policyholders' sales, income and well-being^{39,40}.

Low levels of investment in the sector and a lack of effective policies, combined with high resource demand and limited information to implement solutions (Fig. 3), reinforce the gap between current livestock systems and their future potential. There is limited evidence regarding the scalability of the actions outlined here. As such, the set of interventions and practices that best protect against the combined

hazards and can reliably measure their effectiveness warrants further investigation. The gathering of such evidence must be combined with building the capacities of local and regional organizations, the public sector and producers, to promote and implement adaptation options. Governments need technical support to access finance, implement programmes and report adaptation and mitigation achievements. These challenges apply equally to the private sector and small-scale systems. Large-scale production changes landscapes, and supply and demand shifts can provide major benefits and influence consumer behaviour. In small-scale systems, the value of livestock to livelihoods goes far beyond their productive capacity, and building local knowledge and capacity is vital to achieving transformation.

Methods

We used multiple spatial datasets (Supplementary Tables 3 and 4) to implement a prioritization approach to identify priority geographies (that is, countries and production systems) for livestock system investments and to identify adoption constraints to adaptation and mitigation options. We selected two main challenges related to climate adaptation and mitigation strategies: (1) livestock production and producer livelihoods are threatened by short- and long-term rainfall and temperature variations and their changing predictability; and (2) livestock are responsible for a substantial proportion of national and global GHG emissions. A fully detailed description of the methods is provided in the Supplementary Information.

We used the livestock production system classification of Robinson et al.⁴¹. This classification divides livestock production systems into landless, rangeland and mixed systems. Mixed systems are divided into rainfed and irrigated, giving rise to four broad system categories (that is, landless, rangelands, mixed rainfed and mixed irrigated). Each category was then divided into agroecologies according to temperature and whether they were humid, arid or hyperarid⁴¹. The study area included livestock production systems in all LMICs in mid- and low latitudes identified by The World Bank ($n = 132$ countries).

Analysis of adaptation

We performed a geospatial data analysis to characterize livestock production systems and producer livelihoods and to identify threats related to short- and long-term variations in rainfall and temperature. First, we mapped regions projected to experience climatic hazards and masked them by their adaptive capacity within the livestock systems (Supplementary Fig. 8). Second, we assessed the exposure per hazard within each country and livestock system (Supplementary Fig. 9). Exposure was represented by the total VOP⁴², total rural population⁴³, pasture area⁴⁴ and TLUs. TLU values were computed following Rothman-Ostrow et al.⁴⁵ using data from the Gridded Livestock of the World database (version 3)⁴⁶. VOP data were derived from ref. 42, which used a combination of modelling and FAOSTAT data¹³ to produce geospatial datasets of the value of livestock production.

Sixteen climatic hazard classes were identified from the intersection of rainfall variability, heat stress, drought and flood risk. To characterize rainfall variability, we used the coefficient of variation in annual mean rainfall (15–30% = highly variable; >30% = extremely variable) derived from the Climate Hazards Group InfraRed Precipitation with Stations dataset⁴⁷. Areas of heat stress were defined as having thermal stress (projected for 2030 under the Representative Concentration Pathway 8.5 scenario) with a of ≥ 79 (ref. 48). To define areas at risk of flooding, we used the UN Environment Programme–Division of Early Warning and Assessment–GRID-Europe dataset⁴⁹, in which flooding risk is ranked from 0 (no risk) to 5 (extreme). Areas at risk of drought were defined as having more than 25 d without rain per month on average.

We defined three categories of livestock system adaptive capacity to these climate hazards (low $\geq 25\%$; medium = 10–25%; high $\leq 10\%$) using the national-level poverty headcount ratio of US\$1.90 d⁻¹ (ref. 50) as a proxy. For the exposure analysis, we considered areas with poverty

rates above 10% and their climate dimensions (that is, the rainfall coefficient of variation, heat stress, flood hazard risk and drought).

Analysis of mitigation

We calculated total livestock emissions by geography by summing direct and estimated emissions from deforestation (Supplementary Figs. 10 and 11). We did not include changes in soil carbon stocks due to the difficulty in estimating these reliably at the scale of our study. Across the 132 countries included in this analysis, emissions were estimated to be 2,995 megatons (CO₂e), which is lower than other recently published estimates⁵¹. This difference arises primarily due to the exclusion of direct CO₂ emissions from cropland used for feed (see Supplementary Information) and the focus on LMICs only (compared with global estimates). Despite inherent limitations and uncertainties in the input datasets (see the corresponding references), our sources of emissions data are derived well-established sources (refs. 13,42,52,53) and robust for the purposes of our prioritization analysis. Furthermore, because our analysis does not introduce specific equations or parameters, it is unlikely that it introduces new uncertainties or propagates any original uncertainties present in the input datasets, beyond what would be a simple addition of these original uncertainties.

Direct emissions data come from ref. 42 and are for the year 2010. To update these data to 2019, we multiplied 2010 values by the proportional increase in national livestock emissions between 2000 and 2019, as reported by FAOSTAT^{13,54}. To estimate livestock emissions linked to deforestation, we first combined above- and below-ground carbon biomass data from 2010⁵⁵ and masked these values by areas with tree cover values $\geq 30\%$ ⁵⁶, giving a layer of carbon biomass per hectare of forest. Commodity-linked deforestation data are available for subnational administrative areas⁵². For each area, we calculated the average amount of carbon biomass (using the data from ref. 55) and multiplied this by the mean annual rate of commodity-linked (soy and pasture) forest loss for the period 2010–2015 (data from ref. 52) giving the average rate of carbon loss per area per year. Data from ref. 52 are available only at the subnational administration level; thus, our analysis of indirect emissions was conducted at that spatial scale. Carbon was converted to CO₂ using a factor of 3.67 to account for its atomic weight. We assume that carbon in below-ground biomass is lost on conversion to pastureland or soybeans. We included deforestation due to soybeans as the majority of soybean production is used for livestock feed, but note that a proportion of this production has other fates.

Prioritization

All indicators were normalized by dividing the value of each indicator per country by the maximum value across geographies to rank countries and visualize adaptation and mitigation indicators (Fig. 1). For example, TLUs for each country were normalized using India's TLUs, since it has the highest value worldwide. Each normalized exposure indicator was then weighted and summed, creating a unique adaptation index. For the case of mitigation, each indicator corresponding to emissions was normalized by dividing by the maximum value across geographies.

Constraints for the adoption of climate actions

Based on a literature review⁵⁷ and expert opinion, we identified adaptation and mitigation options available for livestock systems (Fig. 3). To represent the constraints for the adoption of these options, we identified key global indicators reported by The World Bank to quantify the relevance of each constraint for each adaptation and mitigation option per country (that is, accessibility, cost, knowledge, labour and land tenure). We represented the constraints as follows: (1) accessibility was measured using research and development expenditure (percentage of GDP); (2) cost was measured by GDP per capita; (3) knowledge was measured by the total adult literacy rate; (4) labour was measured by employment in agriculture; and (5) land tenure was measured by the

Rule of Law (see Supplementary Table 4). The relative importance of each constraint for each option—not for a specific country but generally—was quantified based on expert opinion. We represent these constraints for the top three countries, by continent, that were identified in the prioritization analysis.

This research was conducted as part of a global research programme on livestock and climate (see <https://www.cgiar.org/initiative/34-livestock-climate-and-system-resilience/>), which is part of the CGIAR consortium of research centres. No involvement of human participants or animal subjects took place as part of this work. To the best of the authors' knowledge, the findings reported herein do not carry any racial, cultural or gender bias.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

The datasets analysed during the current study are available from the livestock prioritization repository https://github.com/CIAT/livestock_prioritization/tree/main/Data. No restrictions on data availability exist other than those related to refs. 42,52, the data from which are not publicly available. For use of these datasets, we strongly recommend contacting the first and/or corresponding authors of the studies. Access links for all datasets are provided in Supplementary Table 3.

Code availability

An interactive markdown is available at https://github.com/CIAT/livestock_prioritization.

References

1. *World Livestock: Transforming the Livestock Sector Through the Sustainable Development Goals* (FAO, 2018).
2. Hong, C. et al. Global and regional drivers of land-use emissions in 1961–2017. *Nature* **589**, 554–561 (2021).
3. Herrero, M. et al. The roles of livestock in developing countries. *Animal* **7**, 3–18 (2013).
4. Megersa, B., Markemann, A., Angassa, A. & Valle Zárate, A. The role of livestock diversification in ensuring household food security under a changing climate in Borana, Ethiopia. *Food Secur.* **6**, 15–28 (2014).
5. Moll, H. A. J., Staal, S. J. & Ibrahim, M. N. M. Smallholder dairy production and markets: a comparison of production systems in Zambia, Kenya and Sri Lanka. *Agric. Syst.* **94**, 593–603 (2007).
6. Gupta, S. Brain food: clever eating. *Nature* **531**, S12–S13 (2016).
7. Thornton, P. K. & Herrero, M. Adapting to climate change in the mixed crop and livestock farming systems in sub-Saharan Africa. *Nat. Clim. Change* **5**, 830–836 (2015).
8. Rojas-Downing, M. M., Nejadhashemi, A. P., Harrigan, T. & Woznicki, S. A. Climate change and livestock: impacts, adaptation, and mitigation. *Clim. Risk Manag.* **16**, 145–163 (2017).
9. Thornton, P., Nelson, G., Mayberry, D. & Herrero, M. Impacts of heat stress on global cattle production during the 21st century: a modelling study. *Lancet Planet. Health* **6**, e192–e201 (2022).
10. Thornton, P. K., van de Steeg, J., Notenbaert, A. & Herrero, M. The impacts of climate change on livestock and livestock systems in developing countries: a review of what we know and what we need to know. *Agric. Syst.* **101**, 113–127 (2009).
11. Godde, C. M., Mason-D'Croz, D., Mayberry, D. E., Thornton, P. K. & Herrero, M. Impacts of climate change on the livestock food supply chain; a review of the evidence. *Glob. Food Sec.* **28**, 100488 (2021).
12. Jarvis, A. et al. *Climate-Informed Priorities for One CGIAR Regional Integrated Initiatives* (CGIAR Research Program on Climate Change, Agriculture and Food Security, 2021).

13. FAOSTAT (FAO, 2021); <http://www.fao.org/faostat/en/#data/GT>
14. *Climate Watch* (World Resources Institute, 2023); <https://www.wri.org/initiatives/climate-watch>
15. Herrero, M. et al. Greenhouse gas mitigation potentials in the livestock sector. *Nat. Clim. Change* **6**, 452–461 (2016).
16. Pendrill, F. et al. Disentangling the numbers behind agriculture-driven tropical deforestation. *Science* **377**, eabm9267 (2022).
17. Mehrabi, Z., Gill, M., Wijk, M., van, Herrero, M. & Ramankutty, N. Livestock policy for sustainable development. *Nat. Food* **1**, 160–165 (2020).
18. Paul, B. K., Butterbach-Bahl, K., Notenbaert, A., Nduah Nderi, A. & Ericksen, P. Sustainable livestock development in low- and middle-income countries: shedding light on evidence-based solutions. *Environ. Res. Lett.* **16**, 011001 (2021).
19. Smith, P. et al. Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? *Glob. Change Biol.* **26**, 1532–1575 (2020).
20. Khoury, C. K. et al. Increasing homogeneity in global food supplies and the implications for food security. *Proc. Natl Acad. Sci. USA* **111**, 4001–4006 (2014).
21. Richards, M. et al. *How Countries Plan to Address Agricultural Adaptation and Mitigation: An Analysis of Intended Nationally Determined Contributions*. CCAFS Info Note (CGIAR Research Program on Climate Change, Agriculture and Food Security, 2015).
22. Wiese, L. et al. Countries' commitments to soil organic carbon in Nationally Determined Contributions. *Clim. Policy* **21**, 1005–1019 (2021).
23. Thornton, P. K. & Herrero, M. Potential for reduced methane and carbon dioxide emissions from livestock and pasture management in the tropics. *Proc. Natl Acad. Sci. USA* **107**, 19667–19672 (2010).
24. Rivera-Ferre, M. G. et al. Re-framing the climate change debate in the livestock sector: mitigation and adaptation options. *WIREs Clim. Change* **7**, 869–892 (2016).
25. Zander, K. K. & Drucker, A. G. Conserving what's important: using choice model scenarios to value local cattle breeds in East Africa. *Ecol. Econ.* **68**, 34–45 (2008).
26. Ainslie, A. The sociocultural contexts and meanings associated with livestock keeping in rural South Africa. *Afr. J. Range Forage Sci.* **30**, 35–38 (2013).
27. Harris, F. Management of manure in farming systems in semi-arid West Africa. *Exp. Agric.* **38**, 131–148 (2002).
28. Uwizeye, A. et al. Nitrogen emissions along global livestock supply chains. *Nat. Food* **1**, 437–446 (2020).
29. Bossio, D. A. et al. The role of soil carbon in natural climate solutions. *Nat. Sustain.* **3**, 391–398 (2020).
30. Griscom, B. W. et al. Natural climate solutions. *Proc. Natl Acad. Sci. USA* **114**, 11645–11650 (2017).
31. Yamamoto, W., Dewi, I. A. & Ibrahim, M. Effects of silvopastoral areas on milk production at dual-purpose cattle farms at the semi-humid old agricultural frontier in central Nicaragua. *Agric. Syst.* **94**, 368–375 (2007).
32. Jose, S. & Dollinger, J. Silvopasture: a sustainable livestock production system. *Agrofor. Syst.* **93**, 1–9 (2019).
33. Nair, P. K. R., Tonucci, R. G., Garcia, R. & Nair, V. D. in *Carbon Sequestration Potential of Agroforestry Systems: Opportunities and Challenges* (eds Kumar, B. & Nair, P.) 145–162 (Springer, 2011).
34. Zeppetello, L. R. V. et al. Consistent cooling benefits of silvopasture in the tropics. *Nat. Commun.* **13**, 708 (2022).
35. Vaughan, C., Hansen, J., Roudier, P., Watkiss, P. & Carr, E. Evaluating agricultural weather and climate services in Africa: evidence, methods, and a learning agenda. *WIREs Clim. Change* **10**, e586 (2019).
36. Hansen, J. W. et al. Climate services can support African farmers' context-specific adaptation needs at scale. *Front. Sustain. Food Syst.* **3**, 21 (2019).
37. Chiputwa, B., Blundo-Canto, G., Steward, P., Andrieu, N. & Ndiaye, O. Co-production, uptake of weather and climate services, and welfare impacts on farmers in Senegal: a panel data approach. *Agric. Syst.* **195**, 103309 (2022).
38. Matere, J. et al. Predictive Livestock Early Warning System (PLEWS): monitoring forage condition and implications for animal production in Kenya. *Weather Clim. Extremes* **27**, 100209 (2020).
39. Matsuda, A., Takahashi, K. & Ikegami, M. Direct and indirect impact of index-based livestock insurance in Southern Ethiopia. *Geneva Pap. Risk Insur. Issues Pract.* **44**, 481–502 (2019).
40. Jensen, N. D., Barrett, C. B. & Mude, A. G. Cash transfers and index insurance: a comparative impact analysis from northern Kenya. *J. Dev. Econ.* **129**, 14–28 (2017).
41. Robinson, T. P. et al. Mapping the global distribution of livestock. *PLoS ONE* **9**, e96084 (2014).
42. Herrero, M. et al. Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proc. Natl Acad. Sci. USA* **110**, 20888–20893 (2013).
43. *WorldPop* (University of Southampton, 2021); <http://www.worldpop.org>
44. Ramankutty, N., Evan, A. T., Monfreda, C. & Foley, J. A. Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Glob. Biogeochem. Cycles* **22**, GB1003 (2008).
45. Rothman-Ostrow, P., Gilbert, W. & Rushton, J. Tropical livestock units: re-evaluating a methodology. *Front. Vet. Sci.* **7**, 556788 (2020).
46. Gilbert, M. et al. Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010. *Sci. Data* **5**, 180227 (2018).
47. Funk, C. et al. The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. *Sci. Data* **2**, 150066 (2015).
48. Thornton, P., Nelson, G., Mayberry, D. & Herrero, M. Increases in extreme heat stress in domesticated livestock species during the twenty-first century. *Glob. Change Biol.* **27**, 5762–5772 (2021).
49. *Flood Risk Maps* (UN Environment Programme, Division of Early Warning and Assessment & GRID-Europe, 2021); <https://uneppgrid.ch/en/activity/1BDE1705>
50. *Poverty Headcount Ratio for \$1.90 a Day at 2011 International Prices* (The World Bank, 2021); <https://maps.worldbank.org/projects>
51. Xu, X. et al. Global greenhouse gas emissions from animal-based foods are twice those of plant-based foods. *Nat. Food* **2**, 724–732 (2021).
52. Goldman, E., Weisse, M. J., Harris, N. & Schneider, M. *Estimating the Role of Seven Commodities in Agriculture-Linked Deforestation: Oil Palm, Soy, Cattle, Wood Fiber, Cocoa, Coffee, and Rubber* (World Resources Institute, 2020).
53. Tubiello, F. N. et al. Pre- and post-production processes increasingly dominate greenhouse gas emissions from agri-food systems. *Earth Syst. Sci. Data* **14**, 1795–1809 (2022).
54. Flammioni, A. et al. Emissions of greenhouse gases from energy use in agriculture, forestry and fisheries: 1970–2019. *Earth Syst. Sci. Data* **14**, 811–821 (2022).
55. Soto-Navarro, C. et al. Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action. *Phil. Trans. R. Soc. B Biol. Sci.* **375**, 20190128 (2020).
56. Hansen, M. C. et al. High-resolution global maps of 21st-century forest cover change. *Science* **342**, 850–853 (2013).

57. Thornton, P. et al. Responding to global change: a theory of change approach to making agricultural research for development outcome-based. *Agric. Syst.* **152**, 145–153 (2017).

Acknowledgements

We thank the CGIAR Livestock and Climate Initiative for support and P. Ericksen for supporting the development of this manuscript. We thank E. Goldman for access to the soybean and pasture-driven forest loss database.

Author contributions

T.S.R. and J.R.-V. conceived of the research. C.B.-C., P.S., T.S.R. and J.R.-V. performed the data acquisition and processing. C.B.-C. and P.S. analysed the data. C.B.-C., P.S., T.S.R., P.T., J.A. and J.R.-V. wrote the manuscript. C.B.-C., P.S., T.S.R., P.T., J.A., M.K. and J.R.-V. edited the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41893-023-01161-1>.

Correspondence and requests for materials should be addressed to Julian Ramirez-Villegas.

Peer review information *Nature Sustainability* thanks Matthew Harrison, Juan-Pablo Inamagua and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Data collection was performed manually from the original sources. In two cases data were provided by authors upon direct contact. The R markdown file contains all data transformations performed, with modifiable options. The GitHub repository (https://github.com/CIAT/livestock_prioritization) contains all input datasets (under the folder Data). These datasets will allow reproducing the entire analysis conducted. All datasets are listed in Supplementary Table S3. Below we provide a list of data with the access web addresses:

- 001 – Gilbert et al. (2018). Exposure: livestock density. Available online at <https://dataverse.harvard.edu/dataverse/glw>
- 002 – Herrero et al. (2013). Exposure: value of livestock production. Dataset provided by author.
- 003 – Ramankutty et al. (2008). Exposure: pasture area. Available at <http://www.earthstat.org/cropland-pasture-area-2000/>
- 004 – World Pop. Exposure: total rural population. Available at https://data.worldpop.org/GIS/Population/Global_2000_2020/2020/0_Mosaicked/ppp_2020_1km_Aggregated.tif
- 005 – Robinson et al. (2014). Exposure: global livestock systems. Available at <https://dataverse.harvard.edu/dataverse/GLPS>
- 006 – Funk et al. (2015). Hazards: coefficient of variation in mean rainfall. Calculated from CHIRPS data at <https://data.chc.ucsb.edu/products/CHIRPS-2.0/>
- 007 – Funk et al. (2015). Hazards: dry days per months. Calculated from CHIRPS data at <https://data.chc.ucsb.edu/products/CHIRPS-2.0/>
- 008 – UNEP/DEWA/GRID-Europe. Hazards: flood risk. <https://preview.grid.unep.ch/index.php?preview=data&events=floods&evcat=5&lang=eng>
- 009 – Thornton et al. (2021). Hazards: heat stress index. Dataset provided by author.
- 010 – World Bank. Adaptive capacity: poverty headcount ratio of US\$1.9 per day. Available at <https://maps.worldbank.org/projects>
- 011 – Herrero et al. (2013). Emissions: direct GHG emissions. Dataset provided by author.
- 012 – FAOSTAT (2021) & Tubiello et al. (2022). Emissions: direct GHG emissions. Available at https://fenixservices.fao.org/faostat/static/bulkdownloads/Emissions_Totals_E_All_Data.zip
- 013 – Soto-Navarro et al. (2020). Emissions: aboveground biomass carbon density. Available at https://datadownload-production.s3.amazonaws.com/WCMC_carbon_tonnes_per_ha.zip
- 014 – Soto-Navarro et al. (2020). Emissions: belowground biomass carbon density. Available at <https://datadownload->

production.s3.amazonaws.com/WCMC_carbon_tonnes_per_ha.zip
 015 – Hansen et al. (2013). Emissions: forest tree cover. Available at <https://storage.googleapis.com/earthenginepartners-hansen/GFC-2021-v1.9/download.html>
 016 – Goldman et al. (2020). Emissions: pasture-driven forest loss. Dataset provided by author.
 017 – Goldman et al. (2020). Emissions: soy-driven forest loss. Dataset provided by author.
 018 – Global Administrative Areas (2021). Administrative regions. Available at https://gadm.org/download_world.html.

Data analysis

An interactive markdown is publicly available at: https://github.com/CIAT/livestock_prioritization

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

The datasets analyzed during the current study are available in the Livestock_Prioritization repository https://github.com/CIAT/livestock_prioritization/tree/main/Data. No restrictions on data availability exist other than related to refs. 42 and 52 (see Main Text), which are not publicly available. For use of these datasets we strongly recommend contacting the first and/or corresponding authors of those studies. Access links for all datasets can be found in Supplementary Table S3.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We synthesized spatial dataset to identify adaptation and mitigation priority geographies for livestock system investments. The study area included livestock production systems in all low-income, low- and middle-income, and upper-middle-income countries in mid and low latitudes (n=132 countries).
Research sample	We analyzed n=132 countries
Sampling strategy	We selected these n=132 countries because they are Low and Middle Income Countries according to the World Bank in mid- and low latitudes.
Data collection	Data were downloaded from existing (public) repositories for all datasets except two, which were provided directly by the authors of those datasets.
Timing and spatial scale	We analyze a combination of historical and future projected conditions (see Methods and Supplementary Table S3).
Data exclusions	No data exclusions were made.
Reproducibility	All code and data are freely available in a GitHub repository. See data and code availability statements on the Main Text.
Randomization	Does not apply.
Blinding	Does not apply.
Did the study involve field work?	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern

Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging