

## Mental Health and Loneliness in University Students During the COVID-19 Pandemic in Germany: A Longitudinal Study

Weber, Maxi; Schulze, Lars; Bolzenkötter, Teresa; Niemeyer, Helen; Renneberg, Babette

Veröffentlichungsversion / Published Version

Zeitschriftenartikel / journal article

### Empfohlene Zitierung / Suggested Citation:

Weber, M., Schulze, L., Bolzenkötter, T., Niemeyer, H., & Renneberg, B. (2022). Mental Health and Loneliness in University Students During the COVID-19 Pandemic in Germany: A Longitudinal Study. *Frontiers in Psychiatry*, 13, 1-11. <https://doi.org/10.3389/fpsyt.2022.848645>

### Nutzungsbedingungen:

Dieser Text wird unter einer CC BY Lizenz (Namensnennung) zur Verfügung gestellt. Nähere Auskünfte zu den CC-Lizenzen finden Sie hier:

<https://creativecommons.org/licenses/by/4.0/deed.de>

### Terms of use:

This document is made available under a CC BY Licence (Attribution). For more information see:

<https://creativecommons.org/licenses/by/4.0>



# Mental Health and Loneliness in University Students During the COVID-19 Pandemic in Germany: A Longitudinal Study

Maxi Weber<sup>1\*</sup>, Lars Schulze<sup>1</sup>, Teresa Bolzenkötter<sup>1</sup>, Helen Niemeyer<sup>2</sup> and Babette Renneberg<sup>1</sup>

<sup>1</sup> Clinical Psychology and Psychotherapy, Freie Universität Berlin, Berlin, Germany, <sup>2</sup> Clinical Psychological Intervention, Freie Universität Berlin, Berlin, Germany

## OPEN ACCESS

### Edited by:

Haibo Yang,  
Tianjin Normal University, China

### Reviewed by:

Melissa Macalli,  
Université de Bordeaux, France  
Jeff Huarcaya,  
Hospital Base Guillermo Almenara  
Irigoyen, Peru

### \*Correspondence:

Maxi Weber  
maxi.weber@fu-berlin.de

### Specialty section:

This article was submitted to  
Public Mental Health,  
a section of the journal  
Frontiers in Psychiatry

Received: 04 January 2022

Accepted: 16 March 2022

Published: 15 April 2022

### Citation:

Weber M, Schulze L, Bolzenkötter T,  
Niemeyer H and Renneberg B (2022)  
Mental Health and Loneliness in  
University Students During the  
COVID-19 Pandemic in Germany: A  
Longitudinal Study.  
Front. Psychiatry 13:848645.  
doi: 10.3389/fpsy.2022.848645

The COVID-19 pandemic and its preventive measures had adverse consequences for mental health. However, knowledge of mental health trajectories across the pandemic is limited. This study investigated the mental health levels and changes among university students during the pandemic and lockdown in Germany, as well as their associated factors. We surveyed students' mental health ( $N = 363$ , 68% female) with the patient health questionnaire (PHQ-8) and the generalized anxiety disorder scale (GAD-7) during the first easing phase (July 2020; time 1) and the second lockdown (November 2020; time 2). Cut-off scores from the GAD-7 and PHQ-8 were used to determine clinically relevant symptoms and to define trajectory groups. Sociodemographic and pandemic-related data were assessed (e.g., coping with academic life, social contacts) as well as loneliness, stress, repetitive negative thinking, quality of life, and perceived social support. Paired  $t$ -test, multiple regression, and repeated-measures ANOVA were applied. Means and prevalence rates for symptoms of depression (38.8%) and anxiety (25.6%) did not differ between time 1 and time 2, and most students were asymptomatic on the PHQ-8 (44.4%) and the GAD-7 (56.3%) across the pandemic. Feelings of loneliness significantly increased from time 1 to time 2,  $d = -0.30$ ,  $[-0.47, -0.13]$ , with higher symptom levels in symptomatic groups at time 2 and greater increases in the asymptomatic groups. Levels of stress, repetitive negative thinking, quality of life, and social support did not differ during the pandemic. At time 1, loneliness and repetitive negative thinking were associated with anxiety and depressive symptoms. Anxiety and depressive symptoms were prevalent among students, and increased levels of loneliness during the pandemic were associated with elevated symptoms and differing trajectories. Further research using representative and larger samples should determine the long-term impact of the pandemic on mental health and loneliness to identify vulnerable students and offer adequate support.

**Keywords:** COVID-19, pandemic, depression, anxiety, loneliness, students, university

## INTRODUCTION

The first wave of the coronavirus disease (COVID-19) pandemic and its lockdown measures negatively affected the mental health of many individuals (1, 2). However, specific subgroups at higher risk for mental health problems were described, including university and college students (3–7). Even before COVID-19, students were exposed to multiple stressors during the emergent adulthood adapting to social and academic life (8–11). With the COVID-19 pandemic, further potential stressors emerged due to closed universities, remote learning formats, and prolonged social distancing measures.

Studies across the globe revealed elevated symptoms of anxiety, depression, stress, and loneliness among university and college students during the pandemic (12–23). However, most of the studies applied cross-sectional designs and mental health impacts should be interpreted with caution (24). Some studies provided longitudinal data comparing the same students before and closely after the first peak of the pandemic to examine how mental health has changed. Compared with pre-pandemic levels, the majority of studies likewise showed increased symptoms of depression, anxiety, and stress (12, 25–32), but not all (33). Mixed findings were reported by Meda et al. (34) showing increased symptoms for depression but not for anxiety among students during the first lockdown in Italy compared to pre-pandemic levels, while another study among medical students in India suggested increased rates for anxiety but not for depression (35).

A few additional longitudinal studies compared different time points during the pandemic to examine how mental health has changed and revealed conflicting results. For example, studies among Chinese college students found increased anxiety and depression rates when the pandemic was under control compared with the acute phase of the pandemic, but not for self-reported stress (22, 36). Two other studies indicated decreased anxiety and depression symptoms (37), and stress levels (38) during the first lockdown compared to the pre-pandemic academic semester before increasing again in the post-lockdown period. In contrast, others reported generally high but declining anxiety and depression symptoms along with reduced daily COVID-cases and eased lockdown measures in Italy (34) and the United States (39). One recent repeated cross-sectional study with a large sample of students and non-students compared anxiety and depressive symptoms at three pandemic time points in France (40). Relative to non-students, students showed higher depressive symptoms during the first national lockdown (19% vs. 36%), comparable rates during the easing phase (21% vs. 27%), and again dramatic increases during the second lockdown (27% vs. 54%). Symptoms of anxiety were likewise more prevalent in students compared with non-students during the pandemic.

In addition to the mixed findings on mental health courses during the pandemic, it is less understood which risk and protective factors co-determine mental health levels and changes among university students during the pandemic. Previous cross-sectional data largely based on the general population suggested that increased symptoms of anxiety and depression were associated with female gender, younger age, living alone,

and financial insecurities (3, 23, 41–47). In addition, these studies implied that adverse coping styles, repetitive negative thinking, boredom, pre-existing mental health conditions, and adverse childhood experiences were associated with worsened mental health, while perceived social support, having social contacts, and self-efficacy were protective for mental health.

In recent years, loneliness has consistently been linked to poorer mental health, symptoms of anxiety and depression. (e.g., (48–50)). With the COVID-19 pandemic and the established social distancing measures, the link between loneliness and mental health was further emphasized (4). In fact, loneliness during the pandemic increased in the general population (51) as well as in university students compared with pre-pandemic levels (12, 29), and this increase was more prevalent in students compared with non-students (52). Moreover, loneliness was largely responsible for the exacerbated course of depressive symptoms in young adults during the pandemic (53).

Overall, knowledge is limited regarding the mental health levels of university students and trajectories after prolonged threats and stressors during the COVID-19 pandemic. Longitudinal research is warranted on how mental health changes and which stressors may be of primary concern to target prevention efforts, particularly following multiple lockdowns. In Germany, the first lockdown started in March 2020 with easing steps from May to June 2020; the second four months later in November 2020 to May 2021. Most shops, restaurants, and universities were closed and gatherings of more than five people were banned. Between the two lockdowns, restaurants and shops re-opened and contact restrictions were eased while universities remained closed. Additional measures were maintained during the easing phase such as a minimum distance of 1.5 m to others, wearing face masks in public transport, and the recommendation to reduce physical contacts whenever possible.

Using a cross-sectional and longitudinal cohort design with measures during the easing phase and the second lockdown in Germany, the current study aimed at 1) investigating the levels and changes of mental health in university students, and 2) identifying associated factors for mental health (i.e., sociodemographic, pandemic-related, and psychological variables). As registered previously (osf.io/na5b6), we expected overall worsened primary anxiety and depressive symptoms in students and worsened psychological outcomes (i.e., loneliness, stress, repetitive negative thinking, quality of life, social support) along with increasing COVID-19 cases and deaths, and re-introduced lockdown measures. To better understand how mental health changed during the pandemic, we examined trajectories based on the clinical cut-off scores for probable anxiety and depression at each time point and their associations with changes in loneliness. Moreover, we tested whether anxiety and depressive symptom levels were associated with sociodemographic variables (e.g., female gender, living alone, socioeconomic status), pandemic-related variables (e.g., coping with daily and academic life, reduced social contacts since the pandemic), and psychological variables (e.g., higher levels of loneliness, ruminative thinking, lower perceived social support, current mental disorder).

## METHODS

### Study Design and Participants

This cross-sectional and longitudinal online survey study was conducted according to the Strengthening the Reporting of Observational Studies in Epidemiology guidelines for observational studies (54). Data was collected 2 months after the first lockdown in Germany (20 July–28 August 2020; time 1), and during the second lockdown when rates of COVID-19 infections and related deaths increased dramatically (10 November–2 December 2020; time 2). Participants older than 18 years studying in Berlin, Germany (no other inclusion or exclusion criteria) were recruited via social media, mailing lists, and the institutional website. The study was approved by the ethics committee at Freie Universität Berlin (032/2020).  $N = 467$  students provided informed consent and initiated the survey via the *Questback* platform. We included  $n = 363$  (77.7%) participants for the cross-sectional analyses at time 1 with complete questions regarding the primary outcomes, of which 343 (94%) completed the whole survey.  $N = 254$  participants agreed to participate in the survey at time 2, of which 157 respondents completed the whole survey. Matched data at time 1 and time 2 was available for 135 respondents. This sample size entailed more than 90% power to observe a small within-effect at the 5% level ( $G^*$ Power 3.1.9.2,  $F$ -test, repeated-measures ANOVA). Participants completing the survey at time 1 were entered into a raffle to receive one of ten 25 € gift cards. Psychology students from Freie Universität Berlin could receive course credits after each wave.

### Measurements

#### Sociodemographic and Pandemic-Related Variables

The questionnaire battery at time 1 comprised data related to age, gender, family status, living situation, highest degree, field of study, students' income, and the socioeconomic status indexed by the degree and profession of the student parents (55).

To measure pandemic-specific experiences, additional items were formulated. Participants rated their perceived wellbeing and their coping abilities in daily life, in academic life, and with a potential future lockdown on a five-point Likert scale, ranging from 1 "good" to 5 "poor" (e.g., *How have you been feeling in general since the pandemic?*). Perceived wellbeing and finances compared to before the pandemic were rated on a five-point Likert scale, ranging from 1 "strongly improved" to 5 "strongly worsened" (e.g., *Has your financial situation changed since the pandemic?*). Finally, participants reported the number of days during the last 2 weeks (0–14 days) that they had social contacts and consumed alcohol, respectively (see [osf.io/na5b6](https://osf.io/na5b6) for study materials used).

#### Primary Mental Health Outcomes

Anxiety and depression symptoms were assessed with the well-validated 7-item Generalized Anxiety Disorder Scale [GAD-7, (56, 57)] and the 8-item Patient Health Questionnaire [PHQ-8, (58)] with equivalent diagnostic accuracy compared to the PHQ-9 (59). Total scores range from 0 to 21 for the GAD-7 and from 0 to 24 for the PHQ-8. Scoring 10 or above indicates moderate-to-severe symptomatology, which typically

represents clinically significant depression and anxiety (58, 60). Both instruments demonstrated excellent internal consistency in the present study (GAD-7,  $\alpha = 0.87$ ; PHQ-8, Cronbach's  $\alpha = 0.86$ ). Open-ended responses toward the most distressing and most positive experiences and perceived changes in academic life during the last 2 weeks were gathered to cross-validate symptoms levels, which will be presented elsewhere in detail.

#### Additional Psychological Variables

Outcomes assessed with reliable and valid questionnaires at the two time points were loneliness [UCLA loneliness scale, ULS-8, (61)] stress [perceived stress scale, PSS-10, items 3, 6, (62, 63)], quality of life [satisfaction with life scale, SWLS, (64, 65)], repetitive negative thinking [perseverative thinking questionnaire, (66)], and social support [brief form of perceived social support questionnaire, (67)]. Single-measure items were applied to measure feelings of boredom, presence of diagnosed mental disorder, and subjective health at time 1 (68). Associated factors with mental health at time 1 included coping strategies [active coping, positive reframing, acceptance, religion, and substance use, Brief-COPE, (69, 70)], self-efficacy [generalized self-efficacy scale; (71)], social anxiety [mini social phobia inventory, (72)], and adverse childhood experiences [ACE, (73)]. The applied scales proved acceptable to excellent internal consistency in this study (Cronbach's  $\alpha = 0.72$ – $0.96$ ), except for the coping subscales religion and active coping ( $\alpha = 0.64, 0.69$ ), which were subsequently removed from further analyses.

#### Data Analysis

The Welch's  $t$ -test, univariate analysis of variance (ANOVA) and Chi-square test of independence were used to test differences between the cross-sectional sample and the longitudinal sample. To determine research question 1) regarding the mental health levels and changes during the pandemic, paired sample  $t$ -tests and their respective effect size estimates using Cohen's  $d$  were used to examine mean changes in variables tested at time 1 and time 2 (i.e., anxiety, depression, loneliness, stress, quality of life, social support, and repetitive negative thinking). Cohen's  $d$  of 0.2, 0.5, and 0.8 indicate a small, medium, and large effect size, respectively (74). Regarding the dependent variables anxiety and depressive symptoms, clinically relevant symptom levels at time 1 and time 2 were determined using the established cut-off score of 10 of the GAD-7 and PHQ-8 (58, 60). To adjust trajectories during the pandemic [e.g., (75)], the cut-off scores at time 1 and time 2 were used, resulting in four potential paths for anxiety and depression: 1) the asymptomatic, 2) the worsened, 3) the symptomatic, and 4) the improved trajectory. To further explore how mental health changed during the pandemic, two-way repeated measures ANOVAs with loneliness as the dependent variable were performed with the factors time and trajectory groups for depression and anxiety. Post hoc analysis using the Tukey's test (76) were applied as well as partial eta squared ( $\eta_p^2$ ) as measures of small (0.01), medium (0.06), and large (0.14) effect sizes (74).

To examine research question 2) on factors associated with increased anxiety and depressive symptoms, we performed multiple linear regression analyses for the two primary

outcomes individually, as others have done (15, 77). The two models tested associations measured at time 1, respectively, with sociodemographic (e.g., age, gender, living situation, socioeconomic status), pandemic-related (e.g., coping with academic life since the pandemic, social contacts), and additional psychological variables (e.g., loneliness, social support, presence of current mental disorder). Associated factors for the two primary outcomes were first examined in univariate linear regression analyses and subsequently entered into a multiple linear regression model adjusting for all other tested sociodemographic, pandemic-related, and psychological factors. The statistical assumptions were tested regarding multicollinearity (i.e., tolerance and VIF factor  $\leq 2$ ). Residual and scatter plots indicated that the assumptions toward normality, linearity, and homoscedasticity were met.

Nine percent of observations at time 1 had missing values in the independent variables and were assessed with multivariate imputation by chained equations following the conditional multiple imputation approach (78). Associated factors from the regression analysis models were included in the imputation model for the dependent variable anxiety symptoms and for depressive symptoms, respectively. Twenty data sets were each imputed and subsequently pooled using Rubin's rules [(79); see (80) for an overview]. Sensitivity analyses were applied to explain any differences between the complete case analysis using list-wise deletion and the multiple imputation approach. All statistical analyses were conducted using R version 4.0.2. (81). *P*-values < .05 indicated statistical significance.

## RESULTS

### Sample Characteristics

Participants at time 1 ( $N = 363$ ) were mostly female (68%) and had a mean age of 26 years ( $SD = 4.27$ ; **Table 1**). About half of the participants were single (49.2%), undergraduate students (46.4%), and the majority lived with others (75.2%). Most of the participants' parents (63%) had a middle economic status, while students themselves had an average income of 700–1000 Euro/month or less (67.1 %). On average, participants reported an overall very good or good health status ( $M = 1.76$ ,  $SD = 0.73$ ). Sixty-two participants (17.1%) self-reported a diagnosed mental disorder; most frequently named were anxiety and depressive disorders. Sociodemographic characteristics at time 1 did not differ between the cross-sectional sample and the longitudinal sample with matched data pairs, but slightly more participants with matched data had reported a mental disorder at time 1,  $\chi^2(2, N = 498) = 9.63, p = .008$  (**Table 1**).

### Pandemic-Related Responses and Changes

Wellbeing since the pandemic was overall perceived as moderate (40%) or somewhat good (27.5%) at time 1, and slightly worse or worse (61%) compared to pre-pandemic levels (**Table 2**). Students stated to cope rather well (38.3%) with their daily lives since the pandemic and that their income at time 1 did not change (52.5%) compared to pre-pandemic levels. At time 1, students slightly agreed or agreed (43.5%) to fear a potential

**TABLE 1** | Sample characteristics.

	Cross-sectional sample n (%) (n = 363)	Longitudinal sample n (%) (n = 135)
<b>Age</b>		
mean (SD)	25.87 (4.69)	25.32 (3.83)
median	25	25
<b>Gender<sup>a</sup></b>		
female	247 (68.0%)	100 (74.1%)
male	116 (32.0%)	35 (25.9%)
<b>Family status</b>		
single	179 (49.2%)	68 (50.4%)
partnership	169 (46.6%)	61 (45.2%)
other	15 (4.1%)	6 (4.4%)
having children	19 (5.2%)	6 (4.4%)
<b>Living situation</b>		
with others	273 (75.2%)	96 (71.1%)
alone	90 (24.8%)	39 (28.9%)
<b>Highest degree</b>		
High school diploma	169 (46.4%)	71 (52.6%)
Bachelor's degree	158 (43.5%)	53 (39.3%)
Master's degree	36 (9.9%)	11 (8.1%)
<b>University in Berlin</b>		
Freie Universität	128 (35.3%)	57 (42.2%)
Technical University	55 (15.2%)	16 (11.9%)
Humboldt University	53 (14.6%)	18 (13.3%)
Other	127 (35.0%)	44 (32.6%)
<b>Field of study</b>		
Social sciences	146 (40.2%)	57 (42.2%)
Humanities and arts	73 (20.1%)	26 (19.1%)
Natural sciences	43 (11.8%)	16 (11.8%)
Engineering	41 (11.3%)	9 (6.6%)
Economics and politics	40 (11.0%)	16 (11.8%)
Other	4 (1.1%)	2 (1.5%)
<b>Income (€)</b>		
<700	101 (27.9%)	38 (28.1%)
700–1000	142 (39.2%)	57 (42.2%)
1001–1300	60 (16.5%)	18 (13.3%)
>1301–1700	60 (16.5%)	22 (16.3%)
<b>Parents' SES</b>		
high	65 (18.6%)	22 (16.7%)
average	220 (63.0%)	87 (65.9%)
low	64 (18.3%)	23 (17.4%)
missing n	13 (3.6%)	3 (2.2%)
<b>Health status</b>		
mean SD	1.76 (0.73)	1.77 (0.67)
missing n	21 (5.6%)	0 (0.0%)
<b>COVID risk group (yes)</b>		
missing n	21 (5.6%)	0 (0.0%)
<b>Reported mental disorder (yes)</b>		
missing n	21 (5.6%)	0 (0.0%)

*SD*, standard deviation; *SES*, socioeconomic status.

<sup>a</sup>None of the participants identified as diverse.

**TABLE 2** | COVID-19 related responses and changes during the pandemic.

Variables	Cross-sectional sample	Longitudinal sample		Paired <i>t</i> test (134)	<i>p</i> value
		Time 1	Time 2		
		Mean (SD)			
Perceived Wellbeing since COVID <sup>a</sup>	2.65 (0.99)	2.65 (0.96)	2.66 (0.98)	−0.06	0.950
Wellbeing since COVID vs. pre-COVID <sup>a</sup>	3.50 (1.01)	3.58 (1.00)	3.40 (1.04)	1.32	0.188
Coping with changes in daily life since COVID <sup>a</sup>	2.48 (0.90)	2.51 (0.93)	2.36 (0.99)	1.26	0.209
Coping with academic life since COVID <sup>a</sup>	2.92 (1.21)	3.01 (1.14)	2.65 (1.17)	2.41	0.017
Changed income since COVID (time 1) <sup>a</sup>	3.44 (0.81)	3.36 (0.76)	-		
Afraid towards potential future lockdown (time 1) <sup>a</sup>	3.14 (1.27)	3.13 (1.34)	-		
Coping with potential future lockdown (time 1) <sup>a</sup>	2.67 (1.01)	2.65 (1.04)	-		
Days/last 2 weeks with social contacts	5.26 (3.72)	4.79 (3.76)	3.17 (2.95)	3.90	<0.001
Days/last 2 weeks with social contacts vs. pre-COVID	2.19 (1.07)	2.04 (1.00)	1.61 (0.84)	3.99	<0.001
Days/last 2 weeks drinking alcohol	3.33 (3.14)	3.08 (3.02)	2.07 (2.29)	3.08	0.002
Days/last 2 weeks drinking alcohol vs. pre-COVID	2.93 (0.98)	2.87 (0.92)	2.60 (0.97)	2.42	0.017

<sup>a</sup> Higher scores indicate poor or worsened outcomes using a five-point Likert scale.

future lockdown, and at the same time, indicated they would cope rather well a second lockdown (46.3%). Students coped moderately well (30.0%) with their academic life's at time 1, which significantly improved from time 1 to time 2. The average number of days during the last 2 weeks having social contacts and drinking alcohol decreased compared to pre-pandemic levels, and decreased further from time 1 to time 2, respectively. All other ratings did not differ between the two assessments (Table 2).

## Mental Health Levels and Trajectories During the Pandemic

Table 3 shows the results for the primary mental health outcomes depressive symptoms and anxiety symptoms for the cross-sectional and the longitudinal sample. Prevalence rates in the cross-sectional sample were 38% and 26.5% for moderate-to-severe depressive and anxiety symptoms, respectively. In the longitudinal sample, prevalence rates did not change for depressive symptoms (43% vs. 43%) and anxiety symptoms (29.6% vs. 28.1%). Between time 1 and time 2, mean scores did not significantly differ for symptoms of depression,  $t(134) = -0.09$ ,  $p = 0.924$ ,  $d = -0.01$ , 95% CI [-0.16, 0.14], and anxiety,  $t(134) = -0.04$ ,  $p = 0.968$ ,  $d = -0.004$ , 95% CI [-0.17, 0.16]. Most participants showed asymptomatic PHQ-8 trajectories from time 1 to time 2 (44.4%), followed by stable symptomatic (30.4%), worsened (12.6%), and improved depressive trajectories (12.6%). Similarly, most GAD-7 trajectories during the pandemic were asymptomatic (56.3%), followed by improved (15.5%), stable symptomatic (14.1%), or worsened trajectories (14.1%).

Regarding the additional psychological outcomes, symptoms of stress and repetitive negative thinking did not significantly differ between time 1 and time 2 (Table 3). In addition, perceived social support and quality of life did not differ during the course of the pandemic. However, feelings of loneliness significantly increased among students between time 1 and time 2,  $t(134) =$

$-2.63$ ,  $p = 0.009$ . The effect size for increased loneliness was small, Cohen's  $d = -0.30$ , 95% CI [-0.47, -0.13]).

To further explore the increases in loneliness during the pandemic, two-way repeated-measures ANOVAs were performed with loneliness across time and between different PHQ-8 and GAD-7 trajectories. Results indicated a large and significant difference in loneliness between the PHQ-8 trajectory groups,  $F(3, 131) = 11.49$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.16$ , and a medium effect of time,  $F(1, 131) = 13.24$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.03$  (Figure 1). In addition, there was a significant and medium interaction effect between increased loneliness during the pandemic and the PHQ-8 trajectory groups  $F(3, 131) = 4.09$ ,  $p = 0.008$ ,  $\eta_p^2 = 0.02$ . Post hoc comparisons revealed that between time 1 and time 2, loneliness significantly increased in the asymptomatic PHQ-8 trajectory group ( $p = 0.002$ ), while increases in the symptomatic and worsened courses did not reach statistical significance. At time 2, compared to asymptomatic PHQ-8 courses loneliness was more pronounced in groups with symptomatic ( $p = 0.007$ ) and worsened courses ( $p = 0.043$ ), but did not differ from the improved trajectory group ( $p = 0.998$ ). Regarding GAD-7 trajectories, effects of group and time were statistically significant and large or medium-sized,  $F(3, 131) = 9.72$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.13$ ;  $F(3, 131) = 12.61$ ,  $p < 0.001$ ,  $\eta_p^2 = 0.02$  (Figure 1). The overall interaction between GAD-7 trajectories and loneliness was not significant ( $p = 0.149$ ,  $\eta_p^2 = 0.01$ ). However, loneliness significantly increased in the asymptomatic GAD-7 trajectory group ( $p = 0.012$ ). At time 2, loneliness was more prevalent in groups with stable symptomatic compared to asymptomatic GAD-7 courses ( $p = 0.014$ ) and did not differ from worsened or improved GAD-7 trajectories ( $p = 0.548$ ;  $p = 0.823$ ).

## Associated Factors for Mental Health Levels and Changes

Table 4 shows the unadjusted and adjusted associations at time 1 with sociodemographic, pandemic-related and psychological variables, separately for the primary outcome of mental health

**TABLE 3** | Primary outcomes and additional psychological variables during the COVID-19 pandemic.

Variables	Cross-sectional sample	Longitudinal sample		Paired <i>t</i> test (134)	<i>p</i>	Cohen's <i>d</i>	95 % CI
		Time 1	Time 2				
Depression (PHQ-8)							
Mean ( <i>SD</i> )	8.12 (5.24)	8.37 (5.52)	8.43 (4.63)	−0.09	0.924	−0.01	[−0.16, 0.14]
Median ( <i>range</i> )	7.00 (0–23)	8.00 (0–23)	8.00 (0–21)				
Moderate-to-severe ( <i>N</i> , %)	141 (38.8%)	58 (43.0%)	58 (43.0%)				
Anxiety (GAD-7)							
Mean ( <i>SD</i> )	7.15 (4.64)	7.50 (4.71)	7.52 (4.30)	−0.04	0.968	−0.004	[−0.17, 0.16]
Median ( <i>range</i> )	6.00 (0–21)	7.00 (0–21)	7.00 (0–20)				
Moderate-to-severe ( <i>N</i> , %)	93 (25.6%)	40 (29.6%)	38 (28.1%)				
Loneliness							
Mean ( <i>SD</i> )	15.61 (5.51)	15.96 (5.93)	17.61 (5.04)	−2.63	0.009	−0.30	[−0.47, −0.13]
Median ( <i>range</i> )	14 (8–30)	14 (8–29)	17 (8–28)				
Stress <sup>a</sup>							
Mean ( <i>SD</i> )	3.91 (1.67)	4.09 (1.78)	4.42 (1.59)	−1.73	0.086	−0.20	[−0.37, −0.02]
Median ( <i>range</i> )	4 (2–8)	4 (2–8)	4 (2–8)				
Satisfaction with life							
Mean ( <i>SD</i> )	23.66 (6.58)	23.82 (6.59)	23.33 (6.17)	0.66	0.509	0.08	[−0.05, −0.21]
Median ( <i>range</i> )	25 (5–34)	25 (6–34)	25 (5–34)				
Perceived social support							
Mean ( <i>SD</i> )	19.83 (3.96)	19.96 (3.81)	19.98 (3.54)	−0.03	0.972	−0.004	[−0.14, −0.12]
Median ( <i>range</i> )	21 (6–24)	21 (7–24)	21 (9–24)				
Repetitive negative thinking							
Mean ( <i>SD</i> )	29.00 (12.59)	30.12 (12.67)	28.58 (13.10)	1.07	0.285	0.11	−0.03, −0.26]
Median ( <i>range</i> )	29 (0–56)	30 (5–57)	29 (0–56)				

*SD*, standard deviation; *PHQ-8*, patient health questionnaire-8; *GAD-7*, generalized anxiety disorder-7.

<sup>a</sup>Stress was measured with two items from the perceived stress scale (Cronbachs'  $\alpha = 0.85$ ).

(depressive symptoms, anxiety symptoms). When adjusting for all other tested variables, increased depressive symptoms at time 1 were associated with worse coping abilities in daily life ( $B = 0.64$ ;  $SE = 0.29$ ) and worse coping with academic life since the pandemic ( $B = 0.74$ ;  $SE = 0.20$ ), with higher levels of loneliness ( $B = 0.24$ ;  $SE = 0.05$ ), social anxiety ( $B = 0.20$ ;  $SE = 0.08$ ), boredom ( $B = 0.46$ ;  $SE = 0.19$ ), and repetitive negative thinking ( $B = 0.11$ ;  $SE = 0.02$ ;  $adjR^2 = 52.3\%$ ,  $p < .001$ ). Regarding anxiety symptoms at time 1, adjusted associations were found for living alone ( $B = -0.91$ ;  $SE = 0.44$ ), worse coping with academic life ( $B = 0.48$ ;  $SE = 0.18$ ), worse coping with a potential future lockdown ( $B = 0.51$ ;  $SE = 0.26$ ), loneliness ( $B = 0.20$ ;  $SE = 0.05$ ), and repetitive negative thinking ( $B = 0.13$ ;  $SE = 0.02$ ;  $adjR^2 = 59.38\%$ ,  $p < .001$  including all variables). Sensitivity analyses with complete data applying list-wise deleted revealed similar conclusions compared to the multiple imputation approach presented for both depressive and anxiety symptoms (see **Supplementary Table 1**).


## DISCUSSION

This study investigated the mental health levels and trajectories of university students during two different stages of the COVID-19 pandemic, i.e., at time 1 during the eased lockdown phase and

at time 2 during the second lockdown in Germany. Contrary to the hypotheses, mean symptoms of depression and anxiety did not significantly change during the pandemic. Students most often had asymptomatic or sustained symptomatic courses of depression (56.3%, 30.4%) and anxiety (44.4%, 10.5%) during the pandemic; fewer worsened or improved between time 1 and time 2. Likewise, mean levels of stress, perceived quality of life, and social support did not change over the course of the pandemic. However, in line with the hypotheses, feelings of loneliness increased from time 1 to time 2. Higher levels of loneliness during the lockdown phase were present in participants with sustained or worsened symptom trajectories, while increases in loneliness were most prevalent in those with asymptomatic courses of depression and anxiety. Moreover, loneliness and repetitive negative thinking were associated with anxiety and depressive symptoms measured at time 1. Here, we discuss the results on mental health in the context of the COVID pandemic and its preventive measures (e.g., social distancing, closure of universities) together with further implications for students in higher education.

## Findings in Context

During the eased lockdown phase in July 2020 at time 1, anxiety (GAD-7) and depression (PHQ-8) scores were more than twice as



**FIGURE 1 |** Feelings of loneliness and trajectories of depressive symptoms (A) and anxiety symptoms (B) during the COVID-19 pandemic.

high as normative and pre-pandemic data for German university students (9, 57, 82). However, mean scores and clinically relevant rates for depressive symptoms (38.8%) in our cross-sectional sample are comparable to pandemic data of German university students assessed between June and August 2020 [37%, (13); 38.5%, 29]. Prevalence rates for anxiety symptoms at time 1 were slightly lower as reported previously during the eased lockdown phase in Germany [25.6% vs. 35.5%; (29)]. Although rates for anxiety and depressive symptoms at time 1 were slightly higher in participants with matched data at time 1 and time 2 compared to the cross-sectional sample, our results fit in with the pooled prevalence rates of anxiety and depression among students from Western countries during the pandemic [e.g., (83, 84)].

In this study, anxiety and depressive symptoms did not change during the course of the pandemic and lockdown, which contradicts our hypotheses as well as recent findings among French students with increased rates between the eased lockdown phase and the second lockdown (40). However, symptom trajectories differed during the pandemic. While most students had asymptomatic courses, approximately four of ten students had a stable symptomatic or worsened depressive course, and three out of ten faced adverse anxiety courses during the pandemic. In addition, and contrary to our hypotheses, levels did not change regarding stress, quality of life, perceived social support, and repetitive negative thinking during the pandemic, reflecting previous mixed longitudinal data [e.g., (22, 36–38)]. First, our findings indicate that most university students reported slightly worse or worse wellbeing at time 1 compared with pre-pandemic levels, which is in line with prior research [e.g., (12, 27, 28, 31)]. These findings generally point to decreased wellbeing during the pandemic, while symptom levels likely persisted between time 1 and time 2. However, coping with academic life improved during the pandemic, which implies that most students adapted to the remote studying formats. Second, this study was conducted at the end of the semester at time 1, and elevated symptoms levels at time 1 may also have developed partly in response to the examination phase (32). The survey at time 2 was conducted shortly after the second lockdown had started, and its long-term consequences were possibly not yet tangible.

Consistent with previous assumptions and data (4, 12, 53) but inconsistent with others (31), loneliness increased among university students during the pandemic. Loneliness particularly

increased in students with asymptomatic trajectories during the pandemic, and the highest levels were present in students with symptomatic or worse trajectories. This mirrors both trajectories and symptom levels during the pandemic among the general population compared to people with pre-existing mental disorders (51). Although aligned with the established social distancing measures, social contacts decreased compared with pre-pandemic levels and further decreased during the pandemic, likely fostering loneliness in the current sample. This finding raises concerns given that loneliness is a crucial risk factor for mental health in general (48) as well as a major reason for increased helpline calls during the pandemic in Germany, and worldwide (85).

Moreover, loneliness was significantly associated with both depressive and anxiety symptoms at time 1 along with repetitive negative thinking, while adaptive coping in daily and academic life was protective for depressive symptoms, and adaptive coping with a potential future lockdown was protective for anxiety symptoms. In addition, these findings generally fit within the literature (41–47), suggesting that the way students appraise the pandemic, as well as their connections with others, may be critical in understanding the mental health during the pandemic.

The current study provides novel evidence on mental health before and during a second lockdown, identified loneliness and repetitive negative thinking as salient risk factors for mental health, and demonstrated diverging trajectories of mental health in a homogenous sample of German university students. The findings on prevalent anxiety and depressive symptoms and increased loneliness during the pandemic may foster immediate preventive actions such as psychoeducation in higher education, but also stimulate research on interventions targeting loneliness among young adults. However, the study also faces limitations. First and most importantly, the sample sizes were small and the response rate at time 2 was low (43.3 %), increasing the risk for inflated data. These numbers are comparable to other studies on university students during the pandemic [e.g., (29)], but results should be replicated with representative and larger sample sizes. Our sample consisted of participants studying at Berlin-based universities, and thus our data may not be generalized to other (student) populations. Second, the current study assessed the mental health levels and changes only twice during the pandemic, as well as the pre-pandemic well-being in



**TABLE 4 |** Associations with the two dependent variables depressive symptoms and anxiety symptoms at time 1 (N = 363) using simple linear (unadjusted models) and multiple linear regression analyses (adjusted models.).

Independent variables	Depressive Symptoms (time 1)						Anxiety Symptoms (time 1)					
	Unadjusted model			Adjusted model <sup>a</sup>			Unadjusted model			Adjusted model <sup>a</sup>		
	Beta	SE	p	Beta	SE	p	Beta	SE	p	Beta	SE	p
<i>Sociodemographic data (time 1)</i>												
Age	-0.08	0.06	0.157	-0.03	0.05	0.568	0.01	0.05	0.918	0.05	0.04	0.279
Gender (male vs.)												
female	1.15	0.58	0.052	0.14	0.44	0.743	1.32	0.52	0.011	0.29	0.40	0.463
Living situation (alone vs.)												
with others	0.45	0.64	0.478	-0.36	0.48	0.460	-0.48	0.56	0.391	-0.91	0.44	0.038
Family status (single vs.)												
partnership	-0.80	0.56	0.158	-0.32	0.44	0.472	0.08	0.50	0.877	0.15	0.40	0.705
other	-1.34	1.41	0.341	0.32	1.05	0.763	-1.85	1.25	0.138	-0.11	0.96	0.909
Parents' SES (low vs.)												
middle	0.65	0.74	0.385	0.07	0.55	0.897	0.08	0.66	0.121	-0.06	0.51	0.907
high status	0.06	0.92	0.949	0.67	0.66	0.315	-0.70	0.83	0.396	0.31	0.65	0.641
<i>Pandemic-related variables (time 1)</i>												
Students' income change	0.95	0.34	0.006	0.25	0.25	0.321	0.90	0.30	0.003	0.26	0.23	0.250
Coping with daily life	2.51	0.28	<0.001	0.64	0.29	0.029	2.16	0.25	<0.001	0.37	0.27	0.162
Coping with academic life	1.63	0.21	<0.001	0.74	0.20	<0.001	1.38	0.19	<0.001	0.48	0.18	0.007
Social contacts	-0.30	0.07	<0.001	-0.06	0.06	0.305	-0.30	0.06	<0.001	-0.08	0.06	0.149
Drinking alcohol	-0.06	0.09	0.515	0.07	0.08	0.341	-0.14	0.08	0.069	0.03	0.07	0.713
Coping future lockdown	1.37	0.27	<0.001	-0.05	0.28	0.863	1.56	0.23	<0.001	0.51	0.26	0.048
Anxiety future lockdown	-0.51	0.22	<0.001	0.19	0.20	0.320	-0.65	0.19	0.001	0.07	0.17	0.676
<i>Psychological variables (time 1)</i>												
Loneliness	0.53	0.42	<0.001	0.24	0.05	<0.001	0.45	0.38	<0.001	0.20	0.05	<0.001
Cope (positive reframing)	-1.07	0.17	<0.001	-0.18	0.15	0.254	-0.86	0.15	<0.001	-0.01	0.14	0.952
Cope (acceptance)	-0.69	0.17	<0.001	0.07	0.14	0.613	-0.81	0.15	<0.001	-0.24	0.13	0.061
Cope (substance use)	0.67	0.17	<0.001	0.22	0.14	0.128	0.23	0.15	0.136	-0.07	0.14	0.626
Social support	-0.48	0.07	<0.001	0.04	0.07	0.538	-0.42	0.06	<0.001	0.00	0.06	0.962
Self-efficacy	-0.44	0.56	<0.001	-0.03	0.05	0.579	-0.35	0.05	<0.001	0.04	0.05	0.451
Social anxiety	0.79	0.92	<0.001	0.20	0.08	0.016	0.58	0.08	<0.001	0.12	0.08	0.125
Boredom	1.77	0.23	<0.001	0.46	0.19	0.016	1.01	0.20	<0.001	-0.06	0.17	0.734
Repetitive negative thinking	0.24	0.19	<0.001	0.11	0.02	<0.001	0.21	0.02	<0.001	0.13	0.02	<0.001
Adverse childhood experiences	0.72	0.15	<0.001	0.19	0.12	0.109	0.54	0.13	<0.001	0.16	0.11	0.147
Current mental disorder (yes)	3.62	0.72	<0.001	0.80	0.56	0.154	2.82	0.65	<0.001	0.57	0.50	0.257
<i>R</i> <sup>2</sup>				0.556		<0.001				0.528		<0.001
<i>adjusted R</i> <sup>2</sup>				0.523		<0.001				0.493		<0.001

SES, socioeconomic status. Positive Beta values indicate a higher risk for depressive and anxiety symptoms.

<sup>a</sup>Adjusted for all other variables listed in the table.

a retrospective manner. In the absence of pre-pandemic data, the consequences of the COVID-19 pandemic in this sample should be interpreted with caution. Third, all measures were self-administered via online surveys, which potentially confounds the validity of the results. We used cut-offs from the PHQ-8 and the GAD-7 to create subgroups with differential symptom trajectories (e.g., to study transitions from uncritical to clinically relevant states), which showed good sensitivity and specificity (58, 60). However, these measures cannot replace a structural clinical interview to diagnose a depression or anxiety disorder.

## CONCLUSION

Symptoms of anxiety and depression overall persisted during the COVID-19 pandemic while trajectories varied and feelings of loneliness significantly increased. Moreover, loneliness was associated with anxiety and depressive symptoms, indicating that preventing loneliness may help to maintain and promote mental health among university students. Representative studies on mental health, loneliness, and other associated factors are needed to fully identify students at high risk. Given that the COVID-19 pandemic and its preventive measures such as social distancing and remote learning continue for an indefinite period, long-term consequences for mental health are likely to occur, and universities should offer adequate support informed by the evidence to mitigate mental health problems and loneliness among university students.

## REFERENCES

- Schafer KM, Lieberman A, Sever AC, Joiner T. Prevalence rates of anxiety, depressive, and eating pathology symptoms between the pre- and peri-COVID-19 eras: A meta-analysis. *J Affect Disord.* (2022) 298:364–72. doi: 10.1016/j.jad.2021.10.115
- Robinson E, Sutin AR, Daly M, Jones A. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J Affect Disord.* (2022) 296:567–76. doi: 10.1016/j.jad.2021.09.098
- Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: immediate psychological responses and associated factors. *IJERPH.* (2020) 17:3165. doi: 10.3390/ijerph17093165
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* (2020) 395:912–20. doi: 10.1016/S0140-6736(20)30460-8
- Skapinakis P, Bellos S, Oikonomou A, Dimitriadis G, Gkikas P, Perdikari E, et al. Depression and its relationship with coping strategies and illness perceptions during the COVID-19 lockdown in Greece: a cross-sectional survey of the population. *Depress Res Treat.* (2020) 2020:3158954. doi: 10.1155/2020/3158954
- van der Velden PG, Contino C, Das M, van Loon P, Bosmans MW. Anxiety and depression symptoms, and lack of emotional support among the general population before and during the COVID-19 pandemic: a prospective national study on prevalence and risk factors. *J Affect Disord.* (2020) 277:540–8. doi: 10.1016/j.jad.2020.08.026
- Xiong J, Lipsitz O, Nasri F, Lui LM, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *J Affect Disord.* (2020) 277:55–64. doi: 10.1016/j.jad.2020.08.001
- Farrer LM, Gulliver A, Bennett K, Fassnacht DB, Griffiths KM. Demographic and psychosocial predictors of major depression and generalised anxiety

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Ethics Committee of the Freie Universität Berlin, Department of Education and Psychology. The participants provided their electronic informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

MW, TB, and BR initiated this project and all authors contributed to designing and implementing the survey. MW analyzed the data and wrote the initial draft. All authors have reviewed and edited earlier versions and approved the final version of this manuscript.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2022.848645/full#supplementary-material>

- disorder in Australian university students. *BMC Psychiatry.* (2016) 16:241. doi: 10.1186/s12888-016-0961-z
- Wörfel F, Gusy B, Lohmann K, Töpitz K, Kleiber D. Mental health problems among university students and the impact of structural conditions. *J Public Health.* (2016) 24:125–33. doi: 10.1007/s10389-015-0703-6
- Lipson SK, Phillips MV, Winquist N, Eisenberg D, Lattie EG. Mental health conditions among community college students: a national study of prevalence and use of treatment services. *Psychiatr Serv.* (2021) 72:1126–33. doi: 10.1176/appi.ps.202000437
- Auerbach RP, Alonso J, Axinn WG, Cuijpers P, Ebert DD, Green JG, et al. Mental disorders among college students in the world health organization world mental health surveys. *Psychol Med.* (2016) 46:2955–70. doi: 10.1017/S0033291716001665
- Elmer T, Mepham K, Stadtfeld C. Students under lockdown: comparisons of students' social networks and mental health before and during the COVID-19 crisis in Switzerland. *PLoS ONE.* (2020) 15:e0236337. doi: 10.1371/journal.pone.0236337
- Kohls E, Baldofski S, Moeller R, Klemm S-L, Rummel-Kluge C. Mental health, social and emotional well-being, and perceived burdens of university students during COVID-19 pandemic lockdown in Germany. *Front Psychiatry.* (2021) 12:643957. doi: 10.3389/fpsy.2021.643957
- Lahav Y. Psychological distress related to COVID-19 - the contribution of continuous traumatic stress. *J Affect Disord.* (2020) 277:129–37. doi: 10.1016/j.jad.2020.07.141
- Lee J, Solomon M, Stead T, Kwon B, Ganti L. Impact of COVID-19 on the mental health of US college students. *BMC Psychol.* (2021) 9:95. doi: 10.1186/s40359-021-00598-3
- Mechili EA, Saliaj A, Kamberi F, Girvalaki C, Peto E, Patelarou AE, et al. Is the mental health of young students and their family members affected during the quarantine period? evidence from the COVID-19 pandemic in Albania. *J Psychiatr Ment Health Nurs.* (2020) 28:317–25. doi: 10.1111/jpm.12672

17. Odriozola-González P, Planchuelo-Gómez Á, Irurtia MJ, Luis-García R de. Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. *Psychiatry Res.* (2020) 290:113108. doi: 10.1016/j.psychres.2020.113108
18. Padrón I, Fraga I, Vieitez L, Montes C, Romero E, A. Study on the psychological wound of COVID-19 in university students. *Front Psychol.* (2021) 12:589927. doi: 10.3389/fpsyg.2021.589927
19. Rudenstine S, McNeal K, Schulner T, Ettman CK, Hernandez M, Gvozdieva K, et al. Depression and anxiety during the COVID-19 pandemic in an urban, low-income public university sample. *J Trauma Stress.* (2021) 34:12–22. doi: 10.1002/jts.22600
20. Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on college students' mental health in the United States: interview survey study. *J Med Internet Res.* (2020) 22:e21279. doi: 10.2196/21279
21. Wang X, Hegde S, Son C, Keller B, Smith A, Sasangohar F. Investigating mental health of US college students during the COVID-19 pandemic: cross-sectional survey study. *J Med Internet Res.* (2020) 22:e22817. doi: 10.2196/22817
22. Wang C, Zhao H, Zhang H. Chinese college students have higher anxiety in new semester of online learning during COVID-19: a machine learning approach. *Front Psychol.* (2020) 11:587413. doi: 10.3389/fpsyg.2020.587413
23. Chen T, Lucock M. The mental health of university students during the COVID-19 pandemic: an online survey in the UK. *PLoS ONE.* (2022) 17:e0262562. doi: 10.1371/journal.pone.0262562
24. Meda N, Slongo I. Caution when linking COVID-19 to mental health consequences. *Brain Behav Immun.* (2020) 87:152. doi: 10.1016/j.bbi.2020.05.018
25. Fruehwirth JC, Biswas S, Perreira KM. The Covid-19 pandemic and mental health of first-year college students: examining the effect of Covid-19 stressors using longitudinal data. *PLoS ONE.* (2021) 16:e0247999. doi: 10.1371/journal.pone.0247999
26. Li HY, Cao H, Leung DY, Mak YW. The psychological impacts of a COVID-19 outbreak on college students in China: a longitudinal study. *Int J Environ Res Public Health.* (2020) 17:3933. doi: 10.3390/ijerph17113933
27. Savage MJ, Hennis PJ, Magistro D, Donaldson J, Healy LC, James RM. Nine months into the COVID-19 pandemic: a longitudinal study showing mental health and movement behaviours are impaired in UK students. *Int J Environ Res Public Health.* (2021) 18:2930. doi: 10.3390/ijerph18062930
28. Wilson OW, Holland KE, Elliott LD, Duffey M, Bopp M. The impact of the COVID-19 pandemic on US college students' physical activity and mental health. *J Phys Act Health.* (2021) 18:1–7. doi: 10.1123/jpah.2020-0325
29. Werner AM, Tibubos AN, Mülder LM, Reichel JL, Schäfer M, Heller S, et al. The impact of lockdown stress and loneliness during the COVID-19 pandemic on mental health among university students in Germany. *Sci Rep.* (2021) 11:22637. doi: 10.1038/s41598-021-02024-5
30. Yang X, Hu H, Zhao C, Xu H, Tu X, Zhang G, et al. Longitudinal study of changes in smart phone addiction and depressive symptoms and potential risk factors among Chinese college students. *BMC Psychiatry.* (2021) 21:252. doi: 10.1186/s12888-021-03265-4
31. Evans S, Alkan E, Bhangoo JK, Tenenbaum H, Ng-Knight T. Effects of the COVID-19 lockdown on mental health, wellbeing, sleep, and alcohol use in a UK student sample. *Psychiatry Res.* (2021) 298:113819. doi: 10.1016/j.psychres.2021.113819
32. Huckins JF, daSilva AW, Wang W, Hedlund E, Rogers C, Nepal SK, et al. Mental health and behavior of college students during the early phases of the COVID-19 pandemic: longitudinal smartphone and ecological momentary assessment study. *J Med Internet Res.* (2020) 22:e20185. doi: 10.2196/20185
33. van Zyl LE, Rothmann S, Zondervan-Zwijenburg MA. Longitudinal trajectories of study characteristics and mental health before and during the COVID-19 lockdown. *Front Psychol.* (2021) 12:635333. doi: 10.3389/fpsyg.2021.635333
34. Meda N, Pardini S, Slongo I, Bodini L, Zordan MA, Rigobello P, et al. Students' mental health problems before, during, and after COVID-19 lockdown in Italy. *J Psychiatr Res.* (2021) 134:69–77. doi: 10.1016/j.jpsychires.2020.12.045
35. Saraswathi I, Saikarthik J, Senthil Kumar K, Madhan Srinivasan K, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study. *PeerJ.* (2020) 8:e10164. doi: 10.7717/peerj.10164
36. Li Y, Zhao J, Ma Z, McReynolds LS, Lin D, Chen Z, et al. Mental health among college students during the COVID-19 pandemic in China: a 2-wave longitudinal survey. *J Affect Disord.* (2021) 281:597–604. doi: 10.1016/j.jad.2020.11.109
37. Li WW, Yu H, Miller DJ, Yang F, Rouen C. Novelty seeking and mental health in Chinese university students before, during, and after the COVID-19 pandemic lockdown: a longitudinal study. *Front Psychol.* (2020) 11:600739. doi: 10.3389/fpsyg.2020.600739
38. Yang Z, Ji L-J, Yang Y, Wang Y, Zhu L, Cai H. Meaning making helps cope with COVID-19: a longitudinal study. *Pers Individ Dif.* (2021) 174:110670. doi: 10.1016/j.paid.2021.110670
39. Kujawa A, Green H, Compas BE, Dickey L, Pegg S. Exposure to COVID-19 pandemic stress: associations with depression and anxiety in emerging adults in the United States. *Depress Anxiety.* (2020) 37:1280–8. doi: 10.1002/da.23109
40. Macalli M, Texier N, Schück S, Côté SM, Tzourio C. A repeated cross-sectional analysis assessing mental health conditions of adults as per student status during key periods of the COVID-19 epidemic in France. *Sci Rep.* (2021) 11:21455. doi: 10.1038/s41598-021-00471-8
41. Cao W, Fang Z, Hou G, Han M, Xu X, Dong J, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* (2020) 287:112934. doi: 10.1016/j.psychres.2020.112934
42. Ellwardt L, Präg P. Heterogeneous mental health development during the COVID-19 pandemic in the United Kingdom. *Sci Rep.* (2021) 11:15958. doi: 10.1038/s41598-021-95490-w
43. Petzold MB, Bendau A, Plag J, Pyrkosch L, Mascarell Maricic L, Betzler F, et al. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Brain Behav.* (2020) 10:e01745. doi: 10.1002/brb3.1745
44. Man J de, Buffel V, van de Velde S, Bracke P, van Hal GF, Wouters E. Disentangling depression in Belgian higher education students amidst the first COVID-19 lockdown (April–May 2020). *Arch Public Health.* (2021) 79:3. doi: 10.1186/s13690-020-00522-y
45. González-Sanguino C, Ausin B, Castellanos MÁ, Saiz J, López-Gómez A, Ugidos C, et al. Mental health consequences during the initial stage of the 2020 coronavirus pandemic (COVID-19) in Spain. *Brain Behav Immun.* (2020) 87:172–6. doi: 10.1016/j.bbi.2020.05.040
46. Sheldon E, Simmonds-Buckley M, Bone C, Mascarenhas T, Chan N, Wincott M, et al. Prevalence and risk factors for mental health problems in university undergraduate students: a systematic review with meta-analysis. *J Affect Disord.* (2021) 287:282–92. doi: 10.1016/j.jad.2021.03.054
47. Ye B, Zeng Y, Im H, Liu M, Wang X, Yang Q. The relationship between fear of COVID-19 and online aggressive behavior: a moderated mediation model. *Front Psychol.* (2021) 12:589615. doi: 10.3389/fpsyg.2021.589615
48. Beutel ME, Klein EM, Brähler E, Reiner I, Jünger C, Michal M, et al. Loneliness in the general population: prevalence, determinants and relations to mental health. *BMC Psychiatry.* (2017) 17:97. doi: 10.1186/s12888-017-1262-x
49. Cacioppo JT, Hughes ME, Waite LJ, Hawkley LC, Thisted RA. Loneliness as a specific risk factor for depressive symptoms: cross-sectional and longitudinal analyses. *Psychol Aging.* (2006) 21:140–51. doi: 10.1037/0882-7974.21.1.140
50. Wang J, Mann F, Lloyd-Evans B, Ma R, Johnson S. Associations between loneliness and perceived social support and outcomes of mental health problems: a systematic review. *BMC Psychiatry.* (2018) 18:156. doi: 10.1186/s12888-018-1736-5
51. Pan K-Y, Kok AA, Eikelenboom M, Horsfall M, Jörg F, Luteijn RA, et al. The mental health impact of the COVID-19 pandemic on people with and without depressive, anxiety, or obsessive-compulsive disorders: a longitudinal study of three Dutch case-control cohorts. *Lancet Psychiatry.* (2021) 8:121–9. doi: 10.1016/S2215-0366(20)30491-0
52. Bu F, Steptoe A, Fancourt D. Who is lonely in lockdown? Cross-cohort analyses of predictors of loneliness before and during the COVID-19 pandemic. *Public Health.* (2020) 186:31–4. doi: 10.1016/j.puhe.2020.06.036
53. Lee CM, Cadigan JM, Rhew IC. Increases in loneliness among young adults during the COVID-19 pandemic and association with increases in mental health problems. *J Adolesc Health.* (2020) 67:714–7. doi: 10.1016/j.jadohealth.2020.08.009

54. Elm E. von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. (2007) 370:1453–7. doi: 10.1016/S0140-6736(07)61602-X
55. Lampert T, Kröll L, Müters S, Stolzenberg H. Messung des sozioökonomischen status in der studie zur gesundheit erwachsener in deutschland (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. (2013) 56:631–6. doi: 10.1007/s00103-012-1663-4
56. Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, et al. Validation and standardization of the generalized anxiety disorder screener (GAD-7) in the general population. *Med Care*. (2008) 46:266–74. doi: 10.1097/MLR.0b013e318160d093
57. Hinz A, Klein AM, Brähler E, Glaesmer H, Luck T, Riedel-Heller SG, et al. Psychometric evaluation of the generalized anxiety disorder screener GAD-7, based on a large German general population sample. *J Affect Disord*. (2017) 210:338–44. doi: 10.1016/j.jad.2016.12.012
58. Kroenke K, Strine TW, Spitzer RL, Williams JB, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord*. (2009) 114:163–73. doi: 10.1016/j.jad.2008.06.026
59. Wu Y, Levis B, Riehm KE, Saadat N, Levis AW, Azar M, et al. Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. *Psychol Med*. (2020) 50:1368–80.
60. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. (2006) 166:1092–7. doi: 10.1001/archinte.166.10.1092
61. Hays RD, DiMatteo MR. A short-form measure of loneliness. *J Pers Assess*. (1987) 51:69–81. doi: 10.1207/s15327752jpa5101\_6
62. Cohen S, Williamson G. *Perceived Stress in a Probability Sample of the United States*. Newbury Park, CA: Sage. (1988).
63. Schneider EE, Schönfelder S, Domke-Wolf M, Wessa M. Measuring stress in clinical and nonclinical subjects using a German adaptation of the perceived stress scale. *Int J Clin Health Psychol*. (2020) 20:173–81. doi: 10.1016/j.ijchp.2020.03.004
64. Diener E, Emmons RA, Larsen RJ, Griffin S. The satisfaction with life scale. *J Pers Assess*. (1985) 49:71–5. doi: 10.1207/s15327752jpa4901\_13
65. Janke S, Glöckner-Rist A. *Deutsche Version der Satisfaction with Life Scale*. (SWLS) Zusammenstellung sozialwissenschaftlicher Items und Skalen (ZIS) (2012) doi: 10.6102/zis147
66. Ehring T, Zetsche U, Weidacker K, Wahl K, Schönfeld S, Ehlers A. The Perseverative Thinking Questionnaire (PTQ): validation of a content-independent measure of repetitive negative thinking. *J Behav Ther Exp Psychiatry*. (2011) 42:225–32. doi: 10.1016/j.jbtep.2010.12.003
67. Kliem S, Mößle T, Rehbein F, Hellmann DF, Zenger M, Brähler E, et al. A brief form of the perceived social support questionnaire (F-SozU) was developed, validated, and standardized. *J Clin Epidemiol*. (2015) 68:551–62. doi: 10.1016/j.jclinepi.2014.11.003
68. Bruin A de, Picavet HS, Nossikov A. *Health interview surveys. Towards International Harmonization of Methods and Instruments (WHO Regional Publications European series, no 58)*. Copenhagen: World Health Organization, Regional Office for Europe. (1996).
69. Carver CS. You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med*. (1997) 4:92–100. doi: 10.1207/s15327558ijbm0401\_6
70. Knoll N, Rieckmann N, Schwarzer R. Coping as a mediator between personality and stress outcomes: a longitudinal study with cataract surgery patients. *Eur J Pers*. (2005) 19:229–47. doi: 10.1002/per.546
71. Jerusalem M, Schwarzer R. *SWE - Skala zur Allgemeinen Selbstwirksamkeitserwartung*. ZPID (Leibniz Institute for Psychology) – Open Test Archive. (2003).
72. Wiltink J, Kliem S, Michal M, Subic-Wrana C, Reiner I, Beutel ME, et al. Mini - social phobia inventory (mini-SPIN): psychometric properties and population based norms of the German version. *BMC Psychiatry*. (2017) 17:377. doi: 10.1186/s12888-017-1545-2
73. Wingenfeld K, Schäfer I, Terfehr K, Grabski H, Driessen M, Grabe H, et al. Reliable, valide und ökonomische erfassung früher traumatisierung: erste psychometrische charakterisierung der deutschen version des adverse Childhood Experiences Questionnaire (ACE). *Psychother Psychosom Med Psychol*. (2011) 61:e10–4. doi: 10.1055/s-0030-1263161
74. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. New York, NY: Routledge Academic (1988).
75. Ben-Ezra M, Hamama-Raz Y, Goodwin R, Leshem E, Levin Y. Association between mental health trajectories and somatic symptoms following a second lockdown in Israel: a longitudinal study. *BMJ Open*. (2021) 11:e050480. doi: 10.1136/bmjopen-2021-050480
76. Tukey JW. *Exploratory Data Analysis*. Boston, MA: Addison-Wesley. (1977).
77. Wang Z-H, Yang H-L, Yang Y-Q, Liu D, Li Z-H, Zhang X-R, et al. Prevalence of anxiety and depression symptom, and the demands for psychological knowledge and interventions in college students during COVID-19 epidemic: a large cross-sectional study. *J Affect Disord*. (2020) 275:188–93. doi: 10.1016/j.jad.2020.06.034
78. van Buuren S, Groothuis-Oudshoorn K. MICE: multivariate imputation by chained equations in R. *J. Stat. Soft.* (2011) 45:1–67. doi: 10.18637/jss.v045.i03
79. Rubin DB. *Multiple Imputation for Nonresponse in Surveys*. New York, NY: Wiley. (1987).
80. White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. *Stat Med*. (2011) 30:377–99. doi: 10.1002/sim.4067
81. R Core Team. *R: A language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing. (2021). Available online at: <https://www.R-project.org/>
82. Kocalevent R-D, Hinz A, Brähler E. Standardization of the depression screener patient health questionnaire (PHQ-9) in the general population. *Gen Hosp Psychiatry*. (2013) 35:551–5. doi: 10.1016/j.genhosppsych.2013.04.006
83. Li Y, Wang A, Wu Y, Han N, Huang H. Impact of the COVID-19 Pandemic on the mental health of college students: a systematic review and meta-analysis. *Front Psychol*. (2021) 12:669119. doi: 10.3389/fpsyg.2021.669119
84. Wang C, Wen W, Zhang H, Ni J, Jiang J, Cheng Y, et al. Anxiety, depression, and stress prevalence among college students during the COVID-19 pandemic: a systematic review and meta-analysis. *J Am Coll Health*. (2021):1–8. doi: 10.1080/07448481.2021.1960849
85. Brühlhart M, Klotzbücher V, Lalive R, Reich SK. Mental health concerns during the COVID-19 pandemic as revealed by helpline calls. *Nature*. (2021) 600:121–6. doi: 10.1038/s41586-021-04099-6

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Weber, Schulze, Bolzenkötter, Niemeyer and Renneberg. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.