Degree Monotone Paths and Graph Operations

Yair Caro Department of Mathematics University of Haifa-Oranim Israel Josef Lauri Department of Mathematics University of Malta Malta

Christina Zarb Department of Mathematics University of Malta Malta

Abstract

A path P in a graph G is said to be a degree monotone path if the sequence of degrees of the vertices of P in the order in which they appear on P is monotonic. The length of the longest degree monotone path in G is denoted by mp(G). This parameter was first studied in an earlier paper by the authors where bounds in terms of other parameters of G were obtained.

In this paper we concentrate on the study of how mp(G) changes under various operations on G. We first consider how mp(G) changes when an edge is deleted, added, contracted or subdivided. We similarly consider the effects of adding or deleting a vertex. We sometimes restrict our attention to particular classes of graphs.

Finally we study $mp(G \times H)$ in terms of mp(G) and mp(H) where \times is either the Cartesian product or the join of two graphs.

In all these cases we give bounds on the parameter mp of the modified graph in terms of the original graph or graphs and we show that all the bounds are sharp.

1 Introduction

Given a graph G, a degree monotone path is a path $v_1v_2...v_k$ such that $deg(v_1) \leq deg(v_2) \leq ... \leq deg(v_k)$ or $deg(v_1) \geq deg(v_2 \geq ... \geq deg(v_k)$. This notion, inspired by the well-known Erdős-Szekeres Theorem [6, 7], was introduced in [5] under the name of uphill and downhill path in relation to domination problems, also studied in [3, 4, 9]. In [5], the authors specifically suggested the study of the parameter mp(G), which denotes the length of the longest degree monotone path in G. This parameter was first studied by the authors in [2]. Links between this parameter and other classical parameters such as the chromatic number and clique number, using in particular the Gallai-Roy Theorem [11] were explored, and lower bounds and upper bounds for mp(G) were established in [2]. The close relation to Turan numbers [1] was also studied and explained in [2]. In this paper we consider another natural question related to the parameter mp(G), that of the effect of graph operations on this parameter. We consider both operations on a single graph G which produce a new graph G', as well as operations applied to two graphs to produce a new single graph.

In the first section, we consider operations involving edges, namely edge addition and deletion, subdivision and edge contraction while in the second section vertex addition and deletion is discussed. For each operation we obtain sharp bounds on mp(G'), and give constructions which achieve these bounds. In some cases, we consider the operation for a particular family of graphs which gives more interesting results.

We then consider the *Cartesian product* and the graph join for two graphs G and H, where again we give sharp bounds and constructions which achieve these bounds.

Any graph theory terms not defined here can be found in [11].

2 Edge Operations

2.1 Edge Addition and Deletion

We now look at the concept of adding/deleting an edge to or from a given graph G, and consider the effect of these operations on mp(G). We add an edge e by connecting two vertices in V(G) which are non-adjacent, and the resulting graph is denoted by G + e, while when we delete an edge e, the resulting graph is denoted by G - e.

Theorem 2.1. Given a graph G,

1.
$$\frac{mp(G)+1}{3} \le mp(G+e) \le 3mp(G).$$

2.
$$\frac{mp(G)}{3} \le mp(G-e) \le 3mp(G) - 1$$

In both cases, the bounds attained are sharp.

Proof. Let us first consider the right hand sides of both inequalities.

1. Let G be a graph with $mp(G) = k \ge 1$. If k = 1 then G is the empty graph with no edges, hence $mp(G + e) = 2 \le 3 = 3mp(G)$. So let us assume that $k \ge 2$.

Suppose that $mp(G+e) = t \ge 3k+1$. Consider $P = v_1, v_2, \ldots, v_t$, a longest degree monotone path in G + e in non-decreasing order. Observe that e must have at least one of its vertices in P, otherwise P was originally in G.

Case 1.1. Let us assume that $e = (v_i, z)$, where z is not on the path. If i = 1, then clearly v_1, \ldots, v_t is also a degree monotone path in G and hence $t \leq mp(G) = k < 3k$, a contradiction.

If i = t then v_1, \ldots, v_{t-1} is a degree monotone path in G and hence $t-1 \le k$ which implies $t \le k+1 < 3k$.

So let us assume 1 < i < t. Let us consider the paths v_1, \ldots, v_{i-1} and v_i, \ldots, v_t — both are non-decreasing degree monotone paths in G and hence together they have length at most 2k, implying that $t \leq 2k < 3k$.

Case 1.2. Now assume $e = (v_i, v_j)$, i < j. Hence, in G + e, all vertices in P have the same degree as they have in G except for v_i and v_j whose degree has increased by one. Consider the paths $P_1 = v_1, \ldots, v_i, P_2 = v_i, \ldots, v_{j-1}$ and $P_3 = v_j, \ldots, v_t$.

 P_3 is clearly a degree monotone path in G since $deg_G(v_j)+1 = deg_{G+e}(v_j) \leq deg_{G+e}(v_{j+1})$ (or j = t), hence $|V(P_3)| \leq k$. Similarly $|V(P_2)| \leq k$. So if $t \geq 3k + 1$, $|V(P_1)| \geq k + 2$ since v_i is both in P_1 and P_2 . But v_1, \ldots, v_{i-1} is a degree monotone path of length k + 1 in G, contradicting the fact the mp(G) = k. Hence $mp(G + e) \leq 3k$.

For sharpness of the bound, consider P_{3k} , to which we add a leaf to vertices v_2 up to v_{3k-2} except vertices v_{k+1} and v_{2k+1} , and we connect v_{3k-1} to a new vertex z to which we add two leaves. Thus the resulting graph G_1^+ has 6k - 2 vertices. Figure 1 shows the construction for k = 4. It is clear that $mp(G_1^+) = k$. Now if we add the edge $e = (v_{k+1}, v_{2k+1})$, these two vertices now have degree 3 also, and hence the path $v_1, v_2, \ldots, v_{3k-1}, z$ is a degree monotone path so $mp(G_1^+ + e) = 3k = 3mp(G_1^+)$.




Figure 1: G_1^+ when k = 4

2. Let G be a graph with $mp(G) = k \ge 1$. If k = 1 then G is the empty graph with no edges, hence we cannot delete edges. Therefore assume $k \ge 2$ and suppose that $mp(G-e) = t \ge 3k$. Let $P = v_1, \ldots, v_t$ be a degree monotone path of maximum length in non-increasing order. Observe that e must have at least one of its vertices in P, otherwise P was originally in G.

Case 2.1. Let us first assume that $e = (v_i, z)$, where z is not on the path. If i = 1, then clearly v_1, \ldots, v_t is also a degree monotone path in G and hence $t \leq mp(G) = k < 3k$, a contradiction.

If i = t then v_1, \ldots, v_{t-1} is a degree monotone path in G and hence $t-1 \le k$ which implies $t \le k+1 < 3k$.

So let us assume 1 < i < t. Let us consider the paths v_1, \ldots, v_{i-1} and v_i, \ldots, v_t — both are non-increasing degree monotone paths in G and hence together they have length at most 2k, implying that $t \leq 2k < 3k$.

Case 2.2. Now assume $e = (v_i, v_j)$, i < j. Hence, in G - e, all vertices in P have the same degree as they have in G except for v_i and v_j whose degree has decreased by one. Consider the paths $P_1 = v_1, \ldots, v_i, v_j, P_2 = v_i, \ldots, v_{j-1}$ and $P_3 = v_i, v_j, \ldots, v_t$.

 P_3 is clearly a degree monotone path in G since v_i and v_j have a larger (by 1) degree in G and $deg(v_i) \ge deg(v_j)$. Similarly $|V(P_2)| \le k$. So if $t \ge 3k$, $|V(P_1)| \ge k + 3$ since v_i and v_j are also on P_2 and P_3 . But v_1, \ldots, v_{i-1} is a degree monotone path of length k+1 in G, contradicting the fact the mp(G) = k. Hence $mp(G - e) \le 3k - 1$.

For sharpness, let us construct the graph G_1^- as follows: we start with P_{3k-1} and add three leaves to v_1 and a leaf to v_2 ; we then connect v_{k+1} to v_{2k+1} — we call this edge e. Figure 2 shows the construction for k = 4. Clearly $mp(G_1^-) = k$ while $mp(G_1^- - e) = 3k - 1$ with the path v_1, \ldots, v_{3k-1} and hence $mp(G_1^- - e) = 3mp(G_1^-) - 1$.



Figure 2: G_1^- when k = 4

Let us now turn to the lower bounds in both cases. These results are derived from those for the upperbounds.

1. Suppose for some graph G, $mp(G + e) < \frac{mp(G)+1}{3}$. Then if we take H = G + e so that G = H - e, we have $mp(H) < \frac{mp(H-e)+1}{3}$, which implies that mp(H - e) > 3mp(H) - 1, contradicting the righthand side of part 2.

For sharpness, let G_2^+ be the graph G_1^- without the edge e. Then $mp(G_2^+ + e) = k$ while $mp(G_2^+) = 3k - 1$ giving $mp(G_2^+ + e) = \frac{mp(G_2^+) + 1}{3}$.

2. Suppose that for some graph G, $mp(G-e) < \frac{mp(G)}{3}$. Then let H = G - e so that G = H + e giving $mp(H) < \frac{mp(H+e)}{3}$ which implies mp(H+e) > 3mp(H), contradicing the upper bound in part 1.

For sharpness of the bound, let G_2^- be the graph G_1^+ with the added edge $e = (v_{k+1}, v_{2k+1})$. Then $mp(G_2^-) = 3k$ and $mp(G_2^- - e) = \frac{mp(G_2^-)}{3}$.

2.2 Subdivision

Given a graph G and $e = (u, v) \in E(G)$, the subdivision of e is the addition of a new vertex w such that (u, w) is an edge and (w, v) is an edge but (u, v)is no longer an edge. Note that the degree of u and v remains the same, and the degree of w is 2. The graph obtained by subdividing e is denoted by G^* . Again, we look at the effect of this operation on the maximum length of a degree monotone path, mp(G).

Theorem 2.2. Let G be a graph and e = (u, v) an edge in G such that G^* is the graph obtained by subdividing e with the vertex w. Then

$$\left\lceil \frac{mp(G)+1}{2} \right\rceil \le mp(G^*) \le mp(G)+1,$$

and both bounds are sharp.

Proof. Let us first consider the upper bound. Let P be a degree monotone path of maximum length in G^* . We consider the following cases:

Case 1. If $w \notin P$, then P is a degree monotone path in G hence $mp(G^*) \leq mp(G)$.

Case 2. If $w \in P$, then either u or v or both are in P otherwise $P = \{w\}$ which is not maximal. We consider these cases separately:

- If both u and v are in P then $P \{w\} \cup (u, v)$ is a degree monotone path in G hence $mp(G^*) \le mp(G) + 1$.
- If only one of u or v, say u, is in P then $P \{w\}$ is a degree monotone path in G and again $mp(G^*) \leq mp(G) + 1$.

For sharpness of the bound, consider $G = P_n$ the path on *n* vertices — then mp(G) = n - 1, and subdividing any edge gives $mp(G^*) = n = mp(G) + 1$.

For the lower bound, let P be a degree monotone path of maximum length in G and let $P = v_1, v_2, \ldots, v_t, v_{t+1}, \ldots, v_k$, such that $u = v_t$ and $v = v_{t+1}$. Let us consider e = (u, v) and the subdividing vertex w.

Case 1. If e = (u, v) is not on the path P, then P is a degree monotone path in G^* hence $mp(G^*) \ge mp(G)$.

Case 2. If the edge e = (u, v) is in P, then as w subdivides e, we get $P^* = v_1, \ldots, v_t, w, v_{t+1}, \ldots, v_k$ a path in G^* , where $v_t = u$ and $v_{t+1} = v$.

We assume without loss of generality that P is a non-decreasing monotone path, and hence $deg(u) \leq deg(v)$. The vertex w has degree 2. Now if deg(u) > 2, the paths v_1, \ldots, v_t and the paths w, v_{t+1}, \ldots, v_k are degree monotone in G^* , while if $deg(u) \leq 2$, the path P^* is degree monotone in G^* . Hence in G^* there is a degree monotone path of length at least $\left\lceil \frac{k+1}{2} \right\rceil = \left\lceil \frac{mp(G)+1}{2} \right\rceil$.

For sharpness of the lower bound, let us take the path \dot{P}_n and add a leaf to the vertices v_2 up to v_{n-1} . Hence mp(G) = n - 1. If we subdivide the edge $(v_{\lceil \frac{n}{2} \rceil}, v_{\lceil \frac{n}{2} \rceil + 1})$, we get $mp(G^*) = \lceil \frac{n}{2} \rceil = \lceil \frac{mp(G) + 1}{2} \rceil$. \Box

2.3 Edge Contraction

In a graph G, contraction of an edge e = (u, v) is the replacement of u and v with a single vertex w adjacent (without multiple edges) to all vertices in $N(u)\setminus v \cup N(v)\setminus u$. The resulting graph $G \cdot e$ has one less vertex than G. In case a vertex $z \in V(G)$ is adjacent to both u and v, the degree of z in $G \cdot e$ decreases by one — otherwise it remains the same as in G. In view of this we first consider triangle-free graphs, in which case the degrees of the neighbours of u and v remain unchanged in $G \cdot e$, and deg(w) = deg(u) + deg(v) - 2.

Theorem 2.3. Let G be a triangle-free graph. Let e = uv be an edge of G which is contracted to form $G \cdot e$ with new vertex w. Then

$$\frac{mp(G)}{3} \le mp(G \cdot e) \le 2mp(G).$$

Proof. Let us first consider the upper bound. Clearly $mp(G) \ge 2$ as if mp(G) = 1, G has no edges and $G \cdot e$ is not defined. Let $P = v_1 v_2 \dots v_k$ be a degree monotone path in non-decreasing order of maximum length in $G \cdot e$. We know that deg(w) = deg(u) + deg(v) - 2. Let us look at all the different possibilities.

Case 1: Assume first that deg(u) = deg(v) = 1 and hence deg(w) = 0. If this was the only edge in G, then mp(G) = 2 while $mp(G \cdot e) = 1$, and the upper bound holds. If there are other edges in G, then $mp(G \cdot e) = mp(G)$, and again the upper bound holds.

Case 2: Now assume 1 = deg(u) < deg(v) — then deg(w) = deg(v) - 1 in $G \cdot e$. We consider the following cases:

- Clearly, if w is not a vertex in P, then P is degree monotone in G too, hence $mp(G \cdot e) \leq mp(G)$.
- If $w = v_1$ in P, then in $G, v_2 \dots v_k$ is a degree monotone path, and hence $mp(G \cdot e) 1 \leq mp(G)$ and hence $mp(G \cdot e) \leq mp(G) + 1 \leq 2mp(G)$.
- If $w = v_k$, then $v_1 \dots v_{k-1}v$ is degree monotone in G and hence $mp(G \cdot e) \le mp(G)$.
- If $w = v_j$ for $2 \le j \le k-1$, then $v_1 \dots v_{j-1}v$ and $v_{j+1} \dots v_k$ are degree monotone in G, and $\min\{\max\{j, k-j\}\} = \lceil \frac{k}{2} \rceil$.

If k is odd, then $mp(G) \ge \frac{k+1}{2}$ and hence $mp(G \cdot e) \le 2mp(G) - 1$.

If k is even, then $mp(G) \ge \frac{k}{2}$ and hence $mp(G \cdot e) \le 2mp(G)$.

Case 3: It remains to consider, without loss of generality, the case

$$2 \le deg(u) \le deg(v) \le deg(w) = deg(u) + deg(v) - 2.$$

Again we consider each possibility.

- If in G, w is not a vertex in P, then P is degree monotone in G too, hence $mp(G \cdot e) \leq mp(G)$.
- If $w = v_1$, then in G, either u or v is adjacent to v_2 since $deg(u) \leq deg(v) \leq deg(w)$ then either $uv_2 \dots v_k$ or $vv_2 \dots v_k$ is a degree monotone path in G, and hence $mp(G \cdot e) + 1 \leq mp(G)$ giving $mp(G \cdot e) \leq mp(G) 1 \leq 2mp(G)$.
- If $w = v_k$, then $v_1 v_2 \dots v_{k-1}$ is still a degree monotone path in G hence $mp(G \cdot e) 1 \leq mp(G)$ that is $mp(G \cdot e) \leq mp(G) + 1 \leq 2mp(G)$.
- If $w = v_j$, $2 \le j \le k 1$, then let us consider the following cases:
 - 1. If in G, u is adjacent to both v_{j-1} and v_{j+1} , and hence v is not adjacent to either of these vertices since G is triangle-free, then $v_1 \ldots v_{j-1}$ and $u, v_{j+1} \ldots v_k$ are degree monotone paths of length j-1 and k-j+1 respectively in G. Again, min $\{\max\{j-1, k-j+1\}\} = \lceil \frac{k}{2} \rceil$. If k is odd, then $mp(G) \ge \frac{k+1}{2}$ and hence $mp(G \cdot e) \le 2mp(G) - 1$. If k is even, then $mp(G) \ge \frac{k}{2}$ and hence $mp(G \cdot e) \le 2mp(G)$.




Figure 3: G_1 when k = 4

2. If in G, u is adjacent to v_{j-1} and v is adjacent to v_{j+1} , then if $deg(v_{j-1}) \leq deg(u)$ then $v_1v_2, \ldots, v_{j-1}, u, v, v_{j+1}, \ldots v_k$ is degree monotone in G and hence $mp(G \cdot e) + 1 \leq mp(G)$ implying that $mp(G \cdot e) \leq mp(G) - 1 \leq 2mp(G)$.

If, on the other hand, $deg(v_{j-1}) > deg(u)$, then $deg(u) \le deg(v) \le deg(w) \le deg(v_{j+1})$ and hence $v_1v_2 \ldots v_{j-1}$ and $u, v, v_{j+1} \ldots v_k$ are degree monotone paths in G of length j-1 and k-j+2 respectively. In this case we consider min $\{\max\{j-1, k-j+2\}\}$. If k is even this is equal to $\frac{k+2}{2}$, and hence $mp(G \cdot e) \le 2mp(G) - 2$.

If k is even this is equal to $\frac{k+2}{2}$, and hence $mp(G \cdot e) \leq 2mp(G) - 2$ If k is odd, we get $\frac{k+1}{2}$, and hence $mp(G \cdot e) \leq 2mp(G) - 1$.

- 3. If in G, v is adjacent to v_{j-1} and u is adjacent to v_{j+1} , then $v_1v_2\ldots v_{j-1}$ and $uv_{j+1}\ldots v_k$ are degree monotone paths of length j-1 and k-j+1respectively in G, and again min $\{\max\{j-1, k-j+1\}\} = \lceil \frac{k}{2} \rceil$. If k is odd, then $mp(G) \ge \frac{k+1}{2}$ and hence $mp(G \cdot e) \le 2mp(G) - 1$. If k is even, then $mp(G) \ge \frac{k}{2}$ and hence $mp(G \cdot e) \le 2mp(G)$.
- 4. If in G, v is adjacent to both v_{j-1} and v_{j+1} , and u is adjacent to neither since G is triangle-free, then $v_1 \ldots v_{j-1}$ and $u, v, v_{j+1} \ldots v_k$ are degree monotone paths in G of length j-1 and k-j+2 respectively. In this case we consider min $\{\max\{j-1, k-j+2\}\}$.

If k is even this is equal to $\frac{k+2}{2}$, and hence $mp(G \cdot e) \leq 2mp(G) - 2$. If k is odd, we get $\frac{k+1}{2}$, and hence $mp(G \cdot e) \leq 2mp(G) - 1$.

This bound is attained by the graph G_1 constructed as follows: consider the path on 2k + 1 vertices — we add a leaf to vertices v_2 up to v_{2k} , and to the vertices v_k and v_{2k} we add a second leaf. Then mp(G) = k. If we contract one of the edges joining v_k and a leaf, then vertex w has degree 3 and hence $v_1v_2 \ldots v_{k-1}wv_{k+2} \ldots v_{2k}$ is degree monotone in $G \cdot e$ and has length 2k, giving $mp(G \cdot e) = 2mp(G)$. Figure 3 shows the construction for k = 4.

We now consider the lower bound. Let $P = v_1 \dots v_k$ be a degree monotone path in non-decreasing order of maximum length in G. Let e = uv be the edge contracted to vertex w in $G \cdot e$, and without loss of generality, we assume $deg(u) \leq deg(v)$. Clearly, if the vertices u and v are not on P, then P is still degree monotone in $G \cdot e$ and hence $mp(G \cdot e) \geq mp(G)$. So let us assume that u and v are on P, and hence $k = mp(G) \geq 2$. If k = 2, that is P = uv, then $mp(G \cdot e) \geq 1 \geq \frac{mp(G)}{2}$. So let us assume that $k \geq 3$ and hence $deg(v) \geq 2$. So let us assume that u and v are on P and $mp(G) \geq 3$.

Case 1. We first consider the case in which either u or v, but not both, are on P. Without loss of generality, Let $u = v_i$ be on P. Then in $G \cdot e$, w is on P, and $v_1 \ldots v_{j-1}$ and $v_{j+1} \ldots v_k$ are degree montone in $G \cdot e$, of length j and k - j respectively. Now min $\{\max\{j, k - j\}\} = \lceil \frac{k}{2} \rceil$.

If k is even then $mp(G \cdot e) \ge \frac{mp(G)}{2} \ge \frac{mp(G)}{3}$. If k is odd then $mp(G \cdot e) \ge \frac{mp(G)+1}{2} \ge \frac{mp(G)}{3}$.

We now consider the case in which u and v are in P but e = uv is Case 2. not in P. Let $u = v_i$ and $v = v_j$ in P; in all cases $j - i \ge 3$, otherwise we have a copy of K_3 in G — we consider the following cases:

- If i = 1 and j = k, then $v_2 \dots v_{k-1} w$ is degree monotone in $G \cdot e$, hence $mp(G \cdot e) \ge mp(G) - 1 \ge \frac{mp(G)}{3}$ since $mp(G) \ge 2$.
- If i = 1 and $4 \leq j < k$, then $v_2 \dots v_{j-1} w$ and $v_{j+1} \dots v_k$ are degree monotone in $G \cdot e$. Hence we need min $\{\max\{j-1, k-j\}\} = \lfloor \frac{k}{2} \rfloor$. Then $mp(G \cdot e) \ge \lfloor \frac{mp(G)}{2} \rfloor \ge \frac{mp(G)}{3}.$
- If i > 1 and j = k, then $v_1 \dots v_{i-1}$ and $v_{i+1} \dots v_{k-1} w$ are degree monotone in $G \cdot e$. Hence we need $\min\{\max\{i-1,k-i\}\} = \lfloor \frac{k}{2} \rfloor$. Then $mp(G \cdot e) \geq k$ $\left|\frac{mp(G)}{2}\right| \ge \frac{mp(G)}{3}.$
- If 1 < i < j < k, then $v_1 \dots v_{i-1} w$, $v_{i+1} \dots v_{j-1} w$ and $v_{j+1} \dots v_k$ are degree monotone in $G \cdot e$, of lengths i, j-1 and k-j respectively. Let k = 3t + r where $0 \le r \le 2$.

If k-j > t, then $k-j \ge t+1 \ge \frac{k}{3}$, and hence $mp(G \cdot e) \ge \frac{mp(G)}{3}$.

So let us assume that $k - j \leq t$ and hence $j \geq k - t = 2t + r$. Now $\max\{i, j-1\} \ge \frac{2t+r}{2} = t + \frac{r}{2}$. If r = 0 then $k \ge 3t$ hence $\max\{i, j-1\} = 1$ $t = \frac{k}{3}$. If r = 1, then k = 3t + 1, then $\max\{i, j - 1\} \ge t + 1 = \frac{k+2}{3} > \frac{k}{3}$ since $\max\{i, j-1\}$ must be an integer. Finally, if r = 2, then $\max\{i, j-1\} \ge 1$ $t+1 = \frac{k+1}{3} > \frac{k}{3}$.

Hence, in each case, $mp(G \cdot e) \geq \frac{mp(G)}{3}$.

Finally, we consider the case in which e = uv is in P. We may Case 3. assume that $mp(G) \ge 3$ since if mp(G) = 2 then clearly $mp(G \cdot e) \ge 1 \ge \frac{mp(G)}{3}$. This results in the following cases:

- If $u = v_1$ and $v = v_2$, then in $G \cdot e, v_3 \dots v_k$ is degree monotone, and hence $mp(G \cdot e) \ge mp(G) 2 \ge \frac{mp(G)}{3}$ for $mp(G) \ge 3$.
- If $u = v_{k-1}$ and $v = v_k$, hence $2 \leq deg(u) \leq deg(v) \leq deg(w)$, then in $G \cdot e, v_1 \dots v_{k-2}w$ is degree monotone since $deg(u) \leq deg(v) \leq deg(w) =$ deg(u) + deg(v) - 2. Hence $mp(G \cdot e) \ge mp(G) - 1 \ge \frac{mp(G)}{3}$ for $mp(G) \ge 3$.
- If $u = v_j$ and $v = v_{j+1}$ for $2 \le j \le k-2$ (so $k \ge 4$ otherwise we have one of the previous two cases), then in $G \cdot e, v_1 \dots v_{j-1} w$ and $v_{j+2} \dots v_k$ are degree monone paths of lengths j and k - j - 1 respectively— hence we must consider $\min\{\max\{j, k - j - 1\}\} = \lceil \frac{k}{2} \rceil$. Hence $mp(G \cdot e) \ge \lceil \frac{k}{2} \rceil \ge \frac{mp(G)}{3}$ as required.

The bound is attained by the following construction to give graph G_3 : let us take the path on 3k + 3 vertices — we add a leaf to the vertices v_2 up to v_{3k+2} except for v_{k+1} and v_{2k+2} , and we connect these two vertices by an edge e. We also attach three leaves to each of the leaves attached to v_{k+2} and v_{2k+1} . Finally we add 3 leaves to vertex v_{3k+3} . Thus $deg(v_1) = 1$, $deg(v_2) = deg(v_3) \dots =$ $deg(v_{3k+2}) = 3$ and $deg(v_{3k+3}) = 4$, hence mp(G) = 3k+3. Now let us contract the edge e. Then $deg(v_1) = 1$, $deg(v_2) = deg(v_3) \dots = deg(v_k) = deg(v_{k+2}) =$ $\dots = deg(v_{2k+1}) = deg(v_{2k+3}) = \dots = deg(v_{3k+2}) = 3$ and $deg(v_{3k+3}) = 4$. Vertex w has degree 4. Hence the possible degree monotone paths in $G \cdot e$ are $v_1 \dots v_k w, v_{k+2} \dots v_{2k+1} w$ and $v_{2k+3} \dots v_{3k+3}$ — all these are of length k + 1and therefore $mp(G_3 \cdot e) = \frac{mp(G_3)}{3}$. Figure 4 shows the construction for k = 3.




Figure 4: G_3 when k = 4

The following proposition shows that if we consider contraction in graphs which are not triangle-free, the situation is very different.

Theorem 2.4. There exist arbitrarily large K_4 -free graphs G on n vertices such that mp(G) = 4 while $mp(G.e) = |V(G \cdot e)| = n - 1$.

Proof. Consider the following construction: G is the graph constructed by taking a path $P = v_1 \dots v_{4k}$ on 4k vertices where $k \ge 2$, and another edge e = uv. We connect the vertices $v_{2i} \in P$, $1 \le i \le 2k$ to both u and v — we then connect $v_{2i-1} \in P$ for $1 \le i \le k$ to u, and $v_{2i-1} \in P$ for $k+1 \le i \le 2k$ to v. Finally we connect v_1 to v_{4k} . Figure 5 gives the construction for k = 2. G is not triangle-free, but it is K_4 -free. One can see that in P, the vertices with even index have degree 4, while the vertices with odd index have degree 3, and deg(u) = deg(v) = 3k. Hence it is clear that mp(G) = 4 for any value of $k \ge 2$. Now if we contract e = uv to a vertex w, all vertices in P have degree 3, while $deg(w) = 4k \ge 3$, and therefore $mp(G \cdot e) = 4k + 1$.

3 Vertex Addition and Deletion

Given a graph G, we can add a vertex v and connect it to at least one vertex in G to give the graph G + v. On the other hand, given a graph G and a vertex $v \in V(G)$, the graph G - v is obtained by deleting the vertex v and all its




Figure 5: G when k = 3

incident edges. The effect on the maximum length of a degree monotone path can be seen through the following observations.

Proposition 3.1. Given a graph G and $v \in V(G)$,

1.
$$2 \le mp(G+v) \le |V(G)| + 1$$

2. $1 \le mp(G - v) \le |V(G)| - 1$

Proof.

1. Consider the complete bipartite graph $G = K_{n,n+1}$ so that mp(G) = 2. We add a vertex v_1 and connect it to all the vertices in the larger part, to give $H = G + v_1 = K_{n+1,n+1}$ — now $mp(G + v_1) = mp(H) = |V(G)| + 1$. Now if we add another vertex v_2 and connect it to all the vertices in one part of the partition, we get $H + v_2 = F = K_{n+1,n+2}$ and again $mp(H + v_2) = mp(F) = 2$.

2. For the upperbound, consider the graph $G = K_n$ — then mp(G) = n, and $G - v = K_{n-1}$ hence mp(G - v) = n - 1 = |V(G)| - 1.

For the lower bound consider $G = K_{1,m}$, $m \ge 1$. Then deleting the vertex of degree m gives $mp(K_{1,m} - v) = mp(G - v) = 1$.

In view of this general result, we consider vertex addition and deletion for the family of trees. The following example shows that if one adds non-leaf vertices, the effect on mp(G) can be quite drastic. Consider the tree T constructed by taking a path on 2k + 1 vertices $(v_1v_2 \dots v_{2k+1}), k \ge 1$, and adding a leaf to the vertices $v_{2i}, 1 \le i \le k$. Clearly this tree has mp(T) = 2. Let us add a vertex v and connect it to the vertices $v_{2i+1}, 0 \le i \le k$. So now all the vertices on the path have degree 3 except for v_1 and v_{2k+1} , and vertex v has degree $k + 1 \ge 2$. Hence there is now a degree monotone path of length 2k + 1, that is mp(T + v) = 2k + 1. Hence, in the sequel, we only consider the addition of leaves, so that the resulting graph is another tree.

Theorem 3.2. Let T be a tree and $v \in V(T)$,

1. If we add a vertex v such that T + v is also a tree, then

$$\frac{mp(T)}{2} \le mp(T+v) \le 2mp(T).$$

2. If $v \in V(T)$ such that T - v is a tree, then

$$\frac{mp(T)}{2} \le mp(T-v) \le 2mp(T).$$

In both cases, the bounds attained are sharp.

Proof. Let us first consider the upper bound for each case.

1. If T + v is a tree then clearly v is a leaf otherwise it would form a cycle. Let $P = v_1, v_2, \ldots, v_k$ be a degree monotone path of maximum length in T + v in non-decreasing order. Observe that $k \ge 2$ as T + v contains at least one edge. So we consider all possible cases.

- If v is not adjacent to any vertex in P then P is a degree monotone path in T hence $mp(T+v) \le mp(T)$.
- If v is adjacent to v_1 , then v, v_1, \ldots, v_k is a degree monotone path in T + v contradicting the maximality of P.
- If $v = v_1$ then v_2, \ldots, v_k is a degree monotone path in T of length k-1 so $k-1 \le mp(T)$ and hence $2mp(t) \ge 2k-2 \ge k = mp(T+v)$ since $k \ge 2$.
- If v is adjacent to v_k , then v_1, \ldots, v_{k-1} is a degree monotne path in T, hence again $2mp(t) \ge 2k - 2 \ge k = mp(T+v)$ since $k \ge 2$.
- If $v = v_k$, this implies that deg(v) = 1 and that k = 2, and since T is not $K_1, mp(T) \ge 2$ hence $2mp(T) \ge 4 > 2 = k = mp(T+v)$. (If $T = K_1$ then trivially $T + v = K_2$ so mp(T+v) = 2 = 2mp(T).)
- Let v be adjacent to some v_j for $2 \le j \le k 1$. Note that v can only be v_1 or v_k in P since it is a leaf. Consider v_1, \ldots, v_{j-1} which is a degree monotone path in T hence $j - 1 \le mp(T)$. Similarly v_j, \ldots, v_k is also a degree monotne path in T with length at most mp(T) hence by adding we get $mp(T + v) \le 2mp(T)$ as required.

For sharpness, consider the tree T_1^+ constructed as follows: take the path on 2k + 1 vertices v_1, \ldots, v_{2k+1} , where $k \ge 2$ — we add a leaf to all the vertices except v_1, v_{k+1} and v_{2k+1} . Then $mp(T_1^+) = k$. If we add a vertex v and connect it to vertex v_{k+1} , then $mp(T_1^+ + v) = 2k = 2mp(T_1^+)$. Figure 6 shows the construction for k = 4.




Figure 6: T_1^+ when k = 4

2. Let $P = v_1, v_2, \ldots, v_k$ be a degree monotone path of maximum length in T - v in non-decreasing order. Not that if k = 1 then T - v has no edges which imples that $T = K_{1,m}$ for some $m \ge 1$, and so mp(T) = 2 = 2mp(T - v).

So let us assume that $k \ge 2$. If in T v is not adjacent to any vertex of P, then P is also a degree monotone path in T, hence $mp(T - v) = k \le mp(T) < 2mp(T)$.

So we consider the case in which v is adjacent to exactly one vertex in P. We consider the different scenarios.

- If v is adjacent to v_1 in T, then v_2, \ldots, v_k is a degree monotone path of length k-1 in T hence $mp(T-v) = k \leq 2(k-1) \leq 2mp(T)$ since $k \geq 2$.
- If v is adjacent to v_k in T then v_1, \ldots, v_k is also a degree monotone path in T, hence again $mp(T v) = k \leq mp(T) < 2mp(T)$.
- If v is adjacent to some v_j for $2 \le j \le k-1$, then consider in T the path v_1, \ldots, v_j this is degree monotone hence $j \le mp(T)$. Now v_{j+1}, \ldots, v_k is also a degree monotone path in T and has length at most mp(T). Adding, we get $mp(T-v) \le 2mp(T)$ as required.

For sharpness consider the graph T_1^- constructed as follows — we take a path on 2k+1 vertices for $k \ge 2$ and we add a leaf to vertex v_k and to vertex v_{2k} . Then $mp(T_1^-) = k$, while if v is the leaf attached to v_k , $mp(T_1^- - v) = 2k = 2mp(T_1^-)$. Figure 7 shows the construction for k = 4.




Figure 7: T_1^- when k = 4

We now consider the lower bounds.

1. Assume $mp(T+v) < \frac{mp(T)}{2}$. Let T' = T + v hence T = T' - v — then $mp(T') < \frac{mp(T'-v)}{2}$ implying that mp(T'-v) > 2mp(T'), contradicting the upper bound in part 2.

For sharpness, let T_2^+ be the path on 2k + 1 vertices, $k \ge 2$, with a leaf added to vertex v_{2k} . Then $mp(T_2^+) = 2k$. We add a vertex v and connect it to the vertex v_k — then $mp(T_2^+ + v) = k = \frac{mp(T_2^+)}{2}$.

2. Assume $mp(T-v) < \frac{mp(T)}{2}$. Let T' = T - v hence T = T' + v — then $mp(T') < \frac{mp(T'+v)}{2}$ implying mp(T'+v) > 2mp(T'), contradicting the upperbound in part 1.

For sharpness, we construct the graph T_2^- as follows: take the path on 2k+1 vertices for $k \ge 2$ and add a leaf to every vertex except the first and the last so that $mp(T_2^-) = 2k$. Consider the leaf v connected to v_{k+1} — if we delete this vertex, $mp(T_2^- - v) = k = \frac{mp(T_2^-)}{2}$.

4 Operations involving two graphs

4.1 Cartesian Product

The Cartesian product $G \square H$ of graphs G and H is a graph such that the vertex set of $G \square H$ is the Cartesian product $V(G) \times V(H)$ and any two vertices (u, u') and (v, v') are adjacent in $G \square H$ if and only if either

- u = v and u' is adjacent with v' in H, or
- u' = v' and u is adjacent with v in G.

Theorem 4.1. Let G and H be two connected graphs. Then

$$mp(G) + mp(H) - 1 \le mp(G \square H) \le mp(G)mp(H)$$

and both bounds are sharp.

Proof. Let us first consider the lower bound. Let $v_1 \ldots v_t$ be a longest degree monotone path in G, and let $u_1 \ldots u_s$ be a longest degree monotone path in H, both in non-decreasing order.

Consider the path in $G \square H$ with vertex coordinates

$$(v_1, u_1)(v_1, u_2) \dots (v_1, u_s)(v_2, u_s)(v_3, u_s) \dots (v_t, u_s).$$

This is clearly a degree monotone path in $G \square H$ with t + s - 1 vertices, and hence $mp(G \square H) \ge mp(G) + mp(H) - 1$.

Now let us consider the degree monotone path in in $G \square H$ of maximum length $r = mp(G \square H)$ with vertices z_i for $1 \le i \le r$. Let us label $z_i = (v_{a_i}, u_{b_i})$. So consider the vertices from z_1 to $z_i = (v_{a_i}, u_{b_i})$: for vertex z_{i+1} , either the vcoordinate or the u coordinate will change but not both.

If the v coordinate changes, then $z_{i+1} = (v_{a_{i+1}}, u_{b_{i+1}})$ where $a_{i+1} > a_i$ and $b_{i+1} = b_i$, hence it follows that $deg(v_{a_{i+1}}) \ge deg(v_{a_i})$.

If the *u* coordinate changes, then $z_{i+1} = (v_{a_{i+1}}, u_{b_{i+1}})$ where $a_{i+1} = a_i$ and $b_{i+1} > b_i$, hence it follows that $deg(u_{b_{i+1}}) \ge deg(u_{b_i})$.

Now let us consider those vertices in which the index $a_{i+1} > a_i$, which implies that the corresponding vertices in G are distinct — it is clear that these vertices form a degree monotone paths in G. Hence if the number of such vertices is $t, t \leq mp(G)$.

Similarly, if we consider the vertices in which the index $b_{i+1} > b_i$, the corresponding vertices in H are distinct and form a degree monotone path in H — if the number of such vertices is s, then $s \leq mp(H)$.

Now since for each move from z_i to z_{i+1} , only one coordinate changes, we have at most st coordinates, hence $mp(G \square H) \leq mp(G)mp(H)$.

Now we look at construction which achieve these bounds. Firstly, let $G = H = K_{1,m}$, where $m \ge 2$, and hence mp(G) = mp(H) = 2. In $G \square H$ there is one vertex of degree 2m, 2m vertices of degree m + 1 which are independent, and m^2 vertices of degree 2 which are independent. Hence the longest degree monotone path has three vertices so $mp(G \square H) = 2 = mp(G) + mp(H) - 1$.

Now for the upper bound, let G be a connected regular graph on t vertices and let H be a graph such that mp(H) = s. Let the vertices of G be v_1, \ldots, v_t , and let the vertices u_1, \ldots, u_s in H be vertices on a longest degree monotone path in H.

Now in $G \square H$, consider the path

 $(v_1, u_1), (v_2, u_1), \dots, (v_t, u_1), (v_t, u_2), (v_{t-1}, u_2), \dots, (v_1, u_2), (v_1, u_3), \dots, (v_t, u_3), \dots$

and we carry on in this fashion until we have used all the *st* vertices. This path is degree monotone and hence $mp(G \Box H) = mp(G)mp(H)$.

4.2 Graph join

The graph join G + H of two graphs G and H with disjoint vertex sets, V(G) and V(H) and disjoint edge sets E(G) and E(H) is the graph such that

- $V(G+H) = V(G) \cup V(H)$
- $E(G+H) = E(G) \cup E(H) \cup \{xy : x \in V(G), y \in V(H)\}$

We now consider degree monotone paths in G + H.

Theorem 4.2. Given two graphs G and H,

$$mp(G) + mp(H) \le mp(G + H) \le |V(G)| + |V(H)|,$$

and both bounds are sharp.

Proof. The upper bound is trivial since for any graph $mp(G) \leq |V(G)|$. So let us consider the lower bound. Let $P = v_1, v_2, \ldots, v_t$ be a degree monotone path of maximum length in G and let $P^* = u_1, u_2, \ldots, u_s$ be a degree monotone path of maximum length in H. Let us rearrange $\{v_1, \ldots, v_t, u_1, \ldots, u_s\}$ in nondecreasing order according to their degrees in G + H, noting that $deg_{G+H}(v) =$ $deg_G(v) + |V(H)|$ while $deg_{G+H}(u) = deg_H(u) + |V(G)|$. Then it is clear that these vertices form a degree monotone path in G + H of length s + t, hence $mp(G + H) \leq mp(G) + mp(H)$.

Let us consider the upper bound. Let G and H be two graphs such that |V(G)| = |V(H)| and G and H have the same degree sequence. We write the vertices of G + H in non-decreasing order of their degrees such that each vertex of the G part is followed by the corresponding vertex in the H part. So for the degree monotone path in G + H, we start with the vertex of smallest degree in G + H and alternately take vertices of this same degree from H and G until all vertices of this degree are included in the path: we then move to the second smallest degree in G and carry out the same procedure for every different degree in the sequence. There is an even number of vertices in G + H of each degree since G and H have the same number of vertices and the same degree sequence, so this alternating path can be continued until all vertices in G + H have been included, which implies that mp(G + H) = |V(G)| + |V(H)| in this case, achieveing the upper bound.

For the lower bound, consider $G = K_{1,m}$ and $H = K_k$. So mp(G) = 2 and mp(H) = k. Then in G + H there are k + 1 vertices of degree k + m, namely the vertices of H and the vertex of degree m in G, and the m vertices of degree k + 1 that are independent. So if we can start the path with a vertex of degree k + 1, and then we must move to a vertex of degree k + m, and all the vertices of this degree can be included in the path. hence mp(G + H) = 1 + k + 1 = 2 + k = mp(G) + mp(H).

5 Conclusion

We have given sharp bounds for mp(G') in terms of mp(G) where G' is obtained from G by the most basic operations involving a single vertex or edge. We have shown that the effect of edge contraction on mp(G) in the case of K_3 -free graphs is bounded (above and below) by a multiplicative factor, while there exist K_4 free graphs for which mp(G) = 4 while $mp(G \cdot e) = |V(G)| - 1$. This leads to the following questions:

- 1. Is there a K_4 -free graph G such that mp(G) = 3 and $mp(G \cdot e) \ge |V(G)| 1$?
- 2. Is there a characterization of K_4 -free graphs for which $mp(G \cdot e)$ is bounded above and below by a multiplicative factor?

We have also obtained sharp bounds for $mp(G \times H)$ in terms of mp(G)and mp(H) where \times is either the Cartesian product of the join of two graphs. Repeating this for other products, such as the direct product, might be interesting.

Finally, the notion of edge addition gives rise to the question of what the minimum number of edges of a graph G on n vertices can be if adding any edge increases mp(G). This leads to a problem analogous to the saturation number of a graph [8, 10], and we shall be considering this in a forthcoming paper.

References

- [1] B Bollobás. *Extremal graph theory*. Courier Dover Publications, 2004.
- [2] Y. Caro, J. Lauri, and C. Zarb. Degree Monotone Paths. ArXiv e-prints, May 2014. submitted.
- [3] J. Deering. Uphill & Downhill Domination in Graphs and Related Graph Parameters. PhD thesis, ETSU, 2013.
- [4] J. Deering, T.W. Haynes, S.T. Hedetniemi, and W. Jamieson. Downhill and uphill domination in graphs. 2013. submitted.
- [5] J. Deering, T.W. Haynes, S.T. Hedetniemi, and W. Jamieson. A polynomial time algorithm for downhill and uphill domination. 2013. submitted.

- [6] M. Eliáš and J. Matoušek. Higher-order erdős-szekeres theorems. Advances in Mathematics, 244:1–15, 2013.
- [7] P. Erdös and G. Szekeres. A combinatorial problem in geometry. *Compositio Mathematica*, 2:463–470, 1935.
- [8] J.R. Faudree, R.J. Faudree, and J.R. Schmitt. A survey of minimum saturated graphs. *The Electronic Journal of Combinatorics*, 1000:DS19–Jul, 2011.
- [9] T. W. Haynes, S. T. Hedetniemi, J. D. Jamieson, and W. B. Jamieson. Downhill domination in graphs. *Discussiones Mathematicae Graph Theory*. accepted.
- [10] L. Kászonyi and Z. Tuza. Saturated graphs with minimal number of edges. Journal of graph theory, 10(2):203–210, 1986.
- [11] D. B. West. Introduction to Graph Theory. Prentice Hall, 2 edition, September 2000.