



## ISTITUTO NAZIONALE DI RICERCA METROLOGICA Repository Istituzionale

Physiological variability in brain electric conductivity: correcting the effect of the age for the detection of pathological alterations

This is the author's accepted version of the contribution published as:

*Original*

Physiological variability in brain electric conductivity: correcting the effect of the age for the detection of pathological alterations / Marmin, Sébastien; Arduino, Alessandro; Cencini, Matteo; Lancione, Marta; Biagi, Laura; Tosetti, Michela; Zilberti, Luca. - (2023), pp. 72-74. (Intervento presentato al convegno Joint workshop of ENBIS and MATHMET Mathematical and Statistical Methods for Metrology tenutosi a Torino nel 30-31 May 2023).

*Availability:*

This version is available at: 11696/76819 since: 2023-06-01T08:11:42Z

*Publisher:*

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# Physiological variability in brain electric conductivity: correcting the effect of the age for the detection of pathological alterations

Sébastien Marmin and Alessandro Arduino and Matteo Cencini and Marta Lancione and Laura Biagi and Michela Tosetti and Luca Zilberti

**Key words:** Electric properties tomography, Quantitative imaging, Magnetic resonance imaging, white matter disorders

## Overview

The aim of this work is to analyze the physiological variability of electric conductivity in the brain measured through electric properties tomography (EPT) using Magnetic Resonance Imaging (MRI). Electric conductivity maps of 27 healthy volunteers and 45 patients with pathologies of the white matter (WM) were estimated based on water content [Michel et al. (2017)] measured using MR Fingerprinting [Cencini et al. (2022)] (Fig. 1A).

For each subject, tissue class segmentation was performed to obtain a white matter mask (Fig. 1B) and the median conductivity within this region was measured. A non-linear mixed effect model was used to identify sources of variability. The inter-subject standard deviation of the median WM conductivity was estimated to 11 mS/m. The results in Fig 1C showed strong dependence of WM conductivity with age.

We reported higher WM conductivity in patients with respect to control subjects. To distinguish pathological changes from physiological variability, we established a threshold value of 22.5 mS/m more than the age-dependent average of observed

---

Sébastien Marmin

Laboratoire national de métrologie et d'essais, 29 avenue Hennequin, 78190 Trappes, France,  
e-mail: [sebastien.marmin@lne.fr](mailto:sebastien.marmin@lne.fr)

Alessandro Arduino and Luca Zilberti

Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce, 91 10135 Torino, Italy e-mail:  
[a.arduino@inrim.it](mailto:a.arduino@inrim.it), [l.zilberti@inrim.it](mailto:l.zilberti@inrim.it)


Matteo Cencini

National Institute for Nuclear Physics (INFN), Largo Bruno Pontecorvo 3, 56127 Pisa, Italy  
e-mail: [matteo.cencini@pi.infn.it](mailto:matteo.cencini@pi.infn.it)

Marta Lancione, Laura Biagi and Michela Tosetti

Fondazione IRCCS Stella Maris, Viale del Tirreno, 331, 56128 Pisa, Italy e-mail:  
[marta.lancione@fsm.unipi.it](mailto:marta.lancione@fsm.unipi.it), [laura.biagi@fsm.unipi.it](mailto:laura.biagi@fsm.unipi.it), [michela.tosetti@fsm.unipi.it](mailto:michela.tosetti@fsm.unipi.it)

in the control group (Fig. 1D). These findings are important for the development of biomarkers and personalized medicine using EPT, and demonstrate the potential of metrology in this field.



**Fig. 1** A. Conductivity map on a transverse section for one subject. B. Segmentation of the white matter. C. Median conductivity in WM for each subject with respect to age (black line); The blue area is the prediction envelope at level 95%. D. For each subject, difference with the age-dependent average calculated from the control group (threshold in purple).

**Acknowledgements** This work was developed within the QUIERO project. This project has received funding from the EMPIR programme co-financed by the Participating States and from the European Union’s Horizon 2020 research and innovation programme. This study was also partially supported by grant RC and the 5x1000voluntary contributions to IRCCS Fondazione Stella Maris, funded by the Italian Ministry of Health.

## References

Cencini, Matteo et al. (2022). “Fast high-resolution Electric Properties Mapping using three-dimensional MR Fingerprinting based water fraction estimation (MRF-EPT)”. In: *Proceedings of the Joint Annual Meeting ISMRM-ESMRMB, London*.

Michel, Eric, Daniel Hernandez, and Soo Yeol Lee (2017). “Electrical conductivity and permittivity maps of brain tissues derived from water content based on T1-weighted acquisition”. In: *Magnetic Resonance in Medicine* 77.3, pp. 1094–1103. DOI: 10.1002/mrm.26193.