## **PhD Thesis**

Investigating the Role of Bone Morphogenetic Protein Receptor IB and Putative Ligands in Reproductive Development and Steroidogenic Regulation in Male Mice

## Ilona Ciller

B.Sc.Hon., (University of New England)

## **Acknowledgments**

I would sincerely like to thank my supervisors A/Prof Jim McFarlane and the late Dr Tim O'Shea for allowing me to undertake this PhD. Your ideas and advice have been vital to almost every aspect of my research. I would also like to extend my thanks to Dr Suresh Kumar A.P. for helping me to learn all aspects of molecular biology and assisting in major gene experiments, Dr Ursula Ciller for teaching and helping me with RIA, cell culture and blood draws, Mr Kim Quinn for teaching the operation of the robot and qPCR cycler, Ms Ibtisam Al-Ali for helping me finish my immunization experiment when I was unable to, and Ms Janelle McFarlane for supplying me with all the mice I needed for my experimentation. Thank you also to all the McClymont Staff and technicians for letting me use resources, equipment and reagents needed to undergo my research project. Finally I would like to give a very special thankyou to friends and family – Khanjilla, Ibtisam, Balkis, Thor, Christopher, Sari, Suresh my husband, my sisters Ursula and Anja, nieces Heidi and Alison and my mom and dad Eugenia and Gerhard for your much-appreciated support, inspiration and good times.

Ch	apter 1	1
1.0	Introduction	2
1.1	Overview of TGF-βs in the Reproductive System	4
1.2	2 Biosynthesis and Structure of BMPs	7
1.3	B TGF-β and BMP Signalling	8
	1.3.1 Dimer Formation	8
	1.3.2 Receptors	9
	1.3.3 Binding Epitopes of Type I and Type II receptors	. 11
	1.3.4 Receptor - Ligand Interactions in Surface Plasmon Resonance	
	Studies	. 11
	1.3.5 Receptor – Ligand Interactions in Cell Lines and Compensation	. 12
	1.3.6 Receptor Autophosphorylation in Transfected Cells	. 14
	<b>1.3.7</b> Receptor Activation Profile for BMP-7 in vitro verses in vivo	. 14
	<b>1.3.8</b> Smads	. 15
	1.3.9 BMP Signalling and Other Kinase Pathways	
1.4	BMP Receptors in Reproduction	. 18
	1.4.1 Naturally Occurring Mutant BMP and BMP Receptors	. 19
1.5	Some Bone Morphogenetic Protein Family Members	. 22
	<b>1.5.1</b> BMP-1	. 23
	<b>1.5.2</b> BMP-2 (BMP-2a)	. 23
	<b>1.5.3</b> BMP-3 (Osteogenin)	. 26
	<b>1.5.4</b> BMP-3b (GDF-10)	. 26
	<b>1.5.5</b> BMP-4 (BMP-2b)	. 27
	<b>1.5.6</b> BMP-5	. 28
	<b>1.5.7</b> BMP-6 (Vgr-1)	. 28
	<b>1.5.8</b> BMP-7 (OP-1)	. 30
	<b>1.5.9</b> BMP-8a (OP-2)	. 30
	<b>1.5.10</b> BMP-8b (OP-3)	. 31
	<b>1.5.11</b> BMP-15 (GDF-9B)	. 31
	<b>1.5.12</b> GDF-9	. 32
	BMP Agonists and Antagonists	
1.7	' Embryology - Primordial Germ Cells	
	<b>1.7.1</b> TGF-βs in the Proliferation of Primordial Germ Cells	. 37
	<b>1.7.2</b> TGF-βs and Male Determination	. 39

<b>1.8</b> BMPs in the Female Reproductive System	40
<b>1.8.1</b> The Hypothalamic – Pituitary – Ovarian Axis	41
1.8.2 Follicular Development	43
<b>1.8.2.1</b> TGF-βs Involved in Follicular Development	44
1.8.2.2 Primordial to Primary Follicle Transition	44
1.8.2.3 Primary to Secondary Follicle Transition	46
1.8.2.4 Secondary to Antral Follicle Transition	49
1.8.3 BMPs in Other Female Reproductive Tissues	54
1.8.4 Ovarian Steroidogenesis	55
1.9 BMPs in the Male Reproductive System	59
<b>1.9.1</b> The Hypothalamic – Pituitary – Testicular Axis	61
1.9.2 Spermatogenesis	62
1.9.3 Leydig Cell Steroidogenesis	65
<b>1.9.4</b> TGF-βs in the Prostate	66
<b>1.9.5</b> TGF-βs in Other Male Reproductive Tissues	68
1.10 Proposed Project	69
Chapter 2	70
2.0 Materials and Methods	71
<b>2.1</b> Animals	
2.2 Antibodies and Reagents	
2.3 Antibody Purification from Egg Yolk and Plasma	73
2.4 Antibody Biotinylation	
2.5 Direct ELISA	75
2.6 Competitive ELISA	79
2.7 RNA Extraction	
<b>2.8</b> RT-PCR	81
<b>2.9</b> PCR	81
<b>2.10</b> QPCR	83
2.11 Agarose Gel Electrophoresis	
2.12 Tissue Culture	84
2.13 Leydig Cell Isolation	85
2.14 Testicular Interstitial Cell Isolation	86
2.15 Steroid Extraction	86

2.16 Testosterone Radioimmunoassay	87
2.17 Histology	87
<b>2.18</b> 3β-HSD Histochemistry	88
2.19 Trypan Blue Viability Test	88
2.17 Analysis	89
Chapter 3	90
3.0 Postnatal Expression of Bone Morphogenetic Proteins and their Receptors	
in the Mouse Testis	91
3.1 Introduction	91
3.2 Experimental Design	94
3.3 Results	95
3.3.1 Body and Reproductive Organ Weight Throughout Postnatal	
Development of Male Mice	95
3.3.2 Comparative Analysis of Testicular BMP Receptor Gene Expression	
During Postnatal Development	99
3.3.3 Comparative Analysis of BMP Gene Expression in Mouse Testis	
During Postnatal Development	100
3.3.4 Distribution of BMP Receptor and BMP Genes in Adult Male	
Reproductive and Steroidogenic Tissues	. 103
3.4 Discussion	105
Chapter 4	. 109
<b>4.0</b> The Effects of <i>in vivo</i> BMPR-IB Immunization on Reproductive	
Characteristics of Pubertal Male Mice	. 110
4.1 Introduction	. 110
4.2 Experimental Design	. 113
4.3 Results	
4.3.1 Body Weight of Pubertal Mice Immunized Against BMPR-IB	
4.3.2 Seminal Vesicle and Testis Weight in BMPR-IB immunized Pubertal	
Mice	. 114
4.3.3 Serum and Intra-testicular Testosterone in BMPR-IB immunized	
Pubertal Male Mice	. 115
4.3.4 3β-Hydroxysteroid Dehydrogenase Staining in BMPR-IB immunized	

Pubertal Mouse Testis	117
4.3.5 The mRNA Expression of Cyp19, Hsd3b1, Hsd3b6 and Hsd17b3 in	1
Pubertal Mouse Testis Immunized Against BMPR-IB	119
4.4 Discussion	120
Chautar F	400
Chapter 5.	
<b>5.0</b> The Effects of <i>in vivo</i> BMPR-IB Immunization on Reproductive	
Characteristics of Adult Male Mice	
5.1 Introduction	
<b>5.2</b> Experimental Design	
5.3 Results	
<b>5.3.1</b> Body Weight of Adult Mice Immunized Against BMPR-IB	
5.3.2 Seminal Vesicle and Testis Weight in BMPR-IB Immunized Adult	
Mice	
5.3.3 Serum and Intra-testicular Testosterone in BMPR-IB Immunized	
Adult Male Mice	
<b>5.3.4</b> 3β-Hydroxysteroid Dehydrogenase Staining in BMPR-IB immunized	
Adult Mouse Testis	130
5.3.5 The mRNA Expression of Cyp19, Hsd3b1, Hsd3b6 and Hsd17b3 in	1
Adult Mouse Testis Immunized Against BMPR-IB	132
5.4 Discussion	133
Chapter 6	136
6.0 Effects of in vitro BMPR-IB Immunization on Testosterone Production in	1
Testicular Culture and Leydig Cell Culture of Male Mice	137
6.1 Introduction	137
6.2 Experimental Design	139
6.3 Results	
6.3.1 Effect of anti-BMPR-IB on Testosterone Secretion by Testicular	ŕ
Tissue in Pubertal and Adult Mice	
6.3.2 Effect of Anti-BMPR-IB on Testosterone Secretion by Leydig Cells	143
6.3.3 Testosterone Secretion by Leydig Cells in Leydig Cell Conditioned	
Media	144

<b>6.3.4</b> Testosterone Secretion by Leydig Cells in Seminiferous Tubule	
Conditioned Media	145
6.3.5 Comparison of the Effect of Anti-BMPR-IB on Leydig Cell	
Testosterone Secretion in Normal, Leydig Cell Conditioned and	
Seminiferous Tubule Conditioned Media	146
6.3.6 The Effects of anti-BMP-2 and anti-BMP-4 on Pubertal and Adult	
Mouse Testicular Interstitial Cell Testosterone Production	148
6.4 Discussion	150
Chapter 7.	154
7.0 General Discussion	155
Reference List	160

## **List of Figures and Tables**

- **Figure 1.1** Diagram showing tetrameric complex formation of type I and type II serine/threonine receptors when activated by BMPs and consequential Smad signal transducer activities.
- **Figure 1.2** Bone morphogenetic protein carboxy-tail domain amino acid sequence map for 10 BMP family members.
- Figure 1.3 Enzymatic pathways to steroid synthesis in the male and female gonad.
- Figure 1.4 The male mouse reporoductive track
- **Figure 1.5** Schematic representation of the hypothalamic-pituitary-testicular axis and feedback mechanisms.
- **Figure 1.6** Stages of spermatogenesis as germ cells undergo mitosis and meiosis.
- Figure 2.1 Absorbance of chicken antibodies against their target ligands
- Figure 2.2 Absorbance of sheep antibodies against their target ligands
- **Figure 2.3** A dose response curve for BMPR-IB antibody JMCK#59.
- Figure 2.4 In vitro comparison of BMPR-IB antibodies JMCK#58 and JMCK#59
- Figure 2.5 A dose response curve for BMP-4 using JMCK#54
- Figure 2.6 A Bmpr1b PCR optimized using a MgCl<sub>2</sub> gradient
- **Figure 3.1** Body weight (g) in male mice from 2 to 8 weeks.
- Figure 3.2 Testis weight (g) in male mice from 2 to 8 weeks.
- **Figure 3.3** Epididymis weight (g) in male mice from 2 to 8 weeks.
- **Figure 3.4** Vas deferens weight (g) in male mice from 2 to 8 weeks.
- **Figure 3.5** Seminal vesicle weight (g) in male mice from 2 to 8 weeks.
- **Figure 3.6** Coagulating gland weight (g) in male mice from 2 to 8 weeks.
- Figure 3.7 Prostate weight (g) in male mice from 2 to 8 weeks.
- **Figure 3.8** Bmpr1a, Bmpr1b and Bmpr2 mRNA expression in 2, 4, 6 and 8 week old mouse testis.
- Figure 3.9 (Panel A) Bmp2, Bmp3, Bmp3b, Bmp4 and Bmp5, (Panel B) Bmp6, Bmp7, Bmp8a, Bmp8b and Bmp15 mRNA levels in 2, 4, 6 and 8 week old mouse testis.
- **Figure 3.10** BMP receptor and BMP mRNA expression in adult male reproductive and steroidogenic tissues.

- **Figure 4.1** (Panel A) seminal vesicle weight (g) and (Panel B) testis weight (g) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of pubertal male mice after 6 days of treatment.
- **Figure 4.2** Serum testosterone (ng/ml) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of pubertal male mice after 6 days of treatment.
- **Figure 4.3** Intra-testicular testosterone (ng/mg) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of pubertal male mice after six days of treatment.
- **Figure 4.4** 3β-HSD staining in (A) Control, (B) BMPR-IB immunized, (C) eCG treated and (D) BMPR-IB immunized plus eCG treated groups of pubertal mouse testis.
- **Figure 4.5** *Cyp19, Hsd3b1, Hsd3b6* and *Hsd17b3* mRNA expression in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treated pubertal mouse testis after 6 days of treatment.
- **Figure 5.1** (Panel A) seminal vesicle weight (g) and (Panel B) testis weight (g) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of mature male mice after 6 days of treatment.
- **Figure 5.2** Serum testosterone (ng/ml) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of mature male mice after 6 days of treatment.
- **Figure 5.3** Intra-testicular testosterone (ng/mg) in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treatment groups of mature male mice after 6 days of treatment.
- **Figure 5.4** 3β-HSD staining in (A) Control, (B) BMPR-IB immunized, (C) eCG treated, and (D) BMPR-IB immunized plus eCG treated groups of mature mice.
- **Figure 5.5** *Cyp19, Hsd3b1, Hsd3b6 and Hsd17b3* mRNA expression in control, BMPR-IB immunized, eCG and BMPR-IB immunized plus eCG treated adult mouse testis.
- **Figure 6.1** Percent testosterone in pubertal testis culture media after treatment with anti-BMPR-IB, eCG and anti-BMPR-IB plus eCG in comparison to no treatment (control).

- **Figure 6.2** Percent testosterone in mature testis culture after treatment with anti-BMPR-IB, eCG and anti-BMPR-IB plus eCG in comparison to no treatment (control).
- **Figure 6.3** Percent testosterone in adult Leydig cell culture media after treatment with anti-BMPR-IB, hCG and anti-BMPR-IB plus hCG in comparison to no treatment (control).
- **Figure 6.4** Percent testosterone in adult Leydig cell culture media after treatment in Leydig cell conditioned media previously treated with anti-BMPR-IB, hCG and anti-BMPR-IB plus hCG in comparison to no treatment (control).
- **Figure 6.5** Percent testosterone in adult Leydig cell culture media after treatment with seminiferous tubule conditioned media previously treated with anti-BMPR-IB, FSH and anti-BMPR-IB plus FSH in comparison to no treatment (control).
- **Figure 6.6** Leydig cell culture media testosterone in ng/10K cells in control, anti-BMPR-IB, gonadotrophin and anti-BMPR-IB plus gonadotrophin treatment groups in normal (unconditioned) media, Leydig cell conditioned media and seminiferous tubule conditioned media.
- **Figure 6.7** Testosterone as percent of the control in pubertal interstitial cell culture media in treatment groups of anti-BMP-2, anti-BMP-4, eCG, eCG plus anti-BMP-2, and eCG plus anti-BMP-4 in comparison to no treatment (control).
- **Figure 6.8** Testosterone as percent of the control in mature interstitial cell culture media in treatment groups of anti-BMP-2, anti-BMP-4, eCG, eCG plus anti-BMP-2, and eCG plus anti-BMP-4 in comparison to no treatment (control).
- Table 1.1 Comparison of Homo sapiens BMP-2 aa Sequence Across Species
- **Table 2.1** Species Comparison of BMPR-IB, BMP-2 and BMP-4 Peptide Sequences
- **Table 2.2** BMPR-IB Peptide Sequence Comparisons
- **Table 2.3** List of BMPR-1B, BMP-2, BMP-4 and non-immune control antibodies purified from plasma or yolk
- **Table 2.4** Primers used for RT-PCR and RT-qPCR
- Table 4.1 Effect of 6-day BMPR-IB Immunization on Body Weight of Pubertal Mice
- Table 5.1 Effect of 6-day BMPR-IB Immunization on Body Weight of Adult Mice