

SIMSUG 2010, Exeter, UK, 28-30 April 2010.

Isotope tracing of nitrate: lessons from Malta

Tim H.E. Heaton¹, Marianne E. Stuart², Manuel Sapiano³, Miriam Micallef Sultana³

¹ British Geological Survey, Nottingham NG12 5GG, England

² British Geological Survey, Wallingford, OX10 8BB, England

³ Malta Resources Authority, Marsa, MRS 9065, Malta

ABSTRACT

Average concentrations of nitrate in Malta's groundwaters are probably the highest among EU member states. This compromises the quality of an important resource - almost 60% of Malta's water supply being provided by groundwater. An $^{15}\text{N}/^{14}\text{N} + ^{18}\text{O}/^{16}\text{O}$ isotope study was undertaken as a core part of wide-ranging investigations into the potential sources of the nitrate pollution, its likely future trends, and possible ameliorative actions. The dual isotope ($^{15}\text{N}/^{14}\text{N} + ^{18}\text{O}/^{16}\text{O}$) approach was important for identifying waters affected by denitrification. Excluding these, groundwater from three physically and hydrologically distinct aquifers, with a very wide range in nitrate concentrations (24 to 410 mg $\text{NO}_3 \text{ L}^{-1}$), had remarkably similar isotope compositions: 90% of samples lying within $\delta^{15}\text{N} \approx +8$ to $+12\text{‰}$, and $\delta^{18}\text{O} \approx +3$ to $+6\text{‰}$. The $\delta^{18}\text{O}$ values are entirely consistent with those expected for microbial nitrification in the presence of surface or groundwaters, and together with $\delta^{15}\text{N}$ values rule out nitrate derivation directly from fertilizers or sewage. In other studies the relatively high $\delta^{15}\text{N}$ values for the waters would probably have been interpreted as indicative of nitrate derived from manure. In Malta, however, cultivated soils have high $\delta^{15}\text{N}$ values, $\approx +6$ to $+11\text{‰}$, very similar to the values for nitrate in the groundwater, and argue for a soil-derived source. The implications of a soil-source of such high nitrate levels are discussed, and the study emphasised the importance of characterising the compositions of soils in addition to other sources – a factor often neglected in isotope studies of nitrate.