



# Singular Anisotropic Problems with Competition Phenomena

Nikolaos S. Papageorgiou<sup>1</sup> · Calogero Vetro<sup>2</sup> · Francesca Vetro<sup>3</sup>

Received: 12 December 2022 / Accepted: 10 February 2023 / Published online: 28 March 2023  
© The Author(s) 2023

## Abstract

We consider a Dirichlet problem driven by the anisotropic  $(p(z), q(z))$ -Laplacian, with a parametric reaction exhibiting the combined effects of singular and concave-convex nonlinearities. The superlinear term may change sign. Using variational tools together with truncation and comparison techniques, we prove a global (for the parameter  $\lambda > 0$ ) existence and multiplicity theorem (a bifurcation-type theorem).

**Keywords** Singular and concave–convex nonlinearities · Anisotropic regularity · Hardy’s inequality · Global existence and multiplicity of solutions · Strong comparison · Truncations

**Mathematics Subject Classification** 35J75

## 1 Introduction

Let  $\Omega \subseteq \mathbb{R}^N$  be a bounded domain with a  $C^2$ -boundary  $\partial\Omega$ . In this paper we study the following singular anisotropic Dirichlet problem

---

✉ Calogero Vetro  
calogero.vetro@unipa.it

Nikolaos S. Papageorgiou  
npapg@math.ntua.gr

Francesca Vetro  
francescavetro80@gmail.com

<sup>1</sup> Department of Mathematics, National Technical University, Zografou campus, 15780 Athens, Greece

<sup>2</sup> Department of Mathematics and Computer Science, University of Palermo, Via Archirafi 34, 90123 Palermo, Italy

<sup>3</sup> Palermo, Italy

$$\begin{cases} -\Delta_{p(z)}u(z) - \Delta_{q(z)}u(z) = \lambda[u(z)^{-\eta(z)} + u(z)^{\tau(z)-1}] + f(z, u(z)) & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, \lambda > 0, u > 0. \end{cases} \tag{P_\lambda}$$

Given  $r \in C(\overline{\Omega})$  with  $1 < \min_{\overline{\Omega}} r$ , by  $\Delta_{r(z)}$  we denote the anisotropic  $r$ -Laplace differential operator defined by

$$\Delta_{r(z)}u = \operatorname{div} (|\nabla u|^{r(z)-2}\nabla u) \text{ for all } u \in W_0^{1,r(z)}(\Omega).$$

In contrast to the isotropic  $r$ -Laplacian (that is,  $r(\cdot)$  is constant), the anisotropic operator is not homogeneous. In  $(P_\lambda)$  the equation is driven by the sum of two such operators with distinct variable exponents  $p(\cdot)$  and  $q(\cdot)$  (double phase problem). Given  $\vartheta \in L^\infty(\Omega)$ , we set  $\vartheta_- = \operatorname{ess\,inf}_\Omega \vartheta$  and  $\vartheta_+ = \operatorname{ess\,sup}_\Omega \vartheta$ . In  $(P_\lambda)$  we assume that  $1 < \tau_- \leq \tau_+ < q_- \leq q_+ < p_- \leq p_+$  and  $0 < \eta_- \leq \eta_+ < 1$ . The perturbation  $f(z, x)$  is a Carathéodory function (that is, for all  $x \in \mathbb{R}$   $z \rightarrow f(z, x)$  is measurable and for a.a.  $z \in \Omega$   $x \rightarrow f(z, x)$  is continuous) which exhibits  $(p_+ - 1)$ -superlinear growth as  $x \rightarrow +\infty$ , but need not satisfy the usual in such cases Ambrosetti–Rabinowitz condition (the AR-condition for short) and may change sign (indefinite perturbation). So, problem  $(P_\lambda)$  in the reaction has the combined effects of singular and concave–convex nonlinearities with two distinguishing features. First the superlinear (convex) term need not satisfy the AR-condition and second this perturbation is in general sign-changing. In the past, anisotropic singular equations were studied without the presence of the concave term  $\lambda u^{\tau(z)-1}$  and with a superlinear perturbation which is positive. We refer to the works of Byun–Ko [2] and Saoudi–Ghanmi [21]. Both deal with equations driven by the anisotropic  $p$ -Laplacian only. More recently, Papageorgiou–Rădulescu–Zhang [19] considered singular anisotropic double phase problems with a superlinear positive perturbation and no concave term.

Closer to our work here is the recent paper of Papageorgiou–Winkert [14], who examined an isotropic version of problem  $(P_\lambda)$  (all the exponents of the problem are constant) with a superlinear positive perturbation. The definite sign of the perturbation allows the authors of [14] to produce an ordered pair of upper and lower solutions, which in turn leads to the nonemptiness of the set of admissible parameters. They prove a global existence and multiplicity result (a bifurcation-type theorem). Our aim in this paper is to extend their result to anisotropic problems with an indefinite superlinear perturbation.

Finally we mention also the recent works on some other classes of anisotropic singular problems of Papageorgiou–Winkert [13, 15] and Papageorgiou–Zhang [16, 17]; for problems in divergence form, some recent results are given in Abdalmonem–Scapellato [1], Ragusa [20] and Wei [23] for parabolic equations.

## 2 Mathematical Background: Hypotheses

The analysis of problem  $(P_\lambda)$  is based on the variable Lebesgue and Sobolev spaces. A comprehensive introduction to the subject can be found in the books of Cruz Uribe–Fiorenza [3] and of Diening–Harjulehto–Hästö–Růžička [4].

We introduce the set

$$E_1 = \left\{ r \in C(\overline{\Omega}) : 1 < \min_{\overline{\Omega}} r \right\}.$$

Recall that for  $r \in C(\overline{\Omega})$ ,  $r_- = \min_{\overline{\Omega}} r$  and  $r_+ = \max_{\overline{\Omega}} r$ . Let  $L^0(\Omega)$  be the space of all measurable functions  $u : \Omega \rightarrow \mathbb{R}$ . As usual we identify two such functions which differ only on a Lebesgue null subset of  $\Omega$ . Given  $r \in E_1$ , the variable Lebesgue space  $L^{r(z)}(\Omega)$  is defined by

$$L^{r(z)}(\Omega) = \left\{ u \in L^0(\Omega) : \rho_r(u) = \int_{\Omega} |u|^{r(z)} dz < +\infty \right\}.$$

We endow this space with the so-called “Luxemburg norm” defined by

$$\|u\|_{r(z)} = \inf \left[ \lambda > 0 : \int_{\Omega} \left( \frac{|u|}{\lambda} \right)^{r(z)} dz \leq 1 \right].$$

With this norm the space  $L^{r(z)}(\Omega)$  becomes a separable and uniformly convex (thus reflexive, see [12], p. 225) Banach space. Let  $r' \in E_1$  be the conjugate variable exponent to  $r(\cdot)$ , defined by

$$r'(z) = \frac{r(z)}{r(z) - 1} \text{ or equivalently } \frac{1}{r(z)} + \frac{1}{r'(z)} = 1 \text{ for all } z \in \overline{\Omega}.$$

We have that

$$L^{r(z)}(\Omega)^* = L^{r'(z)}(\Omega),$$

and the following Hölder-type inequality holds

$$\int_{\Omega} |uv| dz \leq \left[ \frac{1}{r_-} + \frac{1}{r'_-} \right] \|u\|_{r(z)} \|v\|_{r'(z)} \text{ for all } u \in L^{r(z)}(\Omega), \text{ all } v \in L^{r'(z)}(\Omega).$$

If  $r_1, r_2 \in E_1$  and  $r_1(z) \leq r_2(z)$  for all  $z \in \overline{\Omega}$ , then

$$L^{r_2(z)}(\Omega) \hookrightarrow L^{r_1(z)}(\Omega) \text{ continuously.}$$

Using the variable Lebesgue spaces, we can define the corresponding variable Sobolev spaces. So, given  $r \in E_1$ , the variable Sobolev space  $W^{1,r(z)}(\Omega)$  is defined by

$$W^{1,r(z)}(\Omega) = \left\{ u \in L^{r(z)}(\Omega) : |\nabla u| \in L^{r(z)}(\Omega) \right\},$$

with  $\nabla u$  being the weak gradient of  $u(\cdot)$ . This space is equipped with the following norm

$$\|u\|_{1,r(z)} = \|u\|_{r(z)} + \|\nabla u\|_{r(z)} \text{ for all } u \in W^{1,r(z)}(\Omega),$$

with  $\|\nabla u\|_{r(z)} = \| |\nabla u| \|_{r(z)}$ .

By  $C^{0,1}(\overline{\Omega})$  we denote the space of all Lipschitz continuous functions  $u : \overline{\Omega} \rightarrow \mathbb{R}$ . Given  $r \in C^{0,1}(\overline{\Omega}) \cap E_1$ , we define

$$W_0^{1,r(z)}(\Omega) = \overline{C_c^\infty(\Omega)}^{\|\cdot\|_{1,r(z)}}.$$

Both spaces  $W^{1,r(z)}(\Omega)$  and  $W_0^{1,r(z)}(\Omega)$  are separable and uniformly convex (thus reflexive) Banach spaces. Since in the definition of  $W_0^{1,r(z)}(\Omega)$  we assume that the exponent  $r(\cdot)$  is Lipschitz continuous, the Poincaré inequality holds, that is, there exists  $c = c(\Omega) > 0$  such that

$$\|u\|_{r(z)} \leq c \|\nabla u\|_{r(z)} \text{ for all } u \in W_0^{1,r(z)}(\Omega).$$

The Poincaré inequality leads to the following equivalent norm on  $W_0^{1,r(z)}(\Omega)$

$$\|u\| = \|\nabla u\|_{r(z)} \text{ for all } u \in W_0^{1,r(z)}(\Omega).$$

In the sequel we will use this norm on  $W_0^{1,r(z)}(\Omega)$ . For  $r \in E_1$ , we introduce the corresponding critical Sobolev exponent  $r^*(\cdot)$  given by

$$r^*(z) = \begin{cases} \frac{Nr(z)}{N-r(z)} & \text{if } r(z) < N, \\ +\infty & \text{if } N \leq r(z). \end{cases}$$

There is an anisotropic version of the Sobolev embedding theorem.

**Proposition 1** *If  $r \in C^{0,1}(\overline{\Omega}) \cap E_1$ ,  $r_+ < N$ ,  $q \in E_1$  and  $q(z) \leq r^*(z)$  (resp.  $q(z) < r^*(z)$ ) for all  $z \in \overline{\Omega}$ , then  $W_0^{1,r(z)}(\Omega) \hookrightarrow L^{q(z)}(\Omega)$  continuously (resp. compactly).*

There is a close relation between the norm  $\|\cdot\|_{r(z)}$  and the modular function  $\rho_r(u) = \int_{\Omega} |u|^{r(z)} dz$ .

**Proposition 2** *Suppose  $r \in E_1$  and  $\{u_n, u\}_{n \in \mathbb{N}} \subseteq L^{r(z)}(\Omega)$ , then we have:*

(a)  $\|u\|_{r(z)} = \lambda \Leftrightarrow \rho_r\left(\frac{u}{\lambda}\right) = 1$  ( $\lambda > 0$ ).

- (b)  $\|u\|_{r(z)} < 1$  ( resp.  $= 1, > 1$ )  $\Leftrightarrow \rho_r(u) < 1$  ( resp.  $= 1, > 1$ ).
- (c)  $\|u\|_{r(z)} < 1 \Rightarrow \|u\|_{r(z)}^{r_+} \leq \rho_r(u) \leq \|u\|_{r(z)}^{r_-}$ .
- (d)  $\|u\|_{r(z)} > 1 \Rightarrow \|u\|_{r(z)}^{r_-} \leq \rho_r(u) \leq \|u\|_{r(z)}^{r_+}$ .
- (e)  $\|u_n\|_{r(z)} \rightarrow 0$  ( resp.  $\rightarrow +\infty$ )  $\Leftrightarrow \rho_r(u_n) \rightarrow 0$  ( resp.  $\rightarrow +\infty$ ).

We know that

$$W_0^{1,r(z)}(\Omega)^* = W^{-1,r'(z)}(\Omega).$$

Let  $A_r : W_0^{1,r(z)}(\Omega) \rightarrow W^{-1,r'(z)}(\Omega)$  be defined by

$$\langle A_r(u), h \rangle = \int_{\Omega} |\nabla u|^{r(z)-2} (\nabla u, \nabla h)_{\mathbb{R}^N} dz \text{ for all } u, h \in W_0^{1,r(z)}(\Omega).$$

This operator has the following properties (see Gasiński–Papageorgiou [6], Proposition 2.5).

**Proposition 3** *The operator  $A_r(\cdot)$  is bounded (that is, maps bounded sets to bounded sets), continuous, strictly monotone (thus maximal monotone too) and of type  $(S)_+$  (that is, if  $u_n \xrightarrow{w} u$  in  $W_0^{1,r(z)}(\Omega)$  and  $\limsup_{n \rightarrow +\infty} \langle A_r(u_n), u_n - u \rangle \leq 0$ , then  $u_n \rightarrow u$  in  $W_0^{1,r(z)}(\Omega)$ ).*

The anisotropic regularity theory (see Fan [5] and Lieberman [11] for the corresponding isotropic theory) will lead us to the space  $C_0^1(\Omega) = \{u \in C^1(\overline{\Omega}) : u|_{\partial\Omega} = 0\}$ . This is an ordered Banach space with positive (order) cone  $C_+ = \{u \in C_0^1(\overline{\Omega}) : u(z) \geq 0 \text{ for all } z \in \overline{\Omega}\}$ . This cone has a nonempty interior given by

$$\text{int } C_+ = \left\{ u \in C_+ : u(z) > 0 \text{ for all } z \in \Omega, \quad \frac{\partial u}{\partial n} \Big|_{\partial\Omega} < 0 \right\},$$

where  $\frac{\partial u}{\partial n} = (\nabla u, n)_{\mathbb{R}^N}$  with  $n(\cdot)$  being the outward unit normal on  $\partial\Omega$ .

Let  $u_1, u_2 \in L^0(\Omega)$  such that  $u_1(z) \leq u_2(z)$  for a.a.  $z \in \Omega$ . We introduce the following sets:

$$\begin{aligned} [u_1, u_2] &= \{h \in W_0^{1,p(z)}(\Omega) : u_1(z) \leq h(z) \leq u_2(z) \text{ for a.a. } z \in \Omega\}, \\ \text{int}_{C_0^1(\overline{\Omega})} [u_1, u_2] &= \text{interior in } C_0^1(\overline{\Omega}) \text{ of } [u_1, u_2] \cap C_0^1(\overline{\Omega}), \\ [u_1] &= \{h \in W_0^{1,p(z)}(\Omega) : u_1(z) \leq h(z) \text{ for a.a. } z \in \Omega\}. \end{aligned}$$

If  $h_1, h_2 \in L^0(\Omega)$ , then we say that  $h_1 \prec h_2$  if and only if for every  $K \subseteq \Omega$  compact we have

$$0 < c_K \leq h_2(z) - h_1(z) \text{ for a.a. } z \in K.$$

Evidently, if  $h_1, h_2 \in C(\Omega)$  and  $h_1(z) < h_2(z)$  for all  $z \in \Omega$ , then  $h_1 < h_2$ .

Given  $h \in L^0(\Omega)$ , we set

$$h^+(z) = \max\{h(z), 0\} \text{ and } h^-(z) = \max\{-h(z), 0\} \text{ for all } z \in \Omega.$$

We have  $h^\pm \in L^0(\Omega)$ ,  $h = h^+ - h^-$ ,  $|h| = h^+ + h^-$  and if  $h \in W_0^{1,p(z)}(\Omega)$ , then  $h^\pm \in W_0^{1,p(z)}(\Omega)$ .

Let  $X$  be a Banach space and  $\varphi \in C^1(X, \mathbb{R})$ . By  $K_\varphi$  we denote the critical set of  $\varphi(\cdot)$ , that is

$$K_\varphi = \{u \in X : \varphi'(u) = 0\}.$$

We say that  $\varphi(\cdot)$  satisfies the ‘‘C-condition’’, if it has the following property:

‘‘Every sequence  $\{u_n\}_{n \in \mathbb{N}} \subseteq X$  such that  
 $\{\varphi(u_n)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$  is bounded  
 and  $(1 + \|u_n\|_X)\varphi'(u_n) \rightarrow 0$  in  $X^*$  as  $n \rightarrow +\infty$ ,  
 admits a strongly convergent subsequence’’.

The hypotheses on the data of  $(P_\lambda)$  are the following:

- $H_0 : p, q \in C^{0,1}(\overline{\Omega}), \tau \in C(\overline{\Omega}), 1 < \tau_- \leq \tau_+ < q_- \leq q_+ < p_- \leq p_+ < N,$   
 $\eta \in C(\overline{\Omega}), 0 < \eta_- \leq \eta_+ < 1.$
- $H_1 : f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function such that  $f(z, 0) = 0$  for a.a.  $z \in \Omega$ , and
  - (i)  $|f(z, x)| \leq a(z)[1 + x^{r(z)-1}]$  for a.a.  $z \in \Omega$ , all  $x \geq 0$ , with  $a \in L^\infty(\Omega)_+,$   
 $r \in C(\overline{\Omega}), p_+ < r_- \leq r_+ < p_-^*;$
  - (ii) if  $F(z, x) = \int_0^x f(z, s)ds$ , then  $\lim_{x \rightarrow +\infty} \frac{F(z,x)}{x^{p_+}} = +\infty$  uniformly for a.a.  $z \in \Omega;$
  - (iii) if  $e_\lambda(z, x) = \lambda [x^{1-\eta(z)} + x^{\tau(z)}] + f(z, x)x - \lambda p_+ \left[ \frac{1}{1-\eta(z)}x^{1-\eta(z)} + \frac{1}{\tau(z)}x^{\tau(z)} \right] + p_+F(z, x),$   
 then there exists  $\vartheta \in L^1(\Omega)$  such that  $e_\lambda(z, x) \leq e_\lambda(z, y) + \vartheta(z)$  for a.a.  $z \in \Omega,$   
 all  $0 \leq x \leq y;$
  - (iv)  $\lim_{x \rightarrow 0^+} \frac{f(z,x)}{x^{q_+ - 1}} = 0$  uniformly for a.a.  $z \in \Omega$ , there exists  $\delta > 0$  such that  $0 < m_s \leq f(z, x)$  for a.a.  $z \in \Omega$ , all  $0 < s \leq x \leq \delta$ , and for every  $\rho > 0$  there exists  $\widehat{\xi}_\rho > 0$  such that for a.a.  $z \in \Omega$ , the function  $x \rightarrow f(z, x) + \widehat{\xi}_\rho|x|^{p(z)-1}$  is nondecreasing on  $[0, \rho].$

**Remark 1** Since we are looking for positive solutions and the above hypotheses concern the positive semiaxis  $\mathbb{R}_+ = [0, +\infty)$ , without any loss of generality we may assume that  $f(z, x) = 0$  for a.a.  $z \in \Omega$ , all  $x \leq 0$ . Hypotheses  $H_1$  (ii), (iii) imply that

$$\lim_{x \rightarrow +\infty} \frac{f(z, x)}{x^{p_+ - 1}} = +\infty \text{ uniformly for a.a. } z \in \Omega.$$

So,  $f(z, \cdot)$  is  $(p_+ - 1)$ -superlinear, but need not satisfy the AR-condition which is common in the literature when studying superlinear problems (see Willem [24], p. 46). Instead we use the quasimonotonicity condition on  $e_\lambda(z, \cdot)$  (see hypothesis  $H_1$

(iii). This is a slight generalization of a condition used by Li-Yang [10]. If there exists  $M > 0$  such that for a.a.  $z \in \Omega$ ,  $x \rightarrow \frac{f(z,x)}{x^{p_+-1}}$  is nondecreasing on  $[M, +\infty)$ , then hypothesis  $H_1$  (iii) is satisfied. We stress that in contrast to [14], the perturbation here can be sign-changing.

Let  $V : W_0^{1,p(z)}(\Omega) \rightarrow W^{-1,p'(z)}(\Omega)$  be defined by

$$\langle V(u), h \rangle = \int_{\Omega} (|\nabla u|^{p(z)-2} + |\nabla u|^{q(z)-2})(\nabla u, \nabla h)_{\mathbb{R}^N} dz \text{ for all } u, h \in W_0^{1,p(z)}(\Omega).$$

Evidently  $V = A_p + A_q$  and so on account of Proposition 3, we have:

**Proposition 4** *The operator  $V(\cdot)$  is bounded, continuous, strictly monotone (thus maximal monotone too) and of type  $(S)_+$ .*

### 3 An Auxiliary Problem

In this section, we examine the following auxiliary anisotropic Dirichlet problem

$$\begin{cases} -\Delta_{p(z)}u(z) - \Delta_{q(z)}u(z) = \lambda u(z)^{\tau(z)-1} & \text{in } \Omega, \\ u|_{\partial\Omega} = 0, \lambda > 0, u > 0. \end{cases} \tag{Q_\lambda}$$

The solution of this problem will help us bypass the singularity and prove the existence of admissible parameters for problem  $(P_\lambda)$ .

**Proposition 5** *If hypothesis  $H_0$  holds, then for every  $\lambda > 0$  problem  $(Q_\lambda)$  has a unique positive solution  $\bar{u}_\lambda \in \text{int } C_+$  and  $\bar{u}_\lambda \rightarrow 0$  in  $C_0^1(\bar{\Omega})$  as  $\lambda \rightarrow 0^+$ .*

**Proof** First we show the existence of a positive solution for problem  $(Q_\lambda)$ . To this end let  $\sigma_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$  be the  $C^1$ -functional defined by

$$\sigma_\lambda(u) = \int_{\Omega} \frac{1}{p(z)} |\nabla u|^{p(z)} dz + \int_{\Omega} \frac{1}{q(z)} |\nabla u|^{q(z)} dz - \int_{\Omega} \frac{\lambda}{\tau(z)} (u^+)^{\tau(z)} dz$$

for all  $u \in W_0^{1,p(z)}(\Omega)$ . If  $\|u\|, \|u\|_{\tau(z)} \geq 1$ , then we have

$$\sigma_\lambda(u) \geq \frac{1}{p_+} \|u\|^{p_-} - \frac{\lambda c_0}{\tau_-} \|u\|^{\tau_-} \text{ for some } c_0 > 0 \text{ (see Proposition 2).}$$

Since  $\tau_+ < q_- < p_-$ , it follows that

$$\sigma_\lambda(\cdot) \text{ is coercive.}$$

The modular functions are convex continuous, hence sequentially weakly lower semi-continuous. This fact and Proposition 1 (the anisotropic Sobolev embedding theorem) imply that

$\sigma_\lambda(\cdot)$  is sequentially weakly lower semicontinuous.

Then the Weierstrass–Tonelli theorem implies that there exists  $\bar{u}_\lambda \in W_0^{1,p(z)}(\Omega)$  such that

$$\sigma_\lambda(\bar{u}_\lambda) = \inf \left[ \sigma_\lambda(u) : u \in W_0^{1,p(z)}(\Omega) \right]. \tag{1}$$

Let  $u \in W_0^{1,p(z)}(\Omega)$ ,  $u \neq 0$ . Then for  $t \in (0, 1)$  we have

$$\begin{aligned} \sigma_\lambda(tu) &\leq \frac{t^{q_-}}{q_-} [\rho_p(\nabla u) + \rho_q(\nabla u)] - \frac{t^{\tau_+}}{\tau_+} \rho_\tau(u) \\ &\leq c_1 t^{q_-} - c_2 t^{\tau_+} \text{ for some } c_1, c_2 > 0. \end{aligned}$$

Since  $\tau_+ < q_-$ , choosing  $t \in (0, 1)$  even smaller if necessary, we have

$$\begin{aligned} \sigma_\lambda(tu) &< 0, \\ \Rightarrow \sigma_\lambda(\bar{u}_\lambda) &< 0 = \sigma_\lambda(0) \quad (\text{see (1)}), \\ \Rightarrow \bar{u}_\lambda &\neq 0. \end{aligned}$$

From (1) we have

$$\begin{aligned} \langle \sigma'_\lambda(\bar{u}_\lambda), h \rangle &= 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega), \\ \Rightarrow \langle V(\bar{u}_\lambda), h \rangle &= \int_\Omega \lambda(u^+)^{\tau(z)-1} h \, dz \text{ for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned} \tag{2}$$

In (2) we use the test function  $h = -\bar{u}_\lambda^- \in W_0^{1,p(z)}(\Omega)$  and obtain

$$\begin{aligned} \rho_p(\nabla \bar{u}_\lambda^-) &\leq 0, \\ \Rightarrow \bar{u}_\lambda^- &\geq 0, \bar{u}_\lambda^- \neq 0 \quad (\text{see Proposition 2}). \end{aligned} \tag{3}$$

From (2) and (3) it follows that  $\bar{u}_\lambda$  is a positive solution of  $(Q_\lambda)$ . From [19] (Proposition A1), we have that  $\bar{u}_\lambda \in L^\infty(\Omega)$ . Then the anisotropic regularity theory (see Fan [5]) implies that  $\bar{u}_\lambda \in C_+ \setminus \{0\}$ . Finally the anisotropic maximum principle (see [19], Proposition A2) implies that

$$\bar{u}_\lambda \in \text{int } C_+.$$

Next we show that this positive solution of  $(Q_\lambda)$  is unique. For  $\tau_0 \in (\tau_+, q_-)$ , we consider the integral functional  $j : L^1(\Omega) \rightarrow \bar{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$  defined by

$$j(u) = \begin{cases} \int_\Omega \frac{1}{p(z)} |\nabla u|^{1/\tau_0} |p(z)| \, dz + \int_\Omega \frac{1}{q(z)} |\nabla u|^{1/\tau_0} |q(z)| \, dz & \text{if } u \geq 0, u^{1/\tau_0} \in W_0^{1,p(z)}(\Omega), \\ +\infty & \text{otherwise.} \end{cases}$$



Theorem 2.2 of Takáč–Giacomoni [22] implies that  $j(\cdot)$  is convex. Suppose  $\tilde{u}_\lambda$  is another positive solution of  $(Q_\lambda)$ . Again we have  $\tilde{u}_\lambda \in \text{int } C_+$ . Using Proposition 4.1.22, p. 274, of Papageorgiou–Rădulescu–Repovš [18], we have

$$\frac{\bar{u}_\lambda}{\tilde{u}_\lambda} \in L^\infty(\Omega) \text{ and } \frac{\tilde{u}_\lambda}{\bar{u}_\lambda} \in L^\infty(\Omega). \tag{4}$$

Let  $\text{dom } j = \{u \in L^1(\Omega) : j(u) < +\infty\}$  (the effective domain of  $j(\cdot)$ ) and let  $h = (\bar{u}_\lambda^{\tau_0} - \tilde{u}_\lambda^{\tau_0}) \in W_0^{1,p(z)}(\Omega)$ . On account of (4) for  $t \in (0, 1)$  small we have

$$\bar{u}_\lambda^{\tau_0} + th \in \text{dom } j, \quad \tilde{u}_\lambda^{\tau_0} + th \in \text{dom } j.$$

Then since  $j(\cdot)$  is convex, the directional derivatives of  $j(\cdot)$  at  $\bar{u}_\lambda^{\tau_0}$  and at  $\tilde{u}_\lambda^{\tau_0}$  in the direction  $h$  exist and using Green’s identity we have

$$\begin{aligned} j'(\bar{u}_\lambda^{\tau_0})(h) &= \frac{1}{\tau_0} \int_\Omega \frac{-\Delta_{p(z)}\bar{u}_\lambda - \Delta_{q(z)}\bar{u}_\lambda}{\bar{u}_\lambda^{\tau_0-1}} h \, dz \\ &= \frac{1}{\tau_0} \int_\Omega \lambda \bar{u}_\lambda^{\tau(z)-\tau_0} h \, dz, \\ j'(\tilde{u}_\lambda^{\tau_0})(h) &= \frac{1}{\tau_0} \int_\Omega \frac{-\Delta_{p(z)}\tilde{u}_\lambda - \Delta_{q(z)}\tilde{u}_\lambda}{\tilde{u}_\lambda^{\tau_0-1}} h \, dz \\ &= \frac{1}{\tau_0} \int_\Omega \lambda \tilde{u}_\lambda^{\tau(z)-\tau_0} h \, dz. \end{aligned}$$

The convexity of  $j(\cdot)$  implies the monotonicity of  $j'(\cdot)$ . Hence

$$\begin{aligned} 0 &\leq \int_\Omega \lambda [\bar{u}_\lambda^{\tau(z)-\tau_0} - \tilde{u}_\lambda^{\tau(z)-\tau_0}] (\bar{u}_\lambda^{\tau_0} - \tilde{u}_\lambda^{\tau_0}) \, dz \leq 0 \text{ (since } \tau_+ < \tau_0), \\ &\Rightarrow \tilde{u}_\lambda = \bar{u}_\lambda. \end{aligned}$$

This proves the uniqueness of the positive solution  $\bar{u}_\lambda \in \text{int } C_+$  of  $(Q_\lambda)$ . Finally we have

$$\langle V(\bar{u}_\lambda), h \rangle = \int_\Omega \lambda \bar{u}_\lambda^{\tau(z)-1} h \, dz \text{ for all } h \in W_0^{1,p(z)}(\Omega).$$

Using  $h = \bar{u}_\lambda \in W_0^{1,p(z)}(\Omega)$ , we obtain

$$\begin{aligned} \rho_p(\nabla \bar{u}_\lambda) &\leq \lambda \rho_\tau(\bar{u}_\lambda), \\ \Rightarrow \min\{\|\bar{u}_\lambda\|^{p^+}, \|\bar{u}_\lambda\|^{p^-}\} &\leq \lambda \max\{\|\bar{u}_\lambda\|_{\tau(z)}^{\tau_+}, \|\bar{u}_\lambda\|_{\tau(z)}^{\tau_-}\} \\ &\text{(see Proposition 2)} \\ &\leq \lambda c_3 \max\{\|\bar{u}_\lambda\|^{\tau_+}, \|\bar{u}_\lambda\|^{\tau_-}\} \\ &\text{for some } c_3 > 0 \text{ (see Proposition 1)}. \end{aligned}$$

Recall that  $\tau_+ < q_- < p_-$ . So, it follows that

$$\bar{u}_\lambda \rightarrow 0 \text{ in } W_0^{1,p(z)}(\Omega) \text{ as } \lambda \rightarrow 0^+. \tag{5}$$

The anisotropic regularity theory (see Fan [5]) implies that we can find  $\alpha \in (0, 1)$  and  $c_4 > 0$  such that

$$\bar{u}_\lambda \in C_0^{1,\alpha}(\bar{\Omega}), \|\bar{u}_\lambda\|_{C_0^{1,\alpha}(\bar{\Omega})} \leq c_4 \text{ for all } \lambda \in (0, 1]. \tag{6}$$

We know that  $C_0^{1,\alpha}(\bar{\Omega}) \hookrightarrow C_0^1(\bar{\Omega})$  compactly. So, from (6) and (5) we conclude that

$$\bar{u}_\lambda \rightarrow 0 \text{ in } C_0^1(\bar{\Omega}) \text{ as } \lambda \rightarrow 0^+.$$

□

### 4 Positive Solutions

We introduce the following two sets:

$$\begin{aligned} \mathcal{L} &= \{ \lambda > 0 : \text{problem } (P_\lambda) \text{ has a positive solution} \} \\ &\text{(set of admissible parameters),} \\ \mathcal{S}_\lambda &= \{ \text{set of positive solutions of } (P_\lambda) \}. \end{aligned}$$

**Proposition 6** *If hypotheses  $H_0, H_1$  hold, then  $\mathcal{L} \neq \emptyset$  and for all  $\lambda > 0$   $\mathcal{S}_\lambda \subseteq \text{int } C_+$ .*

**Proof** Let  $\delta > 0$  be as postulated by hypothesis  $H_1(iv)$ . On account of Proposition 5, we can find  $\lambda^* > 0$  such that

$$\|\bar{u}_\lambda\|_\infty \leq \delta \text{ for all } \lambda \in (0, \lambda^*]. \tag{7}$$

We fix  $\lambda \in (0, \lambda^*]$  and let  $\bar{u}_\lambda \in \text{int } C_+$  be the unique positive solution of  $(Q_\lambda)$  (see Proposition 5). We introduce the Carathéodory function  $g_\lambda(z, x)$  defined by

$$g_\lambda(z, x) = \begin{cases} \lambda[\bar{u}_\lambda(z)^{-\eta(z)} + \bar{u}_\lambda(z)^{\tau(z)-1}] + f(z, x^+) & \text{if } x \leq \bar{u}_\lambda(z), \\ \lambda[x^{-\eta(z)} + x^{\tau(z)-1}] + f(z, x) & \text{if } \bar{u}_\lambda(z) < x. \end{cases} \tag{8}$$

Let  $G_\lambda(z, x) = \int_0^x g_\lambda(z, s) ds$  and consider the  $C^1$ -functional  $\varphi_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\begin{aligned} \varphi_\lambda(u) &= \int_\Omega \frac{1}{p(z)} |\nabla u|^{p(z)} dz + \int_\Omega \frac{1}{q(z)} |\nabla u|^{q(z)} dz \\ &\quad - \int_\Omega G_\lambda(z, u) dz \text{ for all } u \in W_0^{1,p(z)}(\Omega). \end{aligned}$$

**Claim:**  $\varphi_\lambda(\cdot)$  satisfies the  $C$ -condition.

Consider a sequence  $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(z)}(\Omega)$  such that

$$|\varphi_\lambda(u_n)| \leq c_5 \text{ for some } c_5 > 0, \text{ all } n \in \mathbb{N}, \tag{9}$$

$$(1 + \|u_n\|)\varphi'_\lambda(u_n) \rightarrow 0 \text{ in } W^{-1,p'(z)}(\Omega) \text{ as } n \rightarrow +\infty. \tag{10}$$

From (10) we have

$$\left| \langle V(u_n), h \rangle - \int_\Omega g_\lambda(z, u_n)h \, dz \right| \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|}$$

for all  $h \in W_0^{1,p(z)}(\Omega)$ , with  $\varepsilon_n \rightarrow 0^+$ . (11)

In (11) we choose the test function  $h = -u_n^- \in W_0^{1,p(z)}(\Omega)$ . Using (8) we obtain

$$\begin{aligned} \rho_p(\nabla u_n^-) &\leq \varepsilon_n \text{ for all } n \in \mathbb{N}, \\ \Rightarrow u_n^- &\rightarrow 0 \text{ in } W_0^{1,p(z)}(\Omega) \text{ (see Proposition 2)}. \end{aligned} \tag{12}$$

We define

$$\widehat{f}_\lambda(z, x) = \lambda[x^{-\eta(z)} + x^{\tau(z)-1}] + f(z, x)$$

and  $\widehat{F}_\lambda(z, x) = \int_0^x \widehat{f}_\lambda(z, s) \, ds$ .

From (12), (9) and (8), we have

$$\begin{aligned} \int_\Omega \frac{p_+}{p(z)} |\nabla u_n^+|^{p(z)} \, dz + \int_\Omega \frac{p_+}{q(z)} |\nabla u_n^+|^{q(z)} \, dz - \int_\Omega p_+ \widehat{F}_\lambda(z, u_n^+) \, dz &\leq c_6 \\ \text{for some } c_6 > 0, \text{ all } n \in \mathbb{N}, \\ \Rightarrow \rho_p(\nabla u_n^+) + \rho_q(\nabla u_n^+) - \int_\Omega p_+ \widehat{F}_\lambda(z, u_n^+) \, dz &\leq c_6 \text{ for all } n \in \mathbb{N} \\ \text{(since } p(z) \leq p_+ \text{ for all } z \in \overline{\Omega}\text{)}. \end{aligned} \tag{13}$$

In (11) we choose the test function  $h = u_n^+ \in W_0^{1,p(z)}(\Omega)$  and obtain

$$\begin{aligned} -\rho_p(\nabla u_n^+) - \rho_q(\nabla u_n^+) + \int_\Omega g_\lambda(z, u_n^+)u_n^+ \, dz &\leq \varepsilon_n \text{ for all } n \in \mathbb{N}, \\ \Rightarrow -\rho_p(\nabla u_n^+) - \rho_q(\nabla u_n^+) + \int_\Omega \widehat{f}_\lambda(z, u_n^+)u_n^+ \, dz &\leq c_7 \\ \text{for some } c_7 > 0, \text{ all } n \in \mathbb{N} \text{ (see (8))}. \end{aligned} \tag{14}$$

We add (13) and (14) and obtain

$$\int_\Omega e_\lambda(z, u_n^+) \, dz \leq c_8 \text{ for some } c_8 > 0, \text{ all } n \in \mathbb{N}, \tag{15}$$

(note that  $e_\lambda(z, x) = \widehat{f}_\lambda(z, x)x - p_+ \widehat{F}_\lambda(z, x)$  for all  $z \in \Omega$ , all  $x \geq 0$ ). Using (15) we will show that  $\{u_n^+\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(z)}(\Omega)$ . Arguing by contradiction, suppose that at least for a subsequence we have

$$\|u_n^+\| \rightarrow +\infty \text{ as } n \rightarrow +\infty, \|u_n^+\| \geq 1 \text{ for all } n \in \mathbb{N}. \tag{16}$$

We set  $y_n = \frac{u_n^+}{\|u_n^+\|}$ ,  $n \in \mathbb{N}$ . Then  $y_n \in W_0^{1,p(z)}(\Omega)$ ,  $y_n \geq 0$ ,  $\|y_n\| = 1$  for all  $n \in \mathbb{N}$ . So, we may assume that

$$y_n \xrightarrow{w} y \text{ in } W_0^{1,p(z)}(\Omega), y_n \rightarrow y \text{ in } L^{r(z)}(\Omega), y \geq 0 \text{ (see Proposition 1)}. \tag{17}$$

Suppose  $y \neq 0$ . We set  $\Omega_+ = \{z \in \Omega : y(z) > 0\}$ . From (17) we see that  $|\Omega_+|_N > 0$  (by  $|\cdot|_N$  we denote the Lebesgue measure on  $\mathbb{R}^N$ ). We have

$$u_n^+(z) \rightarrow +\infty \text{ for a.a. } z \in \Omega_+. \tag{18}$$

Then from (18), hypothesis  $H_1(ii)$  and since  $\tau_+ < p_+$ , we see that

$$\frac{\widehat{F}_\lambda(z, u_n^+(z))}{\|u_n^+\|^{p_+}} = \frac{\widehat{F}_\lambda(z, u_n^+(z))}{(u_n^+(z))^{p_+}} y_n(z)^{p_+} \rightarrow +\infty \text{ for a.a. } z \in \Omega_+.$$

Using Fatou’s lemma, we have

$$\lim_{n \rightarrow +\infty} \int_\Omega \frac{\widehat{F}_\lambda(z, u_n^+)}{\|u_n^+\|^{p_+}} dz = +\infty. \tag{19}$$

From (12), (9) and (8), we have

$$\begin{aligned} & -\frac{1}{q_-} [\rho_p(\nabla u_n^+) + \rho_q(\nabla u_n^+)] + \int_\Omega \widehat{F}_\lambda(z, u_n^+) dz \leq c_9 \\ & \text{for some } c_9 > 0, \text{ all } n \in \mathbb{N}, \\ \Rightarrow & \int_\Omega \widehat{F}_\lambda(z, u_n^+) dz \leq c_9 + \frac{1}{q_-} [\rho_p(\nabla u_n^+) + \rho_q(\nabla u_n^+)] \\ & \leq c_{10} [1 + \rho_p(\nabla u_n^+)] \text{ for some } c_{10} > 0, \\ & \leq c_{10} [1 + \|u_n^+\|^{p_+}] \text{ (see (16) and Proposition 2),} \\ \Rightarrow & \int_\Omega \frac{\widehat{F}_\lambda(z, u_n^+)}{\|u_n^+\|^{p_+}} dz \leq c_{10} \left[ \frac{1}{\|u_n^+\|^{p_+}} + 1 \right] \text{ for all } n \in \mathbb{N}. \end{aligned} \tag{20}$$

Comparing (20) and (19), we have a contradiction.

Next suppose that  $y = 0$ . Consider the function

$$\mu_n(t) = \varphi_\lambda(tu_n^+) \text{ for all } t \in [0, 1].$$

The function  $\mu_n(\cdot)$  is continuous and we can find  $t_n \in [0, 1]$  such that

$$\mu_n(t_n) = \max_{0 \leq t \leq 1} \mu_n(t). \tag{21}$$

Let  $\beta > 1$  and set  $v_n = (2\beta)^{1/p_-} y_n, n \in \mathbb{N}$ . From (17) and since we assume that  $y = 0$ , we have

$$\begin{aligned} v_n &\rightarrow 0 \text{ in } L^{r(z)}(\Omega), \\ &\Rightarrow \int_{\Omega} G_{\lambda}(z, v_n) dz \rightarrow 0 \text{ as } n \rightarrow +\infty. \end{aligned} \tag{22}$$

From (16) we see that we can find  $n_0 \in \mathbb{N}$  such that

$$\frac{(2\beta)^{1/p_-}}{\|u_n^+\|} \in (0, 1] \text{ for all } n \geq n_0. \tag{23}$$

Then from (21) and (23) we see that

$$\begin{aligned} \varphi_{\lambda}(t_n u_n^+) &\geq \varphi_{\lambda}((2\beta)^{1/p_-} y_n) = \varphi_{\lambda}(v_n) \text{ for all } n \geq n_0, \\ \Rightarrow \varphi_{\lambda}(t_n u_n^+) &\geq \frac{1}{p_+} \int_{\Omega} (2\beta)^{\frac{p(z)}{p_-}} |\nabla y_n|^{p(z)} dz - \int_{\Omega} G_{\lambda}(z, v_n) dz \\ &\geq \frac{2\beta}{p_+} \rho_p(\nabla y_n) - \int_{\Omega} G_{\lambda}(z, v_n) dz \text{ (recall } \beta > 1) \\ &= \frac{2\beta}{p_+} - \int_{\Omega} G_{\lambda}(z, v_n) dz \text{ for all } n \geq n_0 \\ &\text{(since } \|y_n\| = 1, \text{ see Proposition 2).} \end{aligned}$$

From (22) we see that there exists  $n_1 \in \mathbb{N}, n_1 \geq n_0$ , such that

$$\varphi_{\lambda}(t_n u_n^+) \geq \frac{\beta}{p_+} \text{ for all } n \geq n_1.$$

But  $\beta > 1$  is arbitrary. So, we infer that

$$\varphi_{\lambda}(t_n u_n^+) \rightarrow +\infty \text{ as } n \rightarrow +\infty. \tag{24}$$

We have

$$\begin{aligned} 0 &\leq t_n u_n^+ \leq u_n^+ \text{ for all } n \in \mathbb{N}, \\ \Rightarrow \int_{\Omega} e_{\lambda}(z, t_n u_n^+) dz &\leq \int_{\Omega} e_{\lambda}(z, u_n^+) dz + c_{11} \\ &\text{for some } c_{11} > 0, \text{ all } n \in \mathbb{N} \text{ (see hypothesis } H_1(ii)), \\ \Rightarrow \int_{\Omega} e_{\lambda}(z, t_n u_n^+) dz &\leq c_{12} \text{ for some } c_{12} > 0, \text{ all } n \in \mathbb{N} \text{ (see (15)).} \end{aligned} \tag{25}$$

We set

$$\widehat{e}_\lambda(z, x) = g_\lambda(z, x)x - p_+G_\lambda(z, x) \text{ for all } z \in \Omega, \text{ all } x \geq 0.$$

Then from (8), (7) and hypothesis  $H_1(iv)$ , we see that

$$\widehat{e}_\lambda(z, x) \leq e_\lambda(z, x) + c_{13} \text{ for some } c_{13} > 0, \text{ a.a. } z \in \Omega, \text{ all } x \geq 0. \tag{26}$$

Note that

$$\varphi_\lambda(0) = 0, \varphi_\lambda(u_n^+) \leq c_{14} \text{ for some } c_{14} > 0, \text{ all } n \in \mathbb{N} \text{ (see (9), (12))}. \tag{27}$$

From (24) and (27) it follows that there exists  $n_2 \in \mathbb{N}$  such that

$$t_n \in (0, 1) \text{ for all } n \geq n_2. \tag{28}$$

Then (28) and (21) imply that for all  $n \geq n_2$ , we have

$$\begin{aligned} & \frac{d}{dt}\mu_n(t)\Big|_{t=t_n} = 0, \\ \Rightarrow & \langle \varphi'_\lambda(t_n u_n^+), u_n^+ \rangle = 0 \text{ (by the chain rule),} \\ \Rightarrow & \langle \varphi'_\lambda(t_n u_n^+), t_n u_n^+ \rangle = 0 \text{ for all } n \geq n_2 \text{ (see (28))}. \end{aligned} \tag{29}$$

For  $n \geq n_2$  we have

$$\begin{aligned} \varphi_\lambda(t_n u_n^+) &= \varphi_\lambda(t_n u_n^+) - \frac{1}{p_+} \langle \varphi'_\lambda(t_n u_n^+), t_n u_n^+ \rangle \text{ (see (29))}, \\ \Rightarrow \varphi_\lambda(t_n u_n^+) &\leq \int_\Omega \left[ \frac{1}{p(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{p(z)} dz + \int_\Omega \left[ \frac{1}{q(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{q(z)} dz \\ &+ \frac{1}{p_+} \int_\Omega \widehat{e}_\lambda(z, t_n u_n^+) dz \text{ (see (28))} \\ &\leq \int_\Omega \left[ \frac{1}{p(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{p(z)} dz + \int_\Omega \left[ \frac{1}{q(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{q(z)} dz \\ &+ \frac{1}{p_+} \int_\Omega e_\lambda(z, t_n u_n^+) dz + c_{15} \text{ for some } c_{15} > 0 \text{ (see (26))} \\ &\leq \int_\Omega \left[ \frac{1}{p(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{p(z)} dz + \int_\Omega \left[ \frac{1}{q(z)} - \frac{1}{p_+} \right] |\nabla u_n^+|^{q(z)} dz \\ &+ c_{16} \text{ for some } c_{16} > 0 \text{ (see (25))} \\ &\leq \varphi_\lambda(u_n^+) - \frac{1}{p_+} \langle \varphi'_\lambda(u_n^+), u_n^+ \rangle + c_{17} \text{ for some } c_{17} > 0 \text{ (see (26), (15))} \\ &\leq c_{18} \text{ for some } c_{18} > 0 \text{ (see (9), (10), (12))}. \end{aligned} \tag{30}$$

We compare (24) and (30) and reach a contradiction. Therefore  $\{u_n^+\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(z)}(\Omega)$  is bounded and this combined with (12) implies that  $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(z)}(\Omega)$  is bounded. We may assume that

$$u_n \xrightarrow{w} u \text{ in } W_0^{1,p(z)}(\Omega), \quad u_n \rightarrow u \text{ in } L^{r(z)}(\Omega). \tag{31}$$

In (11) we use the test function  $h = (u_n - u) \in W_0^{1,p(z)}(\Omega)$ , pass to the limit as  $n \rightarrow +\infty$  and use (31). We obtain

$$\begin{aligned} \lim_{n \rightarrow +\infty} \langle V(u_n), u_n - u \rangle &= 0, \\ \Rightarrow u_n &\rightarrow u \text{ in } W_0^{1,p(z)}(\Omega) \text{ (see Proposition 4),} \\ \Rightarrow \varphi_\lambda(\cdot) &\text{ satisfies the } C\text{-condition} \end{aligned}$$

This proves the Claim.

On account of hypotheses  $H_1(i), (iv)$ , given  $\varepsilon > 0$ , we can find  $c_{19} = c_{19}(\varepsilon) > 0$  such that

$$F(z, x) \leq \frac{\varepsilon}{p_+} x^{q_+} + c_{19} x^{r(z)} \text{ for a.a. } z \in \Omega, \text{ all } x \geq 0. \tag{32}$$

Consider  $u \in W_0^{1,p(z)}(\Omega)$  with  $\|u\| \leq 1$  small. We have

$$\begin{aligned} \varphi_\lambda(u) &\geq \frac{1}{p_+} [\rho_p(\nabla u) + \rho_q(\nabla u)] - \int_{\{u \leq \bar{u}_\lambda\}} \lambda [\bar{u}_\lambda^{-\eta(z)} + \bar{u}_\lambda^{\tau(z)-1}] u^+ dz \\ &\quad - \frac{\lambda}{1 - \eta_+} \int_{\{\bar{u}_\lambda < u\}} [u^{1-\eta(z)} - \bar{u}_\lambda^{1-\eta(z)}] dz - \frac{\lambda}{\tau_-} \int_{\{\bar{u}_\lambda < u\}} [u^{\tau(z)} - \bar{u}_\lambda^{\tau(z)}] dz \\ &\quad - \int_\Omega F(z, u^+) dz \text{ (see (8)).} \end{aligned} \tag{33}$$

Let  $\widehat{d}(z) = d(z, \partial\Omega)$  for all  $z \in \overline{\Omega}$ . Then Lemma 14.16, p. 355, of Gilbarg–Trudinger [8] implies that  $\widehat{d} \in C_+ \setminus \{0\}$ . Since  $\bar{u}_\lambda \in \text{int } C_+$  (see Proposition 5), using Proposition 4.1.22, p. 274, of [18], we can find  $c_{20} > 0$  such that

$$c_{20} \widehat{d} \leq \bar{u}_\lambda. \tag{34}$$

Using the anisotropic Hardy’s inequality of Harjulehto–Hästö–Koskenoja [9], we have

$$\begin{aligned} \int_\Omega \left( \frac{|u|}{\bar{u}_\lambda^{\eta(z)}} \right)^{p(z)} dz &= \int_\Omega \left( \frac{|u|}{\bar{u}_\lambda^{1-\eta(z)}} \right)^{p(z)} \left( \frac{|u|}{\bar{u}_\lambda} \right)^{p(z)} dz \\ &\leq c_{21} \int_\Omega \left( \frac{|u|}{\bar{u}_\lambda} \right)^{p(z)} dz \text{ for some } c_{21} > 0 \text{ (since } \bar{u}_\lambda \in \text{int } C_+) \end{aligned}$$

$$\begin{aligned}
 &\leq c_{22} \int_{\Omega} \left( \frac{|u|}{\widehat{d}} \right)^{p(z)} dz \text{ for some } c_{22} > 0 \text{ (see (34))} \\
 &\leq c_{22} \left\| \frac{u}{\widehat{d}} \right\|_{p(z)} \text{ for } \|u\| \leq 1 \text{ small (see Proposition 2 and [9])} \\
 &\leq c_{23} \|u\| \text{ for some } c_{23} > 0.
 \end{aligned}
 \tag{35}$$

Also we have

$$\begin{aligned}
 \frac{\lambda}{1 - \eta_+} \int_{\{\bar{u}_\lambda < u\}} u^{1-\eta(z)} dz &\leq \frac{\lambda}{1 - \eta_+} \int_{\{\bar{u}_\lambda < u\}} \frac{u}{\bar{u}_\lambda^{\eta(z)}} dz \\
 &\leq \lambda c_{24} \int_{\Omega} \frac{|u|}{\widehat{d}} dz \text{ for some } c_{24} > 0 \\
 &\leq \lambda c_{25} \left\| \frac{|u|}{\widehat{d}} \right\|_{p(z)} \text{ for some } c_{25} > 0 \\
 &\text{(since } L^{p(z)}(\Omega) \hookrightarrow L^1(\Omega) \text{ continuously)} \\
 &\leq \lambda c_{26} \|u\| \text{ for some } c_{26} > 0 \\
 &\text{(anisotropic Hardy's inequality, see [9]),}
 \end{aligned}
 \tag{36}$$

and

$$\frac{\lambda}{\tau_-} \int_{\{\bar{u}_\lambda < u\}} |u|^{\tau(z)} dz \leq \frac{\lambda}{\tau_-} \rho_\tau(u) \leq \lambda c_{27} \|u\| \text{ for some } c_{27} > 0 (\|u\| \leq 1 \text{ small}).
 \tag{37}$$

We return to (33) and use (35), (36), (37) and (32). We obtain

$$\begin{aligned}
 \varphi_\lambda(u) &\geq \frac{1}{p_+} \|u\|^{p_+} + \frac{1}{p_+} \left[ \|u\|_{1,q(z)}^{q_+} - \varepsilon c_{28} \|u\|_{1,q(z)}^{q_+} \right] - c_{29} [\lambda \|u\| + \|u\|^{r_-}] \\
 &\text{for some } c_{28}, c_{29} > 0 \text{ (recall } \|u\| \leq 1 \text{ is small).}
 \end{aligned}$$

Choosing  $\varepsilon > 0$  small, we have

$$\varphi_\lambda(u) \geq \left[ \frac{1}{p_+} - c_{29}(\lambda \|u\|^{1-p_+} + \|u\|^{r_- - p_+}) \right] \|u\|^{p_+}.$$

Consider the function

$$\widehat{\gamma}_\lambda(t) = \lambda t^{1-p_+} + t^{r_- - p_+}, \quad t \geq 0.$$

Evidently  $\widehat{\gamma}_\lambda \in C^1(0, \infty)$  and

$$\widehat{\gamma}_\lambda(t) \rightarrow +\infty \text{ as } t \rightarrow 0^+ \text{ and as } t \rightarrow +\infty \text{ (since } 1 < p_+ < r_-).$$



Therefore we can find  $t_0 \in (0, 1)$  such that

$$\begin{aligned} \widehat{\gamma}_\lambda(t_0) &= \min_{t>0} \widehat{\gamma}_\lambda, \\ \Rightarrow \widehat{\gamma}'_\lambda(t_0) &= 0, \\ \Rightarrow \lambda(p_+ - 1)t_0^{-p_+} &= (r_- - p_+)t_0^{r_- - p_+ - 1}, \\ \Rightarrow t_0 = t_0(\lambda) &= \left[ \frac{\lambda(p_+ - 1)}{r_- - p_+} \right]^{\frac{1}{r_- - 1}}. \end{aligned}$$

We have

$$\widehat{\gamma}_\lambda(t_0) = \lambda \left[ \frac{r_- - p_+}{\lambda(p_+ - 1)} \right]^{\frac{p_+ - 1}{r_- - 1}} + \left[ \frac{\lambda(p_+ - 1)}{r_- - p_+} \right]^{\frac{r_- - p_+}{r_- - 1}}.$$

Since  $p_+ < r_-$ , we see that

$$\widehat{\gamma}_\lambda(t_0) \rightarrow 0 \text{ as } \lambda \rightarrow 0^+.$$

Therefore we can find  $\widehat{\lambda}_0 > 0$  such that

$$\begin{aligned} \frac{1}{p_+} - c_{29}\widehat{\gamma}_\lambda(t_0) &\geq \beta_\lambda > 0 \text{ for all } \lambda \in (0, \widehat{\lambda}_0), \\ \Rightarrow \varphi_\lambda(u) &\geq \beta_\lambda > 0 \text{ for all } \lambda \in (0, \widehat{\lambda}_0), \text{ all } \|u\| = t_0(\lambda). \end{aligned} \tag{38}$$

Let  $\overline{B}_\lambda = \{u \in W_0^{1,p(\cdot)}(\Omega) : \|u\| \leq t_0(\lambda)\}$ . The reflexivity of  $W_0^{1,p(\cdot)}(\Omega)$  and the Eberlein–Šmulian theorem imply that  $\overline{B}_\lambda$  is sequentially weakly compact. Also, the sequential weak lower semicontinuity of the modular function and Proposition 1 imply that  $\varphi_\lambda(\cdot)$  is sequentially weakly lower semicontinuous. So, we can find  $u_\lambda \in \overline{B}_\lambda$  such that

$$\varphi_\lambda(u_\lambda) = \inf [\varphi_\lambda(u) : u \in \overline{B}_\lambda]. \tag{39}$$

Recall that  $\overline{u}_\lambda \in \text{int } C_+$ . So, if  $u \in C_+ \setminus \{0\}$ , we can find  $t \in (0, 1)$  small such that

$$0 \leq tu \leq \overline{u}_\lambda, \quad 0 \leq tu(z) \leq \delta \text{ for all } z \in \overline{\Omega} \text{ (see [12], p. 274)}.$$

Using (8) and hypothesis  $H_1(iv)$ , we have

$$\begin{aligned} \varphi_\lambda(tu) &\leq \frac{t^{q_-}}{q_-} [\rho_p(\nabla u) + \rho_q(\nabla u)] - \lambda t \int_\Omega [\overline{u}_\lambda^{-\eta(z)} + u^{\tau(z)-1}] u dz \\ &\leq c_{30}t^{q_-} - \lambda c_{31}t \text{ for some } c_{30}, c_{31} > 0 \text{ (recall } \frac{u}{\overline{u}_\lambda^{\eta(\cdot)}} \in L^1(\Omega)). \end{aligned}$$

Since  $1 < q_-$ , choosing  $t \in (0, 1)$  even smaller if necessary, we have

$$\begin{aligned} \varphi_\lambda(tu) &< 0, \\ \Rightarrow \varphi_\lambda(u_\lambda) &< 0 = \varphi_\lambda(0) \quad (\text{see (39)}), \\ \Rightarrow u_\lambda &\neq 0. \end{aligned}$$

Then from (38) we see that

$$\begin{aligned} 0 &< \|u_\lambda\| < t_0(\lambda), \\ \Rightarrow \langle \varphi'_\lambda(u_\lambda), h \rangle &= 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega), \\ \Rightarrow \langle V(u_\lambda), h \rangle &= \int_\Omega g_\lambda(z, u_\lambda) h dz \text{ for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned} \tag{40}$$

We use the test function  $h = (\bar{u}_\lambda - u_\lambda)^+ \in W_0^{1,p(z)}(\Omega)$ . We have

$$\begin{aligned} &\langle V(u_\lambda), (\bar{u}_\lambda - u_\lambda)^+ \rangle \\ &= \int_\Omega (\lambda[\bar{u}_\lambda^{-\eta(z)} + \bar{u}_\lambda^{\tau(z)-1}] + f(z, u_\lambda^+))(\bar{u}_\lambda - u_\lambda)^+ dz \text{ (see (8))} \\ &\geq \int_\Omega \lambda \bar{u}_\lambda^{\tau(z)-1} (\bar{u}_\lambda - u_\lambda)^+ dz \text{ (see (7) and hypothesis } H_1(iv)) \\ &= \langle V(\bar{u}_\lambda), (\bar{u}_\lambda - u_\lambda)^+ \rangle \text{ (see Proposition 5),} \\ \Rightarrow \bar{u}_\lambda &\leq u_\lambda \text{ (see Proposition 4).} \end{aligned} \tag{41}$$

From (41), (8) and (40), we infer that

$$u_\lambda \text{ is a positive solution of } (P_\lambda).$$

From Proposition A1 of Papageorgiou–Rădulescu–Zhang [19], we know that  $u_\lambda \in L^\infty(\Omega)$ . Then the singular anisotropic regularity theory (see Saoudi–Ghanmi [21] and Giacomoni–Kumar–Sreenadh [7] for the corresponding isotropic theory) implies that  $u_\lambda \in C_+ \setminus \{0\}$ . Let  $\rho = \|u_\lambda\|_\infty$  and let  $\widehat{\xi}_\rho > 0$  be as postulated by hypothesis  $H_1(iv)$ . We have

$$\begin{aligned} &-\Delta_{p(z)}u_\lambda - \Delta_{q(z)}u_\lambda + \widehat{\xi}_\rho u_\lambda^{p(z)-1} - \lambda u_\lambda^{-\eta(z)} \\ &= \lambda u_\lambda^{\tau(z)-1} + f(z, u_\lambda) + \widehat{\xi}_\rho u_\lambda^{p(z)-1} \\ &\geq 0 \text{ in } \Omega, \\ \Rightarrow u_\lambda &\in \text{int } C_+ \text{ (see Proposition A2 of [13]).} \end{aligned}$$

We conclude that

$$\mathcal{L} \neq \emptyset \text{ and } S_\lambda \subseteq \text{int } C_+ \text{ for all } \lambda > 0.$$

□

The next proposition establishes a structural property of the set  $\mathcal{L}$ , namely that it is connected.

**Proposition 7** *If hypotheses  $H_0, H_1$  hold,  $\lambda \in \mathcal{L}$  and  $0 < \mu < \lambda$ , then  $\mu \in \mathcal{L}$ .*

**Proof** Let  $u_\lambda \in S_\lambda \subseteq \text{int } C_+$ . From Proposition 5 we know that  $\bar{u}_\sigma \rightarrow 0$  in  $C_0^1(\bar{\Omega})$  as  $\sigma \rightarrow 0^+$ . So, we can find  $\sigma \in (0, \mu)$  small such that

$$\bar{u}_\sigma \leq \min\{\delta, u_\lambda\} \quad (\text{recall } u_\lambda \in \text{int } C_+).$$

We introduce the Carathéodory function  $\widehat{g}_\mu(z, x)$  defined by

$$\widehat{g}_\mu(z, x) = \begin{cases} \mu[\bar{u}_\sigma(z)^{-\eta(z)} + \bar{u}_\sigma(z)^{\tau(z)-1}] + f(z, \bar{u}_\sigma(z)) & \text{if } x < \bar{u}_\sigma(z), \\ \mu[x^{-\eta(z)} + x^{\tau(z)-1}] + f(z, x) & \text{if } \bar{u}_\sigma(z) \leq x \leq u_\lambda(z), \\ \mu[u_\lambda(z)^{-\eta(z)} + u_\lambda(z)^{\tau(z)-1}] + f(z, u_\lambda(z)) & \text{if } u_\lambda(z) < x. \end{cases} \tag{42}$$

We set  $\widehat{G}_\mu(z, x) = \int_0^x \widehat{g}_\mu(z, s) ds$  and consider the  $C^1$ -functional  $\widehat{\psi}_\mu : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\begin{aligned} \widehat{\psi}_\mu(u) &= \int_\Omega \frac{1}{p(z)} |\nabla u|^{p(z)} dz + \int_\Omega \frac{1}{q(z)} |\nabla u|^{q(z)} dz \\ &\quad - \int_\Omega \widehat{G}_\mu(z, u) dz \text{ for all } u \in W_0^{1,p(z)}(\Omega). \end{aligned}$$

From (42) and Proposition 2, it is clear that  $\widehat{\psi}_\mu(\cdot)$  is coercive. Also, it is sequentially weakly lower semicontinuous (see Proposition 1). Then by the Weierstrass-Tonelli theorem, we can find  $u_\mu \in W_0^{1,p(z)}(\Omega)$  such that

$$\begin{aligned} \widehat{\psi}_\mu(u_\mu) &= \inf \left[ \widehat{\psi}_\mu(u) : u \in W_0^{1,p(z)}(\Omega) \right], \\ \Rightarrow \langle \widehat{\psi}'_\mu(u_\mu), h \rangle &= 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned} \tag{43}$$

In (43) first we use the test function  $h = (\bar{u}_\sigma - u_\mu)^+ \in W_0^{1,p(z)}(\Omega)$ . We have

$$\begin{aligned} &\langle V(u_\mu), (\bar{u}_\sigma - u_\mu)^+ \rangle \\ &= \int_\Omega (\mu[\bar{u}_\sigma^{-\eta(z)} + \bar{u}_\sigma^{\tau(z)-1}] + f(z, \bar{u}_\sigma)) (\bar{u}_\sigma - u_\mu)^+ dz \text{ (see (42))} \\ &\geq \int_\Omega \mu \bar{u}_\sigma^{\tau(z)-1} (\bar{u}_\sigma - u_\mu)^+ dz \text{ (because } 0 \leq \bar{u}_\sigma(z) \leq \delta, z \in \bar{\Omega}, \text{ use } H_1(iv))} \\ &\geq \int_\Omega \sigma \bar{u}_\sigma^{\tau(z)-1} (\bar{u}_\sigma - u_\mu)^+ dz \text{ (since } \sigma < \mu) \\ &= \langle V(\bar{u}_\sigma), (\bar{u}_\sigma - u_\mu)^+ \rangle \text{ (see Proposition 5),} \end{aligned}$$

$$\Rightarrow \bar{u}_\sigma \leq u_\mu.$$

Next in (43) we choose the test function  $h = (u_\mu - u_\lambda)^+ \in W_0^{1,p(z)}(\Omega)$ . We have

$$\begin{aligned} & \langle V(u_\mu), (u_\mu - u_\lambda)^+ \rangle \\ &= \int_\Omega (\mu[u_\lambda^{-\eta(z)} + u_\lambda^{\tau(z)-1}] + f(z, u_\lambda))(u_\mu - u_\lambda)^+ dz \text{ (see (42))} \\ &\leq \int_\Omega (\lambda[u_\lambda^{-\eta(z)} + u_\lambda^{\tau(z)-1}] + f(z, u_\lambda))(u_\mu - u_\lambda)^+ dz \text{ (since } \mu < \lambda) \\ &= \langle V(u_\lambda), (u_\mu - u_\lambda)^+ \rangle \text{ (since } u_\lambda \in S_\lambda), \\ \Rightarrow u_\mu &\leq u_\lambda \text{ (see Proposition 4).} \end{aligned}$$

So, we have proved that

$$u_\mu \in [\bar{u}_\sigma, u_\lambda]. \tag{44}$$

Then from (44), (42) and (43) it follows that

$$u_\mu \in S_\mu \subseteq \text{int } C_+ \text{ and so } \mu \in \mathcal{L}.$$

□

A quick inspection of the above proof reveals that we get, as a useful byproduct of it, the following corollary.

**Corollary 1** *If hypotheses  $H_0, H_1$  hold,  $\lambda \in \mathcal{L}, u_\lambda \in S_\lambda \subseteq \text{int } C_+$  and  $\mu \in (0, \lambda)$ , then  $\mu \in \mathcal{L}$  and there exists  $u_\mu \in S_\mu \subseteq \text{int } C_+$  such that  $u_\mu \leq u_\lambda$ .*

In fact with little additional effort, we can improve the above ‘‘monotonicity’’ property of the solution multifunction  $\lambda \rightarrow S_\lambda$ .

**Proposition 8** *If hypotheses  $H_0, H_1$  hold,  $\lambda \in \mathcal{L}, u_\lambda \in S_\lambda \subseteq \text{int } C_+$  and  $\mu \in (0, \lambda)$ , then  $\mu \in \mathcal{L}$  and there exists  $u_\mu \in S_\mu \subseteq \text{int } C_+$  such that  $u_\lambda - u_\mu \in \text{int } C_+$ .*

**Proof** From Corollary 1 we already know that  $\mu \in \mathcal{L}$  and there exists  $u_\mu \in S_\mu \subseteq \text{int } C_+$  such that

$$u_\mu \leq u_\lambda.$$

Let  $\rho = \|u_\lambda\|_\infty$  and let  $\widehat{\xi}_\rho > 0$  be as postulated by hypothesis  $H_1(iv)$ . We have

$$\begin{aligned} & -\Delta_{p(z)}u_\mu - \Delta_{q(z)}u_\mu + \widehat{\xi}_\rho u_\mu^{p(z)-1} - \lambda u_\mu^{-\eta(z)} \\ & \leq \lambda u_\lambda^{\tau(z)-1} + f(z, u_\lambda) + \widehat{\xi}_\rho u_\lambda^{p(z)-1} \text{ (see hypothesis } H_1(iv)) \\ & = -\Delta_{p(z)}u_\lambda - \Delta_{q(z)}u_\lambda + \widehat{\xi}_\rho u_\lambda^{p(z)-1} - \lambda u_\lambda^{-\eta(z)}. \end{aligned} \tag{45}$$

Since  $u_\mu \in \text{int } C_+$ , we see that  $0 < (\lambda - \mu)u_\mu^{\tau(z)-1}$ . Hence from (45) and Proposition 2.3 of Papageorgiou-Winkert [13], we obtain

$$u_\lambda - u_\mu \in \text{int } C_+.$$

□

From the proof of Proposition 7, we know that for  $\sigma \in (0, \mu)$  small, we have

$$u_\mu \in [\bar{u}_\sigma, u_\lambda] \quad (\text{see (44)}).$$

In fact using Proposition 8, we can improve this.

**Proposition 9** *If hypotheses  $H_0, H_1$  hold,  $\lambda \in \mathcal{L}, u_\lambda \in S_\lambda \subseteq \text{int } C_+$  and  $\mu \in (0, \lambda)$ , then we can find  $u_\mu \in S_\mu \subseteq \text{int } C_+$  and  $\sigma \in (0, \mu)$  small such that  $u_\mu \in \text{int}_{C_0^1(\bar{\Omega})}[\bar{u}_\sigma, u_\lambda]$ .*

**Proof** From Proposition 8, we already know that there exists  $u_\mu \in S_\mu \subseteq \text{int } C_+$  such that

$$u_\lambda - u_\mu \in \text{int } C_+. \tag{46}$$

Also if  $\sigma \in (0, \mu)$  is small, we have  $\bar{u}_\sigma \leq \min\{\delta, u_\mu\}$  (see Proposition 5). Let  $\rho = \|u_\mu\|_\infty$  and let  $\hat{\xi}_\rho > 0$  be as postulated by hypothesis  $H_1(iv)$ . We have

$$\begin{aligned} & -\Delta_{p(z)}\bar{u}_\sigma - \Delta_{q(z)}\bar{u}_\sigma + \hat{\xi}_\rho\bar{u}_\sigma^{p(z)-1} - \mu\bar{u}_\sigma^{-\eta(z)} \\ & \leq \sigma\bar{u}_\sigma^{\tau(z)-1} + \hat{\xi}_\rho\bar{u}_\sigma^{p(z)-1} + f(z, \bar{u}_\sigma) \quad (\text{since } \bar{u}_\sigma \leq \delta, \text{ see } H_1(iv)) \\ & \leq \mu u_\mu^{\tau(z)-1} + \hat{\xi}_\rho u_\mu^{p(z)-1} + f(z, u_\mu) \quad (\text{see } H_1(iv)) \\ & = -\Delta_{p(z)}u_\mu - \Delta_{q(z)}u_\mu + \hat{\xi}_\rho u_\mu^{p(z)-1} - \mu u_\mu^{-\eta(z)}. \end{aligned} \tag{47}$$

Since  $\bar{u}_\sigma \in \text{int } C_+$ , on account of hypothesis  $H_1(iv)$ , we have

$$0 < f(\cdot, \bar{u}_\sigma(\cdot)).$$

So, from (47) and Proposition 2.3 of Papageorgiou-Winkert [13], we infer that

$$u_\mu - \bar{u}_\sigma \in \text{int } C_+. \tag{48}$$

Then (46) and (48) imply that

$$u_\mu \in \text{int}_{C_0^1(\bar{\Omega})}[\bar{u}_\sigma, u_\lambda].$$

□

Let  $\hat{\lambda} = \sup \mathcal{L}$ .

**Proposition 10** *If hypotheses  $H_0, H_1$  hold, then  $\widehat{\lambda} < +\infty$ .*

**Proof** Hypotheses  $H_1(i), (ii), (iv)$  imply that we can find  $\lambda_0 > 0$  such that

$$\lambda_0 x^{\tau(z)-1} + f(z, x) \geq x^{p(z)-1} \text{ for a.a. } z \in \Omega, \text{ all } x \geq 0. \tag{49}$$

Let  $\lambda > \lambda_0$  and suppose that  $\lambda \in \mathcal{L}$ . Then we can find  $u_\lambda \in S_\lambda \subseteq \text{int } C_+$  (see Proposition 6). Let  $\Omega_0 \subseteq \Omega$  be an open subset with  $C^2$ -boundary  $\partial\Omega_0$  and such that  $\overline{\Omega_0} \subseteq \Omega$ . We define

$$0 < m_0 = \min_{\Omega_0} u_\lambda \quad (\text{since } u_\lambda \in \text{int } C_+).$$

For  $\varepsilon > 0$ , let  $m_0^\varepsilon = m_0 + \varepsilon$ . Also, let  $\rho = \max\{\|u_\lambda\|_\infty, m_0^\varepsilon\}$  and take  $\widehat{\xi}_\rho > 0$  as postulated by hypothesis  $H_1(iv)$ . We have

$$\begin{aligned} & -\Delta_{p(z)} m_0^\varepsilon - \Delta_{q(z)} m_0^\varepsilon + \widehat{\xi}_\rho (m_0^\varepsilon)^{p(z)-1} - \lambda (m_0^\varepsilon)^{-\eta(z)} \\ & \leq \widehat{\xi}_\rho (m_0)^{p(z)-1} + \chi(\varepsilon) \text{ with } \chi(\varepsilon) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0^+ \\ & \leq [\widehat{\xi}_\rho + 1] m_0^{p(z)-1} + \chi(\varepsilon) \\ & \leq \lambda_0 m_0^{\tau(z)-1} + f(z, m_0) + \widehat{\xi}_\rho m_0^{p(z)-1} + (\lambda - \lambda_0) m_0^{\tau(z)-1} + \chi(\varepsilon) \text{ (see (49))} \\ & \leq \lambda u_\lambda^{\tau(z)-1} + f(z, u_\lambda) + \widehat{\xi}_\rho u_\lambda^{p(z)-1} \text{ (see } H_1(iv)) \\ & = -\Delta_{p(z)} u_\lambda - \Delta_{q(z)} u_\lambda + \widehat{\xi}_\rho u_\lambda^{p(z)-1} - \lambda u_\lambda^{-\eta(z)} \text{ in } \Omega_0. \end{aligned} \tag{50}$$

For  $\varepsilon > 0$  small, we have

$$0 < \widehat{c} \leq (\lambda - \lambda_0) m_0^{\tau(z)-1} - \chi(\varepsilon).$$

So, from (50) and Proposition 2.3 of [13] (see also Proposition A4 of [19]), we obtain

$$m_0^\varepsilon < u_\lambda(z) \text{ for all } z \in \Omega_0,$$

a contradiction. Therefore  $\widehat{\lambda} \leq \lambda_0 < +\infty$ . □

If  $\lambda \in (0, \widehat{\lambda})$ , then we have multiplicity of positive solutions.

**Proposition 11** *If hypotheses  $H_0, H_1$  hold and  $\lambda \in (0, \widehat{\lambda})$ , then problem  $(P_\lambda)$  has at least two positive solutions  $u_0, \widehat{u} \in \text{int } C_+$ .*

**Proof** Let  $\beta \in (\lambda, \widehat{\lambda})$  and  $\sigma \in (0, \lambda)$  small such that  $\|\overline{u}_\sigma\|_\infty \leq \delta$  (see Proposition 5). From the previous results, we know that for  $u_\beta \in S_\beta \subseteq \text{int } C_+$ , we can find  $u_0 \in W_0^{1,p(z)}(\Omega)$  such that

$$u_0 \in S_\lambda \subseteq \text{int } C_+, \quad u_0 \in \text{int}_{C_0^1(\overline{\Omega})} [\overline{u}_\sigma, u_\beta]. \tag{51}$$

As in the proof of Proposition 7, truncating the reaction at  $\{\bar{u}_\sigma(z), u_\beta(z)\}$  (see (42)) and introducing the corresponding  $C^1$ -energy functional  $\widehat{\psi}_\lambda(\cdot)$ , via the direct method of the Calculus of Variations, we produce  $u_0$  a global minimizer of  $\widehat{\psi}_\lambda(\cdot)$ .

Also, we introduce the following Carathéodory function

$$\widehat{e}_\lambda(z, x) = \begin{cases} \lambda[\bar{u}_\sigma(z)^{-\eta(z)} + \bar{u}_\sigma(z)^{\tau(z)-1}] + f(z, \bar{u}_\sigma(z)) & \text{if } x \leq \bar{u}_\sigma(z), \\ \lambda[x^{-\eta(z)} + x^{\tau(z)-1}] + f(z, x) & \text{if } \bar{u}_\sigma(z) < x. \end{cases} \tag{52}$$

We set  $\widehat{E}_\lambda(z, x) = \int_0^x \widehat{e}_\lambda(z, s) ds$  and introduce the  $C^1$ -functional  $\widehat{\varphi}_\lambda : W_0^{1,p(z)}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\begin{aligned} \widehat{\varphi}_\lambda(u) &= \int_\Omega \frac{1}{p(z)} |\nabla u|^{p(z)} dz + \int_\Omega \frac{1}{q(z)} |\nabla u|^{q(z)} dz \\ &\quad - \int_\Omega \widehat{E}_\lambda(z, u) dz \text{ for all } u \in W_0^{1,p(z)}(\Omega). \end{aligned}$$

From (42) and (52), we see that

$$\widehat{\psi}_\lambda \Big|_{[\bar{u}_\sigma, u_\beta]} = \widehat{\varphi}_\lambda \Big|_{[\bar{u}_\sigma, u_\beta]}.$$

Recall that  $u_0 \in \text{int } C_+$  is a global minimizer of  $\widehat{\psi}_\lambda(\cdot)$ . Then from (51) it follows that

$$\begin{aligned} &u_0 \text{ is a local } C_0^1(\overline{\Omega}) - \text{minimizer of } \widehat{\varphi}_\lambda(\cdot), \\ \Rightarrow &u_0 \text{ is a local } W_0^{1,p(z)}(\Omega) - \text{minimizer of } \widehat{\varphi}_\lambda(\cdot) \\ &\text{(see [13], Proposition A3).} \end{aligned} \tag{53}$$

Using (52) we can easily check that

$$K_{\widehat{\varphi}_\lambda} \subseteq [\bar{u}_\sigma] \cap \text{int } C_+. \tag{54}$$

Then (54) and (52) imply that we may assume that  $K_{\widehat{\varphi}_\lambda}$  is finite or otherwise we already have an infinity of positive smooth solutions of  $(P_\lambda)$  and so we are done. So, we have that  $K_{\widehat{\varphi}_\lambda}$  is finite and this fact together with (53) and Theorem 5.7.6, p. 449, of Papageorgiou–Rădulescu–Repovš [18] imply that we can find  $\rho \in (0, 1)$  small such that

$$\widehat{\varphi}_\lambda(u_0) < \inf [\widehat{\varphi}_\lambda(u) : \|u - u_0\| = \rho] = \widehat{m}_\lambda. \tag{55}$$

On account of hypothesis  $H_1(ii)$ , we see that if  $u \in \text{int } C_+$  then

$$\widehat{\varphi}_\lambda(tu) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{56}$$

Moreover, using (52) and arguing as in the proof of Proposition 7 (see the ‘‘Claim’’), we show that

$$\widehat{\varphi}_\lambda(\cdot) \text{ satisfies the } C\text{-condition.} \tag{57}$$

Then (55), (56) and (57) permit the use of the mountain pass theorem. We can find  $\widehat{u} \in W_0^{1,p(z)}(\Omega)$  such that

$$\begin{aligned} \widehat{u} &\in K_{\widehat{\varphi}_\lambda} \subseteq [\bar{u}_\sigma] \cap \text{int } C_+ \text{ (see (54)),} \\ \widehat{\varphi}_\lambda(u_0) &< m_\lambda \leq \widehat{\varphi}_\lambda(\widehat{u}) \text{ (see (55)).} \end{aligned}$$

So,  $\widehat{u} \neq u_0$ ,  $\widehat{u} \neq 0$  and  $\widehat{u} \in \text{int } C_+$  is the second positive solution of problem  $(P_\lambda)$  with  $\lambda \in (0, \widehat{\lambda})$ . □

Finally we check the admissibility of the critical parameter  $\widehat{\lambda} > 0$ .

**Proposition 12** *If hypotheses  $H_0, H_1$  hold, then  $\widehat{\lambda} \in \mathcal{L}$ .*

**Proof** Let  $\{\lambda_n\}_{n \in \mathbb{N}} \subseteq \mathcal{L}$  be such that  $\lambda_n \uparrow \widehat{\lambda}$ . We can find  $u_n \in S_{\lambda_n} \subseteq \text{int } C_+$  which are minimizers of  $\widehat{\psi}_{\lambda_n}(\cdot)$  (truncation at  $\bar{u}_\sigma$  for  $\sigma \in (0, \lambda_n)$  small and at  $u_\beta \in S_\beta \subseteq \text{int } C_+$  with  $\beta \in (\lambda_n, \widehat{\lambda})$ ) and so

$$\begin{aligned} \widehat{\psi}_{\lambda_n}(u_n) &\leq \widehat{\psi}_{\lambda_n}(\bar{u}_\sigma) \\ &= \int_\Omega \frac{1}{p(z)} |\nabla \bar{u}_\sigma|^{p(z)} dz + \int_\Omega \frac{1}{q(z)} |\nabla \bar{u}_\sigma|^{q(z)} dz \\ &\quad - \int_\Omega \left( \lambda_n [\bar{u}_\sigma^{1-\eta(z)} + \bar{u}_\sigma^{\tau(z)}] + f(z, \bar{u}_\sigma) \bar{u}_\sigma \right) dz \text{ (see (42))} \\ &\leq \rho_p(\nabla \bar{u}_\sigma) + \rho_q(\nabla \bar{u}_\sigma) - \lambda \rho_\tau(\bar{u}_\sigma) - \widehat{\eta} \text{ with } \widehat{\eta} \in (0, +\infty) \\ &\leq \rho_p(\nabla \bar{u}_\sigma) + \rho_q(\nabla \bar{u}_\sigma) - \sigma \rho_\tau(\bar{u}_\sigma) - \widehat{\eta} \text{ (since } \sigma \in (0, \lambda)) \\ &= -\widehat{\eta} < 0 \text{ (see Proposition 5),} \\ \Rightarrow \widehat{\varphi}_{\lambda_n}(u_n) &< 0 \text{ (since } \widehat{\psi}_{\lambda_n} \Big|_{[\bar{u}_\sigma, u_\beta]} = \widehat{\varphi}_{\lambda_n} \Big|_{[\bar{u}_\sigma, u_\beta]} \Big). \end{aligned}$$

Also, we have  $\widehat{\varphi}'_{\lambda_n}(u_n) = 0$  in  $W^{-1,p'(z)}(\Omega)$  for all  $n \in \mathbb{N}$ . Then as in the proof of Proposition 7 (see the ‘‘Claim’’), we obtain

$$u_n \rightarrow u_* \text{ in } W_0^{1,p(z)}(\Omega) \text{ as } n \rightarrow +\infty.$$

We have

$$\begin{aligned} \langle \widehat{\varphi}'_{\lambda_n}(u_n), h \rangle &= 0 \text{ for all } h \in W_0^{1,p(z)}(\Omega), \text{ all } n \in \mathbb{N}, \\ \Rightarrow \langle V(u_n), h \rangle &= \int_\Omega \widehat{e}_{\lambda_n}(z, u_n) h dz, \\ \Rightarrow \langle V(u_*), h \rangle &= \int_\Omega \widehat{e}_{\widehat{\lambda}}(z, u_*) h dz \text{ for all } h \in W_0^{1,p(z)}(\Omega). \end{aligned}$$



Also we have  $\bar{u}_\sigma \leq u_n$  for all  $n \in \mathbb{N}$  and so  $\bar{u}_\sigma \leq u_*$  which means that  $u_* \in \mathcal{S}_{\hat{\lambda}} \subseteq \text{int } C_+$ , hence  $\hat{\lambda} \in \mathcal{L}$ .  $\square$

We have proved that

$$\mathcal{L} = (0, \hat{\lambda}].$$

We can state the following global existence and multiplicity theorem of problem  $(P_\lambda)$  (bifurcation-type theorem).

**Theorem 1** *If hypotheses  $H_0, H_1$  hold, then there exists  $\hat{\lambda} > 0$  such that:*

- (a) *for all  $\lambda \in (0, \hat{\lambda})$  problem  $(P_\lambda)$  has at least two positive solutions  $u_0, \hat{u} \in \text{int } C_+$ ;*
- (b) *for  $\lambda = \hat{\lambda}$  problem  $(P_\lambda)$  has at least one positive solution  $u_* \in \text{int } C_+$ ;*
- (c) *for all  $\lambda > \hat{\lambda}$  problem  $(P_\lambda)$  has no positive solution.*

**Acknowledgements** The authors wish to thank the two knowledgeable referees for their corrections and remarks. The second author is supported by the research fund of University of Palermo: “FFR 2023 Calogero Vetro”.

**Funding** Open access funding provided by Università degli Studi di Palermo within the CRUI-CARE Agreement. The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

**Data Availability** This paper has no associated data and material.

## Declarations

**Conflict of interest** The authors read and approved the final manuscript. The authors have no relevant financial or non-financial interests to disclose.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Abdalmonem, A., Scapellato, A.: Intrinsic square functions and commutators on Morrey–Herz spaces with variable exponents. *Math. Methods Appl. Sci.* **44**, 12408–12425 (2021)
2. Byun, S.-S., Ko, E.: Global  $C^{1,\alpha}$ -regularity and existence of multiple solutions for singular  $p(x)$ -Laplacian equations. *Calc. Var.* **56**, 76 (2017)
3. Cruz-Uribe, D., Fiorenza, A.: *Variable Lebesgue Spaces. Foundations and Harmonic Analysis*, Birkhäuser/Springer, Heidelberg (2013)
4. Diening, L., Harjulehto, P., Hästö, P., Růžička, M.: *Lebesgue and Sobolev Spaces with Variable Exponents. Lecture Notes in Mathematics*, vol. 2017. Springer, Heidelberg (2011)
5. Fan, X.: Global  $C^{1,\alpha}$ -regularity for variable exponent elliptic equations in divergence form. *J. Diff. Equ.* **235**, 397–417 (2007)

6. Gasirski, L., Papageorgiou, N.S.: Anisotropic nonlinear Neumann problems. *Calc. Var.* **42**, 323–354 (2011)
7. Giacomoni, J., Kumar, D., Sreenadh, K.: Sobolev and Hölder regularity results for some singular nonhomogeneous quasilinear problems. *Calc. Var.* **60**, 121 (2021)
8. Gilbarg, D., Trudinger, N.S.: *Elliptic Partial Differential Equations of Second Order*. Springer, Berlin (2001)
9. Harjulehto, P., Hästö, P., Koskenoja, M.: Hardy’s inequality in a variable exponent Sobolev space. *Georgian Math. J.* **12**, 431–442 (2005)
10. Li, G., Yang, C.: The existence of a nontrivial solution to a nonlinear elliptic boundary value problem of  $p$ -Laplacian type without the Ambrosetti-Rabinowitz condition. *Nonlinear Anal.* **72**, 4602–4613 (2010)
11. Lieberman, G.M.: The natural generalization of the natural conditions of Ladyzhenskaya and Ural’tseva for elliptic equations. *Comm. Partial Diff. Equ.* **16**, 311–361 (1991)
12. Papageorgiou, N.S., Winkert, P.: *Applied Nonlinear Functional Analysis*. W. de Gruyter, Berlin (2018)
13. Papageorgiou, N.S., Winkert, P.: Positive solutions for singular anisotropic  $(p, q)$ -equations. *J. Geom. Anal.* **31**, 11849–11877 (2021)
14. Papageorgiou, N.S., Winkert, P.:  $(p, q)$ -Equations with singular and concave-convex nonlinearities. *Appl. Math. Optim.* **84**, 2601–2628 (2021)
15. Papageorgiou, N.S., Winkert, P.: On a class of singular anisotropic  $(p, q)$ -equations. *Rev. Mat. Complut.* **35**, 545–571 (2022)
16. Papageorgiou, N.S., Zhang, C.: Singular  $(p, q)$ -equations with competing perturbations. *Appl. Anal.* **101**, 6151–6171 (2022)
17. Papageorgiou, N.S., Zhang, C.: Global multiplicity for the positive solutions of parametric singular  $(p, q)$ -equations with indefinite perturbations. *Bull. Malays. Math. Sci. Soc.* **46**, 5 (2023)
18. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: *Nonlinear Analysis Theory Methods*. Springer Monographs in Mathematics, Springer, Cham (2019)
19. Papageorgiou, N.S., Rădulescu, V.D., Zhang, Y.: Anisotropic singular double phase Dirichlet problems. *Discret. Contin. Dyn. Syst. S* **14**, 4465–4502 (2021)
20. Ragusa, M.A.: Parabolic Herz spaces and their applications. *Appl. Math. Lett.* **25**, 1270–1273 (2012)
21. Saoudi, K., Ghanmi, A.: A multiplicity results for a singular equation involving the  $p(x)$ -Laplace operator. *Compl. Var. Elliptic Equ.* **62**, 695–725 (2017)
22. Takáč, P., Giacomoni, J.: A  $p(x)$ -Laplacian extension of the Díaz–Saa inequality and some applications. *Proc. R. Soc. Edinb. A* **150**, 205–232 (2019)
23. Wei, M.: A characterization of  $C^{\alpha, \beta}$  via the commutator of Hardy-type operators on mixed Herz spaces. *Appl. Anal.* **101**, 5727–5742 (2022)
24. Willem, M.: *Minimax Theorems*. Birkhäuser, Boston (1996)

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.