

Antifungal activity of chili pepper extract with potential for the control of some major pathogens in grapevine

Marco Vuerich,^a Elisa Petrusa,^{a*}  Antonio Filippi,^b Stéphanie Cluzet,^c  Josep Valls Fonayet,^{c,d} Angela Sepulcri,^a Barbara Piani,^a Paolo Ermacora^a and Enrico Braidot^a



Abstract

BACKGROUND: In recent years, biofungicides have drawn increasing interest in vineyards for a more sustainable integrated and copper-limited pest management. Among alternatives, botanicals could represent valuable tools, being rich sources of biologically active compounds. Conversely to the well-known antioxidant and biological properties in relation to health benefits, investigation on bioactivity of hot pungent *Capsicum* sp. products against fungal phytopathogens in vineyards is still scarce. Therefore, the present study aimed at exploring the biologically active compounds profile of a chili pepper (*Capsicum chinense* Jacq.) pod extract and its antimicrobial properties against some of the major fungal and Oomycetes pathogens of grapevine, including *Botrytis cinerea* Pers., *Guignardia bidwellii* (Ellis) Viala & Ravaz and *Plasmopara viticola* (Berk. & M.A. Curtis) Berl. & De Toni.

RESULTS: The ethyl acetate-extracted oleoresin from the most pungent varieties was rich in capsaicinoids and polyphenols (371.09 and 268.5 $\mu\text{g mg}^{-1}$ dry weight, respectively). Capsaicin and dihydrocapsaicin, hydroxycinnamic and hydroxybenzoic acids and quercetin derivatives were the most abundant, while carotenoids represented only a minor fraction. The oleoresin was efficient to inhibit all three pathogenic fungi and ED₅₀ values were determined, evidencing that *G. bidwellii* was the more sensitive ($0.233 \pm 0.034 \text{ mg mL}^{-1}$).

CONCLUSION: The results suggested a potentiality of chili pepper extract for the control of some important grapevine pathogens, their possible application being helpful for the recommended limitation in extensive use of copper in vineyard. The complex mixture of high amounts of capsaicinoids, associated to specific phenolic acids and other minor bioactive components might contribute to the observed antimicrobial action of chili pepper extract.

© 2023 The Authors. *Pest Management Science* published by John Wiley & Sons Ltd on behalf of Society of Chemical Industry.

Supporting information may be found in the online version of this article.

Keywords: biological control; *Botrytis cinerea*; *Capsicum* spp.; grapevine; *Guignardia bidwellii*; *Plasmopara viticola*

1 INTRODUCTION

In relation to the cultivated area (7.5 billion ha)¹ and economic value, grapevine (*Vitis vinifera* L.) is one of the most important fruit crops worldwide. Due to the wide range of pathogens affecting grapevine, causing direct and indirect damage to the production, an intense use of pesticides is required for the protection of this crop.^{2,3} Together with *Erysiphe necator* (causal agent of powdery mildew), *Botrytis cinerea* Pers. and *Plasmopara viticola* (Berk. & M.A. Curtis) Berl. & De Toni, respectively the causal agents of gray mold and downy mildew, are the pathogens for which most treatments are required.^{2,4}

Copper-based pesticides represent one of the main means of containment of gray mold and downy mildew, especially in organic viticulture where its use is still authorized due to its wide range of action.^{5,6} However, the extensive use of copper can lead to issues at different scales. From an ecological point of view, copper accumulated in the soil affects plants and fauna.^{6,7} Moreover,

the presence of copper and other fungicides residues in grapes and wine can be a threat to human health.⁷

* Correspondence to: E Petrusa, Department of Agricultural, Food, Environmental and Animal Sciences, University of Udine, via delle Scienze 91, 33100, Udine, Italy, E-mail: elisa.petrussa@uniud.it

a Department of Agricultural, Food, Environmental and Animal Sciences, University of Udine, Udine, Italy

b Department of Medicine, University of Udine, Udine, Italy

c Equipe Molécules d'Intérêt Biologique (MIB)—ISVV, University of Bordeaux, Faculté des Sciences Pharmaceutiques, INRAE, Bordeaux INP, Bordeaux Sciences Agro, OENO, UMR 1366, Villenave d'Ornon, France

d MetaboHUB, Bordeaux Metabolome Facility, Villenave d'Ornon, France

In this context, the European Union has been regulating for years to reduce and/or replace the use of copper. Currently, according to EU regulation 2018/1981⁸ the maximum amount of copper allowed is 28 kg ha⁻¹ over a period of 7 years (i.e., 4 kg ha⁻¹ per year). Moreover, in the Farm to Fork (F2F) strategy, the EU targets 25% of organic land by 2030 and also to reduce by 50% the use of chemical pesticides.⁹

One of the strategies to reduce or replace copper and other synthetic pesticides use is the development of grapevine resistant varieties, the use of which has recently been allowed by the EU Commission for the production of wines with protected denominations of origin.^{3,10} In particular, in recent decades numerous efforts have primarily focused on breeding in order to obtain varieties resistant to downy and powdery mildew, and gray mold.¹¹ The use of these cultivars allows the reduction of fungicide applications. However, the onset of secondary diseases previously controlled by phytosanitary treatments occurs as a backlash.³ *Guignardia bidwellii* (Ellis) Viala & Ravaz, the causal agent of black rot disease, has diffused from Northern America and increased in importance in the last few years in northern wine-growing regions in Europe, becoming one of the more serious new emergent grapevine pathogens in Italy and France, favored by both reduction of copper dosage allowed in organic viticulture and for the significant reduction of fungicide treatments on mildew resistant/tolerant varieties, yet black-rot susceptible, in integrated management.^{3,12,13} The latter factors are favoring the increase of black rot outbreaks and leading to relevant yield loss in European viticultural production.^{14,15}

The research on botanicals with bioactive action against phytopathogens has received increasing interest, since biopesticides could represent widely acceptable, sustainable as environment-friendly and potentially lower toxic substitutes to synthetic fungicides in combination with limited copper application in an integrated management program.^{16,17} Recently, the use of the biocontrol agents has been highly endorsed by European Commission's recommendations in the Farm to Fork (F2F) strategy for sustainable food,⁹ under the European Green Deal. Many traditional or novel aromatic and medicinal plants represent an important source of secondary metabolites such as alkaloids, tannins and flavonoids, quinones, glycosides, saponins and terpenoids, conferring a huge range of variable antimicrobial properties.^{7,16,18–20} These bioactive molecules are naturally accumulated as plant defense barriers against phytopathogens²¹ and represent strategies for environmental adaptation to abiotic stresses²² (e.g., UV exposure, dehydration, high temperatures), or as pollinator's attractants.²³ Among the most domesticated crops utilized for centuries as culinary spice, chili peppers (*Capsicum* spp.) have been recognized to be rich in health-beneficial bioactive phytochemicals for the human diet and to possess important medicinal properties.^{24–26} Regarding biochemical composition, the five most widely cultivated chili pepper species (*C. annuum* L., *C. frutescens* L., *C. chinense* Jacq., *C. baccatum* L. and *C. pubescens* Ruiz and Pav.) are recognized as important sources of vitamins and several bioactive compounds.^{27,28} The most important are carotenoids and flavonoids (responsible for the attractive colors of fruits and functioning as natural antioxidants and defenses against biotic and abiotic stresses), and more of 20 capsaicinoids (simple arylalkylamide alkaloids), the principal member of which is capsaicin (*trans*-8-methyl-N-vanillyl-6-nonenamide), synthesized mainly in the placenta, conferring the typical pungent taste.^{24,29} Among the latter, capsaicin and dihydrocapsaicin are the most abundant capsaicinoids found in chili peppers, being

69 and 22%, respectively, followed by nordihydrocapsaicin, homocapsaicin and homodihydrocapsaicin, in a wide range of variance affected by fruit ripening, species, cultivars, and cultivation conditions.^{29,30} Most of these bioactive compounds are already known to exert antimicrobial or antifungal actions both as single molecules or as major components of pepper extracts and oleoresins.^{31–37} A number of reports confirmed also that *Capsicum* sp. extracts, in addition to be proposed as biopesticide against insects,³⁸ were effective against plant pathogens as post-harvest fungal agent in apple³¹ and several agriculture fungi like *Alternaria alternata*, *Fusarium oxysporum*, *Botrytis cinerea*, *Verticillium dahliae*, *Phytophthora capsici*.^{39–43} Also, it has suggested that capsaicinoids accumulation in wild chili peppers may represent an adaptive response to *Fusarium* seed infection in an evolutionary sense.⁴⁰ In some cases, along with an antifungal direct action, some secondary metabolites found in chili peppers were also proven to elicit natural plant induced resistance.^{41,43–45}

Despite the recognized inhibitory actions against several pathogenic fungi of plant crops, to date investigation on antifungal activity of hot pungent *Capsicum* sp.-based products against the most important pathogens of *Vitis vinifera* is still scarce, with the exception of one existing published patent about aqueous solution comprising *Capsicum* for contrasting downy or powdery mildew.⁴⁶

Accordingly, in this research, we examined the potential grapevine fungicide activity of a chili pepper oleoresin, obtained from fruits of the hottest varieties of *C. chinense* and extracted through maceration with a safe and eco-friendly solvent. The profile of the major bioactive compounds was also discussed in relation to their putative action against some of the most detrimental phytopathogens in viticulture production.

2 MATERIALS AND METHODS

2.1 Chemicals

Capsorubin was obtained from CaroteNature Gmbh (Musingen, Switzerland). Capsaicin (99.9% purity), capsanthin, zeaxanthin, β -cryptoxanthin, potato dextrose agar (PDA), β -carotene, lutein, β -Apo-8-carotenal (*trans*), dichloromethane and ethyl acetate were purchased from Merck, Darmstadt (Germany); violaxanthin, anteraxanthin and flavonoids from Extrasynthese, Genais (France); ethanol (analytical grade) from Carlo Erba Reagents, Milan (Italy); acetonitrile and absolute methanol from Thermo Fisher Scientific (Illkirch, France). Fluorescein, 2,2'-azobis (2-methylpropionamide) dihydrochloride (AAPH) and hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), were purchased from Sigma-Aldrich (France). The water used for HPLC–MS was purified with an Elga system (Bucks, U.K.). Formic acid was obtained from Sigma Chemical Company (St Louis, MO). Phenolic acids were purchased from Sigma.

2.2 Preparation of chili pepper oleoresin

Commercial dry powder of fruits and seeds of *Capsicum chinense* Jacq., var. 'Habanero Mustard', 'Habanero Pastel', 'Trinidad Moruga Scorpion Red', 'Trinidad Moruga Scorpion Choco', 'Carolina Reaper', 'White Naga', 'Naga Morich Chocolate' was purchased by means of local producers (Italy). The plant material was chosen on the basis of the highest pungency level [ranging from 350 000 Scoville heat units (SHU) up to more than 1 million SHU],^{47–49} a parameter which is known to be directly related to capsaicinoid content. Since the different chili pepper powders were obtained from plant material available on the internet market, we prefer

to be not strictly dependent on characteristics of a single variety. Indeed, this could be a limit to an easy and quick acquisition of the plant material and so we prefer to make a blend that represented an intermediate option, easy to be replicated.

In the case of all varieties used, the percentage of the powders added to the total mixture was always the same and the oleoresin was extracted by macerating them in 100% ethyl acetate (100 g L⁻¹) for 2 h at 25 °C under stirring. After centrifugation at 28 000 × *g* for 10 min (Beckman Coulter, Avanti J-26S XP, Milan, Italy), the supernatant fraction was collected, and the solvent removed by rotary evaporator (Buchi, Rotavapor® R-100, Cornaredo, Italy). The final extract was diluted in 100% ethanol obtaining a stock solution of approx. 400–500 mg dry weight (DW) mL⁻¹, stored at –20 °C until use.

2.3 GC–MS analysis of chili pepper oleoresin

Oleoresin extract composition was determined by GC–MS (Agilent Technologies GC–MS 5977E, USA) and the amount of each compound was expressed as percentage (area percent method), and also in µg mg⁻¹ DW by external calibration with capsaicin standard.

Oleoresin extract was dried under nitrogen and resuspended in dichloromethane (Merck, Darmstadt, Germany)⁵⁰ leading to a solution of known concentration; the sample was transferred in vial suitable for injection. GC–MS was performed in EI mode (70 eV) with a 5977E MSD system, a single-quadrupole (Agilent Technologies, USA) equipped with 7683A autosampler and automatic split/splitless injector. An aliquot (1 µL) of solutions was injected in splitless mode for the first 1.5 min of the analysis and then in the split mode for the remainder of the run. The separation was carried out on SP-5 ms capillary column (30 m × 0.25 mm × 0.25 µm) (Supelco, Bellefonte, PA). The GC oven program started for 10 min at 60 °C; then, the temperature was ramped up (3 °C min⁻¹) to 95 °C (hold time 5 min), then again to 270 °C at the rate of 3 °C min⁻¹ and to end with 5 min hold time. The total run time was 89.93 min and in post run the temperature was set to 280 °C for 5 min in order to clean the column. The length of the run was necessary to check up to which minute compounds of interest eluted; after 79 min, however, nothing of interest was detected. Helium (6.0) was used as carrier gas with a constant flow rate of 1.0 mL min⁻¹. The temperature value of the ion source and the quadrupole were set to 240 °C and 150 °C, respectively; the transfer line was set to 240 °C. The GC–MS analyses in the full scan mode for a range of 50 to 650 amu and the analysis were performed after a solvent delay of 4 min, with 3 microscan sec⁻¹. Compounds were identified by comparing their mass spectra of NIST 14 Mass spectral Library and by matching the results with those reported in the literature.^{50–52}

2.4 HPLC analysis of principal polyphenols

To 50 mg of a mixture of a fine powder from chili pepper pods, prepared as the one used to obtain oleoresin, 1 mL of 70% methanol was added and the mixture was put into an ultrasonic water bath at 25 °C for 15 min. This extraction step was based on the one developed by Jeong *et al.*⁵³ for polyphenols extraction of pepper. After centrifugation (15 000 rpm, 5 min), the supernatant was collected and placed in a new tube. The powder was extracted a second time in a similar manner. All the supernatants were combined and diluted with water (1:1 v/v) and newly centrifuged previous HPLC-MS injection.

Analysis of polyphenols was performed by HPLC-MS/MS in Multiple Reaction Monitoring (MRM) using a 1260 Infinity UPLC

(Agilent Technologies, Courtaboeuf, France) coupled to a 6430 triple quadrupole mass spectrometer (Agilent Technologies, Courtaboeuf, France) and adapting a previous protocol.⁵⁴ Two µL of sample were injected into an Agilent Zorbax SB-C18 (100 mm × 2.1 mm, 1.8 µm) thermostated at 40 °C, and separation of the compounds was performed at a flow rate of 0.43 mL min⁻¹ with a mobile phase composed of solvent A (distilled water, 0.1% formic acid) and solvent B (acetonitrile, 0.1% formic acid). The gradient was as follows: 0 to 4 min, (1% B-10% B); 4 to 12 min (10% B 8% to 20% B); 12 to 13 min (20% B to 30% B); 13 to 16 min 30% B; 16 to 20 min (30% B to 50% B); 21 to 25 min, 95% B. The source parameters were: capillary voltage, +3000 V; nebulizer pressure, 15 psi; dry gas, 11 L min⁻¹; dry temperature, 350 °C. Specific MRM transitions for each polyphenol were used for identification⁵⁴ and quantification was done with the Mass Hunter Data Analysis software (Agilent, Technologies, Courtaboeuf, France). Polyphenols were determined from calibration curves of pure standards (injected concentrations ranging from 0.08 to 43 µg mL⁻¹). Feruloyl hexoside compounds were tentatively identified by their MRM fragmentation and quantified as equivalent of kaempferol hexoside. Concentrations were expressed in µg mg⁻¹ DW of phenolic compound.

2.5 HPLC analysis of principal carotenoids

Carotenoids such as capsorubin, violaxanthin, capsanthin, antheraxanthin, zeaxanthin, β-cryptoxanthin, β-carotene and lutein were extracted according to Su *et al.*⁵⁵ and to Borguini *et al.*⁵⁶ with some modifications. Briefly, 20 µL of oleoresin and 100 µL of internal standard (IS) β-Apo-8-carotenal (*trans*) (65 mg L⁻¹) were mixed in a Falcon tube, then ethyl acetate (1 mL) and Milli-Q® water (2 mL) were added and the sample was centrifuged at 3000 rpm for 15 min on a Centrifuge 5415 (Eppendorf AG, Germany). Sodium sulfate anhydrous was added to dry and the sample was kept in the dark for about 20 min, then the organic phase was filtered through syringe filter (PTFE 0.22 µm, 25 mm, DTO Servizi Srl, Venice, Italy). Two hundred milliliters of extract were then evaporated to dryness under nitrogen, reconstituted with 200 µL of acetone and filtered through syringe filter. All samples were stored at –20 °C until analysis. The filtrate was transferred into an autosampler vial and finally 20 µL were injected into HPLC. The HPLC system included a Shimadzu LC-20AT pump, a vacuum degasser, a Prominence SPD-M20A photodiode-array detector, a Prominence SIL-20 AC HT autosampler (20 µL loop) and a Prominence CTO-20 AC column oven set at 25 °C (Shimadzu Corporation, Kyoto, Japan). The HPLC separation was achieved using a Spherisorb column ODS 2 (4.6 × 250 mm, 5.0 µm particle size) and a mobile phase of acetonitrile: methanol (40:60) at a flow rate of 1 mL min⁻¹ was used. The optimal wavelength for the detection of the main carotenoid components in the extract was found to be 454 nm and the detector slit width was 4 nm. Full spectra were recorded in the range 190–800 nm. Equipment control, data acquisition and integration were performed with Shimadzu LabSolutions (Ver. 5.54 SP2) Software.

2.6 Total carotenoid quantification by spectrophotometric analysis

The average carotenoid quantification of the dried oleoresin extract was determined according to the spectrophotometric method of Biehler *et al.*⁵⁷ Dry extract was suspended in 0.5 mL acetone and sonicated for 2 min. The absorbance at the mean absorbance maximum (A₄₅₀) of increasing dilutions of the suspension was read in a 1 mL-quartz cuvette by spectrophotometer

(Agilent Technologies, 8453, Milan, Italy). The average carotenoid concentration (c) was calculated by the equation:

$$c \text{ (mol L}^{-1}\text{)} = A_{450} \cdot \text{dilution factor} / 135310$$

and expressed as mg per 100 g of oleoresin by using an average molar mass of 536.88.

2.7 Antioxidant potential assay

The antioxidant capacity of the oleoresin extract was evaluated with the Oxygen Radical Absorbance Capacity (ORAC) method. The assay was made according to a modified method developed by Dávalos *et al.*⁵⁸ by using an automated plate reader (Fluostar Omega; BMG Labtech, Offenburg, Germany). Briefly, oleoresin (20 μ L), Trolox (standard) or phosphate buffer (blank) prepared in 75 mM phosphate buffer (pH 7.4) and 120 μ L of fluorescein solution (70 nM final concentration) were mixed with in a 96-well black plate, then placed at 37 °C during 5 min. After that, 60 μ L of AAPH (12 mM final concentration) were added and the reaction was immediately followed by monitoring fluorescence using 485 nm excitation and 530 nm emission wavelengths at each cycle of 90 min. A standard curve of Trolox was used and its ORAC value was obtained by using the Trolox standard curve $y = 44.231.47x + 140.977.55$ with a $R^2 = 0.89$. The antioxidant activity of the pepper extract was expressed as μ moles Trolox equivalents (TE) g^{-1} DW extract (ORAC). All samples were analyzed five times.

2.8 Pathogen inoculum source

B. cinerea Pers. isolate used in the experiment was isolated from harvested bunches in local vineyards (Udine, Italy) since 2018 and grown on potato dextrose agar (PDA) medium at 21 °C under a 12 h photoperiod; the isolate was multiplied every 2 weeks by hyphal plug or exposed to actinic blue light under a 12 h photoperiod during one more week for inducing sporulation. Conidial suspension was adjusted at the concentration of 4×10^5 conidia mL^{-1} in sterile distilled water, before inoculation on PDA for the growth inhibition assessment.

G. bidwellii (Ellis) Viala & Ravaz isolate was obtained in June 2018 under sterile condition, by selecting single pycnidia from symptomatic leaves of infected grapevines, at the University of Udine. The strain was grown and maintained on PDA at 21 °C under a photoperiod of 12 h light and transferred in fresh medium every 3–4 weeks for maintenance. A plug of 3–4-week-old mycelium (5 mm \varnothing) was aseptically removed and used as fungal inoculation for growth inhibition assessment.

P. viticola (Berk. & M.A. Curtis) Berl. & De Toni sporangia were collected from natural infected leaves in an untreated vineyard in 2019 (University of Udine, Italy) and weekly propagated by infecting 4th or 5th leaf of one-year old potted vines of *Vitis vinifera* cv. Sauvignon Blanc grafted on SO4 rootstock and maintained in controlled conditions at 25 °C under a 12 h photoperiod, 55% of relative humidity. The fresh sporulation was collected the day of the experiment, counted by microscope under visible light and adjusted with sterile distilled water at the final concentration of 4×10^5 sporangia mL^{-1} .

All pathogen isolates were identified on the basis of disease symptoms on plants where they were collected and on microscope observation of characteristic morphological traits of reproductive structures in artificial substrate and/or host surface.

2.9 In vitro antifungal activity against Botrytis cinerea

The inhibition of different concentrations of chili pepper oleoresin on *B. cinerea* growth was analyzed by the diffusion technique on PDA growth medium. Eleven doses of oleoresin (0.44–8.86 mg mL^{-1} in sterile distilled water with 4.9% (v/v) ethanol) were prepared for the assay and the diluted solutions (70 μ L) were spread on 5 mL of solidified PDA medium in Petri dishes (50 mm \varnothing) just before inoculation with a drop of 5 μ L of conidial suspension, adjusted at 4×10^5 conidia mL^{-1} . Sealed Petri dishes were incubated in a growth chamber at 21 °C under a 12 h photoperiod, until full evasion of fungus was reached in control samples. The mycelium growth area was monitored daily and photographs of Petri dishes with mycelia were taken and images analyzed by ImageJ Fiji software⁵⁹ for the assessment of percentage of growth inhibition with respect to the control. Three independent sets of experiments were performed with four replicates each set.

2.10 In vitro antifungal activity against Guignardia bidwellii

Inhibition activity of different concentrations of chili pepper oleoresin against *G. bidwellii* was determined by aseptically incubating a plug of 3–4 week-old mycelium (5 mm \varnothing) on 3 mL of PDA medium in 6 well cell culture microplates for growth test assessment. The diluted solutions (70 μ L at 0.001 to 12.5 mg mL^{-1} in sterile distilled water with 6.9% (v/v) ethanol) were spread on PDA medium just before inoculation. Radial growth of the mycelium was monitored for 3, 6, 9 and 12 days, respectively and weighted to each respective area of the plug at time 0. Photographs of Petri dishes with mycelia were taken and images were then analyzed by ImageJ software for the assessment of the growth area. Three independent sets of experiments were performed with four replicates each set.

2.11 Inhibition of Plasmopara viticola sporulation on leaf discs

The inhibitory effect of different concentrations of chili pepper oleoresin extract against *P. viticola* sporulation was analyzed on leaf discs, according to Corio-Costet *et al.*⁶⁰ Eleven doses of oleoresin (0.2–10 mg mL^{-1}) in sterile distilled water with 3.8% (v/v) ethanol were prepared for the assay the day of experiment. Four leaf discs (21 mm \varnothing) on wet Whatman filter paper were sprayed with each solution (1.25 mL) and incubated for 1 d in the dark before inoculation with three drops (15 μ L) of fresh *P. viticola* inoculum adjusted at 4×10^5 sporangia mL^{-1} . Sealed Petri dishes were incubated in a humid chamber at 21 °C under a 12 h photoperiod until complete evasion of conidiophores. Photographs of Petri dishes with sporangia evasion were taken after 5 days, and images were analyzed by ImageJ software for the assessment of percentage of sporulation to total leaf area, as described in Peresotti *et al.*⁶¹ Three independent experiments were performed with four replicates each set.

2.12 Statistical analysis

Antifungal activity of chili pepper oleoresin was statistically analyzed using R extension package drc⁶² and for each pathogen a regression curve was fitted using dose–response analysis sigmoidal model Log-logistic. The effective dose or concentration (ED_{50}) of oleoresin to inhibit 50% of fungal growth was derived by log-logistic dose–response curve analysis.⁶³

Table 1. GC–MS analysis of chili pepper extract

Name of compound	Formula	%	m/z	Content ($\mu\text{g mg}^{-1}$ DW)	Biological activity
Undecane	C ₁₁ H ₂₄	1.067			Antioxidant ^{66,67}
Butanoic acid, 3-methyl-, hexyl ester	C ₁₁ H ₂₂ O ₂	0.054			
<i>cis</i> -Muurola-3,5-diene	C ₁₅ H ₂₄	0.017			
5-Octen-1-ol, (Z)-	C ₈ H ₁₆ O	0.078			
1,3-Pinanylamine	C ₁₀ H ₁₉ N	0.007			
3,7-Diacetamido-7H-s-triazolo[5,1-c]-s-triazole	C ₇ H ₉ N ₇ O ₂	0.005			
N-[4-Aminobutyl]aziridine	C ₆ H ₁₄ N ₂	0.007			
α -Himachalene	C ₁₅ H ₂₄	0.103			
2,6,6-Trimethyl-bicyclo[3.1.1]hept-3-ylamine	C ₁₀ H ₁₉ N	0.016			
2-Methylbutyl 8-methylnon-6-enoate	C ₁₅ H ₂₈ O ₂	0.113			
Z,Z-2,5-Pentadecadien-1-ol	C ₁₅ H ₂₈ O	0.098			
2-Trifluoroacetoxydodecane	C ₁₄ H ₂₅ F ₃ O ₂	0.102			Antimicrobial ^{68,69}
8,9,9,10,10,11-Hexafluoro-4,4-dimethyl-3,5-dioxatetracyclo[5.4.1.0(2,6)0.0(8,11)]dodecane	C ₁₂ H ₁₂ F ₆ O ₂	0.006			Antimicrobial ⁷⁰
13-Tetradecenal	C ₁₄ H ₂₆ O	0.026			
10-(3-Ethyl-2-oxiranyl)-1-decanol	C ₁₄ H ₂₈ O ₂	0.036			
Imidazole, 2-amino-5-[(2-carboxy)vinyl]-	C ₆ H ₇ N ₃ O ₂	0.033			Antimicrobial ⁷¹
6-Tridecene, (Z)-	C ₁₃ H ₂₆	0.044			
Hexadecen-1-ol, trans-9-	C ₁₆ H ₃₂ O	0.139			Antioxidant ⁷²
γ -Guanidinobutyric acid	C ₅ H ₁₁ N ₃ O ₂	0.010			
unknown		0.012			
1-Octanesulfonyl chloride	C ₈ H ₁₇ ClO ₂ S	0.012			
Cyclopentanone, 2-(1-methylpropyl)-	C ₉ H ₁₆ O	0.013			
Z-2-Dodecenol	C ₁₂ H ₂₄ O	0.011			
l-Guanidinosuccinimide	C ₅ H ₇ N ₃ O ₂	0.003			
<i>E</i> -2-Tetradecen-1-ol	C ₁₄ H ₂₈ O	0.358			
(Z)6-Pentadecen-1-ol	C ₁₅ H ₃₀ O	0.038			Antimicrobial ⁷³
(8Z)-14-Methyl-8-hexadecen-1-ol	C ₁₇ H ₃₄ O	0.059			
2(1H)-Benzocyclooctenone, decahydro-10a-methyl-, trans-	C ₁₃ H ₂₂ O	0.092			
3-Trifluoroacetoxypentadecane	C ₁₇ H ₃₁ F ₃ O ₂	0.091			Antioxidant ⁷⁴
7-Hexadecenal, (Z)-	C ₁₆ H ₃₀ O	0.100			
Methyl 12,13-tetradecadienoate	C ₁₅ H ₂₆ O ₂	0.021			
3-Trifluoroacetoxydodecane	C ₁₄ H ₂₅ F ₃ O ₂	0.046			
10-Bromodecanoic acid, ethyl ester	C ₁₂ H ₂₃ BrO ₂	0.012			
Octadecanal	C ₁₈ H ₃₆ O	0.139			Antimicrobial ⁷⁵
Nerolidol	C ₁₅ H ₂₆ O	0.134			Antioxidant/antibacterial/ antifungal/insecticidal/anti-inflammatory/ defense-related terpenes in <i>Vitis</i> and <i>Camellia</i> ^{76–81}
<i>E</i> -11-Hexadecenal	C ₁₆ H ₃₀ O	0.093			
1,10-Hexadecanediol	C ₁₆ H ₃₄ O ₂	0.005			
unknown		0.015			
Methyl 8-methyl-nonanoate	C ₁₁ H ₂₂ O ₂	0.008			
N-Pentadecylacetamide	C ₁₇ H ₃₅ NO	0.228			
9-Octadecenamide, (Z)-	C ₁₈ H ₃₅ NO	0.198			
γ -Thionodecalactone	C ₁₀ H ₁₈ OS	0.065			
N-Hexadecylacetamide	C ₁₈ H ₃₇ NO	0.095			
unknown		0.029			
9-Octadecenamide, (Z)-	C ₁₈ H ₃₅ NO	0.048			
1-Ethyl-dodecyl acrylate	C ₁₇ H ₃₂ O ₂	0.022			
<i>p</i> -Cresol, 2,2'-methylenebis[6-tert-butyl-	C ₂₃ H ₃₂ O ₂	0.037			
Pelargonic acid vanillylamide (Nonivamide)	C ₁₇ H ₂₇ NO ₃	0.614	137, 151, 293	37.81	Antioxidant/antifungal ^{40,41,82}

Table 1. Continued

Name of compound	Formula	%	m/z	Content ($\mu\text{g mg}^{-1}$ DW)	Biological activity
Tetradecane, 2,6,10-trimethyl- Nordihydrocapsaicin	C ₁₇ H ₃₆	0.239			
	C ₁₇ H ₂₇ NO ₃	0.373	137 , 151, 293	37.53	Antioxidant/antifungal ^{40,41,82}
Dihydro- β -agarofuran Capsaicin	C ₁₅ H ₂₆ O	0.068			
	C ₁₈ H ₂₇ NO ₃	69.436	137 , 152, 305	118.58	Antioxidant/antifungal ^{40,41,82}
Dihydrocapsaicin	C ₁₈ H ₂₉ NO ₃	23.755	137 , 151, 307	64.97	Antioxidant/antifungal ^{40,41,82}
Dihydrocapsaicin, O-acetyl-	C ₂₀ H ₃₁ NO ₄	0.675	137 , 195, 307	37.88	Antioxidant/antifungal ^{40,41,82}
unidentified		0.080	137	37.09	
unidentified		0.122	137	37.23	
Heptacosane	C ₂₇ H ₅₆	0.505			Antibacterial ⁸³
unidentified		0.029			
Octacosane	C ₂₈ H ₅₈	0.206			Antibacterial ⁸⁴
Total Capsaicinoids				371.09 (37.11)*	

Note: Peak area relative amount ratio (%), based on the normalized value (100.0%) and content of the compounds detected from chili pepper oleoresin after chromatographic run in dichloromethane.

*Relative abundance (% wt) on oleoresin DW basis.

3 RESULTS

3.1 GC-MS analysis of chili pepper oleoresin

To investigate the antifungal efficacy of chili pepper extracts against some fungal pathogens in vineyard, a number of the hottest varieties belonging to the highly pungent species *C. chinense* were selected in relation to their high content in carotenoid and capsaicinoids. The choice of a mix of varieties instead of a single one was dictated by the fact that this option could be more easily reproduced from commercial market. Of course, if we would be able to produce by ourselves one of the best varieties this would be the best choice.

The powder mixture of each of these varieties was macerated in equal proportion in ethyl acetate, chosen as a very good organic solvent for achieving high capsaicinoid recovery in oleoresins^{64,65} and GC-MS analysis was then performed for quantification and characterization of these compounds.

Comparing GC-MS chromatograms of pure capsaicin (Supporting Information, Fig. S1, panel A), used as standard, with that of oleoresin sample (Supporting Information, Fig. S1, panel B), it can be observed that its peak overlapped quite similarly with the corresponding one identified in the oleoresin sample, with a retention time of 74.096 and 74.218 min, respectively. Among the 79 detected compounds in the oleoresin under analysis, 48 were identified by matching their spectra with the NIST library and reported in Table 1, in relation to their relative peak area (as a percentage of the whole chromatogram) and their potential bioactivity. The benzyl cation fragment at m/z 137, commonly found in all capsaicinoids of pepper extracts, has been used for their specific identification in SIM mode of GC-MS analysis, along with m/z 305, 307, 293.⁵¹

Among capsaicinoids, the most abundant principal compounds in relation with their % of total peak area were represented by capsaicin and dihydrocapsaicin, and, at minor extent, O-acetyl dihydrocapsaicin, nonivamide and nordihydrocapsaicin, followed by two unidentified capsaicinoids. The results obtained also evidenced that capsaicin and dihydrocapsaicin contribute for the highest amount (69.44 and 23.76% of the total compounds analyzed by GC-MS, respectively) in chili pepper extract, whereas the other minor capsaicinoids (nonivamide and nordihydrocapsaicin) accounted only for low percentage (0.987%). As expected, total capsaicinoids were well represented, reaching approximately 95% of the total amount of various compounds detected by GC-MS analysis in oleoresin.

In term of abundance, other most represented compounds identified were alkanes (Undecane, Heptacosane and Octacosane, 1.07, 0.51 and 0.21% of the total, respectively), essential oils (*E*-2-Tetradecen-1-ol and Hexadecen-1-ol, trans-9-, 0.36 and 0.139%, respectively), terpenes (2,6,10-Trimethyltetradecane, Nerolidol, α -himachalene, 0.24, 0.13, and 0.10%), oleic acid derivative (9-Octadecenamide, (*Z*)-, 0.2%), and volatile fatty aldehydes (Octadecanal, 0.14%). Some minor components were acids (as 3-Methylbutanoic acid), alcohols (5-Octen-1-ol, (*Z*)-), aldehydes (*E*-11-Hexadecenal and 13-Tetradecenal), esters (as Butanoic acid, 3-methyl-, hexyl ester and 2-Methylbutyl 8-methylnon-6-enoate). By means of linear regression equation obtained from calibration with the internal standard of capsaicin, the concentration of the more abundant capsaicinoid found in oleoresin reached 118.58 $\mu\text{g mg}^{-1}$ DW and total content of capsaicinoids accounted for the value of 371.09 $\mu\text{g mg}^{-1}$ DW.

Table 2. Polyphenol concentration and biological activity in chili pepper extract

Compounds	Content ($\mu\text{g mg}^{-1}$ DW)	Biological activity
Phenolic acids		
Hydroxycinnamic acids		
Caffeic acid	1.5 \pm 0.1 (0.58)*	
Vanillic acid	174.0 \pm 9.4 (64.91)*	Antifungal ⁸⁵
Ferulic acid	14.4 \pm 0.6 (5.38)*	Antifungal ⁸⁶
Feruloyl hexoside isomer	1.4 \pm 0.3 (0.52)*	
Feruloyl hexoside isomer	3.6 \pm 0.2 (1.34)*	
Feruloyl hexoside isomer	1.3 \pm 0.2 (0.48)*	
Feruloyl hexoside isomer	1.8 \pm 0.2 (0.68)*	
Hydroxybenzoic acids		
Protocatechuic acid	35.8 \pm 0.6 (13.35)*	Antifungal ⁸⁷
Flavonoids		
Flavonols		
Quercetin 3-glucuronide	0.5 \pm 0.1 (0.20)*	
Quercetin 3-rhamnoside	13.2 \pm 1.6 (4.94)*	Antifungal ⁸⁸
Quercetin 3-rutinoside	1.2 \pm 0.4 (0.46)*	
Kaempferol 3-glucoside	<LOD	
Flavanones		
Naringenin	1.3 \pm 0.1 (0.47)*	
Naringenin glucoside	16.0 \pm 1.7 (5.98)*	Antifungal ^{89,90}
Flavanols		
Catechin	<LOD	
Epicatechin	<LOD	
B1	<LOD	
B2	<LOD	
Total polyphenols	268.5 \pm 15.4 (100)*	

Note: Values are means \pm standard error of triplicate analysis.
*Relative abundance (% wt) on oleoresin DW basis.

Among the identified metabolites, some capsaicinoids and nerolidol were described to have potential antifungal property.

3.2 HPLC analysis of polyphenols in chili pepper oleoresin

The polyphenolic analysis of chili pepper oleoresin was obtained after a double extraction of pepper powder in 100 and 70% methanol, according to Jeong *et al.*⁵³ which allowed identification of 13 different polyphenols, such as phenolic acids components among cinnamic and benzoic derivatives, three quercetin derivatives as flavonols, and naringenin and its glucoside as flavanones, while flavone and flavanol components were not detected or present under limit of detection (Table 2). The most abundant compound on a DW basis was represented by vanillic acid (65%), followed by protocatechuic acid (13%), being hydroxycinnamic derivatives the most important fraction in terms of quantity (74%) in respect to the other polyphenol compounds. The total concentration of polyphenols in oleoresin accounted for 268.5 \pm 15.4 $\mu\text{g mg}^{-1}$ DW, quite similar to the amount reached by total fractions of capsaicinoids.

3.3 HPLC analysis of carotenoids and antioxidant potential of chili pepper oleoresin

The main carotenoid compounds present in chili pepper oleoresin were identified by comparing retention times (Rt) and UV-Vis spectra obtained by HPLC analysis with the corresponding parameters exhibited by the standards (Supporting Information, Fig. S2 and Table 3).

Oleoresin from the red and orange-yellow pigmented chili pepper varieties contained large abundance of both major yellow and red carotenoids, as β -carotene (0.59 $\mu\text{g mg}^{-1}$ DW and 12.5% of relative peak area), capsanthin (0.22 $\mu\text{g mg}^{-1}$ DW and 8.5%), zeaxanthin (0.06 $\mu\text{g mg}^{-1}$ DW and 3.7%), yellow xanthophyll β -cryptoxanthin (0.08 $\mu\text{g mg}^{-1}$ DW and 3.1%), and epoxy-xanthophyll antheraxanthin (0.04 $\mu\text{g mg}^{-1}$ DW and 2.9%) (Supporting Information, Fig. S2 and Table 3). Capsorubin, a carotenoid exclusively found only in red *Capsicum* species as well as the above mentioned capsanthin, and yellow epoxy-xanthophyll violaxanthin contributed only to a minor level, being under the Limit of Detection (LOD). A number of unidentified compounds were also detected at different retention times, contributing to almost 30% of the total peak area.

Table 3. HPLC analysis of principal carotenoids in chili pepper oleoresin

Compounds	Rt	Content in Oleoresin ($\mu\text{g mg}^{-1}$ DW)	LOD (mg L^{-1})	Biological activity
Capsorubin	4.140	<LOD	0.01	
Violaxanthin	4.458	<LOD	0.06	Antioxidant ⁹¹
Capsanthin	5.023	0.22 \pm 0.02 (21.18)*	0.01	Antioxidant ⁹¹
Antheraxanthin	5.472	0.04 \pm 0.00 (4.05)*	0.01	Antioxidant ⁹¹
Lutein	6.129	0.03 \pm 0.00 (2.80)*	0.01	Antioxidant ⁹¹
Zeaxanthin	6.508	0.06 \pm 0.01 (6.32)*	0.01	Antioxidant ⁹¹
IS	8.390	-		
β -Cryptoxanthin	13.296	0.08 \pm 0.01 (7.76)*	0.03	Antioxidant ^{91,92}
β -Carotene	30.077	0.59 \pm 0.04 (57.44)*	0.02	Antimicrobial ^{33,91,93} Antioxidant ⁹⁴
Total Carotenoids		1.02 \pm 0.07 (0.1) [†]		

Note: Retention time (Rt), relative content, Limit of Detection (LOD) and biological activity of the principal carotenoids detected in chili pepper oleoresin after HPLC analysis in acetonitrile: methanol (40:60). IS, internal standard.
*Relative abundance (% wt) on total identified carotenoids.
[†] Relative abundance (% wt) on oleoresin DW basis.




Figure 1. Antifungal activity of chili pepper oleoresin. Log-logistic dose–response curve illustrating the growth inhibitory effect of increasing pepper oleoresin concentrations on radial growth of *G. bidwellii* (A) and *B. cinerea* (B) after 6 and 2 days of incubation at 21 °C, respectively and on sporulation on grapevine leaf discs for *P. viticola* (C). Data are expressed as mean \pm SD ($n = 3$). Lack-of-fit P value > 0.05 .

By comparison with their relative internal standards, HPLC analysis showed that the total amount of the main carotenoids accounted for a very limited concentration ($1.02 \mu\text{g mg}^{-1}$ DW) compared to capsaicinoids, contributing only to approximately 0.1% on oleoresin DW basis. This result was also confirmed by basic spectrophotometric method for carotenoid quantification on acetone-resuspended oleoresin, determining an almost similar amount of $0.692 \pm 0.074 \mu\text{g mg}^{-1}$ DW ($n = 3$). The most represented pigments of the yellow and red fraction of carotenoids were β -Carotene and capsanthin (57 and 21% of the total carotenoids), respectively.

Finally, the overall reducing power of chili pepper oleoresin was assessed, accounting for the average value of $3704.48 \pm 192.21 \mu\text{moles Trolox equivalents (TE) g}^{-1}$ DW extract (ORAC).

3.4 Antifungal activity of chili pepper oleoresin on grapevine pathogens

After chemical characterization, antifungal activity of chili pepper extract at different concentrations was tested by *in vitro* assays

against three grapevine pathogens (Fig. 1). Overall, the oleoresin exhibited significant inhibition activity against growth and sporulation of isolates from the hemibiotrophic ascomycete *G. bidwellii*, the necrotrophic *B. cinerea* and the biotrophic oomycete *P. viticola*, over a large range of concentrations, ranging from 0.001 to 12.5 mg mL⁻¹.

In particular, the inhibition of *G. bidwellii* mycelial growth on culture medium was measured at day 6, starting from sterilized 1 month-old plug, when it reached 89% at the highest dose of 12.5 mg mL^{-1} (Supporting Information, Fig. S3).

Further concentrations of the extract were not tested, due to difficulties in achieving homogenous dispersion of the oleoresin in water at higher concentrations and for the necessity to avoid excess of ethanol in the assay. With diluted different concentrations of oleoresin, it was possible to interpolate the curve shown in Fig. 1 panel A, and thus to calculate the concentration that inhibited 50% of *G. bidwellii* mycelium growth, *i.e.* ED₅₀ of $0.233 \pm 0.034 \text{ mg mL}^{-1}$.

In the case of *B. cinerea*, inhibition of radial mycelium growth starting from sporulating conidia at day 2 was measured at several dilutions of extract (Supporting Information, Fig. S4).

In this case, chili pepper extract almost completely inhibited the pathogen at a lower dose of 8.8 mg mL⁻¹, however exhibiting a lower ED₅₀ (0.445 ± 0.091 mg mL⁻¹) if compared to *G. bidwellii* (Fig. 1(B)). This difference could be related to differential sensitivities and/or detoxifying capacities possessed by the two fungal pathogens or their specific growth stages towards the bioactive compounds present in oleoresin extract.

Similarly, as for the previous pathogens, oleoresin was also effective in reducing the sporulation and evasion of the obligate biotrophic *P. viticola*, reaching 100% of inhibition at 12.5 mg mL⁻¹ (Supporting Information, Fig. S5). Conversely, it seemed that its efficacy was less strong at lower doses, since higher value of ED₅₀ of 1.535 ± 0.157 mg mL⁻¹ was reported (Fig. 1(C)). It has to be noted that, even at the highest dosage, no apparent symptoms of phytotoxicity of pepper oleoresin on leaf discs were observed.

Overall, these findings could be very promising for the high efficacy shown by chili pepper oleoresin over a range of fungal diseases which pose increasing concern and challenges for grapevine pest sustainable management. Among the pathogens tested, the oleoresin could be particularly valuable for its potential bioactivity against *G. bidwellii*, since it showed the highest efficacy in terms of ED₅₀ and it could be used in biocontrol of this new emerging disease.

4 DISCUSSION

4.1 Extraction of crude chili pepper oleoresin by eco-safe solvent

Reduction of fungicide dosage and use of conventional pesticides in vineyards is now recognized as a priority by government authorities, which encourage the development and adoption of sustainable alternatives. Besides agricultural practices and cultivation of resistant genotypes for limiting the risks of fungal pathogens spreading, the implementation of alternative treatments, including resistance inducers and biofungicides of both botanical^{95,96} or bacterial origin,⁹⁷ is additionally required.

In this regard, chili peppers are rich in different polar bioactive compounds and several solvents could be used to extract them efficiently.^{64,65} Therefore, a preliminary issue has been the best choice of solvent for oleoresin preparation. Accordingly, previous reports highlighted that solvent chemical properties such as polarity may determine variations on content of capsaicinoids, carotenoids and phenolic compounds,⁶⁵ and they influenced also chili pepper oleoresin antioxidant activity since the latter was strongly correlated with the content of bioactive compounds.⁶⁴ As mid-polar solvent, ethyl acetate has been still utilized in oleoresin extraction from hot and non-pungent peppers, attaining the highest levels of total capsaicinoids in comparison to other organic solvents, better than or following only to what resulted by extraction with hexane.^{65,98} Lastly, considering that capsaicinoids are partially soluble in hexane and avoiding the use of highly toxic and inflammable solvents such as methanol and acetone, ethyl acetate was considered to be the best compromise for obtaining a 'green' hot chili pepper oleoresin extract with high levels of bioactive molecules. In accordance, ethyl acetate is scored as recommended in guidelines for solvent selection, given that it can be sourced renewably.⁹⁹

4.2 Bioactive compounds of chili pepper oleoresin and antioxidant potential

From the results of this study, the oleoresin achieved from a mixture of hot pungent varieties of *C. chinense* fruits contained particularly high amounts of capsaicinoids and polyphenols (371 and 268 µg mg⁻¹ DW, respectively, see Tables 1 and 2). In comparison, in fresh whole fruits from 18 different accessions of Habanero peppers the concentration of acetonitrile-extracted capsaicinoids ranged from 10 to 60 mg g⁻¹,¹⁰⁰ similarly to the average content found in dry fruits.¹⁰¹ Moreover, *C. chinense* has several times higher capsaicinoid content and hottest pungency compared to other *Capsicum* spp.,¹⁰² being 'Carolina Reaper' the variety with the highest capsaicinoid content at 73.34 mg g⁻¹ DW by methanolic extraction.⁴⁹

Concerning polyphenols, *C. chinense* displayed on average 1350 mg 100 g⁻¹ DW of total phenolic content by colorimetric method in whole dry fruits,⁴⁹ being even higher if placenta was considered with respect to other parts of the pericarp.¹⁰³ Interestingly, several authors stated a strong correlation between high polyphenols and capsaicinoids in *Capsicum* sp.,^{49,104,105} which was mainly explained for that the phenylpropanoid and capsaicinoid biosynthetic pathways are converging during pepper fruit ripening.^{104,106}

Carotenoids represent the third class of important bioactive metabolites we have investigated in this study, though in terms of abundance it remained a minor fraction, reaching only 1.02 µg mg⁻¹ on DW basis (Table 3). Nonetheless, they contribute, together with capsaicinoids and polyphenols, to the total antioxidant potential of chili pepper oleoresins.^{105,107} Wahyuni *et al.*^{24,108} reported an average total carotenoid content of approximately 40 mg 100 g⁻¹ fresh weight (FW) in different *Capsicum* accessions, whereas others found large variation for carotenoid concentration and composition.^{109,110} In dry fruit, carotenoids content ranged from 0.1 to 3.2 g 100 g⁻¹.²⁸ Chili peppers, as 'Habanero' and 'Naga Morich', were also particularly rich in β-carotene,¹⁰⁷ which explained the major contribution of this yellow pigment on the oleoresin quantified in the present study, followed by capsanthin, characteristic of *Capsicum* sp., which generally contributes to a large fraction of carotenoids in most of the varieties.²⁸

In this study, all the previous major phytochemicals mentioned may contribute to the total antioxidant activity of the oleoresin with a mean ORAC value of 370 448 ± 19 221 µmoles TE 100 g⁻¹ DW, which is very similar to or even higher than the values reported for cayenne pepper^{111,112} and quite lower than *C. frutescens* extracts.¹¹³ A reliable comparison on literature reports is however very difficult to be achieved, due to high variation in the ORAC values in reliance on *Capsicum* variety, origin, fruit processing and solvent used for product extraction. Moreover, it has to be underlined that the oleoresin used in the present study was originating from an assortment of seven different varieties of *C. chinense*, whose particular contribution to the total antioxidant value of the oleoresin is difficult to disentangle. Nonetheless, it is proposed that a significant component of the antioxidant activity in the chili pepper oleoresin used in this study could be attributed mainly to the large abundance of capsaicinoids and polyphenols, given that this activity in *Capsicum* has been positively related to the latter compounds.^{65,103,104,114,115}

4.3 Antimicrobial activity of chili pepper oleoresin

Apart from the high antioxidant value of the oleoresin, the results of *in vitro* pathogen growth assays evidenced that the extract is

effective, at different levels, on mycelial growth or sporulation of three important pathogens of grapevine (Fig. 1). In particular, comparison of dose–response curve patterns and ED₅₀ of pathogens considered highlights that the highest doses of oleoresin (in the range of 4.5–12.5 mg mL⁻¹) did not cause a 100% inhibition of mycelial growth in the case of black rot agent, *G. bidwellii*, differently from what observed for *P. viticola* and *B. cinerea*. This feature might suggest that oleoresin exerted more a fungistatic than a fungitoxic activity on this Ascomycete. Hence, using a pharmacology definition, the oleoresin exhibited a higher ‘efficacy’ in inhibiting *P. viticola* and *B. cinerea* mycelial growth (potential fungitoxic action) at higher concentrations, but a stronger ‘potency’ against *G. bidwellii* since the ED₅₀ is reached at a lower dose.¹¹⁶ However, this fungus appeared to be particularly sensitive at lesser concentrations of chili pepper extract, being its radial mycelial growth decreased by 25% even at a low dose of 0.05 mg mL⁻¹. This observation was confirmed by the lowest calculated value of ED₅₀ (0.233 mg mL⁻¹), which doubled in *B. cinerea*, followed by the far higher level assessed for *P. viticola* sporulation.

To the best of our knowledge, this is the first report of fungicidal effect by *Capsicum*-based extract on the black rot of grapevine, since there is no available information from literature about specific fungicidal efficacy by the specific capsaicinoids or phenolic compounds we have identified so far in the chili pepper oleoresin. In fact, in searching of natural alternatives for organic wine-production, black rot agent was only shown to be sensitive to saponin-containing botanicals on *in vitro* experiments, where a high level of inhibition of conidial germination was provided by *Primula* and *Hedera helix* root extract and other species at concentrations below 0.5% (w/v aqueous or ethanolic solution).¹³ Moreover, both protective and curative efficacy of the extract was also demonstrated by greenhouse tests on potted vines under controlled conditions in the same study. The bioactive compounds mainly responsible for inhibition of conidia germination are ascribed to saponins, which unfortunately exhibited poor efficacy under field conditions, due to the high water-solubility of their formulation. Detergent properties of saponins are believed to explain the possible lytic and fungicidal action on fungal spores.¹¹⁷ Saponin identification in the pepper oleoresin was out of the scope of the present study, however, interestingly, Lucca et al.¹¹⁸ isolated a potent fungicidal saponin from dry ground *Capsicum* sp. fruit, effective against *Aspergillus* sp. germinating conidia.

Therefore, without any additional information, we could only suppose that the most abundant compounds detected in oleoresin, i.e., capsaicinoids (capsaicin and dihydrocapsaicin) and hydroxycinnamic acids (mainly vanillic) and protocatechuic acid among polyphenols, might be responsible for fungicidal action on *G. bidwellii*, although we cannot exclude *a priori* that also other minor bioactive components could contribute in a concerted manner with the above mentioned phytochemicals.⁸⁵

Regarding the efficacy of the oleoresin on conidia sporulation and radial mycelial growth of the necrotrophic agent *B. cinerea*, the pathogen was found to be susceptible to increasing oleoresin doses, showing a quite low value of ED₅₀ and displaying maximal activity at just the concentration of 8.8 mg mL⁻¹. Similarly, also *P. viticola* sporulation was inhibited completely by pepper oleoresin, without any evident phyto-toxicity, though at higher concentrations it showed quite high ED₅₀ value. In accordance, some authors reported that *B. cinerea* and some oomycetes species were highly susceptible to capsaicinoids,^{39,41,43} suggesting that the lateral chain of capsaicinoids compared to the phenolic

one could explain high fungal inhibition, due to osmotic stress and phospholipid membrane damage.^{43,119} Moreover, several data from literature are consistent with the efficacy of phenolic acids, and specifically ferulic and vanillic acid, in inhibiting filamentous fungi.⁸⁶ Protocatechuic acid, from both bacteria or plant origin, displayed strong inhibition against some major post-harvest pathogenic fungi as well as on *B. cinerea* colonization of strawberry fruits, probably by causing fungus membrane permeability impairment.⁸⁷

Interestingly, GC–MS analysis of chili pepper oleoresin run in dichloromethane has identified other minor bioactive components, known from literature to exert some antimicrobial and antifungal activities apart from capsaicinoids, such as essential oils (Table 1). In particular, among volatile compounds, nerolidol displayed remarkable antifungal action against most important phytopathogens (*Fusarium*, *Phytophthora*, *Colletotrichum*, *Alternaria*, *Rhizopus*, *Penicillium*, *Sclerotium* and *Rhizoctonia* species).^{76,77,120} This volatile compound (3,7,11-trimethyl-1,6,10-dodecatrien-3-ol) is a sesquiterpene naturally present in many plant species, with large application in food and cosmetic industries.¹²¹ In this regard, the single compound and its derivatives displayed bactericidal and antifungal properties: in particular, *trans,trans*-farnesol against *Phomopsis obscura*⁷⁸ and nerolidol are effective against *in vitro* *Sclerotium cepivorum* growth, ranging from 2.0 to 5.0 µg per disc.^{77,79} Although very raw, a comparison among these data and our results allows evidence that at the highest dosage of 12.5 mg mL⁻¹ of oleoresin used against *G. bidwellii*, the ascomycete encountered approximately 1.68 µg per fungal disc, giving a very plausible contribution to the oleoresin antifungal efficacy. They reduced sclerotia formation and caused strong fungal growth inhibition by alterations of mycelial morphology and membrane permeability, possibly through their insertion into lipid membranes due to their high hydrophobicity. Consistently, a number of published studies have demonstrated that terpene synthase activity increased and monoterpene and sesquiterpene accumulation took place in response to pathogen's attack^{80,122} and recently (*E*)-nerolidol has been reported to act also as a volatile signal enhancing the natural plant's immune system in tea plant.⁸¹ Taken together, all of this evidence might suggest that also some other bioactive compounds, although represented in minor quantity in the oleoresin, may be related to the high antifungal potential exerted against the three pathogens of grapevine. Furthermore, the co-presence of several bioactive compounds with high antioxidative properties in the complex mixture of the oleoresin might even represent an advantage if compared to the use of a single, isolated active compound. For example, the industrial utilization of capsaicin has been limited by its low water solubility shown by one-single formulation and by its degradation under extreme environmental conditions, thus requiring its incorporation into specific matrices or nanoformulates.^{123,124}

5 CONCLUSION

These results, though preliminary, demonstrated that the oleoresin from chili peppers is able to inhibit the growth of some of the most harmful pathogens in grapevine. At least, capsaicinoids and phenolic acids found at high content, associated to other biologically active components are all likely responsible for the activity against all three phytopathogens investigated.

Noteworthy, the high potential shown by the pepper oleoresin to specifically contrast *G. bidwellii*, the causal agent of black rot of grapevine, may deserve more investigations to produce it as new alternative or complementary bio-fungicide. The diffusion

of the latter disease has recently emerged as one of the major concerns in modern viticulture, due to the combination of restricted limitations in extensive use of copper by EU and introduction of mildew-resistant hybrids in organic vineyard, making it more vulnerable to infections from secondary fungal pathogens and ongoing yield losses. Nonetheless, it is still required to gain more information about the mechanism of action of this extract and to further explore the suitability of this material in vineyard, as well as to improve and achieve the best formulation for its application to vine plant.

Regarding pepper oleoresin manufacturing for pest management, it has to be said that the increase in the total dry pepper production by 30% during the last decade (2011–2021) (FAOSTAT 2022) is an attempted response to the enhanced request for its wide industrial applications in pharmaceutical and cosmetic sectors. Hence, the implementation in industrial production of habanero peppers, together with the development of new breeding programs of F1 hybrids very rich in bioactive compounds,¹²⁵ could really make the oleoresin production more sustainable, answering to the required necessity of new eco-friendly products for pest management. Other technological approaches aiming at increasing the sustainability of the pepper oleoresin application would be the exploration for the optimized industrial size process such as carbon dioxide supercritical extraction systems¹²⁶ or the use of chili pepper by-products such as discarded vegetative parts, peduncles⁸⁹ or fruit waste generated by seed production (coat debris, small seeds.) as source material, in the framework of a circular economy strategy.

ACKNOWLEDGMENTS

This work was supported by the Bordeaux Metabolome Facility and the MetaboHUB (ANR11-INBS-0010) project. Open Access Funding provided by Università degli Studi di Udine within the CRUI-CARE Agreement.

CONFLICT OF INTEREST STATEMENT

All the authors confirmed that there is no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

SUPPORTING INFORMATION

Supporting information may be found in the online version of this article.

REFERENCES

- 1 FAO. Land use statistics and indicators. *Land Use Stat Indic Stat Glob Reg Ctry Trends* **28**:1990–2019 (2021).
- 2 Pertot I, Caffi T, Rossi V, Mugnai L, Hoffmann C, Grando MS *et al.*, A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for the implementation of IPM in viticulture. *Crop Prot* **97**:70–84 (2017).
- 3 Pirrello C, Mizzotti C, Tomazetti TC, Colombo M, Bettinelli P, Prodanutti D *et al.*, Emergent Ascomycetes in viticulture: an interdisciplinary overview. *Front Plant Sci* **10**:1394–1423 (2019).
- 4 Aziz A, Trotel-Aziz P, Dhuciq L, Jeandet P, Couderchet M and Vernet G, Chitosan oligomers and copper sulfate induce grapevine defense reactions and resistance to gray mold and downy mildew. *Phytopathology* **96**:1188–1194 (2006).
- 5 Jacometti MA, Wratten SD and Walter M, Review: alternatives to synthetic fungicides for *Botrytis cinerea* management in vineyards. *Aust J Grape Wine Res* **16**:154–172 (2010).
- 6 Bavaresco L, Squeri C and Vercesi A, Field evaluation of new plant protection products against *Plasmopara viticola*. *BIO Web Conf* **12**:01007 (2019).
- 7 Gabaston J, Richard T, Cluzet S, Palos Pinto A, Dufour M-C, Corio-Costet M-F *et al.*, Pinus pinaster knot: a source of polyphenols against *Plasmopara viticola*. *J Agric Food Chem* **65**:8884–8891 (2017).
- 8 EU 2018/1981, Commission Implementing Regulation (EU) 2018/1981—of 13 December 2018—renewing the approval of the active substances copper compounds, as candidates for substitution, in accordance with Regulation (EC) No 1107 / 2009 of the European Parliament and of the Council concerning the placing of plant protection products on the market, and amending the Annex to Commission Implementing Regulation (EU) No 540/2011, 5 (2018).
- 9 Farm to Fork Strategy, Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. A Farm to Fork Strategy for a fair, healthy and environmentally-friendly food system (2020).
- 10 EUR-Lex—2021/2117, <https://eur-lex.europa.eu/eli/reg/2021/2117/oj> [13 June 2022].
- 11 Pedneault K and Provost C, Fungus resistant grape varieties as a suitable alternative for organic wine production: benefits, limits, and challenges. *Sci Hortic* **208**:57–77 (2016).
- 12 Wicht B, Petrini O, Jermini M, Gessler C and Broggini GAL, Molecular, proteomic and morphological characterization of the ascomycete *Guignardia bidwellii*, agent of grape black rot: a polyphasic approach to fungal identification. *Mycologia* **104**:1036–1045 (2012).
- 13 Koch E, Enders M, Ullrich C, Molitor D and Berkelmann-Löhnertz B, Effect of *Primula* root and other plant extracts on infection structure formation of *Phyllosticta ampellicida* (asexual stage of *Guignardia bidwellii*) and on black rot disease of grapevine in the greenhouse. *J Plant Dis Prot* **120**:26–33 (2013).
- 14 Malheiro A, Santos J, Fraga H and Pinto J, Climate change scenarios applied to viticultural zoning in Europe. *Climate Res* **43**:163–177 (2010).
- 15 Rex F, Fechter I, Hausmann L and Töpfer R, QTL mapping of black rot (*Guignardia bidwellii*) resistance in the grapevine rootstock “Börner” (*V. riparia* Gm183 × *V. cinerea* Arnold). *Theor Appl Genet* **127**:1667–1677 (2014).
- 16 Dagostin S, Schärer H-J, Pertot I and Tamm L, Are there alternatives to copper for controlling grapevine downy mildew in organic viticulture? *Crop Prot* **30**:776–788 (2011).
- 17 Yoon M-Y, Cha B and Kim J-C, Recent trends in studies on botanical fungicides in agriculture. *Plant Pathol J* **29**:1–9 (2013).
- 18 Castillo F, Hernández D, Gallegos G, Rodríguez R and Aguilar CN, *Antifungal Properties of Bioactive Compounds from Plants, Fungicides for Plant and Animal Diseases*, ed. by Dhanasekaran. InTech (2012). ISBN: 978-953-307-804-5
- 19 Gahukar RT, Evaluation of plant-derived products against pests and diseases of medicinal plants: a review. *Crop Prot* **42**:202–209 (2012).
- 20 Gabaston J, Cantos-Villar E, Biais B, Waffo-Teguo P, Renouf E, Corio-Costet M-F *et al.*, Stilbenes from *Vitis vinifera* L. waste: a sustainable tool for controlling *Plasmopara viticola*. *J. Agric. Food Chem* **65**: 2711–2718 (2017). <https://doi.org/10.1021/acs.jafc.7b00241>.
- 21 Rejeb IB, Pastor V and Mauch-Mani B, Plant responses to simultaneous biotic and abiotic stress: molecular mechanisms. *Plants* **3**: 458–475 (2014).
- 22 Akula R and Ravishankar GA, Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal Behav* **6**:1720–1731 (2011).
- 23 Wink M, Plant secondary metabolites modulate insect behavior—steps toward addiction? *Front Physiol* **9**:364–372 (2018).
- 24 Wahyuni Y, Ballester A-R, Sudarmonowati E, Bino RJ and Bovy AG, Secondary metabolites of *capsicum* species and their importance in the human diet. *J Nat Prod* **76**:783–793 (2013).
- 25 Chopan M and Littenberg B, The association of hot red chili pepper consumption and mortality: a large population-based cohort study. *PLoS One* **12**:e0169876 (2017).
- 26 Saleh BK, Omer A and Teweldemedhin B, Medicinal uses and health benefits of chili pepper (*capsicum* spp.): a review. *MOJ Food Process Technol* **6**:325–328 (2018).
- 27 Duranova H, Valkova V and Gabriny L, Chili peppers (*capsicum* spp.): the spice not only for cuisine purposes: an update on current knowledge. *Phytochem Rev* **21**:1379–1413 (2021).
- 28 Arimboor R, Natarajan RB, Menon KR, Chandrasekhar LP and Moorkoth V, Red pepper (*Capsicum annum*) carotenoids as a

- source of natural food colors: analysis and stability—a review. *J Food Sci Technol* **52**:1258–1271 (2015).
- 29 Antonio AS, Wiedemann LSM and Veiga Junior VF, The genus *capsicum*: a phytochemical review of bioactive secondary metabolites. *RSC Adv* **8**:25767–25784 (2018).
 - 30 Lu M, Ho C-T and Huang Q, Extraction, bioavailability, and bioefficacy of capsaicinoids. *J Food Drug Anal* **25**:27–36 (2017).
 - 31 Bautista-Baños S, DeLucca AJ and Wilson C, Evaluation of the antifungal activity of natural compounds to reduce postharvest blue mould of apples during storage. *Mex J Phytopathol* **22**:362–369 (2004).
 - 32 Soumya SL and Nair BR, Antifungal efficacy of *Capsicum frutescens* L extracts against some prevalent fungal strains associated with groundnut storage. **12** (2012).
 - 33 Hayashi M, Nakukool S, Hayakawa S, Ogawa M and Ni'matulah A-BA, Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and β -carotene. *Food Chem* **130**:541–546 (2012).
 - 34 Bello I, Boboye BE and Akinyosoye FA, Phytochemical screening and antibacterial properties of selected Nigerian long pepper (*Capsicum frutescens*) fruits. *Afr J Microbiol Res* **9**:2067–2078 (2015).
 - 35 Levono and Prasad M, GC-MS profiling of capsaicinoids present in *Capsicum chinense* Jacq. cv. (Naga king Chili) and evaluation of its antifungal activity. *Asian J Chem* **29**:2674–2678 (2017).
 - 36 Melgar-Lalanne G, Hernández-Álvarez AJ, Jiménez-Fernández M and Azuara E, Oleoresins from *capsicum* spp.: extraction methods and bioactivity. *Food Bioprocess Technol* **10**:51–76 (2017).
 - 37 Ahmad R, Alqathama A, Aldholmi M, Riaz M, Abdalla AN, Mostafa A *et al.*, Gas chromatography-mass spectrometry (GC-MS) metabolites profiling and biological activities of various *Capsicum annuum* cultivars. *Plan Theory* **11**:1022 (2022).
 - 38 Koleva-Gudeva L, Mitrev S, Maksimova V and Spasov D, Content of capsaicin extracted from hot pepper (*Capsicum annuum* ssp. *microcarpum* L.) and its use as an ecopesticide. *Hem Ind* **67**:671–675 (2013).
 - 39 Wilson CL, Solar JM, El Ghaouth A and Wisniewski ME, Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*. *Plant Dis* **81**:204–210 (1997). <https://doi.org/10.1094/PDIS.1997.81.2.204>.
 - 40 Tewksbury JJ, Reagan KM, Machnicki NJ, Carlo TA, Haak DC, Peñañoza ALC *et al.*, Evolutionary ecology of pungency in wild chilies. *Proc Natl Acad Sci* **105**:11808–11811 (2008).
 - 41 Veloso J, Prego C, Varela MM, Carballeira R, Bernal A, Merino F *et al.*, Properties of capsaicinoids for the control of fungi and oomycetes pathogenic to pepper. *Plant Biol Stuttg Ger* **16**:177–185 (2014).
 - 42 Rodríguez-Maturino A, Troncoso-Rojas R, Sánchez-Estrada A, González-Mendoza D, Ruiz-Sánchez E, Zamora-Bustillos R *et al.*, Efecto antifúngico de extractos fenólicos y de carotenoides de chiltepín (*Capsicum annuum* var. *glabrusculum* en *Alternaria alternata* y *Fusarium oxysporum*). *Rev Argent Microbiol* **47**:72–77 (2015).
 - 43 García T, Veloso J and Díaz J, Properties of vanillyl nonanoate for protection of pepper plants against *Phytophthora capsici* and *Botrytis cinerea*. *Eur J Plant Pathol* **150**:1091–1101 (2018).
 - 44 Song GC, Ryu SY, Kim YS, Lee JY, Choi JS and Ryu C-M, Elicitation of induced resistance against *Pectobacterium carotovorum* and *Pseudomonas syringae* by specific individual compounds derived from native Korean plant species. *Molecules* **18**:12877–12895 (2013).
 - 45 Malo I, De Bastiani M, Arevalo P and Bernacchia G, Natural extracts from pepper, wild rue and clove can activate defenses against pathogens in tomato plants. *Eur J Plant Pathol* **149**:89–101 (2017).
 - 46 Neumann RE, Capsicum based mildew killing solution and method of use US20100159042A1 (2010).
 - 47 Bosland PW, Coon D and Reeves G, “Trinidad Moruga scorpion” pepper is the world’s hottest measured Chile pepper by more than two million scoville heat units. *HortTechnology* **22**:534–538 (2012).
 - 48 Muñoz-Ramírez LS, Peña-Yam LP, Avilés-Viñas SA, Canto-Flick A, Guzmán-Antonio AA and Santana-Buzzy N, Behavior of the hottest chili peppers in the world cultivated in Yucatan, Mexico. *HortScience* **53**:1772–1775 (2018).
 - 49 Zamljen T, Jakopić J, Hudina M, Veberić R and Slatnar A, Influence of intra and inter species variation in chilies (*capsicum* spp.) on metabolite composition of three fruit segments. *Sci Rep* **11**:4932 (2021).
 - 50 Wesołowska A, Grzeszczuk M, and Jadczyk D, GC-MS analysis of essential oils isolated from fruits of chosen hot pepper (*Capsicum annuum* L.) cultivars, (2015).
 - 51 Antonious GF and Jarret RL, Screening capsicum accessions for Capsaicinoids content. *J Environ Sci Health, Part B* **41**:717–729 (2006).
 - 52 Wesołowska A, Jadczyk D, and Grzeszczuk M, Chemical composition of the pepper fruit extracts of hot cultivars *Capsicum annuum* L. (2011).
 - 53 Jeong WY, Jin JS, Cho YA, Lee JH, Park S, Jeong SW *et al.*, Determination of polyphenols in three *Capsicum annuum* L. (bell pepper) varieties using high-performance liquid chromatography-tandem mass spectrometry: their contribution to overall antioxidant and anticancer activity. *J Sep Sci* **34**:2967–2974 (2011).
 - 54 Loupit G, Prigent S, Franc C, De Revel G, Richard T, Cookson SJ *et al.*, Polyphenol profiles of just pruned grapevine canes from wild *Vitis* accessions and *Vitis vinifera* cultivars. *J Agric Food Chem* **68**:13397–13407 (2020).
 - 55 Su Q, Rowley K, Itsiopoulos C and O’Dea K, Identification and quantification of major carotenoids in selected components of the Mediterranean diet: green leafy vegetables, figs and olive oil. *Eur J Clin Nutr* **56**:1149–1154 (2002).
 - 56 Borguini RG, Pacheco S, Chávez DWH, Couto GA, Wilhelm AE, Santiago MCP d A *et al.*, Carotenoid extraction using edible vegetable oil: an enriched provitamin a product. *Plant Dis* **81**:204–210 (2020).
 - 57 Biehler E, Mayer F, Hoffmann L, Krause E and Bohn T, Comparison of 3 spectrophotometric methods for carotenoid determination in frequently consumed fruits and vegetables. *J Food Sci* **75**:C55–C61 (2010).
 - 58 Dávalos A, Gómez-Cordovés C and Bartolomé B, Extending applicability of the oxygen radical absorbance capacity (ORAC—fluorescein) assay. *J Agric Food Chem* **52**:48–54 (2003). <https://doi.org/10.1021/jf0305231>.
 - 59 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682 (2012).
 - 60 Corio-Costet M-F, Dufour M-C, Cigna J, Abadie P and Chen W-J, Diversity and fitness of *Plasmopara viticola* isolates resistant to QoI fungicides. *Eur J Plant Pathol* **129**:315–329 (2011).
 - 61 Peressotti E, Duchêne E, Merdinoglu D and Mestre P, A semi-automatic non-destructive method to quantify grapevine downy mildew sporulation. *J Microbiol Methods* **84**:265–271 (2011).
 - 62 Ritz C, Baty F, Streibig JC and Gerhard D, Dose-response analysis using R. *PLoS One* **10**:e0146021 (2015).
 - 63 Guidelines for accurate EC50/IC50 estimation—Sebaugh. *Pharmaceut Stat* **10**:128–134 (2011). <https://doi.org/10.1002/pst.426>.
 - 64 Bae H, Jayaprakasha GK, Jifon J and Patil BS, Variation of antioxidant activity and the levels of bioactive compounds in lipophilic and hydrophilic extracts from hot pepper (*capsicum* spp.) cultivars. *Food Chem* **134**:1912–1918 (2012).
 - 65 Bae H, Jayaprakasha GK, Crosby K, Jifon JL and Patil BS, Influence of extraction solvents on antioxidant activity and the content of bioactive compounds in non-pungent peppers. *Plant Foods Hum Nutr* **67**:120–128 (2012).
 - 66 Kang JR, Lee MK and Kang SM, Antioxidant property and tyrosinase inhibition activity of various extracts from plants in Compositae plants. **321–328**:321–328 (2008).
 - 67 Kim MK, Nam P-W, Lee S-J and Lee K-G, Antioxidant activities of volatile and non-volatile fractions of selected traditionally brewed Korean rice wines. *J Inst Brewing* **120**:537–542 (2014).
 - 68 Mohan Das N, Sivakama Sundari S, Karuppusamy S, Mohan VR and Parthipan B, GC-MS analysis of leaf and stem bark of *Cleidion nitidum* (Muell.-Arg.) Thw. ex Kurz. (Euphorbiaceae). *Asian J Pharm Clin Res* **7**:41–47 (2014).
 - 69 Ganesh M and Mohankumar M, Extraction and identification of bioactive components in *Sida cordata* (Burm.f.) using gas chromatography-mass spectrometry. *J Food Sci Technol* **54**:3082–3091 (2017).
 - 70 Sumathi BM and Uthayakumari F, GC MS Analysis of Leaves of *Jatropha maheswarii* Subram & Nayar. **7** (2014).
 - 71 Jawad Kadhim M, Mohammed GJ and Hadi Hameed I, *In vitro* antibacterial, antifungal and phytochemical analysis of methanolic extract of fruit *Cassia fistula*. *Orient J Chem* **32**:1329–1346 (2016).
 - 72 Ghathe N, Das A, Chaudhuri D, Panja S and Mandal N, Sundew plant, a potential source of anti-inflammatory agents, selectively induces G2/M arrest and apoptosis in MCF-7 cells through upregulation of p53 and Bax/Bcl-2 ratio. *Cell Death Discov* **2**:15062 (2016).
 - 73 Erukainure OL, Zaruwa MZ, Msaik AM, Muhammad A, Adoga JO, Ogunyemi IO *et al.*, Suppression of phagocytic oxidative burst, cytotoxic effect, and computational prediction of oral toxicity of dietary fatty acids of *Clerodendrum volubile* stem. *Comp Clin Pathol* **26**:663–671 (2017).

- 74 Hussein HM, Hameed IH and Ibraheem OA, Antimicrobial activity and spectral chemical analysis of methanolic leaves extract of *Adiantum capillus-veneris* using GC-MS and FT-IR spectroscopy. **17** (2016).
- 75 Rad JS, Alfatemi MH, Rad MS and Jyoti D, Phytochemical and antimicrobial evaluation of the essential oils and antioxidant activity of Aque www.ajadd.co.uk. *Am J Adv Drug Delivery* **1**:1–10 (2013).
- 76 Rahman A, Al-Reza SM and Kang SC, Antifungal activity of essential oil and extracts of *Piper chaba* Hunter against phytopathogenic fungi. *J Am Oil Chem Soc* **88**:573–579 (2011).
- 77 Pontin M, Bottini R, Burba JL and Piccoli P, *Allium sativum* produces terpenes with fungistatic properties in response to infection with *sclerotium cepivorum*. *Phytochemistry* **115**:152–160 (2015).
- 78 Krist S, Banovac D, Tabanca N, Wedge DE, Gochev VK, Wanner J *et al.*, Antimicrobial activity of nerolidol and its derivatives against airborne microbes and further biological activities. *Nat Prod Commun* **10**:143–148 (2015).
- 79 Chan W-K, Tan LT-H, Chan K-G, Lee L-H and Goh B-H, Nerolidol: a sesquiterpene alcohol with multi-faceted pharmacological and biological activities. *Molecules* **21**:529 (2016).
- 80 Escoriza G, García Lampasona S, Gomez Talquenca S and Piccoli P, *In vitro* plants of *Vitis vinifera* respond to infection with the fungus *Phaeoacremonium parasiticum* by synthesizing the phytoalexin nerolidol. *Plant Cell, Tissue Organ Cult* **138**:459–466 (2019).
- 81 Chen S, Zhang L, Cai X, Li X, Bian L, Luo Z *et al.*, (E)-Nerolidol is a volatile signal that induces defenses against insects and pathogens in tea plants. *Hortic Res* **7**:1–15 (2020).
- 82 Rosa A, Appendino G, Melis MP, Deiana M, Atzeri A, Alessandra I *et al.*, Protective effect and relation structure-activity of nonivamide and iododerivatives in several models of lipid oxidation. *Chem Biol Interact* **180**:183–192 (2009).
- 83 Mihailovi V, Vukovi N, Maškovi P and Stankovi MS, Studies on the antimicrobial activity and chemical composition of the essential oils and alcoholic extracts of *Gentiana asclepiadea* L. **11** (2011).
- 84 Khatua S, Pandey A and Biswas SJ, Phytochemical evaluation and antimicrobial properties of *Trichosanthes dioica* root extract. **4** (2016).
- 85 Lattanzio V, Lattanzino VMT and Cardinali A, Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. **23–67** (2006).
- 86 Martínez-Fraca J, de la Torre-Hernández ME, Meshoulam-Alamilla M and Plasencia J, In search of resistance against *Fusarium* Ear Rot: ferulic acid contents in maize pericarp are associated with antifungal activity and inhibition of fumonisin production. *Front Plant Sci* **13**: 852257–852267 (2022).
- 87 Nguyen XH, Naing KW, Lee YS, Moon JH, Lee JH and Kim KY, Isolation and characteristics of protocatechuic acid from *Paenibacillus elgii* HOA73 against *Botrytis cinerea* on strawberry fruits. *J Basic Microbiol* **55**:625–634 (2015).
- 88 Sudheeran PK, Ovidia R, Galsarker O, Maoz I, Sela N, Maurer D *et al.*, Glycosylated flavonoids: fruit's concealed antifungal arsenal. *New Phytol* **225**:1788–1798 (2020).
- 89 Al Aboody MS and Mickymaray S, Anti-fungal efficacy and mechanisms of flavonoids. *Antibiotics* **9**:45 (2020).
- 90 Joaquín-Ramos ADJ, López-Palestina CU, Pinedo-Espinoza JM, Altamirano-Romo SE, Santiago-Saenz YO, Aguirre-Mancilla CL *et al.*, Phenolic compounds, antioxidant properties and antifungal activity of jarilla (*Barkleyanthus salicifolius* [Kunth] H. Rob & Brettell). *Chil J Agric Res* **80**:352–360 (2020).
- 91 Kaulmann A and Bohn T, Carotenoids, inflammation, and oxidative stress—implications of cellular signaling pathways and relation to chronic disease prevention. *Nutr Res* **34**:907–929 (2014).
- 92 Wang L, Lu W, Li J, Hu J, Ding R, Lv M *et al.*, Optimization of ultrasonic-assisted extraction and purification of zeaxanthin and lutein in corn gluten meal. *Molecules* **24**:2994 (2019).
- 93 Abdulhadi SY, Gergees RN and Hasan GQ, Molecular identification, antioxidant efficacy of phenolic compounds, and antimicrobial activity of beta-carotene isolated from fruiting bodies of *Suillus* sp. *Karbala Int J Mod Sci* **6**:365–374 (2020).
- 94 Ghosh A, Hazra U and Dutta D, Role of β -cryptoxanthin as an antioxidant and its ability to bind with transferrin. *Int J Biosci Biochem Bioinf* **9**:258–264 (2019).
- 95 Schnee S, Queiroz EF, Voinesco F, Marcourt L, Dubuis P-H, Wolfender J-L *et al.*, *Vitis vinifera* canes, a new source of antifungal compounds against *Plasmopara viticola*, *Erysiphe necator*, and *Botrytis cinerea*. *J Agric Food Chem* **61**:5459–5467 (2013).
- 96 Thuerig B, James EE, Schärer H-J, Langat MK, Mulholland DA, Treutwein J *et al.*, Reducing copper use in the environment: the use of larixol and larixyl acetate to treat downy mildew caused by *Plasmopara viticola* in viticulture. *Pest Manage Sci* **74**:477–488 (2018).
- 97 Puopolo G, Cimmino A, Palmieri MC, Giovannini O, Evidente A and Pertot I, *Lysobacter capsici* AZ78 produces cyclo(L-Pro-L-Tyr), a 2,5-diketopiperazine with toxic activity against sporangia of *Phytophthora infestans* and *Plasmopara viticola*. *J Appl Microbiol* **117**: 1168–1180 (2014).
- 98 Restrepo Gallego M, Llanos Ríos N and Fonseca Echeverri CE, Composition of oleoresins from two kinds of chili pepper (habanero and tabasco) obtained by lixiviation with organic solvents. *Rev Lasallista Investig* **4**:14–19 (2007).
- 99 Byrne FP, Jin S, Paggiola G, Petchey THM, Clark JH, Farmer TJ *et al.*, Tools and techniques for solvent selection: green solvent selection guides. *Sustainable Chem Process* **4**:7 (2016).
- 100 Canto-Flick A, Balam-Uc E, Bello-Bello JJ, Lecona-Guzmán C, Solís-Marroquín D, Avilés-Viñas S *et al.*, Capsaicinoids content in Habanero pepper (*Capsicum chinense* Jacq.): hottest known cultivars. *HortScience* **43**:1344–1349 (2008).
- 101 Pino J, González M, Ceballos L, Centurión-Yah AR, Trujillo-Aguirre J, Latournerie-Moreno L *et al.*, Characterization of total capsaicinoids, colour and volatile compounds of Habanero chilli pepper (*Capsicum chinense* Jack.) cultivars grown in Yucatan. *Food Chem* **104**:1682–1686 (2007).
- 102 Guillen NG, Tito R and Mendoza NG, Capsaicinoids and pungency in *Capsicum chinense* and *Capsicum baccatum* fruits. *Pesqui Agropecuária Trop* **48**:237–244 (2018).
- 103 Castro-Concha LA, Tuyub-Che J, Moo-Mukul A, Vazquez-Flota FA and Miranda-Ham ML, Antioxidant capacity and total phenolic content in fruit tissues from accessions of *Capsicum chinense* Jacq. (Habanero pepper) at different stages of ripening. *Sci World J* **2014**:e809073 (2014).
- 104 Materska M and Perucka I, Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L.). *J Agric Food Chem* **53**:1750–1756 (2005).
- 105 Sora GTS, Haminiuk CWI, da Silva MV, Zielinski AAF, Gonçalves GA, Bracht A *et al.*, A comparative study of the capsaicinoid and phenolic contents and *in vitro* antioxidant activities of the peppers of the genus *Capsicum*: an application of chemometrics. *J Food Sci Technol* **52**:8086–8094 (2015).
- 106 Sukrasno N and Yeoman MM, Phenylpropanoid metabolism during growth and development of *Capsicum frutescens* fruits. *Phytochemistry* **32**:839–844 (1993).
- 107 Giuffrida D, Dugo P, Torre G, Bignardi C, Cavazza A, Corradini C *et al.*, Characterization of 12 *Capsicum* varieties by evaluation of their carotenoid profile and pungency determination. *Food Chem* **140**: 794–802 (2013).
- 108 Wahyuni Y, Ballester A-R, Sudarmonowati E, Bino RJ and Bovy AG, Metabolite biodiversity in pepper (*Capsicum*) fruits of thirty-two diverse accessions: variation in health-related compounds and implications for breeding. *Phytochemistry* **72**:1358–1370 (2011).
- 109 Acunha T d S, Crizel RL, Tavares IB, Barbieri RL, Pereira de Pereira CM, Rombaldi CV *et al.*, Bioactive compound variability in a Brazilian *Capsicum* pepper collection. *Crop Sci* **57**:1611–1623 (2017).
- 110 Padilha HKM, dos Pereira E, Munhoz PC, Vizzotto M, Valgas RA and Barbieri RL, Genetic variability for synthesis of bioactive compounds in peppers (*Capsicum annuum*) from Brazil. *Food Sci Technol* **35**:516–523 (2015).
- 111 Sinisgalli C, Faraone I, Vassallo A, Caddeo C, Bisaccia F, Armentano MF *et al.*, Phytochemical profile of *Capsicum annuum* L. cv Senise, incorporation into liposomes, and evaluation of cellular antioxidant activity. *Antioxidants* **9**:428 (2020).
- 112 Sudjaroen Y, Evaluation for nutritive values and antioxidant activities of Bang Changs Cayenne pepper (*Capsicum annuum* var. *acuminatum*). *Sci Res Essays* **9**:844–850 (2014).
- 113 Manikharda, Takahashi M, Arakaki M, Yonamine K, Asikin Y, Takara K *et al.*, Physical properties, flavor characteristics and antioxidant capacity of Shimatogarashi (*Capsicum frutescens*). *Food Sci Technol Res* **23**:427–435 (2017).
- 114 Sandoval-Castro CJ, Valdez-Morales M, Oomah BD, Gutiérrez-Dorado R, Medina-Godoy S and Espinosa-Alonso LG, Bioactive compounds and antioxidant activity in scalded Jalapeño pepper industrial byproduct (*Capsicum annuum*). *J Food Sci Technol* **54**:1999–2010 (2017).

- 115 Loizzo MR, Pugliese A, Bonesi M, Menichini F and Tundis R, Evaluation of chemical profile and antioxidant activity of twenty cultivars from *Capsicum annuum*, *Capsicum baccatum*, *Capsicum chacoense* and *Capsicum chinense*: a comparison between fresh and processed peppers. *LWT--Food Sci Technol* **64**:623–631 (2015).
- 116 Mwamula AO, Kabir MF and Lee D, A review of the potency of plant extracts and compounds from key families as an alternative to synthetic nematicides: history, efficacy, and current developments. *Plant Pathol J* **38**:53–77 (2022).
- 117 Segal R and Schlösser E, Role of glycosidases in the membranolytic, antifungal action of saponins. *Arch Microbiol* **104**:147–150 (1975).
- 118 Lucca AJ d, Bland JM, Vigo CB, Cushion M, Selitrennikoff CP, Peter J et al., CAY-1, a fungicidal saponin from *Capsicum* sp. fruit. *Med Mycol* **40**:131–137 (2002).
- 119 Kurita S, Kitagawa E, Kim C-H, Momose Y and Iwahashi H, Studies on the antimicrobial mechanisms of capsaicin using yeast DNA microarray. *Biosci Biotechnol Biochem* **66**:532–536 (2002).
- 120 Znini M, Cristofari G, Majidi L, El Harrak A, Paolini J and Costa J, In vitro antifungal activity and chemical composition of *Warionia saharae* essential oil against 3 apple phytopathogenic fungi. *Food Sci Biotechnol* **22**:113–119 (2013).
- 121 Lapczynski A, Bhatia SP, Letizia CS and Api AM, Fragrance material review on nerolidol (isomer unspecified). *Food Chem Toxicol* **46**:S247–S250 (2008).
- 122 Toffolatti SL, Maddalena G, Passera A, Casati P, Bianco PA and Quaglino F, Role of terpenes in plant defense to biotic stress, in *Biocontrol Agents and Secondary Metabolites*, ed. by Jogaiah S. Woodhead Publishing, Sawston, Cambridge, Ch. 16, pp. 401–417 (2021).
- 123 Kopec SE, DeBellis RJ and Irwin RS, Chemical analysis of freshly prepared and stored capsaicin solutions: implications for tussigenic challenges. *Pulm Pharmacol Ther* **15**:529–534 (2002).
- 124 Sánchez-Arreguin A, Carriles R, Ochoa-Alejo N, López MG and Sánchez-Segura L, Generation of BSA-capsaicin nanoparticles and their hormesis effect on the *Rhodotorula mucilaginosa* yeast. *Molecules* **24**:2800 (2019).
- 125 Barik S, Ponnampalnam N, Reddy AC, LR DC, Saha K, Acharya GC et al., Breeding peppers for industrial uses: progress and prospects. *Ind Crops Prod* **178**:114626 (2022).
- 126 Uribe JAR, Perez JIN and Mercado CAR, Cost estimation for CO₂ supercritical extraction systems and manufacturing cost for Habanero pepper. *J Supercrit Fluids* **93**:38–41 (2014).