



UNIVERSITÀ
DEGLI STUDI
DI UDINE

Università degli studi di Udine

Modulation of Extra Virgin Olive Oil Digestibility through Oleogelation

Original

Availability:

This version is available <http://hdl.handle.net/11390/1235164> since 2022-10-24T19:18:20Z

Publisher:

Published

DOI:

Terms of use:

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

Publisher copyright

(Article begins on next page)


Background



The development of a **SUSTAINABLE FOOD SYSTEM**, to favor the **TRANSITION TO HEALTHIER AND MORE SUSTAINABLE DIETS**, is one of the major challenges of the modern food industry [1].



The use of extra virgin olive oil (**EVOO**) as a **FUNCTIONAL INGREDIENT** would be particularly interesting due to its recognized health-promoting capacity [2]. However, the direct addition of EVOO to food is challenging due to its liquid state.




Liquid oil conversion into a solid-like material through **OLEOGELATION** could enlarge its possible applications, increasing the technological performances, while reducing hard stock fat content in food [3].

AIM


To understand the digestibility of EVOO triglycerides in oleogels obtained by different oleogelators

Material & Methods




MICROSCOPIC APPEARANCE

PLM showed **needle-like crystals** in MG, SW, and RW of different sizes. PS is not recordable with PLM being a fibrillar network.



MECHANICAL BEHAVIOUR


Both firmness and critical stress indicated that PS was the strongest gel, followed by SW, RW, and MG. These results were attributed to **the microstructure and the nature of the network**, i.e., fibrillar (PS) or crystalline (MG, RW, and SW).




Results & Discussion

DIGESTION BEHAVIOUR

The kinetics of FFAs released differed among samples. All oleogels presented a **lower lipid digestibility** compared to unstructured EVOO. Different structures led to changes in lipid digestibility: MG had the lowest while PS had the highest FFA release values. CLSM highlighted the effect of different gelators on the **formation of mixed micelles** upon intestinal in-vitro digestion. Larger oil droplets were observed in EVOO and PS, whereas MG, RW, and SW enabled the formation of smaller and more dispersed micelles.



Sample	FFA Released (%)
MG	42.88 ± 3.11
RW	50.72 ± 2.92
SW	50.86 ± 2.33
PS	59.10 ± 0.78
Liquid EVOO	67.90 ± 5.55



Conclusions

All **gelators** (MG, RW, SW, and PS) **successfully structured EVOO into oleogels** with peculiar physical characteristics. This can enlarge the possible applications of **EVOO in food formulations**.

The entrapment of liquid oil into oleogel networks based on different microstructures allowed **modulating FFA release** during in vitro digestion. Oleogels can thus represent a **promising strategy to tailor lipid digestibility**.

References

- [1] European Commission. *Farm to Fork Strategy*; 2020.
- [2] A. Romani et al., "Health effects of phenolic compounds found in extra virgin olive oil, by-products, and leaf of olea europaea L.," *Nutrients*, vol. 11, no. 8, 2019.
- [3] Marangoni AG, Garti N. *Edible Oleogels: Structure and Health Implications*. Elsevier Inc; 2011.
- [4] Brodkorb A, Egger L, Alminger M, et al. INFOGEST static in vitro simulation of gastrointestinal food digestion. *Nature Protocols*. 2019;14(4):991-1014.
- [5] Ahmed K, Li Y, McClements DJ, Xiao H. Nanoemulsion- and emulsion-based delivery systems for curcumin: Encapsulation and release properties. *Food Chemistry*. 2012;132(2):799-807.