

1 CBP manuscript 16469

2

3 **Geographical variation in thermal tolerance within Southern Ocean marine ectotherms.**

4

5 Simon A. Morley ^{a,*}, Timo Hirse ^b, Hans-Otto Pörtner ^b, Lloyd S. Peck ^a

6

7 ^a British Antarctic Survey, Natural Environment Research Council, Cambridge, CB30ET, UK

8 ^b Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 27515, Germany.

9

10 Corresponding author. Tel.: +44 1223 221554; fax: +441223 221256.

11 *E-mail address:* smor@bas.ac.uk (S.A. Morley).

12

13 Short title: geographic thermal tolerance

14

15 **Abstract**

16 Latitudinal comparisons of the Southern Ocean limpet, *Nacella concinna*, and clam, *Laternula*
17 *elliptica*, acclimated to 0.0°C, were used to assess differences in thermal response to two
18 regimes, 0.0, 5.1 to 10.0°C and 2.5, 7.5 to 12.5°C, raised at 5.0°C per week. At each temperature,
19 tissue energy status was measured through a combination of O₂ consumption, intracellular pH,
20 cCO₂, citrate synthase (CS) activity, organic acids (succinate, acetate, propionate), and adenylylates
21 (ATP, ADP, AMP, ITP, PLA (phospho-L-arginine)) and heart rate. *L. elliptica* from Signy
22 (60°S) and Rothera (67°S) had the same lethal limits (7.5-10.0°C), but small yet consistent
23 indicators (O₂, CS activity, PLA, ATP, organic acids and cCO₂) suggest that Rothera *L. elliptica*
24 had lower critical and *pejus* (=getting worse) limits than Signy *L. elliptica*. *N. concinna*, which
25 experiences a wider thermal regime, had higher lethal limits (10.0-12.5°C). However, at their
26 Northern geographic limit *N. concinna* living in a warmer environment (South Georgia, 54°S),
27 had a lower critical limit (5.1-10.0°C; O₂, PLA and organic acids) than Rothera and Signy *N.*
28 *concinna* (10.0-12.5°C). This lower limit indicates that South Georgia *N. concinna* have different
29 biochemical responses to temperatures close to their thermal limit, which may make them more
30 vulnerable to future warming trends.

31

32 **Key words:** *Nacella concinna*; *Leternula elliptica*; Southern Ocean; tissue biochemistry; *pejus*
33 limits; critical limits; lethal limits; latitudinal comparisons;

34

35 **1. Introduction**

36 Applying the principles of symmorphosis (Weibel, 2000) to the oxygen supply cascade
37 provides a conceptual framework to explain how temperature affects oxygen availability to
38 tissues, and hence tissue energy status. This has developed towards a unifying principle that the
39 physiological thermal limits of aquatic ectotherms are defined by oxygen availability, the
40 balance between supply and demand (Pörtner, 2002; Pörtner, 2006; Peck et al., 2007). Within
41 their optimal range ectotherms are expected to have the physiological flexibility to cope with
42 normally experienced seasonal variation and tissue energy status should be maximal. However,
43 in the face of extreme conditions, or towards range limits, aerobic scope is reduced, as more of
44 the available energy is required to cope with routine metabolic processes and oxygen supply
45 mechanisms become sub-optimal. There is growing evidence that the effects of thermal stress act
46 hierarchically, from high to low complexity, i.e. whole body processes are affected before
47 biochemical reactions and cells (Pörtner, 2002; Pörtner and Knust, 2007; Pörtner et al., 2007).
48 Improving our understanding of how functionality is lost and energy is partitioned between
49 processes will improve our understanding of the likely impact of climate change on species
50 distributions and resulting ecosystem shifts.

51 Thermal sensitivity can be assessed at a number of levels from whole animal activity
52 through to tissue energy status and finally mortality. Characterising changes in tissue
53 biochemistry has been developed as a test for *critical* thermal limits in aquatic ectotherms which
54 indicate the transition to anaerobic metabolism (*cf* Pörtner, 2002). Within the window bounded
55 by critical temperatures, an even narrower window is bordered by *pejus* limits (*pejus*: getting
56 worse) which indicate when temperatures move beyond the optimal range and functionality starts
57 to be lost, which is paralleled by a shortfall in oxygen supply and demand until critical limits are
58 reached and anaerobic metabolic pathways are recruited. Species tested this way include the
59 boreal freshwater gadoid, *Lota lota* (Hardewig et al., 2004), the worms, *Sipunculus nudus*
60 (Zielinski and Pörtner, 1996), several populations of *Arenicola marina* (Sommer et al., 1997),
61 and several Antarctic species (e.g. *Limopsis marionensis*; Pörtner et al., 1999) including both *L.*
62 *elliptica* (Peck et al., 2002) and *N. concinna* (Pörtner et al., 1999). Biochemical parameters were

63 selected that measure the integrity of cellular systems and identify the temperature at which there
64 is a departure from homeostasis (pHi and cCO₂), with the main focus on those that measure the
65 aerobic status of the animal, circulation (heart rate), aerobic metabolism (O₂ consumption and
66 citrate synthase activity), tissue energy status (balance of adenylates) and recruitment of
67 anaerobic pathways (build up of organic acids). We aim to utilise these techniques to compare
68 *pejus* limits, *critical* and *lethal* limits of *N. concinna* and *L. elliptica* from different locations
69 within the Antarctic. These measures will be used to look for adaptive differences between
70 locations that relate to experienced thermal environment.

71 We exploit access to a unique latitudinal gradient within the Southern Ocean, Rothera
72 (67°S), Signy (60°S) and South Georgia, (54°S) which enable us to separate latitude and
73 temperature (Barnes et al. 2006). Our highest latitude site at Rothera Point, Adelaide Island, has
74 an annual seawater temperature ranging from -2°C in winter to +1°C in the height of summer (at
75 15m depth; Barnes et al., 2006). Signy Island, within the South Orkney Islands (60°C) is on the
76 edge of the Weddell Sea gyre and despite its more northerly location has the same annual
77 seawater temperature range to that of Rothera (Clarke and Leakey, 1996). However the sub-
78 Antarctic Island of South Georgia (54°S), although still within the Southern Ocean (south of the
79 Antarctic circum-polar current) and therefore the biological Antarctic (Barnes et al., 2005), has
80 almost twice the annual range due to a higher summer maximum (-1 to +4°C at 10-20m; Barnes
81 et al., 2006). In response to the constant cold at Rothera benthic marine ectotherms are generally
82 stenothermal, they have a limited temperature range, upper lethal limits typically occurring
83 between 5 and 10°C (Peck, 2005) but for some species are considerably lower than these, e.g. the
84 brittle star *Ophionotus victoriae* which cannot survive even 1 month at 2°C (Peck et al, *in press*).
85 Furthermore, limits for critical activities are often considerably lower than the lethal limits (2 to
86 4°C; Peck et al., 2004). The +4°C maximum summer seawater temperature at South Georgia
87 therefore has the potential to impact these species, unless they have adaptive capacity to cope
88 with this thermal range. It is interesting to note that the range of *O. victoriae* does not stretch as
89 far as South Georgia.

90 Symmorphosis (Weibel, 2000) and the climate variability hypothesis (Stevens, 1989)
91 both predict that species, which experience a wider range of temperature, will have greater
92 thermal flexibility. From this, sub-littoral species living at South Georgia should be able to cope
93 with higher temperatures than the same species from Rothera. Habitat temperatures would also

94 predict that the thermal response of individuals from Signy would be closer to those from
95 Rothera than South Georgia. Although littoral species will experience a much higher range of
96 temperatures during low water emersion, habitat temperatures still predict the warmest
97 environment at South Georgia. The highest recorded littoral temperature is 12.3°C at Rothera
98 (Waller et al., 2006), 10.7°C at Signy (Barnes et al., 1996) and 15.8°C at South Georgia
99 (Davenport, 1997). The littoral limpet, *Nacella concinna* (Strebel 1908), would therefore be
100 expected to have a higher thermal capacity than the sub-littoral clam, *Laternula elliptica* (King
101 and Broderip 1831) and higher thermal limits in *N. concinna* and *L. elliptica* would be predicted
102 from South Georgia, than Signy and Rothera. However, species living at their range limit, where
103 they experience temperatures close to their maxima may have a reduced ability to cope with
104 further temperature increases (Tomanek, 2005; Deutsch et al., 2008); adaptations that enable
105 ectotherms to live in warmer environments may occur at the expense of further acclimatory
106 capacity (Stillman, 2003).

107 Through sampling these two species from 3 locations within the Southern Ocean, at the
108 same time of year, we aim to conduct the first latitudinal comparison of thermal tolerance within
109 the Southern Ocean. Environmental history determines thermal response (e.g.. Osovitz and
110 Hofmann, 2005) and so all animals were acclimated to the same temperature (0.0°C) and
111 maintained under as near identical conditions as possible for 2 months before thermal responses
112 were tested. Any differences in thermal response should therefore indicate adaptive differences
113 between locations.

114

115 **2. Materials and Methods**

116 *2.1. Collection and culture*

117 Limpets, *Nacella concinna* and the soft shelled clam, *Laternula elliptica* were collected
118 during cruise JR109 on the RRS James Clark Ross and concurrently at Rothera, the British
119 Antarctic Survey Research Station at Adeleide Island on Antarctic Peninsula, during early
120 summer as detailed in table 1. Except for *N. concinna* from Signy, which were collected from the
121 lowest part of the intertidal, all animals were collected by SCUBA divers.

122 After collection all animals were maintained in recirculating aquaria at $0.0\pm 0.3^{\circ}\text{C}$, for 63
123 days until the start of experiments. Water quality was maintained through biological filtration,
124 U.V. sterilisation, protein skimmers and daily water changes. During this period animals were

125 not fed to reduce any metabolic increment due to feeding (SDA), which can last up to 25 days in
126 Antarctic benthic ectotherms (Peck, 1998). Nonetheless, *N. concinna* were observed grazing on
127 biofilms on the tank walls and *L. elliptica* were seen with siphons open, so both species are likely
128 to have continued low level feeding.

129 At the start of incubation experiments animals were transferred to two 200 litre jacketed
130 acrylic tanks (Engineering Design and Plastics, Cambridge, Cambs.) attached to LTD20G
131 thermocirculators (Grant instruments Ltd, Shepreth, Cambs.), which maintained temperatures to
132 an accuracy of $\pm 0.2^{\circ}\text{C}$. The method was similar to that that used by Morley et al. (2007) and
133 Peck et al. (2002; 2004; 2007 and 2008). Animals were allowed 48 hours to recover from
134 movement stress before temperatures were altered or measurements taken. Temperature was then
135 raised at approximately 0.1°C per hour until the required temperatures were reached and then
136 animals were allowed 48 hours to recover from any metabolic overshoot (Peck et al., 2002). Due
137 to time constraints animals were separated into two groups, which were taken through different
138 5.0°C temperature steps, tank 1) 0.0 , 5.1 and 10.0°C , tank 2) 2.5 , 7.5 and 12.5°C . Following this
139 protocol animals were tested over approximately 3 weeks in each tank.

140

141 2.2. Oxygen consumption

142 Routine metabolic rate was measured in closed cell respirometers, following the
143 methodology of Peck (1989), except that oxygen concentration was measured with a Fibox-3
144 oxygen meter (Presens GmbH, Regensburg, Germany; e.g. Morley et al., 2007). Oxygen
145 sensitive foils were calibrated before each measurement using 5% w/w sodium dithionite for
146 0% and fully aerated water for 100%. During trials oxygen concentration was not allowed to fall
147 below 70% of air saturation, which is above the threshold for oxy-regulation for *L. elliptica*
148 (Peck et al., 2002). Two empty chambers (controls) were run with each trial to account for
149 background oxygen consumption, which was routinely less than 10% of the animal's
150 consumption. The basal metabolic rate of 4-5 individuals was measured for three consecutive
151 days after which data were analysed for significant differences in oxygen consumption between
152 days. When significant differences were found, daily measurements continued until three stable
153 readings were obtained and then temperature was raised. The volume of each animal was
154 measured through displacement and subtracted from the volume of water in the respirometer. At
155 the end of the experiment dry weight (constant weights at 60°C) and ash free dry weight (dry

156 weight minus the weight of ash remaining after ignition at 475°C for 24h) of each whole animal
157 was measured.

158 At each temperature the circulatory system of *L. elliptica* was monitored with the
159 impedance-based heart rate monitoring system (Buchan et al., 1988). Impedance systems record
160 the frequency of heart rate through variations in resistance between two electrodes. Methods
161 used were similar to those of Peck et al. (2002). Two 1-mm holes were drilled through the shell
162 either side of the heart, and 2mm of tinned 18 standard gauge copper wire electrodes, with the
163 coating removed from the tips, were inserted and glued in place with cyanoacrylate gel adhesive.
164 4 Signy and 4 Rothera *L. elliptica* had electrodes implanted in each of the two temperature tanks.

165 At the end of each temperature step the foot muscle from 4 animals was freeze clamped
166 in liquid nitrogen and stored at -80°C for biochemical analysis at the Alfred Wegener Institute in
167 Germany (except South Georgia and Signy *N. concinna* at 10°C for which only 3 animals were
168 available). Foot muscle was the chosen tissue as it performs a similar function in both *N.*
169 *concinna* and *L. elliptica*. Tissue was ground in a pestle and mortar, under liquid nitrogen, and a
170 number of extracts produced for biochemical analysis.

171 Temperature steps continued until 50% mortality occurred in the tanks. A further set of
172 20 *N. concinna* and *L. elliptica* from each location was maintained in the flow through aquarium,
173 at ambient temperature, for the duration of the experiment, with no mortalities.

174

175 2.3. Biochemical analysis

176 Intracellular pH (pH_i) and cCO_2 could only be measured in tissues of *L. elliptica* as *N.*
177 *concinna* foot muscle has large calcium carbonate stores, which would interfere with
178 homogenate acidity (Pörtner et al., 1999). Extracts were prepared in a media containing
179 160mmol.l^{-1} potassium fluoride and 1.0mmol.l^{-1} nitrilotriacetic acid following the methods of
180 Pörtner et al. (1990). pH_i was measured in the supernatant of the centrifuged homogenates at
181 animal incubation temperature, using needle type fibre optic pH sensors attached to a pH11 micro
182 pH meter (Presens GmbH, Regensburg, Germany) whose analogue output signal was boosted
183 and digitised through a Powerlab system (AD instruments). Due to the high sensitivity of the
184 optode system to solution ionic strength, KCl was added to each calibration buffer to match *L.*
185 *elliptica* tissue ionic strength of 0.16mol L^{-1} . Due to the small change in tissue pH, calibration
186 was conducted using three buffers covering the pH range 6.865 to 7.413 (at 25°C) whose pH

187 change with temperature had already been calculated. pH sensors were re-calibrated using
188 buffers cooled to each incubation temperature.

189 To measure $c\text{CO}_2$ 100 μl of this supernatant extract was injected into a 10ml sealed
190 sample vial, containing 1.5ml of nitrogen gassed HCl (0.05M). The sample was rotated for 10
191 minutes and then 50 μl of this gas phase was injected into an Agilent technologies 6890N
192 Network gas chromatogram through a G1888 network headspace sampler. $c\text{CO}_2$ was measured
193 with a HP-PLOTQ column and a TCD detector. Triplicate peak areas were converted to $c\text{CO}_2$ by
194 plotting a calibration curve for standard samples between 0.5 mM and 10.0 mM dissolved
195 inorganic carbon ($r^2 = 0.998$).

196 Tissue citrate synthase activity was measured from tissue extract at a ratio of 1g of tissue
197 to 9ml of extraction buffer, following the methods of Sidell et al. (1987). Enzyme activity was
198 measured at animal incubation temperature with a Beckman UV-DU 7400 spectrophotometer
199 fitted with a temperature controlled cuvette holder and diode array detector.

200 To measure both the build up of the end products of anaerobic metabolism (organic
201 acids) and tissue energy status (the adenylates) tissue homogenate was prepared using 0.6mol.l⁻¹
202 perchloric acid (PCA) (modified from Hardewig et al., 1998). Diluted PCA extracts were
203 injected into an ICS 2000 ion chromatography system fitted with a Gilson Diluter 401
204 autosampler. Organic acids were separated on an ion exclusion column (Dionex ICE-AS 1) at
205 60°C fitted with an AMMS-ICE 4mm micro-membrane suppressor cell, with 0.20mM
206 heptafluorobutyric acid (HFBA) as eluent at a flow rate of 1 ml.min⁻¹ and 5mM
207 tetrabutylammonium hydroxide (TBAOH) as the regenerant. The signal was detected on a DS6
208 conductivity detector. Duplicate sample peaks were converted into concentrations using
209 calibration curves with standard solutions, succinate (12 μM to 212 μM , $r^2 = 0.997$) acetate
210 (14 μM to 296 μM , $r^2 = 0.997$) and propionate (12 μM to 189 μM , $r^2 = 0.999$).

211 Adenylates were measured on a Beckman PACE/MDQ capillary electrophoresis system
212 following the adapted methods of Casey et al. (1999). The capillary was uncoated fused silica,
213 50 μm ID, 50 cm long. The pH of homogenates was set to 9.5, diluted in separation buffer
214 (40mM tetraborate buffer and 10mM sodium chloride (pH 9.7)) and 50 μl of 4 mM uric acid was
215 added to 200 μl of each standard and sample as an internal standard. The absorbance signals of
216 separation were simultaneously detected at different wavelengths with a photo diode array
217 detector: at 254 nm, adenosine, AMP, ATP, ADP and ITP; at 200 nm, arginine and PLA; at 290

218 nm, internal standard uric acid. Duplicate peaks were converted into concentrations using the
219 following calibration curves: adenosine (71.5 μM to 572 μM , $r^2 = 0.995$), AMP (70.5 μM to 564
220 μM , $r^2 = 0.988$), ATP (80.4 μM to 643 μM , $r^2 = 0.995$), ADP (71.3 μM to 570 μM , $r^2 = 0.966$),
221 ITP (70.2 μM to 561 μM , $r^2 = 0.992$), arginine (82.0 μM to 656 μM , $r^2 = 0.986$) and PLA (68.8
222 μM to 550 μM , $r^2 = 0.996$).

223

224 2.4. Analysis

225 Conducting the experiment in two tanks simultaneously created a mixed design where,
226 within tanks, some individuals were sampled more than once (e.g. for oxygen consumption), but
227 different individuals were sampled between tanks. As there was a small, non significant, effect of
228 individual (as a random factor in a GLM analysis) a simple analyses was conducted, ignoring
229 repeated observations on some individuals (P. Rothery, pers. coms.). ANOVAS with Tukey post-
230 hoc tests were used to test for differences between locations and temperatures for each species.
231 Where ANOVAS indicated only significant differences between locations a separate analysis
232 was conducted for each location. Significant differences was accepted as $P < 0.05$ throughout.
233 Oxygen consumption was standardised to that of a 0.373g dry mass *N. concinna* and a 10.6g dry
234 mass *L. elliptica* (the mean mass of animal used) using scaling coefficients of 0.82 (Fraser et al.,
235 2002) and 0.73 (Peck et al., 2002) respectively.

236

237 3. Results

238 3.1. Animal size

239 The shell length of *L. elliptica* sampled for biochemistry was not significantly different
240 between locations, table 2. However, due to logistic constraints on collections by SCUBA divers,
241 limpets from Rothera were smaller than those from Signy and South Georgia. *N. concinna*
242 collected inter-tidally from Signy were not significantly different in size than *N. concinna*
243 collected sub-tidally from South Georgia.

244

245 3.2. *L. elliptica*

246 More than 50% of *L. elliptica* died during the transition from 5.0 to 10°C and from 7.5 to
247 12.5°C so the experiment was stopped at this stage and the upper thermal limit recorded as
248 between 7.5 and 10.0°C. Although the sample size was too small to detect significant effects,

249 heart rate increased with temperature between 0.0 and 5.1°C. 50% of animals with implanted
250 impedance wires died at 7.5°C so heart rate was not measured at this temperature.

251 Temperature had a significant effect on every biochemical parameter, there were a few
252 significant differences between Rothera and Signy *L. elliptica* (Table 3) and some consistent
253 patterns that indicate differences between *L. elliptica* from the two locations. Although generally
254 not significant, at 0.0°C, Rothera *L. elliptica* had higher oxygen consumption, CS activity and
255 higher concentrations of high energy phosphates (PLA, ITP and ATP) than Signy *L. elliptica*
256 suggesting an overall higher metabolism at 0.0°C. Between 0.0 and 2.5°C there was also a, non-
257 significant, but consistently greater magnitude change in PLA and ATP (reduction), cCO₂
258 (increase) and organic acids (increase of succinate and acetate) in Rothera than Signy *L. elliptica*
259 (Fig 1, Table 3).

260 The oxygen consumption of *L. elliptica* rose between 0.0 and 2.5°C for specimens from
261 both sites, but no oxygen consumption was significantly different from that at 0.0°C. There was,
262 however, a significant drop in oxygen consumption between 2.5 and 7.5°C for both Signy and
263 Rothera *L. elliptica*. Citrate synthase activity at 2.5, 5.1 and 7.5°C was not significantly different
264 from that at 0.0°C. However, in a similar fashion to oxygen consumption, *L. elliptica* from both
265 locations had a peak in enzyme activity at 2.5°C, which was significantly higher than activity at
266 both 5.1 and 7.5°C.

267 The energy status of the high energy phosphates fell as the concentration of their low
268 energy forms, arginine, AMP and ADP, increased at 7.5°C in *L. elliptica* from both locations.
269 PLA the buffer that allows rapid rephosphorylation of ADP yielding ATP was also reduced at
270 7.5°C. The response of the triphosphates, ITP and ATP, to temperature was slightly different
271 between *L. elliptica* from Signy and Rothera. The ATP levels of Rothera *L. elliptica* were lower
272 at both 5.1 and 7.5°C than at 0.0°C, whilst the ATP levels of Signy *L. elliptica* were only
273 reduced at 7.5°C. The ITP concentration of Rothera *L. elliptica* generally fell upon warming but
274 was only significantly lower than 0.0°C at 7.5°C, whilst the ITP concentration of Signy *L.*
275 *elliptica* was lower at both 0.0°C and 7.5°C than 5.1°C.

276 The concentration of all three organic acids, succinate, acetate and propionate was
277 elevated at 2.5°C in Rothera *L. elliptica* and at 5.0°C in both, but was only significantly higher at
278 7.5°C. There were no significant differences in acetate and propionate accumulation patterns

279 between locations except that succinate rose to a significantly higher level in Signy than Rothera
280 *L. elliptica* at 7.5°C.

281 Intracellular pH was significantly higher at 0°C than at 2.5, 5.1 or 7.5°C for *L. elliptica*
282 from both locations. There was also a significant drop in pH_i between 5.1 and 7.5°C. Tissue
283 cCO₂ was highly variable and there was only one significantly elevated value, in Rothera *L.*
284 *elliptica* at 5.1°C.

285

286 3.3. *N. concinna*

287 More than 50% of *N. concinna* died during the transition from 7.5 to 12.5°C and 10.0 to
288 15.0°C so the experiment was stopped at this point. There was no significant effect of location or
289 temperature on arginine but there were complex, significant effects on the other adenylates, with
290 higher concentrations of both low (AMP and ADP) and high (ATP) energy adenylates at 10.0°C
291 (Fig. 2). At 10.0°C AMP was significantly higher in Rothera *N. concinna* than all treatments
292 except Signy *N. concinna* at 2.5°C and South Georgia at 10.0°C. At 10.0°C ADP was higher in
293 *N. concinna* from all locations. Although there was no significant difference in ATP
294 concentration from that at 0.0°C, at 10.0°C the concentration of ATP was higher at all locations
295 than at 2.5°C. However, despite this significant increase in ATP for all locations, the increase for
296 South Georgia *N. concinna* appeared to be less (Fig. 2). The PLA concentration of South Georgia
297 *N. concinna* was also lower at 10.0°C than at either 0.0 or 5.1°C.

298 Despite a general increase in the organic acids with increasing temperature, succinate,
299 acetate and proprionate were only significantly elevated in South Georgia *N. concinna* at 10.0°C
300 (Fig.3). The large inter-individual variation in organic acids at 10°C is typical of biochemical
301 responses around physiological limits as some individuals reach their limits before others. For *N.*
302 *concinna* from all locations, citrate synthase activity determined in tissue samples collected at
303 each temperature generally increased upon warming but was only significantly higher at 7.5 and
304 10.0°C (Fig.4).

305 Oxygen consumption was very different between *N. concinna* from different locations.
306 Oxygen consumption increased in a linear fashion with temperature (T) for Rothera ($O_2\text{cons} =$
307 $0.14T + 1.58$, $R^2=0.27$, $p<0.05$, $F=6.8$) and Signy ($O_2\text{cons} = 0.10T + 0.92$, $R^2=0.57$, $p<0.01$,
308 $F=22.9$) *N. concinna* (Fig 4). Oxygen consumption for South Georgia *N. concinna* reached a
309 similar maximum value but was best described by a curvilinear relationship ($O_2\text{cons} = 0.64T -$

310 $0.06T^2 + 0.76$, $R^2=0.45$, $p<0.01$, $F=6.6$). The calculated maximum (the point of inflexion of the
311 differential equation) occurred at 5.3°C , but the highest oxygen consumption was recorded at
312 2.5°C (Fig. 4).

313

314 **4. Discussion**

315 Measuring a suite of physiological and biochemical parameters gave a comprehensive
316 picture of how tissue energy status changed in response to a 5.0°C per week temperature
317 elevation. It also allowed comparisons of the responses of *L. elliptica*, and *N. concinna*, collected
318 from different geographic locations. As all animals were collected at the same time, were held
319 under nearly identical conditions for approximately 2 months before thermal sensitivity was
320 tested and they were tested over a maximum of 3 weeks, any consistent differences between
321 locations are most likely due differences in response between animals from these locations.

322

323 *4.1. L. elliptica*

324 The critical limit, which was confirmed by a drop in oxygen consumption, a drop in
325 energy status of the adenylates, an increase in anaerobic end products of metabolism (the organic
326 acids) and a drop in intracellular pH, occurred between 5.1 and 7.5°C , which is comparable to
327 that measured previously, over a similar rate of temperature change (6°C ; Peck et al., 2002).
328 Anaerobic metabolism is generally considered one of the protective mechanisms that allow time-
329 limited survival above *critical* temperatures (Pörtner, 2006). However, molluscs in particular can
330 have high anaerobic capacities that can be utilised to survive hypoxia for days or weeks,
331 particularly when accompanied by metabolic depression (e.g. Holmes and Miller, 2006; Morley
332 et al., 2007; Long et al., 2008). Reductions in citrate synthase activity and metabolic rate suggest
333 that thermal limitation may have started at a lower temperature, with *pejus* limits below 5.1°C .
334 Succinate and volatile fatty acid formation set in somewhat earlier in Rothera than the Signy *L.*
335 *elliptica*. This would indicate that, as a consequence of severe oxygen limitation, the critical
336 temperature is lower in the Rothera than the Signy population. The greater, but non-significant,
337 biochemical changes between 0.0 and 2.5°C for Rothera *L. elliptica*, coupled with significant
338 accumulation of carbon dioxide in the haemolymph, which shows that Rothera *L. elliptica* have a
339 lower capacity for gas exchange at 5.0°C (cf Peck et al., 2002), also suggest that *pejus* limits are
340 lower in Rothera than Signy *L. elliptica*. After 2 months at 0.0°C , there were also signs of

341 differences in cold acclimation of; CS activity, oxygen consumption and the turnover of high
342 energy phosphates were consistently (but not significantly) higher in Rothera than Signy *L.*
343 *elliptica* at 0.0 °C.

344 The lethal limit in the current study also occurred at a similar temperature, between 7.5
345 and 10.0°C (within 15 days), to that recorded previously (9°C within 25 days; Peck et al., 2002).
346 The similarity of lethal limits between Rothera and Signy *L. elliptica*, is consistent with the
347 hypothesis that thermal capacities will be matched to the experienced environment (both -2 to
348 +1°C), although the lower *pejus* limits of Rothera *L. elliptica* compared to Signy indicate subtle
349 differences in thermal tolerance between locations. Clearly, studies incorporating *L. elliptica*
350 from South Georgia, which is at the northern limit of its range, are needed to quantify how close
351 it is living to its upper thermal limits and therefore with reduced capacity to cope with further
352 temperature increases (Stillman, 2003). Such studies are crucial to our understanding of species
353 wide thermal capacity and therefore the ability of species to cope with environmental change.
354 Acclimations of *L. elliptica* at 3-4°C would indicate if South Georgia *L. elliptica* have
355 adaptations to cope with higher maximum experienced summer temperature.

356

357 4.2. *N. concinna*

358 Due to the constraints on geographical sampling within the Southern Ocean, there were
359 sampling differences between *N. concinna* collected at different locations. *N. concinna* were
360 collected from the bottom of the inter-tidal from Signy, compared to sub-tidal animals collected
361 from South Georgia and Rothera, however, all *N. concinna* were collected from depths shallower
362 than 6m. A recent study found small, non-consistent, genetic differences between inter-tidal and
363 15m *N. concinna* from Potter Cove, King George Island, but not between inter-tidal and 6m *N.*
364 *concinna* (de Aranzamendi et al., 2008). Different biochemical responses to emersion were also
365 evident between 15m and intertidal *N. concinna*, also from King George Island (Weihe and
366 Abele, *In Press*) was also evident This suggests that comparisons of 6m (Rothera), 2-3m (South
367 Georgia) and inter-tidal (Signy) *N. concinna* will not be affected by population differences
368 between animals collected from different depths. *N. concinna* collected in the middle of summer
369 and tested within days of collection had no significant ($t=0.57$, $p=0.57$) differences in upper
370 lethal thermal limits (1°C per day) between those collected in the inter-tidal (11.6°C) or sub-
371 tidal (6m, 10.8°C; SAM, unpublished data). The lack of any clear differences between Rothera

372 sub-tidal and Signy inter-tidal *N. concinna*, in the present study suggests that they respond
373 similarly to temperature, after a 2 month acclimation to 0.0°C.

374 Rothera *N. concinna* were also smaller than those from the other two locations, which
375 could introduce complicating factors due to growth and age differences as well as wider thermal
376 windows of small specimens (e.g. Pörtner 2002; Peck et al., 2007; Gsottbauer et al., 2007).
377 However, broad scale metabolic differences (see below), in the response to temperature, were
378 found between South Georgia and both Signy and Rothera *N. concinna*, which are unlikely to be
379 caused by size differences. Accepting these caveats, the present geographic comparisons within
380 the Southern Ocean, show for the first time specific responses of the adenylates, organic acids
381 and oxygen consumption to temperature, which suggest differences in critical limits and the
382 metabolic mechanisms employed to cope with acute temperatures close to these limits. There
383 was little evidence to suggest that Rothera or Signy *N. concinna* were temperature limited over
384 the range 0.0 to 10.0°C with measures of aerobic metabolism (oxygen consumption, citrate
385 synthase activity and ATP concentration) all increasing with temperature above 2.5°C. This
386 contrasts strongly with the pattern in *L. elliptica* which didn't show a reduction in these metrics
387 at temperatures above 2.5°C. This indicates they were suffering heat stress and have a narrower
388 thermal window than *N. concinna*.

389 In South Georgia *N. concinna*, citrate synthase activity and therefore the capacity of
390 aerobic metabolism in the mitochondria, increased with temperature in a similar fashion to
391 limpets from the other locations. However, in contrast to Rothera and Signy *N. concinna*, two
392 indicators of metabolism decreased in South Georgia *N. concinna* at higher temperatures.
393 Oxygen consumption peaked somewhere between 2.5 and 5.3°C and then declined whilst the
394 PLA pool, for buffering rapid ATP production, also declined above 5.1°C. Both of these suggest
395 a lower critical limit between 5.1 and 10.0°C (but perhaps as low as 2.5°C) for South Georgia *N.*
396 *concinna*, which was supported by the sharp increase in organic acids between 7.5 and 10.0°C
397 and therefore the recruitment of anaerobic pathways.

398 The wider thermal tolerance of *N. concinna* compared to *L. elliptica* is not surprising
399 considering it ranges into the intertidal and can therefore be exposed to a more variable thermal
400 environment. However, the difference in maximum littoral temperatures recorded so far, 12.3°C
401 at Rothera (Waller et al., 2006) 10.7°C at Signy (Barnes et al., submitted) and 15.8°C at South
402 Georgia (Davenport, 1997) was not reflected in any difference in the upper lethal limits for *N.*

403 *concinna* from different locations after 2 months acclimation to 0.0°C (10.0 to 12.5°C).
404 Davenport reported a short term survival limit of 15.6°C (12 hour exposure) for *N. concinna* at
405 South Georgia, higher than our medium term, 22 day, temperature limit of between 10.0 and
406 12.5°C. This is not unexpected as different thermal limits and protective mechanisms might be
407 expected if different rates of heating or cooling are used (Terblanche et al., 2007; Barnes et al.,
408 2008; Peck et al., *in press*). This difference may also have been because *N. concinna* tested by
409 Davenport were acclimated to summer conditions which could have raised their lethal limit
410 above that of animals acclimated to 0.0°C. However, it should also be noted that due to
411 behavioural adaptations, such as utilisation of cryptic habitats and evaporative cooling (Branch,
412 1981), body temperatures of limpets in the field are unlikely to reach these temperatures. The
413 maximum foot temperature of a Rothera *N. concinna* exposed on a sunny day has been shown to
414 be 8.8°C, although the average was much lower, 3.3 to 3.8°C, ranging from 1.5 to 9.1°C below
415 air temperature (Clark et al., 2008). If this relationship holds for South Georgia *N. concinna* then
416 foot temperature is unlikely to exceed their short-term (daily exposure) lethal limit during low
417 water emersion but might approach their medium term (weekly exposure) limit. Although *N.*
418 *concinna* are regularly exposed in the intertidal zone at Rothera and Signy, at South Georgia, it is
419 mainly a sub-tidal species (Davenport, 1997) and only rarely found above mean low water spring
420 tide level, reducing the risk of exposure to damaging high temperatures. The few inter-tidal
421 individuals found at South Georgia inhabit sheltered microhabitats and not exposed surfaces,
422 which are colonised by a siphonariid limpet, *Kerguelenella lateralis*. At its northern geographic
423 limit differences in habitat utilisation may be key to reducing the likelihood of *N. concinna* being
424 exposed to air temperatures close to its upper limit. They may therefore have evolved a high
425 anaerobic capacity to allow short-term, passive, survival during less frequent exposure (Pörtner,
426 2002). This fits with a growing body of evidence which shows that adaptation to life at range
427 limits may actually make animals less able to cope with further temperature increases (Stillman,
428 2003, Tomanek, 2005, Deutsch et al., 2008).

429 The passive survival strategy of *N. concinna* appears to be effective over several days, but
430 is unlikely to be effective over longer time scales (Pörtner, 2006). After initial shock, animals
431 may switch from acute mechanisms that allow short-term “passive” survival to longer-term
432 acclimatory mechanisms with associated changes in tissue biochemistry and structure. The
433 greater thermal flexibility of *N. concinna* than *L. elliptica* was further evidenced by the increase

434 in adenylate concentration with temperature which suggests that there was an acclimatory
435 increase of the adenylates over the 2 week duration of the experiment, in an attempt to meet the
436 increase in energy demand with temperature.

437 Two month long incubations of *N. concinna* from South Georgia, Signy and Rothera,
438 long enough for seasonal acclimation, led to a reduction in the surface area of mitochondrial
439 cristae per unit muscle fibre volume (Morley et al., submitted). Latitudinal comparisons showed
440 that acclimated *N. concinna* from South Georgia had a higher citrate synthase activity than
441 Rothera *N. concinna*, which was negatively correlated with mitochondrial density across
442 locations. This strongly suggested an adaptation for fewer, more efficient mitochondria in
443 acclimated *N. concinna* from the northern limit of their range (Morley et al., submitted). Tissue
444 samples of *N. concinna* acclimated to 3.0°C are currently being analysed to investigate if
445 changes in aerobic pathways were accompanied by a change of anaerobic scope. These
446 differences in the mechanisms underlying, and the limits of, thermal tolerance between *N.*
447 *concinna* from South Georgia and both Signy and Rothera *N. concinna* are supported by genetic
448 differences, which suggest South Georgia *N. concinna* are a separate population (Beaumont and
449 Wei, 1991).

450 The Scotia arc region of the Southern Ocean has one of the fastest warming climates on
451 the planet (Meredith and King, 2005). If this rate of warming is faster than the rate of
452 evolutionary adaptation, then the acclimatory ability of individuals, populations and species is
453 likely to be a key determinate of their survival (e.g. Peck, 2008). Species with wider geographic
454 ranges are thought to be more resistant to extinction (Jablonski, 1986), which may in part be due
455 to the wider range of environments they experience. This should in turn lead to wider genetic
456 variation and therefore an increased physiological capacity to cope with environmental
457 perturbations (the climate variability hypothesis; Stevens, 1989). The current study has found
458 differences in thermal response of marine ectotherms between locations on the Scotia Arc and
459 Antarctic Peninsula. Investigations of a wider range of species will confirm if these differences
460 are an adaptive response to differences in experienced environment. Latitudinal comparisons
461 testing for differences in capacity between animals from different locations within the Southern
462 Ocean could be a key tool for predicting the vulnerability of species to the current trend of
463 warming.

464

465 **Acknowledgements**

466 This study was funded by NERC core funding to the BAS bioreach program and
467 Antarctic Funding Initiative as well as by the MarCoPoll program of the AWI. Thanks go to the
468 crew of RRS James Clarke Ross and the dive teams both on JCR and at Rothera, especially
469 Andrew Miller who also maintained animals at Rothera. Diving was supported by the NERC
470 NSDF at SAMS. We also thank David Hoffmannbeck and Christian Polleichtner for assisting
471 with biochemical analysis at the Alfred Wegener Institute.

472

473 **References**

- 474 de Aranzamendi M.C., Sahada R., Tatián M., Chiappero M.B. 2008. Genetic differentiation between
475 morphotypes in the Antarctic limpet *Nacella concinna* as revealed by inter-simple sequence repeat
476 markers. *Mar. Biol.* 154, 875-885.
- 477 Barnes D.K.A., Rothery P., Clarke A. 1996. Colonisation and development in encrusting communities
478 form the Antarctic intertidal and sublittoral. *J. Exp. Mar. Biol. Ecol.* 196, 251-265.
- 479 Barnes D.K.A., Linse K., Waller C., Morley S., Enderlein P., Fraser K.P.P., Brown M. 2005. Shallow
480 benthic fauna communities of South Georgia Island. *Polar Biol.* 29, 223-228.
- 481 Barnes, D.K.A., Fuentes, V., Clarke, A., Schloss, I.R., Wallace, M.I. 2006. Spatial and temporal variation
482 in sweater temperatures around Antarctica. *Deep-Sea Res. II* 53, 853-865.
- 483 Barnes, D.K.A., Peck, L.S., Morley, S.A. *submitted*. Acute temperature sensitivity of Antarctic
484 invertebrates determines colonisation potential, biogeography and resilience to environmental change.
- 485 Beaumont, A.R., Wei, J.H.C. 1991. Morphological and genetic variation in the Antarctic limpet *Nacella*
486 *concinna* (Strebel, 1908). *J. Moll. Stud.* 57, 443-450.
- 487 Branch, G.M. 1981. The biology of limpets: physical factors, energy flow, and ecological interactions.
488 *Oceanogr. Mar. Biol. Ann. Rev.* 19, 235-380.
- 489 Buchan, P., Peck, L.S., Tublitz, N.J. 1988. A light, portable apparatus for the assessment of invertebrate
490 heartbeat rate. *J. Exp. Biol.* 136, 495-498.
- 491 Casey, T.M., Dufall, K.G., Arthur, P.G. 1999. An improved capillary electrophoresis method for
492 measuring tissue metabolites associated with cellular energy state. *Eur. J. Biochem.* 261, 740-745.
- 493 Clark, M.S., Geissler, P., Waller, C.L., Fraser, K.P.P., Barnes, D.K.A., Peck, L.S. 2008. Low heat shock
494 thresholds in wild Antarctic inter-tidal limpets (*Nacella concinna*). *Cell Stress Chaperon.* 13, 51-58.
- 495 Clarke, A. and Leakey, R.J.G. 1996. The seasonal cycle of phytoplankton, macronutrients, and the
496 microbial community in a nearshore Antarctic marine ecosystem. *Limnol. and Oceanogr.* 4, 1281-
497 1294.

498 Davenport, J. 1997. Comparisons of the biology of the intertidal subAntarctic limpets *Nacella concinna*
499 and *Kerguelenella lateralis*. J. Moll. Stud. 63, 39-48.

500 Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., Martin, P.R.
501 2008. Impacts of climate warming on terrestrial ectotherms across latitude. PNAS 105, 6668-6672.

502 Fraser, K.P.P., Peck, L.S., Clarke, A. 2002. Feast and famine in Antarctica: seasonal physiology in the
503 limpet *Nacella concinna*. Mar. Ecol. Prog. Ser. 242, 169-177.

504 Gsottbauer, C., Steinbacher, P., Stoiber, W., Obermayer, A., Haslett, J., Sanger, A., Philipp, E. 2007.
505 Mitochondrial density as a parameter of physiological fitness in young and old queen scallops
506 (*Aequipecten opercularis*). Comp. Biochem. Physiol. A 146, S181.

507 Hardewig, I., van Dijk, P.L.M., Portner, H.O. 1998. High-energy turnover at low temperatures: recovery
508 from exhaustive exercise in Antarctic and temperate eelpouts Am. J. Physiol. 274, R1789-R1796.

509 Hardewig, I., Portner, H.O., van Dijk, P.L.M. 2004. How does the cold stenothermal gadoid *Lota lota*
510 survive high water temperatures during summer? J. Comp. Biochem. Physiol. B 174, 149-156.

511 Holmes, S.P., Miller, N. 2006. The hypoxic tolerance of the Protobranch bivalve *Nucula sulcata* Bronn. J.
512 Shellfish Res. 25, 865-867.

513 Jablonski, D. 1986. Larval ecology and macroevolution in marine invertebrates. Bull. Mar. Sci. 39, 565-
514 587.

515 Long, C.W., Brylawski, B.J., Seitz, R.D. 2008. Behavioural effects of low dissolved oxygen on the
516 bivalve *Macoma balthica*. J. Exp. Mar. Biol. Ecol. 359, 34-39.

517 Meredith, M.P., King, J.C. 2005. Rapid climate change in the ocean west of the Antarctic Peninsula
518 during the second half of the 20th century. Geophys. Res. Lett. 32, L19604.

519 Morley, S.A., Peck, L.S., Miller, A.J., Portner, H.O. 2007. Hypoxia tolerance associated with activity
520 reduction is a key adaptation for *Laternula elliptica* seasonal energetics. Oecologia. 153, 29-36.

521 Morley, S.A., Lurman, G.L., Skepper, J.N., Portner, H.O., Peck, L.S. (*submitted*). Thermal plasticity of
522 mitochondria? A latitudinal comparison between Southern Ocean fish and molluscs.

523 Osovitz C.J., Hofmann G.E. 2005. Thermal history-dependent expression of the *hsp 70* gene in purple sea
524 urchins: Biogeographic patterns and the effects of temperature acclimation. J. Exp. Mar. Biol. Ecol.
525 327, 134-143.

526 Peck, L.S. 1989. Temperature and basal metabolism in two Antarctic marine herbivores. J. Exp. Mar.
527 Biol. Ecol. 127, 1-12.

528 Peck, L.S. 1998. Feeding, metabolism and metabolic scope in Antarctic ectotherms. Society for
529 Experimental Biology Seminar Series 66: 365-389.

530 Peck, L.S. 2005. Prospects for survival in the Southern Ocean: vulnerability of benthic species to
531 temperature change. Ant. Sci. 17, 497-507.

532 Peck, L.S. 2008. Brachiopods and climate change. Earth and environmental science transactions of the
533 Royal Society of Edinburgh. 98, 451-456.

534 Peck, L.S., Pörtner, H.O., Hardewig, I. 2002. Metabolic demand, oxygen supply and critical temperatures
535 in the Antarctic bivalve *Laternula elliptica* Physiol. Biochem. Zool. 75, 123-133.

536 Peck, L.S., Ansell, A.D., Webb, K.E., Hepburn, L., Burrows, M.T. 2004. Movements and burrowing
537 activity in the Antarctic bivalve molluscs *Laternula elliptica* and *Yoldia eightsi*. Polar Biol. 27, 357-
538 267.

539 Peck, L.S., Morley, S.A., Pörtner, H.O., Clark, M.S. 2007. Thermal limits of burrowing capacity are
540 linked to oxygen availability and size in the Antarctic clam *Laternula elliptica*. Oecologia 154, 479-
541 484.

542 Peck L.S., Webb, K.E., Miller, A., Clark, M.S., Hill, T. 2008. Temperature limits to activity, feeding and
543 metabolism in the Antarctic starfish *Odontaster validus*. Mar. Ecol. Prog. Ser. 358, 181-189.

544 Peck, L.S., Clark, M.S., Morley, S.A., Massey, A., Rossetti, H. (submitted) Animal temperature limits,
545 size, activity and rates of change. Func. Ecol.

546 Peck, L.S., Massey, A., Thorne, M.A.S., Clark, M.S. *In Press* Lack of acclimation in *Ophionotus*
547 *victoriae* Brittlestars are not fish. Polar Biol.

548 Pörtner, H.O. 2002. Physiological basis of temperature-dependent biogeography: trade-offs in muscle
549 design and performance in polar ectotherms. J. Exp. Biol. 205, 2217-2230.

550 Pörtner, H.O. 2006. Climate-dependent evolution of Antarctic ectotherms: An integrative analysis. Deep
551 Sea Res. II 53, 1071-1104.

552 Pörtner, H.O., Knust, R. 2007. Climate change affects marine fishes through the oxygen limitation of
553 thermal tolerance. Science 315, 95-97.

554 Pörtner, H.O., Boutilier, R.G., Tang, Y., Toews, D.P. 1990. Measurement of intracellular pH and PCO₂
555 after metabolic inhibition by fluoride and nitrilotriacetic acid. Resp. Physiol. 81, 255-274.

556 Pörtner, H.O., Peck, L.S., Zielinski, S., Conway, L.Z. 1999. Intracellular pH and energy metabolism in
557 the highly stenothermal Antarctic bivalve *Limopsis marioensis* as a function of ambient temperature
558 Polar Biol. 22, 17-30.

559 Pörtner, H.O., Peck, L.S., Somero, G. 2007. Thermal limits and adaptation in marine ectotherms: an
560 integrative view. Phil. Trans. R. Soc. B 362, 2233-2258.

561 Sidell, B.D., Driedzic, W.R., Stowe, D.B., Johnston, I.A. 1987. Biochemical correlations of power
562 development and metabolic fuel preferenda in fish hearts. Physiol. Zool. 60, 221-232.

563 Sommer, A., Klein, B., Pörtner, H.O. 1997. Temperature induced anaerobiosis in tow populations of the
564 polychaete worm *Arenicola marina* (L.). J. Comp. Physiol. B 167, 25-35.

565 Stevens, G.C. 1989. The latitudinal gradient in geographical range: how so many species coexist in the
566 tropics. *Am. Nat.* 133, 240-256.

567 Stillman, J.H. 2003. Acclimation capacity underlies susceptibility to climate change. *Science* 301, 65.

568 Terblanche, J.S., Deere, J.A., Clusells-Trullas, S., Janion, C., Chown, S.L. 2007. Critical thermal limits
569 depend on methodological context. *Proc. Roy. Soc. B.* 274, 2935-2942.

570 Tomanek, L. 2005. Two dimensional gel analysis of the heat-shock response in marine snails (genus
571 *Tegula*): interspecific variation in protein expression and acclimation ability. *J. Exp. Biol.* 208, 3133-
572 3143.

573 Waller, C.L., Barnes, D.K.A., Convey, P. 2006. Ecological contrasts across an Antarctic land-sea
574 interface. *31*, 656-666.

575 Weihe, E., Abele, D. *In Press* Differences in the physiological response of inter- and subtidal Antarctic
576 limpets (*Nacella concinna*) to aerial exposure. *Aquatic Biol.*

577 Weibel, E.R. 2000. Symmorphosis on form and function in shaping life (John M Prather Lectures).
578 Harvard University Press, Cambridge

579 Zielinski, S., Pörtner, H.O. 1996. Energy metabolism and ATP free-energy change of the intertidal worm
580 *Sipunculus nudus* below a critical temperature. *J. Comp. Physiol. B* 166, 492-500.

581

582 Table 1. Site details where limpets, *N. concinna* and the soft shelled clam *L. elliptica* were
583 collected during cruise JR109 and at Rothera research station.

584

Species	Site	Depth/m	Co-ordinates	Location
<i>N. concinna</i>	South Cove	6	67°34.25'S, 68°08.00'W	Rothera Point, Adelaide Island
	Borge Bay	0-1	60°42.16'S, 45°35.45'W	Signy, South Orkney Islands
	King Edward Point	2-3	54°17.03'S, 36°39.30'W	South Georgia
<i>L. elliptica</i>	Hangar Cove	10-20	67°33.92'S, 68°07.67'W	Rothera Point, Adelaide Island
	Borge Bay	10-20	60°42.16'S, 45°35.45'W	Signy, South Orkney Islands

585
586

586 Table 2. Shell lengths for limpets, *N. concinna* and the mud clam *L. elliptica* sampled for tissue
 587 biochemistry. Mean \pm sem (number of individuals in brackets). Within each species shell lengths
 588 not sharing a common superscript are significantly different ($p < 0.05$).

Species	Location	Shell length/mm
<i>N. concinna</i>	Rothera	27.7 \pm 0.7 (20) ^a
	Signy	33.9 \pm 0.8 (20) ^b
	South Georgia	33.2 \pm 0.7 (20) ^b
<i>L. elliptica</i>	Rothera	74.0 \pm 3.9 (16) ^c
	Signy	84.2 \pm 3.7 (16) ^c

589

590

591 Table 3 The concentration of selected biochemical measures in *Laternula elliptica* foot muscle
 592 as temperature was increased in 5.0°C steps, after 1-week incubation at each temperature. Mean
 593 \pm sem. Within each parameter, means not sharing a common superscript were significantly
 594 different ($p < 0.05$).

595

Location	Parameter	Temperature			
		0.0	2.5	5.1	7.5
Rothera	O ₂ consumed	73.3 \pm 12.1 ^{ab}	83.4 \pm 16.4 ^a	85.4 \pm 16.0 ^{ab}	45.8 \pm 3.6 ^b
Signy	O ₂ consumed	49.6 \pm 4.8 ^{ab}	94.9 \pm 21.7 ^a	64.2 \pm 14.3 ^{ab}	34.2 \pm 2.6 ^b
Rothera	Heart rate	4.5 \pm 0.4	5.1	6.1	-
Signy	Heart rate	4.1 \pm 0.4	5.3 \pm 0.9	7.2	-
Rothera	CS activity	0.57 \pm 0.05 ^{ab}	0.65 \pm 0.06 ^a	0.40 \pm 0.14 ^b	0.44 \pm 0.10 ^b
Signy	CS activity	0.38 \pm 0.07 ^{ab}	0.61 \pm 0.05 ^a	0.37 \pm 0.08 ^b	0.30 \pm 0.02 ^b
Rothera	Arginine	3.0 \pm 1.2	3.1 \pm 0.4	1.5 \pm 0.2	8.3 \pm 1.2 ^a

Signy	Arginine	1.5±2.6	1.4±0.2	3.1±0.7	7.4±0.6 ^a
Rothera	AMP	0.08±0.01	0.07±0.005	0.03±0.006	0.22±0.07 ^a
Signy	AMP	0.04±0.007	0.03±0.003	0.07±0.02	0.53±0.22 ^a
Rothera	ADP	0.60±0.02	0.53±0.02	0.41±0.02	0.90±0.2 ^a
Signy	ADP	0.44±0.03	0.42±0.02	0.56±0.03	0.79±0.07 ^a
Rothera	ITP	0.11±0.007 ^a	0.10±0.00 ^{ac}	0.090±0.008 ^{ab}	0.067±0.006 ^{bc}
Signy	ITP	0.06±0.01 ^b	0.08±0.004 ^{ab}	0.10±0.005 ^{ac}	0.06±0.007 ^b
Rothera	Acetate	0.0±0.0	87.0±48	62.1±27.2	1182.7±324 ^a
Signy	Acetate	0.0±0.0	0.0±0.0	52.9±15	1251±267 ^a
Rothera	Propionate	0.0±0.0	0.0±0.0	0.0±0.0	540.7±141 ^a
Signy	Propionate	0.0±0.0	0.0±0.0	0.0±0.0	586±45 ^a
Rothera	pHi	7.6±0.02 ^a	7.4±0.01 ^b	7.5±0.03 ^b	7.1±0.03 ^c
Signy	pHi	7.6±0.02 ^a	7.4±0.02 ^b	7.5±0.01 ^b	7.0±0.09 ^c

596
597
598

598 **Figure Legends**

599 Fig. 1. The concentration of selected biochemical measures in *Laternula elliptica* foot muscle as
600 temperature was increased in 5.0°C steps, after 1-week incubation at each temperature; PLA
601 (phospho-L-arginine), ATP, cCO₂, (tissue carbon dioxide) and succinate. Filled bars, Rothera;
602 Open bars, Signy. Within each parameter, means not sharing a common superscript were
603 significantly different ($p < 0.05$).


604
605 Fig. 2. The concentration of adenylates in *Nacella concinna* foot muscle as temperature was
606 increased in 5.0°C steps, after 1-week incubation at each temperature. * indicates that AMP
607 concentrations for 10.0°C Rothera *N. concinna* were significantly different from all locations
608 except Signy *N. concinna* at 2.5°C and South Georgia *N. concinna* at 10.0°C ($p < 0.05$). a
609 indicates that ADP levels were higher at 10.0°C than at other temperatures. Superscripts a and b
610 indicate that the PLA (phospho-L-arginine) concentration for South Georgia *N. concinna* was
611 lower at 10.0°C than either 0.0 or 5.1°C. Superscript c indicates that ATP concentration was
612 higher at 10.0 than 2.5°C. Mean \pm SEM.

613
614 Fig 3. The concentration of organic acids in *Nacella concinna* foot muscle as temperature was
615 increased in 5.0°C steps, after 1 week incubation at each temperature. Points with superscript a
616 are significantly different from all other treatments ($p < 0.05$). * indicates that the succinate
617 concentration of South Georgia *N. concinna* at 10.0°C was significantly higher than all
618 treatments except Rothera 7.5°C. Mean \pm SEM.

619
620 Fig. 4. The CS activity of foot muscle and the oxygen consumption of *Nacella concinna* as
621 temperature were increased in 5.0°C steps, after 1 week incubation at each temperature.
622 Superscripts indicate that for each location CS activity was significantly higher at 7.5 and 10.0°C
623 than at lower temperatures ($p < 0.05$). The complex significant differences in oxygen consumption
624 between temperatures and locations are explained in the text. Mean \pm SEM.


625

625 Fig. 1




626

627 Fig. 2




628

629 Fig. 3



630

631 Fig.4
632



633