

# Non-trivial self-similar extinction solutions for a 3D Hele-Shaw suction problem

**Citation for published version (APA):**

Prokert, G., & Vondenhoff, E. (2007). *Non-trivial self-similar extinction solutions for a 3D Hele-Shaw suction problem*. (CASA-report; Vol. 0719). Technische Universiteit Eindhoven.

**Document status and date:**

Published: 01/01/2007

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

# Non-trivial self-similar extinction solutions for a 3D Hele-Shaw suction problem

G. Prokert  
E. Vondenhoff

*Department of Mathematics and Computer Science  
Technische Universiteit Eindhoven  
P.O. Box 513, 5600 MB Eindhoven, The Netherlands  
E-mail: g.prokert@tue.nl, e.vondenhoff@tue.nl*

## Abstract

We show the existence of noncircular, self-similar solutions to the three-dimensional Hele-Shaw suction problem with surface tension regularisation up to complete extinction. In an appropriate scaling, these solutions are found as bifurcation solutions to a nonlocal elliptic equation of order three. The bifurcation parameter is the ratio of the suction speed and the surface tension coefficient.

**AMS subject classification:** 35R35, 76D27, 47J15

**Key words:** Hele-Shaw flow, bifurcation solution

## 1 Introduction

The problem of Hele-Shaw flow with suction in one point and surface tension regularisation consists in finding a family of moving domains  $t \mapsto \Omega(t)$  in  $\mathbb{R}^3$  and functions  $p(\cdot, t) : \Omega(t) \rightarrow \mathbb{R}$  such that

$$\Delta p = \mu \delta \quad \text{in } \Omega(t), \quad (1.1)$$

$$p = -\gamma \kappa \quad \text{on } \Gamma(t) := \partial\Omega(t). \quad (1.2)$$

Here  $\kappa$  stands for the mean curvature of the moving boundary  $t \mapsto \Gamma(t)$  (taken negative if  $\Omega(t)$  is convex) and  $\gamma > 0$  is the surface tension coefficient. The parameter  $\mu > 0$  denotes the suction speed and  $\delta$  is the delta distribution.

As usual, the evolution of the boundary is given by

$$v_n = -\frac{\partial p}{\partial n}, \quad (1.3)$$

where  $v_n$  is the normal velocity of the boundary.

Besides liquid flow in a Hele-Shaw cell [2], the model and variations of it describe the growth of tumours [1] and porous media flow [3, 4].

Trivial solutions are given by balls around the origin whose volume decreases linearly in time, with rate  $\mu$ . The evolution of the domain that is initially the unit ball  $\mathbb{B}^3$  is therefore

$$\Omega(t) = \alpha(t)\mathbb{B}^3,$$

with  $\alpha : [0, \frac{4\pi}{3\mu}) \rightarrow [0, 1)$  given by

$$\alpha(t) := \sqrt[3]{1 - \frac{3\mu t}{4\pi}}.$$

In [9] it is shown that if  $\mu/\gamma \leq \zeta_2 := 32\pi/5$  then this solution is nonlinearly stable with respect to perturbations that do not change the volume and the center of mass of the initial domain.

In this note we prove the existence of non-trivial solutions with the property

$$\Omega(t) = \alpha(t)\Omega(0),$$

for  $\mu/\gamma$  near the values

$$\zeta_k := 4\pi \frac{k^3 + k^2 - 2k}{k + 3}, \quad k = 2, 3, 4, \dots$$

This communication is organized as follows: in Section 2 we introduce a rescaled evolution equation (2.2) in a way that the trivial solutions described above are represented by trivial stationary solutions. In turn, non-trivial stationary solutions of (2.2) correspond to non-trivial self-similar extinction solutions of the original Hele-Shaw problem. We repeat some results from [9, 10] which form the framework for our considerations here. Because of the rescaling, the evolution operator depends on time. For the problem in  $\mathbb{R}^3$ , the operator scales in such a way that this time dependence occurs simply as a multiplication by a function of time.

In Section 3 we will apply a well known result on “bifurcation from a simple eigenvalue”. To ensure that the eigenvalue under consideration is simple, we have to restrict our basic space, thereby introducing a symmetry breaking. This approach is also used (for other free boundary problems) in [5], [6] and [7].

We want to point out here that the result depends crucially on the space dimension 3. This is due to the fact that only in this dimension the fundamental solution for the Laplacian has the same scaling behavior with respect to dilations as the curvature.

## 2 The evolution problem

Let  $\mathbb{H}^s(\mathbb{S}^2)$  be the Sobolev space of order  $s$  of functions on the unit sphere  $\mathbb{S}^2$  in  $\mathbb{R}^3$ . We recall some constructions and propositions from [9] which form the basis for the results given in Section 3.

We restrict ourselves to domain evolutions that can be described by a continuous function  $R : \mathbb{S}^2 \times [0, \infty) \rightarrow (-1, \infty)$  such that

$$\Omega(t) = \Omega_{R(\cdot, t)} := \left\{ \xi \in \mathbb{R}^3 \setminus \{0\} : |\xi| < 1 + R\left(\frac{\xi}{|\xi|}, t\right) \right\} \cup \{0\}.$$

Besides  $R$  we introduce  $r : \mathbb{S}^2 \times [0, \infty) \rightarrow (-1, \infty)$  by

$$r(x, t) + 1 = \frac{1 + R(x, t)}{\alpha(t)}.$$

This definition of  $r$  is equivalent to

$$\Omega_{r(\cdot, t)} = \alpha(t)^{-1} \Omega_{R(\cdot, t)}. \quad (2.1)$$

Let  $\Omega_{R(t)} = \alpha(t) \Omega_{r(t)}$  be a solution to the Hele-Shaw problem (1.1)-(1.3). Then  $r(t)$  satisfies an evolution equation of the form

$$\frac{\partial r}{\partial t} = \frac{1}{\alpha(t)^3} (\gamma \mathcal{F}_1(r) - \mu \mathcal{F}_2(r)), \quad (2.2)$$

with smooth  $\mathcal{F}_1 : \mathcal{U} \rightarrow \mathbb{H}^{s-3}(\mathbb{S}^2)$  and  $\mathcal{F}_2 : \mathcal{U} \rightarrow \mathbb{H}^{s-1}(\mathbb{S}^2)$ , where  $s > 5$  and  $\mathcal{U}$  is a certain neighbourhood of the origin in  $\mathbb{H}^s(\mathbb{S}^2)$ . For the precise structure of  $\mathcal{F}_1$  and  $\mathcal{F}_2$  and a derivation of (2.2) we refer to [9]. (Hölder spaces are used there instead of Sobolev spaces. Formally, however, the arguments are identical.) Here it is sufficient to investigate the first Fréchet derivatives of  $\mathcal{F}_1$  and  $\mathcal{F}_2$ .

For the equation (2.2),  $r \equiv 0$  is a stationary solution, corresponding to the shrinking ball  $\Omega(t) = \alpha(t) \mathbb{B}^3$ . On the other hand, it is clear that any stationary solution, i.e. any time independent  $r$  satisfying

$$\gamma \mathcal{F}_1(r) - \mu \mathcal{F}_2(r) = 0 \quad (2.3)$$

corresponds to a solution  $\Omega(t) = \alpha(t) \Omega_r$ , i.e. a self-similar solution of the original problem (1.1)-(1.3) which exists up to time  $4\pi/(3\mu)$  when complete extinction of the domain takes place.

We introduce the Dirichlet-to-Neumann operator  $\mathcal{N} : \mathbb{H}^\sigma(\mathbb{S}^2) \rightarrow \mathbb{H}^{\sigma-1}(\mathbb{S}^2)$ ,  $\sigma > 1$ , as the operator that maps a function  $h$  to  $\mathcal{N}h := \frac{\partial u}{\partial n}$ , where  $u$  satisfies

$$\begin{aligned} \Delta u &= 0 && \text{on } \mathbb{B}^3 \\ u &= h && \text{on } \mathbb{S}^2. \end{aligned}$$

This is a first order pseudodifferential operator on  $\mathbb{S}^2$  whose spectrum consists of the nonnegative integers. The eigenfunctions are the spherical harmonics of corresponding degree.

The linearisations of  $\mathcal{F}_1$  and  $\mathcal{F}_2$  around  $r \equiv 0$  are given by

$$\mathcal{F}'_1(0)h = \mathcal{N}(-\mathcal{N}^2 h - \mathcal{N}h + 2h) \quad (2.4)$$

and

$$\mathcal{F}'_2(0)h = -\frac{1}{4\pi} (\mathcal{N}h + 3h). \quad (2.5)$$

For this we refer to [10] Lemma 2.12 and [9] Lemma 2.5.

### 3 Non-trivial stationary solutions via bifurcation

Let  $S_l$  be the space of spherical harmonics of order  $l$ . Define for  $\sigma \geq 0$  the subspace  $\mathbb{H}_\times^\sigma(\mathbb{S}^2)$  of  $\mathbb{H}^\sigma(\mathbb{S}^2)$  consisting of those functions that are invariant with respect to rotations around the  $z$ -axis. It is well known that

$$S_l \cap H_\times^\sigma(\mathbb{S}^2) = \langle Y_l^0 \rangle,$$

where  $Y_l^0$  are the zonal harmonics given by

$$Y_l^0(\theta) = P_l(\cos \theta),$$

where  $\theta$  denotes the polar angle coordinate on  $\mathbb{S}^2$  and  $P_l$  are the Legendre polynomials.

The mappings  $\mathcal{F}_1$  and  $\mathcal{F}_2$  respect rotational symmetries. Therefore, on a suitable neighbourhood  $\mathcal{U}_\times$  of zero in  $\mathbb{H}_\times^s(\mathbb{S}^2)$ , we have a smooth mapping  $\mathcal{F}_{\times, \mu} : \mathcal{U}_\times \rightarrow \mathbb{H}_\times^{s-3}(\mathbb{S}^2)$  given by

$$\mathcal{F}_{\times, \mu} = (\gamma \mathcal{F}_1 - \mu \mathcal{F}_2)|_{\mathcal{U}_\times}.$$

We shall now state the main result of this note. We keep  $s$  and  $\gamma$  fixed and denote by  $X_k$  the orthoplement of  $\langle Y_k^0 \rangle$  in  $\mathbb{H}_\times^s(\mathbb{S}^2)$ . Moreover, we write  $\mu_k := \gamma \zeta_k$ .

**Theorem 3.1.** *Let  $k \geq 2$  be an integer. There exists a  $\delta > 0$  and a  $\mathcal{C}^1$ -curve  $(f, m) : (-\delta, \delta) \rightarrow X_k \times \mathbb{R}$  such that  $(f(0), m(0)) = (0, \mu_k)$  and for all  $\tau \in (-\delta, \delta)$  we have*

$$\mathcal{F}_{\times, m(\tau)}(\tau Y_k^0 + \tau f(\tau)) = 0. \quad (3.1)$$

Furthermore, there is a neighborhood of  $(0, \mu_k)$  such that any zero of  $(r, \mu) \mapsto \mathcal{F}_{\times, \mu}(r)$ , is either of the form  $(\tau Y_k^0 + \tau f(\tau), m(\tau))$  or of the form  $(0, \mu)$ .

This theorem ensures the existence of non-trivial stationary solutions to (2.3). In particular,  $Y_k^0$  gives the direction in which these solutions bifurcate from the trivial one, see Figure 1.

The proof of Theorem 3.1 uses the following lemma. To simplify notation here, we define  $A_k \in \mathcal{L}(\mathbb{H}_\times^s(\mathbb{S}^2), \mathbb{H}_\times^{s-3}(\mathbb{S}^2))$  by

$$A_k := \mathcal{F}'_{\times, \mu_k}(0).$$

**Lemma 3.2.** *Let  $k \geq 2$  be an integer. We have  $\ker A_k = \langle Y_k^0 \rangle$  and  $R(A_k)$  has codimension one.*

*Proof.* The zonal harmonics form a complete orthogonal system in  $\mathbb{H}_\times^s(\mathbb{S}^2)$ . Consequently, we get from (2.4), (2.5), and the fact that  $\mathcal{N}Y_l^0 = lY_l^0$

$$A_k h = \sum_{l \geq 0} g_l(k) \|Y_l^0\|_0^{-2} (h, Y_l^0)_0 Y_l^0, \quad h \in \mathbb{H}_\times^s(\mathbb{S}^2),$$

where  $(\cdot, \cdot)_0$  denotes the usual inner product on  $\mathbb{L}_2(\mathbb{S}^2)$ ,  $\|\cdot\|_0$  the corresponding norm, and

$$g_l(k) = -\gamma(l^3 + l^2 - 2l) + \frac{\mu_k}{4\pi}(l + 3).$$

As  $g_l(k) = 0$  if and only if  $l = k$  and  $g_l(k) \sim -\gamma l^3$  for large  $l$ , both statements follow immediately.  $\square$




Figure 1: The domains  $\Omega_{cY_2^0}$ , for several values of  $c \in \mathbb{R}$ . For  $|c|$  small but nonzero, these domains approximate non-trivial stationary solutions of the rescaled problem, i.e. shapes of self-similarly vanishing domains.

The proof of Theorem 3.1 follows if we combine Lemma 3.2, [8] Theorem 13.5 and the fact that

$$\partial_\mu(\mathcal{F}'_{\times, \mu}(0))|_{\mu=\mu_k} Y_k^0 = -\mathcal{F}'_2(0)Y_k^0 = \frac{k+3}{4\pi}Y_k^0 \notin R(A_k).$$

Our analysis does not provide any strict results concerning the more complicated question of stability of the solutions we found. At least for  $k > 2$ , instability is to be expected because of the linear instability of the trivial solution for  $\mu = \mu_k$ .

## Acknowledgements

The second author has been supported by NWO grant number 613.000.433.

## References

- [1] CHEN, X., AND FRIEDMAN, A. A free boundary problem for an elliptic-hyperbolic system: an application to tumor growth. *SIAM J. Math. Anal.* 35, 4 (2003), 974–986 (electronic).
- [2] ELLIOTT, C. M., AND OCKENDON, J. R. *Weak and variational methods for moving boundary problems*, vol. 59 of *Research Notes in Mathematics*. Pitman (Advanced Publishing Program), Boston, Mass., (1982).
- [3] ESCHER, J., AND SIMONETT, G. Maximal regularity for a free boundary problem. *NoDEA Nonlinear Differential Equations Appl.* 2, 4 (1995), 463–510.

- [4] ESCHER, J., AND SIMONETT, G. Analyticity of the interface in a free boundary problem. *Math. Ann.* 305, 3 (1996), 439–459.
- [5] FONTELOS, M. A., AND FRIEDMAN, A. Symmetry-breaking bifurcations of charged drops. *Arch. Ration. Mech. Anal.* 172, 2 (2004), 267–294.
- [6] FRIEDMAN, A., AND HU, B. Bifurcation for a free boundary problem modeling tumor growth by Stokes equation. *SIAM J. Math. Anal.* 39, 1 (2007), 174–194.
- [7] FRIEDMAN, A., AND REITICH, F. Symmetry-breaking bifurcation of analytic solutions to free boundary problems: an application to a model of tumor growth. *Trans. Amer. Math. Soc.* 353, 4 (2001), 1587–1634 (electronic).
- [8] SMOLLER, J. *Shock waves and reaction-diffusion equations*, vol. 258 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science]*. Springer-Verlag, New York, (1983).
- [9] VONDENHOFF, E. Asymptotic behaviour of injection and suction for Hele-Shaw flow in  $\mathbb{R}^3$  with surface tension near balls. *CASA-Report 06-42, TU Eindhoven*, <ftp://ftp.win.tue.nl/pub/rana/rana06-42.pdf> (2006).
- [10] VONDENHOFF, E. Long-time behaviour of classical Hele-Shaw flows with injection near expanding balls. *CASA-Report 06-19, TU Eindhoven*, <ftp://ftp.win.tue.nl/pub/rana/rana06-19.pdf> (2006).