

Reconstructing a multicolor random scenery seen along a random walk path with bounded jumps

Citation for published version (APA):

Löwe, M., Matzinger, H., & Merkl, F. (2001). *Reconstructing a multicolor random scenery seen along a random walk path with bounded jumps*. (Report Eurandom; Vol. 2001030). Eurandom.

Document status and date:

Published: 01/01/2001

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Reconstructing a Multicolor Random Scenery seen along a Random Walk Path with Bounded Jumps

Matthias Löwe¹, Heinrich Matzinger², and Franz Merkl³

Abstract

Kesten [11] noticed that the scenery reconstruction method proposed by Matzinger [16] relies heavily on the skip-free property of the random walk. He asked if one can still reconstruct an i.i.d. scenery seen along the path of a non-skip-free random walk. In this article, we positively answer this question. We prove that if there are enough colors and if the random walk is recurrent with at most bounded jumps, and if it can reach every integer, then one can almost surely reconstruct almost every scenery up to translations and reflections. Our reconstruction method works if there are more colors in the scenery than possible single steps for the random walk.

Contents

1	Introduction and Result	2
2	Some notation	4
2.1	Conventions concerning constants	6
3	Skeleton of the Reconstruction Procedure	7
4	Proofs concerning the Skeleton Structure	12
5	Heart of the Reconstruction Procedure: Definition of the Algorithm Alg^n	21
6	Playing Puzzle: Correctness of the Algorithm Alg^n	24
6.1	Combinatorics concerning $E_{\text{xi does it}}^n$	25
6.2	Combinatorics concerning $E_{\text{all pieces ok}}^n$	30
6.3	Probabilistic estimates for basic events	41
7	How to find back: Correctness of the stopping times T_f	48
8	Getting started: The first stopping times	51
8.1	The stopping times T^0	51
8.2	The stopping times T^1	55
A	Appendix: The first return time to the origin	64

¹Department of Mathematics, University of Nijmegen, Toernooiveld 1, NL-6525 ED Nijmegen, The Netherlands. loewe@sci.kun.nl

²Eurandom, PO Box 513, NL-5600 MB Eindhoven, The Netherlands. matzinger@eurandom.tue.nl

³Universität Bielefeld, Fakultät für Mathematik, Postfach 100131, D-33501 Bielefeld, Germany, and Eurandom, PO Box 513, NL-5600 MB Eindhoven, The Netherlands. merkl@mathematik.uni-bielefeld.de

1 Introduction and Result

A (one dimensional) *scenery* is a coloring ξ of the integers \mathbb{Z} with C_0 colors $\{1, \dots, C_0\}$. Two sceneries ξ, ξ' are called *equivalent*, $\xi \approx \xi'$, if one of them is obtained from the other by a translation or reflection. Let $(S(t))_{t \geq 0}$ be a recurrent random walk on the integers. Observing the scenery ξ along the path of this random walk, one sees the color $\xi(S(t))$ at time t . The *scenery reconstruction problem* is concerned with trying to retrieve the scenery ξ , given only the sequence of observations $\chi := (\xi(S(t)))_{t \geq 0}$. Quite obviously retrieving a scenery can only work up to equivalence. Work on the scenery reconstruction problem started by Kesten's question, whether one can recognize a single defect in a random scenery. Kesten [10] answered this question in the affirmative in the case of four colors. He takes the colors to be i.i.d. uniformly distributed. In his Ph.D. thesis [16], see also [17] and [19], Matzinger proved that typical sceneries can be reconstructed: He takes the sceneries as independent uniformly distributed random variables, too. He showed that almost every scenery can be almost surely reconstructed. In [11], Kesten noticed that this proof in [16] heavily relies on the skip-free property of the random walk. He asked whether the result might still hold in the case of a random walk with jumps. This article gives a positive answer to Kesten's question: If the random walk can reach every integer with positive probability and is recurrent with bounded jumps, and if there are strictly more colors than possible single steps for the random walk, then one can almost surely reconstruct almost every scenery up to equivalence.

More formally: Let $\mathcal{C} = \{1, \dots, C_0\}$ denote the set of colors. Let μ be a probability measure over \mathbb{Z} supported over a finite set $\mathcal{M} := \text{supp } \mu \subseteq \mathbb{Z}$. With respect to a probability measure P , let $S = (S(k))_{k \in \mathbb{N}}$ be a random walk starting in the origin and with independent increments having the distribution μ . We assume that $E[S(1)] = 0$; thus S is recurrent. Furthermore we assume that $\text{supp } \mu$ has the greatest common divisor 1, thus S can reach every $z \in \mathbb{Z}$ with positive probability. Let $\xi = (\xi(j))_{j \in \mathbb{Z}}$ be a family of i.i.d. random variables, independent of S , uniformly distributed over \mathcal{C} . We prove:

Theorem 1.1 *If $|\mathcal{C}| > |\mathcal{M}|$, then there exists a measurable map $\mathcal{A} : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ such that*

$$P[\mathcal{A}(\chi) \approx \xi] = 1. \tag{1.1}$$

Research on random sceneries started by work by Keane and den Hollander [9], [3]. They thoroughly investigated ergodic properties of a color record seen along a random walk. These questions were motivated among others by the work of Kalikow [8] and den Hollander, Steif [2], in ergodic theory.

A preform of the scenery reconstruction problem is the problem of distinguishing two given sceneries. It has been investigated by Benjamini and Kesten in [1] and [10]. Howard in a series of articles [7], [6], [5] also contributed to this area; see below. The scenery distinguishing problem is the following: Given two different sceneries ξ, ξ' and observations $(\tilde{\xi}(S(j)))_{j \geq 0}$, where $\tilde{\xi}$ equals either ξ or ξ' , the question is: Can we distinguish whether $\tilde{\xi} = \xi$ or $\tilde{\xi} = \xi'$? Benjamini and Kesten [1] showed that one can almost surely distinguish almost all pairs of sceneries ξ, ξ' , if they are drawn independently with i.i.d. entries. Their result even holds in the two dimensional case. This result is not beaten by a reconstruction result: the reconstruction method in two dimensions by Löwe and Matzinger [15] holds only when we have many colors. When ξ and

ξ' differ in precisely one point, the distinguishing problem was examined by Kesten [10] and Howard [5]. Kesten proved that almost all pairs of those sceneries (ξ, ξ') can be distinguished in the 5-color case. He assumes the sceneries to be i.i.d. Howard proved that all periodic sceneries can be distinguished.

As mentioned above, it is in general not possible to reconstruct ξ ; one can at most expect a reconstruction *up to equivalence*. As a matter of fact, even this is impossible: By a theorem of Lindenstrauss [13], there exist non-equivalent sceneries that cannot be distinguished. Of course, they also cannot get reconstructed.

For sceneries that can be reconstructed Benjamini asked whether the reconstruction works also in polynomial time. This question was positively answered by Matzinger [18] in the case of a two color scenery and a simple random walk with holding. Löwe and Matzinger [14] proved that reconstruction works in many cases even if the scenery is not i.i.d., but has some correlations. For the setting of our article den Hollander asked if the finite bound on the length of the jumps is necessary for scenery reconstruction.

In a way a result by Lenstra and Matzinger complements the present paper. If the random walk might jump more than distance 1 only with very small probability and if the tail of the distribution of the jumps decays sufficiently fast, Lenstra and Matzinger [12] proved that scenery reconstruction is still possible.

Let us explain how this article is organized. In order to avoid getting lost among the many details of the rather complex proof, this article is ordered in a “top-down” approach: In order to show the global structure of the reconstruction procedure in a compact but formal way, we start with a section called “Skeleton”. This section collects the main theorems and main definitions of the reconstruction method, using “lower level” procedures as black boxes. In the “Skeleton” section, we only show how these theorems fit together to yield a proof of the reconstruction theorem 1.1; all proofs of the “ingredient” theorems are postponed to later sections. Although this approach is more abstract than a “bottom-up” structure would be, we hope that it allows the reader to more quickly see the global structure.

Overview on some steps for the reconstruction procedure The reconstruction starts with an ergodicity argument: It suffices to consider only sceneries which produce a very untypical initial piece of observations; in particular we may condition on a large but finite initial piece of the observations to be constant. We apply a reconstruction procedure, which works only in this untypical situation, again and again to the observations with larger and larger initial pieces dropped, disregarding all instances that do not produce the prescribed “untypical” outcome. Finally we will see even the prescribed “untypical situation” sufficiently frequent to successfully reconstruct the scenery. The “untypical initial piece” serves to identify locations close to the origin at later times again, at least up to a certain time horizon.

The reconstruction procedure consists in a hierarchy of partial reconstruction procedures; these try to reconstruct larger and larger pieces of the scenery around the origin. The hierarchy of partial reconstruction procedures is defined recursively.

To reconstruct a large piece in the $(m + 1)$ st hierarchical level, we need some information where the random walker is located while producing its color records. This information is

encoded in stopping times, which stop the random walk with high probability sufficiently close to the origin, at least up to a certain time horizon.

The stopping times for the $(m + 1)$ st hierarchical level are built using the m th level partial reconstruction procedure: Given a reconstructed piece around the origin from the m th level, one starts the whole m th level partial reconstruction procedure again at a later “candidate time”. Whenever the piece of scenery obtained in this way has a sufficiently high overlap with the reconstructed piece around the origin, then one has a high chance that the random walk is close to the origin at the “candidate time”.

The global structure of this recursive construction is formally described in the “Skeleton” Sections 3 and 4, and we prove in Sections 7 and 8 that the stopping times fulfill their specification.

The heart of the reconstruction procedure, i.e. the construction of the partial reconstruction algorithm given the stopping times, is described in Section 5 and proven to be correct in Section 6. Roughly speaking, to reconstruct a piece of scenery of size 2^n , we collect a “puzzle” of words of size proportional to n , i.e. logarithmically in the size of the piece to be reconstructed. The puzzle contains (with high probability) all correct subwords of the given size in the “true” piece of scenery to be reconstructed, but also some “garbage” words. We play a kind of puzzle game with these pieces: starting with seed words, we reconstruct larger and larger pieces by adjoining more and more pieces of the puzzle that fit to the growing piece.

Although the actual construction is much more complicated than the idea described now, let us describe an (over)simplified version of how to collect pieces in the puzzle: Suppose we have two “characteristic signals” A and B in the scenery, which occur only once in the scenery. Suppose that the distance between A and B is a multiple of the maximal step size l_{\rightarrow} of the random walk to the right. Then we can almost surely identify the whole “ladder” word read while stepping from A to B with step size l_{\rightarrow} , as follows: Look at all occurrences of A and B in the color record with minimal distance. The words occurring in the color record between those A and B should (a.s.) be always the same in the whole record, and it is the “ladder” word we are looking for. Of course, by ergodicity there are almost surely no (bounded) signals A and B in the scenery that occur only once; this is why the simple idea described here cannot be applied without considerable refinement.

The “pieces of puzzle” obtained are l_{\rightarrow} -spaced pieces; not pieces with spacing 1. This is why our puzzle game leads to reconstructions of modulo classes of the scenery modulo l_{\rightarrow} only. In order to successfully reconstruct the whole scenery, we need to arrange these modulo classes correctly, using some “neighborship” relation between pieces of the puzzle. Unfortunately, the correct arrangement of modulo classes is a technically intricate step in the reconstruction procedure.

2 Some notation

We collect some globally used nonstandard notations and conventions in this section.

Sets, functions, and integers: For functions f and sets D the notation $f \upharpoonright D$ means the restriction of f to the set D . D need not be contained in the domain of f ; thus $f \upharpoonright D$ is defined on $D \cap \text{domain}(f)$. If f and g are functions, the notation $f \subseteq g$ means that f is a restriction of g ; this notation is consistent with the set theoretic definition of functions. By convention,

$0 \in \mathbb{N}$. The integer part of a real number r is denoted by $[r] := \max\{z \in \mathbb{Z} \mid z \leq r\}$; similarly $\lceil r \rceil := \min\{z \in \mathbb{Z} \mid z \geq r\}$.

Integer intervals: Unless explicitly stated otherwise, intervals are taken over the integers, e.g. $[a, b] = \{n \in \mathbb{Z} : a \leq n \leq b\}$, $]a, b[= \{n \in \mathbb{Z} : a < n < b\}$. Given a fixed number C_0 , we define the *set of colors* $\mathcal{C} := [1, C_0] = \{1, \dots, C_0\}$, $|\mathcal{C}| = C_0$.

In the rest of this section I will denote an arbitrary subset of \mathbb{Z} unless otherwise specified.

Sceneries and equivalence: By definition, a *scenery* is an element of $\mathcal{C}^{\mathbb{Z}}$. If $I \subseteq \mathbb{Z}$, then the elements of \mathcal{C}^I are called *pieces of scenery*. The *length* $|\zeta|$ of a piece of scenery $\zeta \in \mathcal{C}^I$ is the cardinality $|I|$ of its index set. $\zeta^{\leftrightarrow} := (\zeta_{-i})_{i \in -I}$ denotes the reflection of a piece of scenery $\zeta \in \mathcal{C}^I$ at the origin. Two pieces of scenery $\zeta \in \mathcal{C}^I$ and $\zeta' \in \mathcal{C}^{I'}$ are called *strongly equivalent*, $\zeta \equiv \zeta'$, if ζ is obtained by some translation of ζ' , i.e. $I' = I + b$ for some $b \in \mathbb{Z}$, and $\zeta = (\zeta'_{i+b})_{i \in I}$. ζ and ζ' are called *equivalent*, $\zeta \approx \zeta'$, if ζ is obtained by some translation or reflection of ζ' , i.e. $I' = aI + b$ for some $a \in \{\pm 1\}$, $b \in \mathbb{Z}$, and $\zeta = (\zeta'_{ai+b})_{i \in I}$. If $T : \mathbb{Z} \rightarrow \mathbb{Z}$, $T(z) = az + b$, denotes this translation or reflection, then $T[\zeta] := \zeta'$ denotes the transport of ζ' by T ; the same notation is used for the domains: $T[I] = I'$. By definition, $\zeta \preceq \zeta'$ means that $\zeta \approx \zeta'[J$ for some $J \subseteq I'$. If additionally such a subset $J \subseteq I'$ and its reading direction (i.e. either $\zeta \equiv \zeta'[J$ or $\zeta \equiv (\zeta'[J]^{\leftrightarrow})$) is unique, we write $\zeta \preceq_1 \zeta'$. Similarly $\zeta \sqsubseteq \zeta'$ (in words: “ ζ occurs in ζ' ”) means that $\zeta \equiv \zeta'[J$ for some $J \subseteq I'$.

Words: The elements of $\mathcal{C}^* := \bigcup_{n \in \mathbb{N}} \mathcal{C}^n = \bigcup_{n \in \mathbb{N}} \mathcal{C}^{\{0, \dots, n-1\}}$ are called *words* (over \mathcal{C}). We identify \mathcal{C} with \mathcal{C}^1 . The concatenation of two words $w_1 \in \mathcal{C}^n$ and $w_2 \in \mathcal{C}^m$ is denoted by $w_1 w_2 \in \mathcal{C}^{n+m}$.

Probability distributions: The law of a random variable X with respect to a probability measure P is denoted by $\mathcal{L}_P(X)$. The n -fold convolution of a probability distribution μ over \mathbb{R} is denoted by μ^{*n} .

Random sceneries and random walks: As mentioned before, let μ be a probability measure over \mathbb{Z} supported over a finite set $\mathcal{M} = \text{supp } \mu \subseteq \mathbb{Z}$. Let $\Omega_2 \subseteq \mathbb{Z}^{\mathbb{N}}$ denote the set of all paths with starting point $S(0) = 0$ and jump sizes $S(t+1) - S(t) \in \mathcal{M}$, $t \in \mathbb{N}$. Let Q_0 denote the law of a random walk $S = (S(k))_{k \in \mathbb{N}}$ with start in $0 \in \mathbb{Z}$ and with independent increments having the distribution μ . Furthermore, let $\xi = (\xi_j)_{j \in \mathbb{Z}}$ be a family of i.i.d. random variables, independent of S , with uniform distribution $\mathcal{L}(\xi_j) = \nu$ over \mathcal{C} . We realize (ξ, S) as canonical projections of $\Omega = \mathcal{C}^{\mathbb{Z}} \times \Omega_2$ endowed with its canonical product σ -algebra and the probability measure $P := \nu^{\mathbb{Z}} \otimes Q_0$. (The restriction of the random walk paths not to have forbidden jumps even on null sets is technically convenient.) We assume that $E[S(1) - S(0)] = 0$ ($k \in \mathbb{N}$); thus S is recurrent. Furthermore we assume that $\text{supp } \mu$ has the greatest common divisor 1, thus S eventually reaches every $z \in \mathbb{Z}$ with probability one. For fixed sceneries $\xi \in \mathcal{C}^{\mathbb{Z}}$, we set $P_\xi := \delta_\xi \otimes Q_0$, where δ_ξ denotes the Dirac measure at ξ . Thus P_ξ is the “canonical” version of the conditioned probability $P[\cdot \mid \xi]$. We use the notations P_ξ and $P[\cdot \mid \xi]$ as synonyms; i.e. we will never work with a different version of the conditioned measure $P[\cdot \mid \xi]$ than P_ξ .

Filtrations: We define the filtration $\mathcal{F} := (\mathcal{F}_n)_{n \in \mathbb{N}}$, $\mathcal{F}_n := \sigma(\xi, (S(k))_{k=0, \dots, n})$ over Ω . We further introduce the filtration $\mathcal{G} := (\mathcal{G}_n)_{n \in \mathbb{N}}$ over $\mathcal{C}^{\mathbb{N}}$, where \mathcal{G}_n is the σ -algebra generated by the projection map $\mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[0, n]}$, $\chi \mapsto \chi \upharpoonright [0, n]$.

Observations of the scenery along the random walk and shifts: Let $\chi = (\chi_n)_{n \in \mathbb{N}} := (\xi_{S(n)})_{n \in \mathbb{N}}$. We sometimes write simply $\chi = \xi \circ S$; this is to be understood in the sense $\chi(\omega) = \xi(\omega) \circ S(\omega)$ for all $\omega \in \Omega$. Let $\mathcal{H} = (\mathcal{H}_n)_{n \in \mathbb{N}}$, $\mathcal{H}_n := \sigma(\chi_k, 0 \leq k \leq n)$ denote the filtration obtained by observing the scenery along initial pieces of the random walk. We define the shift operations $\theta : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{N}}$, $(\chi_n)_{n \in \mathbb{N}} \mapsto (\chi_{n+1})_{n \in \mathbb{N}}$, and $\Theta : \Omega \rightarrow \Omega$, $(\xi, S) \mapsto ((\xi_{n+S(1)})_{n \in \mathbb{Z}}, (S(k+1) - S(1))_{k \in \mathbb{N}})$; thus $\chi \circ \Theta = \theta \circ \chi$. Intuitively, Θ spatially shifts both the scenery and the random walk by the location $S(1)$ of the random walk after one step, and it drops the first time step. One observes $\xi \approx \xi \circ \Theta$.

Admissible paths: A piece of path $\pi = (\pi_i)_{i \in I} \in \mathbb{Z}^I$ over an integer interval I is called *admissible* if $\pi_{i+1} - \pi_i \in \mathcal{M}$ for all $\{i, i+1\} \subseteq I$. For finite $I \neq \emptyset$, $\pi_{\min I}$ and $\pi_{\max I}$ are called starting point and end point of π , respectively. We set $\text{TimeShift}(\pi) := (\pi_{i-1})_{i \in I+1}$. By definition, the length $|\pi|$ of the path π is the cardinality $|I|$. For $x, t > 0$ let $\text{AdPaths}(x, t)$ denote the set of all admissible pieces of path $\pi \in [-x, x]^{[0, t]}$.

Ladder intervals and ladder paths: Let $l_{\rightarrow} := \max \mathcal{M}$, $l_{\leftarrow} := |\min \mathcal{M}|$; thus l_{\rightarrow} and l_{\leftarrow} are the maximal possible jump sizes of S to the right and to the left, respectively. We abbreviate $l := \max\{l_{\rightarrow}, l_{\leftarrow}\}$ and $h := l|\mathcal{M}|$. By definition, d -spaced intervals ($d \in \mathbb{N}$) are sets of the form $I \cap (a + d\mathbb{Z})$ with a bounded interval I and a modulo class $a + d\mathbb{Z} \in \mathbb{Z}/d\mathbb{Z}$. l_{\rightarrow} -spaced intervals are also called right ladder intervals. Similarly, l_{\leftarrow} -spaced intervals are called left ladder intervals. By definition, a right ladder path is a piece of path that steps through the points of some right ladder interval in increasing order. Similarly, a left ladder path is a piece of path that steps through the points of some left ladder interval in decreasing order.

Reading words from pieces of sceneries: For $I = \{i_0, \dots, i_{n-1}\} \subset \mathbb{Z}$ with $i_0 < \dots < i_{n-1}$ and a piece of scenery $\zeta \in \mathcal{C}^I$, we define $\zeta_{\rightarrow} := (\zeta_{i_k})_{k=0, \dots, n-1} \in \mathcal{C}^n$ and $\zeta_{\leftarrow} := (\zeta_{i_{n-1-k}})_{k=0, \dots, n-1} \in \mathcal{C}^n$; thus ζ_{\rightarrow} and ζ_{\leftarrow} are the words obtained by reading ζ from the left to the right and from the right to the left, respectively. The right ladder word of a scenery ξ over a right ladder interval I is defined to be $(\xi \upharpoonright I)_{\rightarrow}$; similarly one defines left ladder words $(\xi \upharpoonright J)_{\leftarrow}$ over left ladder intervals J .

2.1 Conventions concerning constants

Four fixed ‘‘sufficiently large’’ positive integer parameters c_2 , c_1 , α , and n_0 globally play a role. The meaning of these parameters is explained below at the location of their occurrence; at this point we only describe their mutual dependence:

- $c_2 \in \mathbb{N}$ is chosen first sufficiently large; say $c_2 \geq c_2^{\min}(|\mathcal{C}|, \mu)$.
- Then $c_1 \in 2\mathbb{N}$ is chosen to be even and sufficiently large; say $c_1 \geq c_1^{\min}(c_2, |\mathcal{C}|, \mu)$.
- Then $\alpha \in \mathbb{N}$ is chosen to be sufficiently large; say $\alpha \geq \alpha^{\min}(c_1, |\mathcal{C}|, \mu)$.

- Finally $n_0 \in 2\mathbb{N}$ is chosen to be even and sufficiently large; say $n_0 \geq n_0^{\min}(c_1, \alpha, |\mathcal{C}|, \mu)$.

We do not specify explicitly here how large the allowed lower bounds c_2^{\min} , c_1^{\min} , α^{\min} and n_0^{\min} actually need to be; but we emphasize that the constructions below will work if they are sufficiently large.

All other positive constants are denoted by “ c_i ” with a counting index $i > 2$; they keep their meaning globally during the whole article. Unless explicitly stated otherwise, these constants may depend only on the number of colors $|\mathcal{C}|$ and on the jump distribution μ of the random walk; in particular they may depend on the upper bound l of the jump size, but not on n_0 .

3 Skeleton of the Reconstruction Procedure

Our first “ingredient” theorem reduces the problem of almost surely reconstructing sceneries to the following simpler one: We only need to find an auxiliary reconstruction procedure \mathcal{A}_B which may fail to give an answer, and it may sometimes even give the wrong answer, if only giving the correct answer is more probable than giving a wrong one. Roughly speaking, we apply the auxiliary reconstruction procedure \mathcal{A}_B repeatedly to the observations with initial pieces dropped, taking the answer of the majority as our result; here ergodicity of the observations plays a key role.

Theorem 3.1 *If there exists a measurable map $\mathcal{A}_B : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}} \cup \{\text{fail}\}$ with*

$$P[\mathcal{A}_B(\chi) \neq \text{fail}, \mathcal{A}_B(\chi) \approx \xi] > P[\mathcal{A}_B(\chi) \neq \text{fail}, \mathcal{A}_B(\chi) \not\approx \xi], \quad (3.1)$$

then there exists a measurable map $\mathcal{A} : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ such that

$$P[\mathcal{A}(\chi) \approx \xi] = 1. \quad (3.2)$$

The auxiliary reconstruction procedure \mathcal{A}_B gives the output “fail” if one does not see a long block of 1’s in the initial piece of the observations. Thus failure of \mathcal{A}_B is a very frequent event; however, non-failure still occurs with a positive but small probability, and conditioned on this event the most probable answer will be the correct one. Roughly speaking, when we apply \mathcal{A}_B again and again to the observations with initial pieces dropped, we will finally see sufficiently many long blocks of 1’s to make the whole procedure work correctly.

The required long block of 1’s in the initial piece should have length n_0^{20} for some sufficiently large but fixed even number $n_0 \in 2\mathbb{N}$. The parameter n_0 , which parametrizes the size of this required block, is chosen fixed but large enough (see Subsection 2.1).

Definition 3.2 *With the abbreviation $J_1 = [-2ln_0^{20}, 2ln_0^{20}]$, we define the following events:*

$$E_B(k) := \{\chi_n = 1 \text{ for all } n \leq k\} \quad \text{for } k \in \mathbb{N}, \quad (3.3)$$

$$\text{BigBlock} := \left\{ \text{There is an integer interval } J_0 \subseteq J_1 \text{ with } |J_0| \geq n_0^4 \text{ such that} \right. \\ \left. \xi|_{J_0} = (1)_{j \in J_0} \text{ is a constant piece of scenery with value 1.} \right\}. \quad (3.4)$$

Let P_B denote the image of the conditioned law $P[\cdot | E_B(n_0^{20})]$ with respect to the shift $\Theta^{n_0^{20}}$. Furthermore, we define the conditioned law

$$\tilde{P} := P_B[\cdot | \text{BigBlock}]. \quad (3.5)$$

The event $E_B(n_0^{20})$ occurs when we see a large block of 1's in an initial piece of the observations, while BigBlock occurs when there is a large block of 1's close to the origin in the (unobservable) real scenery ξ . The next lemma tells us that such a large block in the real scenery is very probable whenever we see a large initial block of 1's in the observations:

Lemma 3.3 *There exists $c_3 > 0$ such that $P_B[\text{BigBlock}] \geq 1 - e^{-c_3 n_0^{12}}$.*

We describe the intuitive meaning of \tilde{P} : After having seen a large initial block of 1's in the observations, we drop this initial piece and take the present point as our new starting point. Since then a large block of 1's close to the origin in the unobservable real scenery ξ is typical, it does not change much when we even condition on this (unobservable) event.

Almost all the proofs using the measure \tilde{P} will not explicitly use its definition (3.5), but only the following properties of \tilde{P} (and n_0):

Lemma 3.4 *The probability measure \tilde{P} fulfills:*

1. ξ and S are independent with respect to \tilde{P} ;
2. The common distributions of $(S, \xi[(\mathbb{Z} \setminus J_1)])$ with respect to \tilde{P} and with respect to P coincide.
3. With respect to \tilde{P} , the restriction $\xi[J_1]$ is independent of $\xi[(\mathbb{Z} \setminus J_1)]$.
4. $\tilde{P}[\text{BigBlock}] = 1$.

The next theorem shows that whenever we have a reconstruction procedure \mathcal{A}' that works sufficiently probably with respect to the modified measure \tilde{P} , then there exists the auxiliary reconstruction procedure \mathcal{A}_B that we needed above:

Theorem 3.5 *Assume that there exists a measurable map $\mathcal{A}' : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ with*

$$\tilde{P}[\mathcal{A}'(\chi) \approx \xi] \geq \frac{2}{3}. \quad (3.6)$$

Then there exists a measurable map $\mathcal{A}_B : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}} \cup \{\text{fail}\}$ such that

$$P[\mathcal{A}_B(\chi) \neq \text{fail}, \mathcal{A}_B(\chi) \approx \xi] > P[\mathcal{A}_B(\chi) \neq \text{fail}, \mathcal{A}_B(\chi) \not\approx \xi]. \quad (3.7)$$

The reconstruction function \mathcal{A}' required by the last theorem is built by putting together a hierarchy of partial reconstruction algorithms \mathcal{A}^m , $m \geq 1$. The partial reconstruction algorithms \mathcal{A}^m try to reconstruct longer and longer pieces around the origin; the relevant length scale in the m -th hierarchy is given by 2^{n_m} , where n_m is defined as follows:

Definition 3.6 *We define recursively a sequence $(n_m)_{m \in \mathbb{N}}$: n_0 was already chosen above; we set*

$$n_{m+1} := 2^{\lfloor \sqrt{n_m} \rfloor}. \quad (3.8)$$

The partial reconstruction algorithms may sometimes, but not too frequently, give the wrong answer:

Theorem 3.7 *Assume that there exists a sequence $(\mathcal{A}^m)_{m \geq 1}$ of measurable maps $\mathcal{A}^m : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{n_m}, 5 \cdot 2^{n_m}]}$ such that*

$$\tilde{P} \left[\bigcup_{m=1}^{\infty} (E^m)^c \right] \leq \frac{1}{3}, \quad (3.9)$$

where

$$E^m := \{\xi \in [-2^{n_m}, 2^{n_m}] \preceq \mathcal{A}^m(\chi) \preceq \xi \in [-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]\}. \quad (3.10)$$

Then there exists a measurable map $\mathcal{A}' : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ such that the following holds :

$$\tilde{P}[\mathcal{A}'(\chi) \approx \xi] \geq \frac{2}{3}. \quad (3.11)$$

Before describing it formally, let us intuitively explain how the hierarchy of partial reconstruction algorithms \mathcal{A}^m is constructed: The \mathcal{A}^m are built recursively in a “zig-zag” way simultaneously with a hierarchy of stopping times:

These stopping times have the task to estimate times when the random walk S is sufficiently close back to the origin, at least up to a certain time horizon. For this estimation, one may use only an initial piece of the color record χ . To find “higher level” stopping times, we try to reconstruct a piece of scenery both at the present candidate location and at the starting point, using a “lower level” partial reconstruction algorithm. If the two obtained pieces of scenery have a high overlap with each other, then there is a good chance that the candidate location and the starting point are close to each other. This is the “zig” part of the “zig-zag” recursion.

The “zag” part of the recursion uses the stopping times as follows to construct a “higher level” partial reconstruction algorithm \mathcal{A}^m : Whenever the stopping times indicate that one might be sufficiently close to the origin, one collects “typical signals” which one expects to be characteristic of the local environment in the scenery. The data obtained in this way are then matched together similarly to playing a puzzle game. This procedure is the heart of the whole reconstruction method.

To get the whole construction started, one needs some initial stopping times which indicate that one might be sufficiently close to the origin. A simple way to get such times is the following: Whenever one observes a sufficiently long block of 1’s in the color record, then one has a high chance to be close to the origin. (Remember: We conditioned on seeing a long block of 1’s at an initial piece of the color record.) This is the reason why we introduce the modified measure \tilde{P} , since with respect to \tilde{P} one can be (almost) sure to have a big block of 1’s in the scenery close to the origin. However, the such constructed stopping times are not reliable enough to base the first partial reconstruction algorithm on them. Instead, these stopping times are used as ingredients to construct more reliable stopping times.

We treat the “zig” part and the “zag” part of the recursion separately, starting with the formal specification of the “zig” part: Given an abstract partial reconstruction algorithm f , we build stopping times out of it:

The specification of the stopping times depends on a fixed, sufficiently large parameter $\alpha \in \mathbb{N}$. Informally speaking, α influences how many stopping times in each step should be valuable, and what the time horizon for the m -th partial reconstruction algorithm in the hierarchy should be. The parameter α is chosen fixed but large enough; recall Subsection 2.1.

Definition 3.8 Let $m \geq 1$. Let a function $f : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{2n_m}, 5 \cdot 2^{2n_m}]}$ be given. Assume that $f(\chi)$ depends only on $\chi[[0, 2 \cdot 2^{12\alpha n_m}]$. We define the random set

$$\mathbb{T}_f(\chi) := \{t \in [0, 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m} \mid \exists w \in \mathcal{C}^{2 \cdot 2^{2n_m}} : w \preceq f(\chi) \text{ and } w \preceq f(\theta^t(\chi))\}. \quad (3.12)$$

We define a sequence $T_f = (T_{f,k})_{k \geq 0}$ of \mathcal{G} -adapted stopping times with values in $[0, 2^{12\alpha n_{m+1}}]$: Let $t(0) < \dots < t(|\mathbb{T}_f(\chi)| - 1)$ be the elements of $\mathbb{T}_f(\chi)$ arranged in increasing order. For $k \in \mathbb{N}$, we set

$$T_{f,k}(\chi) := \begin{cases} t(2 \cdot 2^{2n_{m+1}}k) + 2 \cdot 2^{12\alpha n_m} & \text{if } 2 \cdot 2^{2n_{m+1}}k < |\mathbb{T}_f(\chi)|, \\ 2^{12\alpha n_{m+1}} & \text{otherwise.} \end{cases} \quad (3.13)$$

Observe that the stopping times $T_f(\chi)$ depend only on $\chi[[0, 2^{12\alpha n_{m+1}}]$.

In the next definition, we introduce events $E_{\text{stop},\tau}^m$; they specify what the stopping times should fulfill: There should be sufficiently many of them, they should be separated by at least $2 \cdot 2^{2n_m}$, and they should stop the random walk sufficiently close to the origin. Furthermore, given any abstract partial reconstruction algorithm f , we define an event $E_{\text{reconst},f}^m$; it measures whether f correctly reconstructs a piece of the scenery around the origin.

Definition 3.9 Let $m \in \mathbb{N}$.

1. Given a sequence $\tau = (\tau_k)_{k \in \mathbb{N}}$ of \mathcal{G} -adapted stopping times, we define

$$E_{\text{stop},\tau}^m := \bigcap_{k=0}^{2^{\alpha n_m}} \{ \tau_k(\chi) < 2^{12\alpha n_m}, |S(\tau_k(\chi))| \leq 2^{n_m}, \tau_j(\chi) + 2 \cdot 2^{2n_m} \leq \tau_k(\chi) \text{ for } j < k \}. \quad (3.14)$$

2. We set for $f : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{2n_m}, 5 \cdot 2^{2n_m}]}$:

$$E_{\text{reconst},f}^m := \{ \xi[[-2^{n_m}, 2^{n_m}] \preceq f(\chi) \preceq \xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]] \}. \quad (3.15)$$

Roughly speaking, the following theorem states: there are stopping times “to get started” which solve their task with high probability:

Theorem 3.10 There exists a sequence of \mathcal{G} -adapted stopping times $T^1 = (T_k^1)_{k \in \mathbb{N}}$ with values in $[0, 2^{12\alpha n_1}]$ and a constant $c_4 > 0$, such that

$$\tilde{P} \left[(E_{\text{stop},T^1}^1)^c \right] \leq e^{-c_4 n_0}. \quad (3.16)$$

The next theorem states that the “zig”-part of the construction works correctly with high probability. As a premise, the “zig”-part needs the underlying “lower level” partial reconstruction algorithm f to work correctly when f is applied at the beginning. Furthermore, the “zig”-part needs f to have a sufficiently high probability to work correctly on the given scenery ξ whenever it is applied again. Informally speaking, the reason is: In the “zig”-part we can only reconstruct, if we know where we are. The idea is to start the whole lower-level reconstruction procedure again whenever we want to find out whether we are close to the origin. As mentioned before, if the result has a large overlap with the piece we have already reconstructed, we can be rather sure that we are close to the origin.

Theorem 3.11 *Under the assumptions of Definition 3.8, we have that*

$$P \left[(E_{\text{stop}, T_f}^{m+1})^c \cap E_{\text{reconst}, f}^m \cap \left\{ P \left[E_{\text{reconst}, f}^m \mid \xi \right] \geq \frac{1}{2} \right\} \right] \leq e^{-n_{m+1}}. \quad (3.17)$$

We remark: in the “zig part” (Theorem 3.11) we work with the event $E_{\text{stop}, T_f}^{m+1}$, while in the “zag part” (Theorem 3.12 below) we work with E_{stop, T_f}^m .

Intuitively, in order to successfully recognize locations close to the origin, we need not only the “lower level” reconstruction to work correctly the first time (i.e. $E_{\text{reconst}, f}^m$ needs to hold), but also the scenery must be such that whenever one starts the “lower level” reconstruction again, one has a sufficiently high chance to reconstruct again a correct piece; this is why we need the event “ $P[E_{\text{reconst}, f}^m \mid \xi] \geq 1/2$ ”.

Finally the heart of the reconstruction algorithm consists of the “zag”-part: there are partial reconstruction algorithms Alg^{n_m} which take an initial piece of the color record as input data, and abstract “lower level” stopping times τ as “argument procedures”. Intuitively, the following theorem states that the algorithms Alg^{n_m} reconstruct correctly with high probability, provided the “argument procedures” τ fulfill their specification $E_{\text{stop}, \tau}^m$.

Theorem 3.12 *For every $m \in \mathbb{N}$, there is a map*

$$\text{Alg}^{n_m} : [0, 2^{12\alpha n_m}]^{\mathbb{N}} \times \mathcal{C}^{2 \cdot 2^{12\alpha n_m}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{n_m}, 5 \cdot 2^{n_m}]} \quad (3.18)$$

such that for every vector $\tau = (\tau_k)_{k \in \mathbb{N}}$ of \mathcal{G} -adapted stopping times with values in $[0, 2^{12\alpha n_m}]$ one has

$$P \left[(E_{\text{reconst}, \text{Alg}^{n_m}(\tau, \cdot)}^m)^c \cap E_{\text{stop}, \tau}^m \right] \leq c_5 e^{-c_6 n_m} \quad (3.19)$$

for some positive constants c_6 and c_5 , where $\text{Alg}^{n_m}(\tau, \cdot) : \chi \mapsto \text{Alg}^{n_m}(\tau(\chi), \chi \upharpoonright [0, 2 \cdot 2^{12\alpha n_m}])$.

To motivate the allowed range for the abstract arguments τ in this theorem, recall that $T_{f,k}(\chi)$ in (3.13) take their values in $[0, 2^{12\alpha n_{m+1}}]$.

Note that Theorems 3.11 and 3.12 use the original probability measure P , while Theorem 3.10 uses the modified probability measure \tilde{P} .

An algorithm Alg^n is defined in the next Section 5, but its correctness, i.e. Theorem 3.12, is proven in Section 6, below. Theorems 3.11 and 3.10 are proven below in separate Sections 7 and 8, respectively. Right now we show how to use these three theorems: Provided these three theorems are true, the hypothesis of Theorem 3.7 holds, i.e. there exists a sequence of measurable maps $\mathcal{A}^m : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{n_m}, 5 \cdot 2^{n_m}]}$ such that (3.9) is valid. We take the maps Alg^{n_m} and the sequences of stopping times T^1, T_f from Theorems 3.10, 3.11, and 3.12 to define recursively maps \mathcal{A}^m . Then we prove: the properties guaranteed by Theorems 3.10, 3.11, and 3.12 imply that the sequence of maps $(\mathcal{A}^m)_{m \geq 1}$ satisfies (3.9). We are ready to describe the “zig-zag”-recursion formally:

Definition 3.13 *We define $\mathcal{A}^m : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{[-5 \cdot 2^{n_m}, 5 \cdot 2^{n_m}]}$ and sequences $T^m = (T_k^m)_{k \in \mathbb{N}}$ of \mathcal{G} -adapted stopping times by simultaneous recursion over $m \geq 1$:*

- T^1 is chosen using Theorem 3.10.
- $\mathcal{A}^m(\chi) := \text{Alg}^{n_m}(T^m(\chi), \chi \upharpoonright [0, 2 \cdot 2^{12\alpha n_m}])$, with Alg^{n_m} taken from Theorem 3.12.

- $T^{m+1} := T_{\mathcal{A}^m}$, with the notation of Definition 3.8.

Recall Definition (3.10) of the events E^m . From now on, we use our specific choice for \mathcal{A}^m from Definition 3.13. Using (3.15), we rewrite (3.10) in the form

$$E^m = E_{\text{reconst}, \mathcal{A}^m}^m. \quad (3.20)$$

Theorem 3.14 *For the sequence $(\mathcal{A}^m)_{m \geq 1}$ as defined in Definition 3.13 and $(E^m)_{m \in \mathbb{N}}$ as in (3.20), the bound (3.9) is valid.*

All theorems of this section together yield the proof of our main theorem:

Proof of Theorem 1.1. By Theorem 3.14, (3.9) holds; then (3.11) holds by Theorem 3.7; moreover (3.7) holds by Theorem 3.5; finally Theorem 3.1 implies the claim (1.1) of Theorem 1.1. ■

4 Proofs concerning the Skeleton Structure

Lemma 4.1 *The shift $\Theta : \Omega \rightarrow \Omega$, $(\xi, S) \mapsto (\xi(\cdot + S(1)), S(\cdot + 1) - S(1))$ is measure-preserving and ergodic with respect to P .*

Proof. The shift Θ is measure-preserving: Since the distribution of ξ is invariant under (deterministic) translations, and since $S(1)$ is independent of ξ , we get: $\xi(\cdot + S(1))$ has the same distribution as ξ . Furthermore, $(S(t+1) - S(1))_{t \in \mathbb{N}}$ has the same distribution as S . Since ξ , $S(1)$ and $(S(t+1) - S(1))_{t \in \mathbb{N}}$ are independent, $\xi(\cdot + S(1))$ and $(S(t+1) - S(1))_{t \in \mathbb{N}}$ are independent, too. Consequently $\Theta(\xi, S)$ has the same distribution as (ξ, S) .

For the ergodicity part, we condition first on deterministic ξ : Recall from Section 2 that for fixed $\xi \in \mathbb{C}^{\mathbb{Z}}$, $P_\xi = \delta_\xi \otimes Q_0$ denotes the “canonical” version of the conditioned measure $P[\cdot | \xi]$. We claim first that the shift Θ is ergodic (but in general not measure-preserving) with respect to P_ξ . To prove this claim, note that the standard shift

$$\hat{\Theta} : \mathbb{Z}^{\mathbb{N}} \rightarrow \mathbb{Z}^{\mathbb{N}}, \quad (s(t))_{t \in \mathbb{N}} \mapsto (s(t+1))_{t \in \mathbb{N}} \quad (4.1)$$

is ergodic (but not measure-preserving) with respect to the probability measure P_S induced by S . Consider the measurable map

$$f_\xi : \mathbb{Z}^{\mathbb{N}} \rightarrow \Omega, \quad f_\xi(s) := (\xi(\cdot + s(0)), s - s(0)); \quad (4.2)$$

then $f_\xi \circ \hat{\Theta} = \Theta \circ f_\xi$, and P_ξ is the image measure of P_S with respect to f_ξ . Thus Θ is ergodic with respect to P_ξ , since $\hat{\Theta}$ is ergodic with respect to P_S .

Let $A \subseteq \Omega$ be measurable and shift-invariant: $\Theta^{-1}[A] = A$. According to the above we have for every $\xi \in \mathcal{C}^{\mathbb{Z}}$ that $P_\xi[A] \in \{0, 1\}$. Consider the set

$$M := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P_\xi[A] = 1 \right\}. \quad (4.3)$$

We claim that for all $a \in \mathbb{Z}$ holds $\xi \in M$ if and only if $\xi(\cdot + a) \in M$. To prove this claim, let $\xi \in M$ and choose $N \in \mathbb{N}$ such that $P[S(N) = a] > 0$. Then the image measure of

$P_\xi[\cdot \mid S(N) = a]$ with respect to Θ^N equals $P_{\xi(\cdot+a)}$. Assume $\xi \in M$. Then $1 = P_\xi[A \mid S(N) = a] = P_\xi[\Theta^{-N}A \mid S(N) = a] = P_{\xi(\cdot+a)}[A]$; this shows that $\xi \in M$ implies $\xi(\cdot+a) \in M$. The same argument, applied to the translated scenery $\xi' = \xi(\cdot+a)$ and $a' = -a$, shows that $\xi(\cdot+a) \in M$ implies $\xi \in M$; hence M is a translation invariant set.

By the ergodicity of the translation operator on sceneries, M has measure $P[\xi \in M] = 0$ or $P[\xi \in M] = 1$. If $P[\xi \in M] = 0$, then $P[P_\xi[A] = 0] = 1$; in this case Fubini's theorem yields $P[A] = 0$. Otherwise $P[P_\xi[A] = 1] = 1$; thus $P[A] = 1$, again by Fubini's theorem. ■

Proof of Theorem 3.1. The idea of this proof is to apply the reconstruction function \mathcal{A}_B to all the shifted observations $\theta^k(\chi)$ for each $k \in \mathbb{N}$. Every time one does this, one gets either a scenery or the state **fail** as result.

Given $\mathcal{A}_B : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}} \cup \{\mathbf{fail}\}$ as in the hypothesis of the theorem, we define measurable functions $\mathcal{A}_B^k : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$, $k \in \mathbb{N}$, as follows:

- If there exists $j \in [0, k[$ such that $\mathcal{A}_B(\theta^j(\chi)) \neq \mathbf{fail}$ and

$$\begin{aligned} & \left| \left\{ j' \in [0, k[\mid \mathcal{A}_B(\theta^{j'}(\chi)) \neq \mathbf{fail}, \mathcal{A}_B(\theta^{j'}(\chi)) \approx \mathcal{A}_B(\theta^j(\chi)) \right\} \right| \\ & > \left| \left\{ j' \in [0, k[\mid \mathcal{A}_B(\theta^{j'}(\chi)) \neq \mathbf{fail}, \mathcal{A}_B(\theta^{j'}(\chi)) \not\approx \mathcal{A}_B(\theta^j(\chi)) \right\} \right|, \end{aligned} \quad (4.4)$$

then let j_0 be the smallest j with this property, and define $\mathcal{A}_B^k(\chi) := \mathcal{A}_B(\theta^{j_0}(\chi))$.

- Else define $\mathcal{A}_B^k(\chi)$ to be the constant scenery $(1)_{j \in \mathbb{Z}}$.

Finally define the measurable function $\mathcal{A} : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ by

$$\mathcal{A}(\chi) := \begin{cases} \lim_{k \rightarrow \infty} \mathcal{A}_B^k(\chi) & \text{if this limit exists pointwise,} \\ (1)_{j \in \mathbb{Z}} & \text{otherwise.} \end{cases} \quad (4.5)$$

We check that the such defined function \mathcal{A} fulfills the claim (1.1) of the Theorem 3.1:

Let us give the general idea: by the hypothesis (3.1) and an ergodicity argument, "on the long run" the proportion of sceneries $\mathcal{A}_B(\theta^k(\chi))$ (for $k \in \mathbb{N}$) which are equivalent to ξ is strictly bigger than the proportion of sceneries which are not equivalent to ξ . More formally, define for $k \in \mathbb{Z}$ the Bernoulli variables X_{sce}^k and $X_{\text{wrong sce}}^k$: we set X_{sce}^k equal to 1 iff $\mathcal{A}_B(\theta^k(\chi)) \neq \mathbf{fail}$ and $\mathcal{A}_B(\theta^k(\chi)) \approx \xi$. Similarly, $X_{\text{wrong sce}}^k$ is equal to 1 iff $\mathcal{A}_B(\theta^k(\chi)) \neq \mathbf{fail}$ and $\mathcal{A}_B(\theta^k(\chi)) \not\approx \xi$. Define

$$Y_{\text{sce}}^k := \frac{1}{k} \sum_{i=0}^{k-1} X_{\text{sce}}^i \quad \text{and} \quad Y_{\text{wrong sce}}^k := \frac{1}{k} \sum_{i=0}^{k-1} X_{\text{wrong sce}}^i. \quad (4.6)$$

Observe that if $Y_{\text{sce}}^k > Y_{\text{wrong sce}}^k$ holds, then $\mathcal{A}_B^k(\chi) \approx \xi$. As a consequence of Lemma 4.1, the sequences $(X_{\text{sce}}^k)_{k \geq 0}$ and $(X_{\text{wrong sce}}^k)_{k \geq 0}$ are stationary and ergodic, since they can be viewed as a measurable function of the sequence $k \mapsto \Theta^k(\xi, S)$. Note that $\xi \approx \xi(\cdot + S(k))$. By the ergodic theorem, we have almost surely:

$$Y_{\text{sce}}^k \xrightarrow{k \rightarrow \infty} P \left[\mathcal{A}_B(\theta^k(\chi)) \neq \mathbf{fail}, \mathcal{A}_B(\chi) \approx \xi \right], \quad (4.7)$$

$$Y_{\text{wrong sce}}^k \xrightarrow{k \rightarrow \infty} P \left[\mathcal{A}_B(\theta^k(\chi)) \neq \mathbf{fail}, \mathcal{A}_B(\chi) \not\approx \xi \right]. \quad (4.8)$$

Thus by the assumption (3.1) there exists a.s. a (random) k_0 such that for all $k \geq k_0$ we have $Y_{\text{sce}}^k > Y_{\text{wrong sce}}^k$ and hence $\mathcal{A}_B^k(\chi) = \mathcal{A}_B^{k_0}(\chi) \approx \xi$; recall that we chose the smallest possible j_0 in the definition of \mathcal{A}_B^k . Thus a.s. $\mathcal{A}(\chi) \approx \xi$. ■

Definition 4.2 For $k, \kappa \in \mathbb{N}$, let $\Xi_{\text{Block}}(k, \kappa)$ be the event of sceneries

$$\Xi_{\text{Block}}(k, \kappa) := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \left| \begin{array}{l} \text{There is an integer interval } J_0 \subseteq [-lk, lk] \text{ with } |J_0| \geq \kappa \text{ such} \\ \text{that } \xi \upharpoonright_{J_0} = (1)_{j \in J_0} \text{ is a constant piece of scenery with value 1.} \end{array} \right. \right\}. \quad (4.9)$$

In this section, only the case $k = n_0^{20}$, $\kappa = n_0^4$ is relevant. In Section 8 below, another case is used as well.

Lemma 4.3 If $\kappa \in 2\mathbb{N}$ is large enough, $k \geq \kappa^2$, $k \in \mathbb{N}$, and if ξ is a scenery with $\xi \notin \Xi_{\text{Block}}(k, \kappa)$, then $P[E_B(k) | \xi] \leq e^{-c_7 k / \kappa^2}$ with some constant $c_7 > 0$. As a consequence, $P[E_B(k) | \xi \notin \Xi_{\text{Block}}(k, \kappa)] \leq e^{-c_7 k / \kappa^2}$.

Proof. Let $\xi \in \mathcal{C}^{\mathbb{Z}} \setminus \Xi_{\text{Block}}(k, \kappa)$. The idea of the proof is to split the time interval $[0, k]$ into pieces of size κ^2 . Let us examine at first one of these time intervals of size κ^2 : A typical length scale for the distance that the random walks travels in this time interval is κ ; in particular the probability that it travels farther than distance κ is bounded away from 0, at least if κ is large enough. If the random walk travels that far, it gets close by a point not colored with “1”, assuming that $\xi \notin \Xi_{\text{Block}}(k, \kappa)$. (Note that the random walk does not leave the interval $[-lk, lk]$ up to the time horizon k .) But once the random walk is close enough to a point not colored with “1”, the probability to hit this point a few steps later is bounded away from 0, too. Thus in every κ^2 -sized interval the random walk has a probability bounded away from 0 to see not only the color 1. There are roughly k/κ^2 such intervals in $[0, k]$; thus the probability to see only 1’s up to the time horizon k is exponentially small in k/κ^2 .

Formally, we proceed as follows: We define stopping times $(\tau_j)_{j=0, \dots, \lfloor \kappa^{-2}k \rfloor - 1}$:

$$\tau_j := \inf \{ t \in [j\kappa^2, (j+1/2)\kappa^2] \mid \xi \upharpoonright_{[S(t), S(t) + l_{\rightarrow}]} \text{ is not constant 1} \}. \quad (4.10)$$

In other words, τ_j is the smallest time in the interval $[j\kappa^2, (j+1/2)\kappa^2]$ when there is a point sufficiently close to the right of the location of the random walker which is not colored with “1”. If no such time exists, $\tau_j = \infty$. We claim: For some constant $c_8 > 0$ holds

$$P[\tau_j < \infty \mid \xi, S \upharpoonright_{[0, j\kappa^2]}] \geq c_8. \quad (4.11)$$

This means: Uniformly in $\xi \in \mathcal{C}^{\mathbb{Z}} \setminus \Xi_{\text{Block}}(k, \kappa)$ and in the history of the random walker up to time $j\kappa^2$, the chance that the random walk will get sufficiently close to a point not colored by “1” during the next $\kappa^2/2$ steps is bounded from below by a positive constant.

To prove (4.11), we observe by the Markov property, $\Delta_j := S((j+1/2)\kappa^2) - S(j\kappa^2)$ has the distribution $\mu^{*\kappa^2/2}$ with respect to the conditioned law $P'_j := P[\cdot \mid \xi, S \upharpoonright_{[0, j\kappa^2]}]$; recall that κ is even. Since $\mu^{*\kappa^2/2}$ has the standard deviation $c_9\kappa$ for some constant c_9 , the Central Limit Theorem implies

$$P[\Delta_j \geq \kappa] \geq c_{10} \quad (4.12)$$

for large enough κ . Here c_{10} denotes a fixed positive constant less than $P[X \geq c_9^{-1}]$, and X is a standard normal random variable. Observe that whenever $\xi \in \mathcal{C}^{\mathbb{Z}} \setminus \Xi_{\text{Block}}(k, \kappa)$ and $\Delta_j \geq \kappa$ hold (i.e. in the interval of interest the random walk S moves at least the distance κ to the right), then τ_j is finite (i.e. the random walk passes close to a point which is not colored with 1). This is true since $\xi \in \mathcal{C}^{\mathbb{Z}} \setminus \Xi_{\text{Block}}(k, \kappa)$ implies that $\xi[S((j+1/2)\kappa^2), S((j+1/2)\kappa^2) + \kappa]$ cannot be a constant piece 1, and since the random walk does not perform jumps to the right larger than l_{\rightarrow} . Since the jump distribution μ is not supported on a strict sublattice $\gamma\mathbb{Z}$ of \mathbb{Z} , there is a fixed $L \in \mathbb{N}$ such that $[0, l_{\rightarrow}] \subseteq \bigcup_{\ell=0}^L \text{supp}(\mu^{*\ell})$. If the event $\{\tau_j < \infty\}$ holds, then ξ is not constant 1 on the interval $[S(\tau_j), S(\tau_j) + l_{\rightarrow}]$. Let A_j denote the event that $\chi[[j\kappa^2, (j+1)\kappa^2]$ is constant 1. Then we have for some constant $1 > c_{11} > 0$ and $\kappa^2/2 \geq L$:

$$\begin{aligned} P'_j[A_j^c] &\geq P'_j[\tau_j < \infty] P'_j[\exists \ell \in [0, L] : \chi(\tau_j + \ell) \neq 1 \mid \tau_j < \infty] \\ &\geq c_8 P'_j[\exists \ell \in [0, L] : \chi(\tau_j + \ell) \neq 1 \mid \tau_j < \infty] \geq c_{11}. \end{aligned} \quad (4.13)$$

Hence we obtain by the Markov property:

$$P[E_B(k) \mid \xi] \leq P \left[\bigcap_{j=0}^{\lfloor \kappa^{-2}k \rfloor - 1} A_j \mid \xi \right] = E \left[\prod_{j=0}^{\lfloor \kappa^{-2}k \rfloor - 1} P'_j[A_j] \mid \xi \right] \leq (1 - c_{11})^{\lfloor \kappa^{-2}k \rfloor}. \quad (4.14)$$

This proves Lemma 4.3 ■

Proof of Lemma 3.3. Let $\sigma^2 := \text{Var}[S(1)]$ be the variance of the single step distribution μ . Consider the integer interval $I := [-2\sigma n_0^{10}, 2\sigma n_0^{10}] \cap \mathbb{Z}$; then

$$P[\xi[I = (1)_{j \in I}]] = |\mathcal{C}|^{-|I|} \geq |\mathcal{C}|^{-4\sigma n_0^{10} - 1}. \quad (4.15)$$

The first submartingale inequality states $P[\max_{0 \leq j \leq t} X_j > \lambda] \leq E[X_t]/\lambda$, for $\lambda > 0$ and nonnegative submartingales X_t . Recall that S^2 is a submartingale, since $x \mapsto x^2$ is convex. Applying the submartingale inequality yields:

$$P[\exists j \in [0, n_0^{20}] : S(j) \notin I] = P \left[\max_{0 \leq j \leq n_0^{20}} S(j)^2 > 4\sigma^2 n_0^{20} \right] \leq (4\sigma^2 n_0^{20})^{-1} E[S(n_0^{20})^2] = \frac{1}{4}. \quad (4.16)$$

If $S(j) \in I$ is valid for all $j \in [0, n_0^{20}]$ and if $\xi[I = (1)_{j \in I}]$ holds, then $\chi[[0, n_0^{20}] = (1)_{j \in [0, n_0^{20}]}$. Thus (4.15) and (4.16) and the independence of S and ξ imply

$$P[E_B(n_0^{20})] \geq \frac{3}{4} |\mathcal{C}|^{-4\sigma n_0^{10} - 1}. \quad (4.17)$$

Hence we get for some constant $c_3 > 0$, using Lemma 4.3 and the abbreviations $\Xi_{\text{Block}}^c = \mathcal{C}^{\mathbb{Z}} \setminus \Xi_{\text{Block}}(n_0^{20}, n_0^4)$ and $E_B = E_B(n_0^{20})$:

$$P[\xi \in \Xi_{\text{Block}}^c \mid E_B] \leq \frac{P[E_B \mid \xi \in \Xi_{\text{Block}}^c]}{P[E_B]} \leq \frac{4}{3} |\mathcal{C}|^{4\sigma n_0^{10} + 1} e^{-c_7 n_0^{12}} \leq e^{-c_3 n_0^{12}}. \quad (4.18)$$

The shift operation $\Theta^{n_0^{20}}$ applied to (ξ, S) cannot shift the scenery ξ by more than ln_0^{20} steps, and every shift of the interval $[-ln_0^{20}, ln_0^{20}]$ by not more than ln_0^{20} steps is contained in J_1 . Thus the shifted event $\Theta^{-n_0^{20}} \text{BigBlock}$ occurs whenever the event $\xi \in \Xi_{\text{Block}}$ holds; thus (4.18) implies $P[\Theta^{-n_0^{20}} \text{BigBlock}^c \mid E_B] \leq e^{-c_3 n_0^{12}}$, which is equivalent to the claim of Lemma 3.3. ■

Proof of Lemma 3.4. We abbreviate $k := n_0^{20}$.

1. We observe first that $\xi \circ \Theta^k$ and the event $E_B(k)$ are both measurable with respect to the σ -field $\sigma(\xi, (S(j))_{j \leq k})$, and $S \circ \Theta^k$ is measurable with respect to $\sigma((S(j) - S(k))_{j > k})$. Since $\sigma(\xi, (S(j))_{j \leq k})$ and $\sigma((S(j) - S(k))_{j > k})$ are independent with respect to P , this implies that $\xi \circ \Theta^k$ and $S \circ \Theta^k$ are independent with respect to $P[\cdot | E_B(k)]$. Hence ξ and S are independent with respect to the image measure $P_B = (P[\cdot | E_B(k)]) \circ (\Theta^k)^{-1}$. Since $\text{BigBlock} \in \sigma(\xi)$, this implies part 1.
2. By the independence proven in 1., it suffices to show the two claims $\mathcal{L}_{\tilde{P}}(S) = \mathcal{L}_P(S)$ and $\mathcal{L}_{\tilde{P}}(\xi[(\mathbb{Z} \setminus J_1)]) = \mathcal{L}_P(\xi[(\mathbb{Z} \setminus J_1)])$:

- With respect to P , $S \circ \Theta^k$ and S both have i.i.d. μ -distributed increments and the starting point 0; thus their distributions coincide. By the above argument, $S \circ \Theta^k$ and $E_B(k)$ are independent with respect to P . Hence the laws of $S \circ \Theta^k$ with respect to P and with respect to $P[\cdot | E_B(k)]$ coincide with the law $\mathcal{L}_P(S)$ of S with respect to P . Hence $\mathcal{L}_P(S) = \mathcal{L}_{P[\cdot | E_B(k)]}(S \circ \Theta^k) = \mathcal{L}_{P_B}(S)$. Since ξ and S are independent with respect to P_B , and since $\text{BigBlock} \in \sigma(\xi)$, we obtain the first claim $\mathcal{L}_{\tilde{P}}(S) = \mathcal{L}_P(S)$.

- We condition on fixed values of $\xi[[-lk, lk]]$ and $S[[0, k]]$:

We know that $\xi \circ \Theta^k$ is a translation of ξ by $S(k)$ steps, which is not more than kl ; this translation maps $[-lk, lk]$ to a subset of J_1 . Thus $(\xi \circ \Theta^k)[(\mathbb{Z} \setminus J_1)]$ is obtained by translating a $(S(k)$ -dependent) subpiece of $\xi[(-lk, lk)]$. Thus by our i.i.d. and independence assumptions for ξ and S we get: $(\xi \circ \Theta^k)[(\mathbb{Z} \setminus J_1)]$ has the distribution $\mathcal{L}_P(\xi[(\mathbb{Z} \setminus J_1)]) = \nu^{\mathbb{Z} \setminus J_1}$ with respect to $P[\cdot | \xi[[-lk, lk]], S[[0, k]]]$. Furthermore, $(\xi \circ \Theta^k)[(\mathbb{Z} \setminus J_1)]$ and $(\xi \circ \Theta^k)[J_1]$ are independent with respect to $P[\cdot | \xi[[-lk, lk]], S[[0, k]]]$. Since $E_B(k)$ depends only on $\xi[[-lk, lk]]$ and $S[[0, k]]$, this implies

$$\mathcal{L}_{P_B}(\xi[(\mathbb{Z} \setminus J_1)]) = \mathcal{L}_{P[\cdot | E_B(k)]}((\xi \circ \Theta^k)[(\mathbb{Z} \setminus J_1)]) = \nu^{\mathbb{Z} \setminus J_1}, \quad (4.19)$$

and $\xi[(\mathbb{Z} \setminus J_1)]$ is independent of $\xi[J_1]$ with respect to P_B . Since the event BigBlock depends only on $\xi[J_1]$, this independence implies

$$\mathcal{L}_{\tilde{P}}(\xi[(\mathbb{Z} \setminus J_1)]) = \mathcal{L}_{P_B}(\xi[(\mathbb{Z} \setminus J_1)]) = \nu^{\mathbb{Z} \setminus J_1} = \mathcal{L}_P(\xi[(\mathbb{Z} \setminus J_1)]); \quad (4.20)$$

recall our choice of \tilde{P} . This proves our second claim.

3. We have just seen: $\text{BigBlock} \in \sigma(\xi[J_1])$, and the random pieces $\xi[(\mathbb{Z} \setminus J_1)]$ and $\xi[J_1]$ are mutually independent with respect to P_B . These two facts imply part 3.
4. This is an immediate consequence of the definition $\tilde{P} = P_B[\cdot | \text{BigBlock}]$.

■

Proof of Theorem 3.5. Assume $\mathcal{A}' : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ is a measurable map satisfying (3.6):

$$P_B[\mathcal{A}'(\chi) \approx \xi | \text{BigBlock}] \geq \frac{2}{3}. \quad (4.21)$$

So,

$$P_B[\{\mathcal{A}'(\chi) \approx \xi\} \cap \text{BigBlock}] \geq \frac{2}{3} P_B[\text{BigBlock}]. \quad (4.22)$$

By Lemma 3.3 it follows, since n_0 is large enough (see Subsection 2.1):

$$P_{\mathbb{B}}[\mathcal{A}'(\chi) \approx \xi] \geq \frac{2}{3} \left(1 - e^{-c_3 n_0^{12}}\right) > \frac{1}{2}. \quad (4.23)$$

Now, by definition of $P_{\mathbb{B}}$,

$$P_{\mathbb{B}}[\mathcal{A}'(\chi) \approx \xi] = P \left[\mathcal{A}'(\chi \circ \Theta^{n_0^{20}}) \approx \xi \circ \Theta^{n_0^{20}} \mid E_{\mathbb{B}}(n_0^{20}) \right]. \quad (4.24)$$

Obviously $\xi \circ \Theta^{n_0^{20}} \approx \xi$. Thus

$$P \left[\mathcal{A}'(\chi \circ \Theta^{n_0^{20}}) \approx \xi \mid E_{\mathbb{B}}(n_0^{20}) \right] > \frac{1}{2}. \quad (4.25)$$

We define $\mathcal{A}_{\mathbb{B}} : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}} \cup \{\text{fail}\}$:

$$\mathcal{A}_{\mathbb{B}}(\chi) := \begin{cases} \mathcal{A}'(\chi \circ \Theta^{n_0^{20}}) & \text{if } E_{\mathbb{B}}(n_0^{20}) \text{ holds,} \\ \text{fail} & \text{otherwise;} \end{cases} \quad (4.26)$$

this is well defined since $E_{\mathbb{B}}(n_0^{20}) \in \sigma(\chi)$. By (4.25), the such defined $\mathcal{A}_{\mathbb{B}}$ satisfies (3.7). ■

Lemma 4.4 *For all events $E \subseteq \Omega$ we have*

$$\tilde{P}(E) \leq |\mathcal{C}|^{4ln_0^{20}+1} P(E). \quad (4.27)$$

Proof of Lemma 4.4. Define $\Omega' := \mathcal{C}^{\mathbb{Z} \setminus J_1} \times \Omega_2$ and write $\Omega = \mathcal{C}^{\mathbb{Z}} \times \Omega_2 = \mathcal{C}^{J_1} \times \mathcal{C}^{\mathbb{Z} \setminus J_1} \times \Omega_2 = \mathcal{C}^{J_1} \times \Omega'$. Then by definition of the measure P and by Lemma 3.4 we have $P = \nu^{J_1} \otimes P_{\Omega'}$ and $\tilde{P} = \tilde{P}_{J_1} \otimes P_{\Omega'}$ where $P_{\Omega'}$ and \tilde{P}_{J_1} , respectively, are the marginal distributions of \tilde{P} on Ω' and \mathcal{C}^{J_1} , respectively. Thus we have for all measurable cylinder-sets of the form $E = \{e_1\} \times E_2 \subset \Omega$, where $e_1 \in \mathcal{C}^{J_1}$ and $E_2 \subseteq \Omega'$:

$$\tilde{P}[E] = \tilde{P}_{J_1}[\{e_1\}] P_{\Omega'}[E_2] \leq |\mathcal{C}|^{4ln_0^{20}+1} \nu^{J_1}[\{e_1\}] P_{\Omega'}[E_2] = |\mathcal{C}|^{4ln_0^{20}+1} P[E] \quad (4.28)$$

where the inequality follows because ν is the uniform distribution on \mathcal{C} , $|J_1| = 4ln_0^{20} + 1$, and \tilde{P}_{J_1} is bounded from above by one. Since \mathcal{C}^{J_1} is finite, every measurable subset of Ω can be written as a finite disjoint union of sets of the above form $\{e_1\} \times E_2$ with $e_1 \in \mathcal{C}^{J_1}$ and $E_2 \subseteq \Omega'$. This proves the result. ■

Proof of Theorem 3.7. For pieces of scenery ψ, ϕ , we define the piece of scenery $\Phi(\psi, \phi)$ as follows: If $\psi \preceq_1 \phi$, then $\Phi(\psi, \phi)$ denotes the unique piece of scenery with $\Phi(\psi, \phi) \approx \phi$ such that $\psi \subseteq \Phi(\psi, \phi)$; otherwise we set $\Phi(\psi, \phi) := \phi$. Let \mathcal{A}^m as in the hypothesis of the theorem and $\chi \in \mathcal{C}^{\mathbb{N}}$. With the abbreviation $\xi^m := \mathcal{A}^m(\chi)$, we define recursively

$$\zeta^1 := \xi^1, \quad (4.29)$$

$$\zeta^{m+1} := \Phi(\zeta^m, \xi^{m+1}), \quad (4.30)$$

$$\mathcal{A}'(\chi) := \begin{cases} \lim_{m \rightarrow \infty} \zeta^m & \text{if this limit exists pointwise on } \mathbb{Z}, \\ (1)_{j \in \mathbb{Z}} & \text{else.} \end{cases} \quad (4.31)$$

(By convention, a sequence $(\zeta^m)_{m \in \mathbb{N}}$ of pieces of sceneries converges pointwise to a scenery ζ if the following holds: $\liminf_{m \rightarrow \infty} \text{domain}(\zeta^m) = \mathbb{Z}$, and for every $z \in \mathbb{Z}$ there is $m_z > 0$ such that

for all $m \geq m_z$ one has $\zeta^m(z) = \zeta(z)$.) Being a pointwise limit of measurable maps, the map $\mathcal{A}' : \mathcal{C}^{\mathbb{N}} \rightarrow \mathcal{C}^{\mathbb{Z}}$ is measurable. For the purpose of the proof, we abbreviate $\underline{\xi}^m := \xi[[-2^{n_m}, 2^{n_m}]]$ and $\bar{\xi}^m := \xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]$ and we define the events

$$E_{\text{1fit}}^m := \left\{ \underline{\xi}^m \preceq_1 \bar{\xi}^{m+1} \right\}. \quad (4.32)$$

We claim:

1. $\liminf_{m \rightarrow \infty} E_{\text{1fit}}^m$ holds \tilde{P} -a.s.,
2. If the event $\liminf_{m \rightarrow \infty} E_{\text{1fit}}^m \cap \bigcap_{m=1}^{\infty} E^m$ occurs, then $\mathcal{A}'(\chi) \approx \xi$.

These two statements together with the hypothesis (3.9) imply the claim (3.11) of the theorem.

Proof of claim 1.: By Lemma 4.4 we may replace “ \tilde{P} -a.s.” in the claim by “ P -a.s.”. If $I_1 \neq I_2$ are fixed integer intervals with $|I_1| = |I_2|$, then $P[\xi[I_1] \approx \xi[I_2]] \leq 2c_{12}e^{-c_{13}|I|}$ holds for some constants $c_{12}, c_{13} > 0$, even if I_1 and I_2 are not disjoint. (See also the similar Lemma 6.33, in particular estimate (6.66), below. The factor 2 makes the notation consistent with this lemma; recall the binary choice: $\xi[I_1] \approx \xi[I_2]$ means $\xi[I_1] \equiv \xi[I_2]$ or $\xi[I_1] \equiv (\xi[I_2])^{\leftrightarrow}$.) We apply this for $I_1 = [-2^{n_m}, 2^{n_m}]$ and all integer intervals $I_2 \subseteq [-9 \cdot 2^{n_{m+1}}, 9 \cdot 2^{n_{m+1}}]$ with $|I_1| = |I_2| = 2 \cdot 2^{n_m} + 1$, $I_1 \neq I_2$; there are at most $18 \cdot 2^{n_{m+1}}$ choices of I_2 . We obtain $P[(E_{\text{1fit}}^m)^c] \leq 18 \cdot 2^{n_{m+1}} \cdot 2c_{12}e^{-2c_{13}2^{n_m}}$, which is summable over m ; recall $n_{m+1} = o(2^{n_m})$ as $m \rightarrow \infty$. Hence $(E_{\text{1fit}}^m)^c$ occurs P -a.s. only finitely many times by the Borel-Cantelli lemma; this proves claim 1.

Next we prove the second claim: By the assumption made there, there is a (random) M such that the events E_{1fit}^m and E^m hold for all $m \geq M$. Let $m \geq M$. In the considerations below, we use several times the following rule: For pieces of sceneries $\alpha, \beta, \gamma, \delta$:

$$\text{If } \alpha \preceq \beta \preceq \gamma \preceq \delta \text{ and } \alpha \preceq_1 \delta, \text{ then } \beta \preceq_1 \gamma. \quad (4.33)$$

In particular, this applies to

$$\underline{\xi}^m \preceq_1 \bar{\xi}^{m+1} \quad \text{and} \quad \underline{\xi}^m \preceq \xi^m \preceq \bar{\xi}^m \preceq \underline{\xi}^{m+1} \preceq \xi^{m+1} \preceq \bar{\xi}^{m+1}; \quad (4.34)$$

we obtain $\xi^m \preceq_1 \xi^{m+1}$. By the definition of ζ^m and Φ , we know $\zeta^m \approx \xi^m$; hence we obtain $\zeta^m \preceq_1 \xi^{m+1}$. Using the definition of Φ again, we see $\zeta^m \subseteq \Phi(\zeta^m, \xi^{m+1}) = \zeta^{m+1}$. Using (4.33), (4.34), $\zeta^m \approx \xi^m$, $\zeta^{m+1} \approx \xi^{m+1}$ again, we get

$$\zeta^m \preceq_1 \bar{\xi}^m \preceq_1 \zeta^{m+1} \preceq_1 \bar{\xi}^{m+1} \quad \text{and} \quad \zeta^m \preceq_1 \bar{\xi}^{m+1}. \quad (4.35)$$

Let $h^m : \mathbb{Z} \rightarrow \mathbb{Z}$, $m \geq M$, denote the unique translation or reflection that maps ζ^m onto a subpiece of $\bar{\xi}^m$. As a consequence of $\zeta^m \subseteq \zeta^{m+1}$, $\bar{\xi}^m \subseteq \bar{\xi}^{m+1}$, and (4.35) we see that h^m does not depend on m for $m \geq M$. Hence h^m maps $\zeta := \bigcup_{m \geq M} \zeta^m$ to a subpiece of $\xi = \bigcup_{m \geq M} \bar{\xi}^m$; thus $\zeta \preceq \xi$. In fact the domain of ζ is \mathbb{Z} ; to see this we observe that $\text{domain}(\zeta)$ contains all $(h^m)^{-1}[\text{domain}(\bar{\xi}^m)] = (h^m)^{-1}[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]$, which cover all of \mathbb{Z} . To summarize, we have shown that $(\zeta^m)_{m \geq M}$ converges pointwise to a scenery $\zeta \approx \xi$; thus $\mathcal{A}'(\chi) = \zeta \approx \xi$ by the definition of $\mathcal{A}'(\chi)$. This finishes the proof of the second claim and also the proof of Theorem 3.7. ■

Definition 4.5 We define events of sceneries

$$\Xi_I := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P \left[(E_{\text{stop}, T^1}^1)^c \mid \xi \right] \leq e^{-c_4 n_0 / 2} \right\}, \quad (4.36)$$

$$\begin{aligned} \Xi_{II} &:= \bigcap_{m=1}^{\infty} \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid \text{If } P[E^m \mid \xi] \geq \frac{1}{2}, \text{ then } P \left[(E_{\text{stop}, T^{m+1}}^{m+1})^c \cap E^m \mid \xi \right] \leq e^{-n_{m+1}/2} \right\} \\ &= \bigcap_{m=1}^{\infty} \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P \left[(E_{\text{stop}, T^{m+1}}^{m+1})^c \cap E^m \cap \left\{ P[E^m \mid \xi] \geq \frac{1}{2} \right\} \mid \xi \right] \leq e^{-n_{m+1}/2} \right\}, \end{aligned} \quad (4.37)$$

$$\Xi_{III} := \bigcap_{m=1}^{\infty} \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P \left[(E^m)^c \cap E_{\text{stop}, T^m}^m \mid \xi \right] \leq c_5^{1/2} e^{-c_6 n_m / 2} \right\}, \quad (4.38)$$

$$\Xi := \Xi_I \cap \Xi_{II} \cap \Xi_{III}, \quad (4.39)$$

where c_5 and c_6 are taken from Theorem 3.12 and c_4 is taken from Theorem 3.10.

Note the similarity between these events and the bounds in (3.16), (3.17) and (3.19). The following lemma provides a link between bounds with and without conditioning on the scenery ξ :

Lemma 4.6 Let A be an event, $r \geq 0$, and Q be a probability measure on Ω such that $Q[A] \leq r^2$. Then

$$Q[Q[A|\xi] > r] \leq r. \quad (4.40)$$

Proof of Lemma 4.6. This follows directly from

$$r^2 \geq Q[A] \geq \int_{\{Q[A|\xi] > r\}} Q[A|\xi] dQ \geq rQ[Q[A|\xi] > r]. \quad (4.41)$$

■

Lemma 4.7 For some constant $c_{14} > 0$ it holds:

$$\tilde{P}[\xi \notin \Xi] \leq e^{-c_{14} n_0}. \quad (4.42)$$

Proof of Lemma 4.7. Using the bound (3.16), Lemma 4.6 for $Q = \tilde{P}$, the fact $\tilde{P}[\cdot \mid \xi] = P[\cdot \mid \xi]$, and the definition (4.36) of Ξ_I , we obtain for a sufficiently small constant $c_{14} > 0$

$$\tilde{P}[\xi \notin \Xi_I] \leq e^{-c_4 n_0 / 2} \leq \frac{e^{-c_{14} n_0}}{3}; \quad (4.43)$$

recall that n_0 was chosen large enough, see Subsection 2.1. As a consequence of the bounds (3.17) and (3.19) we know

$$P \left[(E_{\text{stop}, T^{m+1}}^{m+1})^c \cap E^m \cap \left\{ P[E^m \mid \xi] \geq \frac{1}{2} \right\} \right] \leq e^{-n_{m+1}}, \quad (4.44)$$

$$P \left[(E^m)^c \cap E_{\text{stop}, T^m}^m \right] \leq c_5 e^{-c_6 n_m}. \quad (4.45)$$

We obtain by the bound (4.44), Lemmas 4.4 and 4.6 with $Q = P$, and (4.37):

$$\tilde{P}[\xi \notin \Xi_{\text{II}}] \leq |\mathcal{C}|^{4ln_0^{20}+1} P[\xi \notin \Xi_{\text{II}}] \leq |\mathcal{C}|^{4ln_0^{20}+1} \sum_{m=1}^{\infty} e^{-n_{m+1}/2} \leq \frac{e^{-c_{14}n_0}}{3}. \quad (4.46)$$

Here we used again that n_0 is large, and that $(n_m)_{m \in \mathbb{N}}$ grows fast; see Definition 3.6. The same argument yields, this time using (4.45) and (4.38):

$$\tilde{P}[\xi \notin \Xi_{\text{III}}] \leq |\mathcal{C}|^{4ln_0^{20}+1} P[\xi \notin \Xi_{\text{III}}] \leq |\mathcal{C}|^{4ln_0^{20}+1} \sum_{m=1}^{\infty} c_5^{1/2} e^{-c_6 n_m/2} \leq \frac{e^{-c_{14}n_0}}{3}. \quad (4.47)$$

The combination of (4.43), (4.46), (4.47), and (4.39) proves Lemma 4.7. ■

Lemma 4.8 *For all $\xi \in \Xi$ and all $m \in \mathbb{N}$ the following holds for some constants $c_{15} > 0$, $c_{16} > 0$:*

$$P[E^m \mid \xi] \geq 1 - \sum_{k=0}^m c_{16} e^{-c_{15}n_k} \geq \frac{1}{2} \quad (4.48)$$

and

$$P[E^m \setminus E^{m+1} \mid \xi] \leq c_{16} e^{-c_{15}n_{m+1}}. \quad (4.49)$$

Proof of Lemma 4.8. Let $\xi \in \Xi$. We prove (4.48) and (4.49) simultaneously by induction over m : For $m = 1$ we obtain, since $\xi \in \Xi_{\text{I}}$ and $\xi \in \Xi_{\text{III}}$; see (4.36) and (4.38):

$$\begin{aligned} P[E^1 \mid \xi] &\geq P[E_{\text{stop}, T^1}^1 \mid \xi] - P[(E^1)^c \cap E_{\text{stop}, T^1}^1 \mid \xi] \\ &\geq 1 - e^{-c_4 n_0/2} - c_5^{1/2} e^{-c_6 n_1/2} \geq 1 - \sum_{m=0}^1 c_{16} e^{-c_{15}n_m} \geq \frac{1}{2}; \end{aligned} \quad (4.50)$$

for some constants c_{16}, c_{15} ; recall that $n_1 \geq n_0$ and n_0 is large enough by Subsection 2.1. Thus (4.48) holds for $m = 1$. Let $m \geq 1$. Using $\xi \in \Xi_{\text{II}}$, (4.37), and our induction hypothesis (4.48), we see $P[(E_{\text{stop}, T^{m+1}}^{m+1})^c \cap E^m \mid \xi] \leq e^{-n_{m+1}/2}$. Hence we obtain (4.49), using $\xi \in \Xi_{\text{III}}$ and (4.38):

$$\begin{aligned} P[E^m \setminus E^{m+1} \mid \xi] &\leq P\left[(E^{m+1})^c \cap E_{\text{stop}, T^{m+1}}^{m+1} \mid \xi\right] + P\left[(E_{\text{stop}, T^{m+1}}^{m+1})^c \cap E^m \mid \xi\right] \\ &\leq c_5^{1/2} e^{-c_6 n_{m+1}/2} + e^{-n_{m+1}/2} \leq c_{16} e^{-c_{15}n_{m+1}}. \end{aligned} \quad (4.51)$$

Consequently we get, using our induction hypothesis (4.48) again:

$$P[E^{m+1} \mid \xi] \geq P[E^m \mid \xi] - P[E^m \setminus E^{m+1} \mid \xi] \geq 1 - \sum_{k=0}^{m+1} c_{16} e^{-c_{15}n_k} \geq \frac{1}{2}; \quad (4.52)$$

this completes our induction step. ■

Lemma 4.9 *For some constant $c_{17} > 0$ and for all $\xi \in \Xi$,*

$$\tilde{P}\left[\bigcup_{m=1}^{\infty} (E^m)^c \mid \xi\right] \leq e^{-c_{17}n_0}. \quad (4.53)$$

Proof of Lemma 4.9. By Lemma 4.8 we have for $\xi \in \Xi$:

$$P \left[\bigcup_{m=1}^k (E^m)^c \mid \xi \right] \leq P[(E^1)^c \mid \xi] + \sum_{m=1}^k P[E^m \setminus E^{m+1} \mid \xi] \leq \sum_{m=0}^k c_{16} e^{-c_{15} n_m} \leq e^{-c_{17} n_0}, \quad (4.54)$$

where $c_{17} < c_{15}$ is a small positive constant; recall that n_0 is large. In the limit as $k \rightarrow \infty$, this yields the result (4.53). ■

Proof of Theorem 3.14. Using Lemma 4.7 we have

$$\begin{aligned} \tilde{P} \left[\bigcup_{m=1}^{\infty} (E^m)^c \right] &\leq \tilde{P}[\xi \notin \Xi] + \tilde{P} \left[\{\xi \in \Xi\} \cap \bigcup_{m=1}^{\infty} (E^m)^c \right] \\ &\leq e^{-c_{14} n_0} + \int_{\{\xi \in \Xi\}} \tilde{P} \left[\bigcup_{m=1}^{\infty} (E^m)^c \mid \xi \right] d\tilde{P} \\ &\leq e^{-c_{14} n_0} + \sup_{\xi \in \Xi} \tilde{P} \left[\bigcup_{m=1}^{\infty} (E^m)^c \mid \xi \right]. \end{aligned} \quad (4.55)$$

We bound the argument of the last supremum, using Lemma 4.9:

$$\tilde{P} \left[\bigcup_{m=1}^{\infty} (E^m)^c \mid \xi \right] = P \left[\bigcup_{m=1}^{\infty} (E^m)^c \mid \xi \right] \leq e^{-c_{17} n_0}. \quad (4.56)$$

The combination of (4.55) and (4.56) yields, since n_0 is large (by Subsection 2.1):

$$P \left[\bigcup_{m=1}^{\infty} (E^m)^c \right] \leq e^{-c_{14} n_0} + e^{-c_{17} n_0} \leq \frac{1}{3}. \quad (4.57)$$

■

5 Heart of the Reconstruction Procedure: Definition of the Algorithm Alg^n

This section contains the heart of the reconstruction procedure: for every $n \in \mathbb{N}$, we define an algorithm Alg^n ; it is designed to reconstruct long pieces of scenery with high probability. In Section 6 below we show that it fulfills the formal specification given in Theorem 3.12.

Informally speaking, the observation χ allows us to collect many pieces of “puzzle words”. These puzzle words are chosen to have size $c_1 n$ with a fixed parameter c_1 ; recall subsection 2.1. To obtain them, we collect triples of words (w_1, w_2, w_3) which occur in sequence in the observations χ soon after a stopping time $\tau(k)$; an initial piece of χ is represented below by a formal argument η . We put those words w_2 into our puzzle which are already uniquely determined by w_1 and w_3 . This means that w_1 and w_3 should be very “characteristic signals”; if w_1 and w_3 could be read at very different locations in the scenery close to a stopping time, then it is improbable that they will enclose always the same word w_2 . Frequently, w_2 turns out

to be a ladder word: Whenever one reads a w_2 in the context $w_1w_2w_3$ along a non-ladder path sufficiently close to the origin, one reads with high probability a different word w'_2 in the context $w_1w'_2w_3$, too, along a different path with the same starting point and the same end point; but then w_2 is not collected as a puzzle word.

Here is the formal construction: We take input data $\tau \in [0, 2^{12\alpha n}]^{\mathbb{N}}$ and $\eta \in \mathcal{C}^{2 \cdot 2^{12\alpha n}}$. A side remark: although for formal reasons there are infinitely many $\tau(k)$ given in the input data, the construction below actually uses only the first $2^{\alpha n}$ of them.

Definition 5.1 *We define for $m \geq 0$ the random sets:*

$$\text{PrePuzzle}^n(\tau, \eta) := \{(w_1, w_2, w_3) \in (\mathcal{C}^{c_1 n})^3 \mid \exists k \in [0, 2^{\alpha n}[: w_1w_2w_3 \sqsubseteq \eta[\tau(k), \tau(k) + 2^{2n}]\}, \quad (5.1)$$

$$\text{Puzzle}_I^n(\tau, \eta) := \{(w_1, w_2, w_3) \in \text{PrePuzzle}^n(\tau, \eta) \mid \forall (w_1, w'_2, w_3) \in \text{PrePuzzle}^n(\tau, \eta): w'_2 = w_2\}, \quad (5.2)$$

$$\text{Puzzle}_{II}^n(\tau, \eta) := \{w_2 \in \mathcal{C}^{c_1 n} \mid \exists w_1, w_3 \in \mathcal{C}^{c_1 n}: (w_1, w_2, w_3) \in \text{Puzzle}_I^n(\tau, \eta)\}. \quad (5.3)$$

Let us explain the idea behind the following constructions: Although many of the words w_2 in “Puzzle_{II}” turn out to be ladder words of a central piece in the true scenery ξ , some of them are not: There are “garbage words” in the puzzle. We play a “puzzle-game” with the words in “Puzzle_{II}”: We try to fit larger and larger pieces together. In order to distinguish “real” pieces from “garbage” pieces, we need some “seed words” which are guaranteed (with high probability) not to be garbage words; every piece that fits to a piece containing a seed word has a high chance not to be garbage, too. This is what the set Seed_{II} defined below is good for. We identify “seed” words as “puzzle” words that occur in the observations almost immediately after a stopping time $\tau(k)$, when we expect the random walk to be close to the origin.

Recall the abbreviation $h = l|\mathcal{M}|$. Formally, we proceed as follows:

Definition 5.2

$$\text{Seed}_I^n(\tau, \eta) := \left\{ (w_1, w_2, w_3) \in \text{Puzzle}_I^n(\tau, \eta) \mid \begin{array}{l} \exists k \in [0, 2^{\alpha n}[\exists j \in [0, 7c_1 n l] : \\ w_1w_2w_3 \equiv \eta[\tau(k) + j + [0, 3c_1 n]] \end{array} \right\}, \quad (5.4)$$

$$\text{Seed}_{II}^n(\tau, \eta) := \{w_2 \in \mathcal{C}^{c_1 n} \mid (w_1, w_2, w_3) \in \text{Seed}_I^n(\tau, \eta)\}, \quad (5.5)$$

$$\text{Seed}_{III}^n(\tau, \eta) := \left\{ u \in \text{Seed}_{II}^n(\tau, \eta) \mid \begin{array}{l} \exists v \in \text{Seed}_{II}^n(\tau, \eta) : \\ (u[\llbracket [0, c_2 n l \leftarrow] \cap l \leftarrow \mathbb{Z} \rrbracket \rrbracket \rightarrow = (v[\llbracket [0, c_2 n l \rightarrow] \cap l \rightarrow \mathbb{Z} \rrbracket \rrbracket \leftarrow \end{array} \right\}, \quad (5.6)$$

$$\text{Neighbors}^n(\tau, \eta) := \{(w_1, w_2) \in (\mathcal{C}^{c_1 n})^2 \mid \exists k \in [0, 2^{\alpha n}[, w \in \mathcal{C}^{h-1} : w_1ww_2 \sqsubseteq \eta[\tau(k), \tau(k) + 2^{2n}]\}. \quad (5.7)$$

Let us explain what “Seed_{III}” is intended for: We need to identify the orientation of the pieces (whether they are to be read “forward” or “backward”). This task consists of two problems:

The identification of the relative orientation of two pieces with respect to each other, and the identification of the absolute orientation with respect to the “true” scenery ξ . Of course, we have no chance to identify the absolute orientation if the random walk is symmetric; we even bother about identifying the absolute orientation only in the very unsymmetric case $l_{\rightarrow} \neq l_{\leftarrow}$. The set Seed_{III} helps us to identify the absolute orientation in this case: Suppose we read every l_{\rightarrow} -th letter in a word from the left to the right, and every l_{\leftarrow} -th letter in the same word from the right to the left; then every $l_{\rightarrow}l_{\leftarrow}$ -th letter appears in both words, when at least one letter is read both times. This turns out to be characteristic enough to identify the reading directions “left” and “right” in the case $l_{\rightarrow} \neq l_{\leftarrow}$. The fixed parameter c_2 influences the length of the sample pieces in this procedure.

The relation “Neighbors” serves as an estimation for the geometric neighborhood relation between ladder words: ladder words that occur closely together in the observation χ are expected to occur on geometrically neighboring intervals in the “true” scenery ξ . The next definition defines a “true” geometric neighborhood relation \triangleright_n . We try to reconstruct the corresponding “true” neighborhood relation for ladder words in a piece of ξ using only the “estimated” neighborhood relation “Neighbors”.

Recall that μ^{*k} denotes the k -fold convolution of μ ; in particular

$$\text{supp } \mu^{*k} := \left\{ \sum_{i=1}^k s_i \mid \forall i : s_i \in \text{supp } \mu \right\}. \quad (5.8)$$

Definition 5.3 *Let I, J be right ladder intervals. By definition, $I \triangleright_n J$ means $|I| = |J| = c_1 n$ and $\min J - \max I \in \text{supp } \mu^{*h}$. Similarly for I', J' being left ladder intervals, $I' \triangleleft_n J'$ means $|I'| = |J'| = c_1 n$ and $\max J' - \min I' \in \text{supp } \mu^{*h}$.*

The next definition is the heart of our method: We describe how to obtain reconstructed pieces of sceneries. All pieces of scenery $w \in \mathcal{C}^{[-5 \cdot 2^n, 5 \cdot 2^n]}$ are tested as candidates in a sequence of “Filters”: Reconstructed ladder words should be in “Puzzle $_{\text{II}}$ ”, the “estimated” and the “reconstructed” neighborhood relation should be consistent with each other, the reconstructed pieces should contain “Seed $_{\text{III}}$ ” words, and no piece of the puzzle should be used twice.

Only candidate pieces that pass all Filters are considered as a solution of the partial reconstruction problem.

Definition 5.4 *Let $\text{Filter}_i^n(\tau, \eta)$, $i = 1, \dots, 5$, denote the set of all $w \in \mathcal{C}^{[-5 \cdot 2^n, 5 \cdot 2^n]}$ which fulfill the following condition 1., . . . , 5., respectively:*

1. *For every right ladder interval $I \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$, $|I| = c_1 n$, one has $(w \upharpoonright I)_{\rightarrow} \in \text{Puzzle}_{\text{II}}^n(\tau)$.*
2. *For all right ladder intervals $I, J \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$:
if $I \triangleright_n J$, then $((w \upharpoonright I)_{\rightarrow}, (w \upharpoonright J)_{\rightarrow}) \in \text{Neighbors}^n(\tau, \eta)$.*
3. *For all right ladder intervals $I, J \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$, $|I| = |J| = c_1 n$:
if $((w \upharpoonright I)_{\rightarrow}, (w \upharpoonright J)_{\rightarrow}) \in \text{Neighbors}^n(\tau, \eta)$, then there is $q \in \mathbb{N}$ such that $I \triangleright_n J + ql_{\rightarrow}$.*
4. *For every right modulo class $Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$ there exists a right ladder interval $I \subseteq Z \cap [-2 \cdot 2^n, 2 \cdot 2^n]$ such that $(w \upharpoonright I)_{\rightarrow} \in \text{Seed}_{\text{III}}^n(\tau, \eta)$.*
5. *For all right ladder intervals $I, J \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$, $|I| = |J| = c_1 n$:
if $(w \upharpoonright I)_{\rightarrow} = (w \upharpoonright J)_{\rightarrow}$, then $I = J$.*

We set

$$\text{SolutionPieces}^n(\tau, \eta) := \bigcap_{i=1}^5 \text{Filter}_i^n(\tau, \eta). \quad (5.9)$$

The output of the algorithm Alg^n could be any of these pieces $w \in \text{SolutionPieces}^n(\tau, \eta)$; we choose one of them, if it exists.

Definition 5.5 *We define $\text{Alg}^n(\tau, \eta)$ as follows:*

- If $\text{SolutionPieces}^n(\tau, \eta)$ is nonempty, then we define $\text{Alg}^n(\tau, \eta)$ to be its lexicographically smallest element.
- Otherwise $\text{Alg}^n(\tau, \eta)$ is defined to be the constant scenery $(1)_{j \in [-5 \cdot 2^n, 5 \cdot 2^n]}$.

We could have equally well taken any element of $\text{SolutionPieces}^n(\tau, \eta)$ in Definition 5.5; we choose the lexicographically smallest one just for definiteness.

6 Playing Puzzle: Correctness of the Algorithm Alg^n

In this section we prove Theorem 3.12 by showing that the Algorithm Alg^n defined in Definition 5.5 fulfills the specification described by this theorem: Let $n = n_m$, $m \in \mathbb{N}$. A remark concerning notation: Events defined in this section are labeled with an upper index n , not m , since the “hierarchy level” m plays no role here, in contrast to the “Skeleton” section. Only events that also occur in the “Skeleton” section keep their old index m . Hopefully, this should not cause any confusion.

Let $\tau = (\tau_k)_{k \in \mathbb{N}}$ denote a fixed vector of \mathcal{G} -adapted stopping times with values in $[0, 2^{12\alpha n}]$. We abbreviate $\text{Input} := (\tau(\chi), \chi[[0, 2 \cdot 2^{12\alpha n}]])$.

Definition 6.1 *We define the following events:*

$$E_{\text{xi does it}}^n := \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{SolutionPieces}^n(\text{Input})\}, \quad (6.1)$$

$$E_{\text{all pieces ok}}^n := \left\{ \begin{array}{l} \forall w \in \text{SolutionPieces}^n(\text{Input}) : \\ \xi[[-2^n, 2^n] \preceq w \preceq \xi[[-9 \cdot 2^n, 9 \cdot 2^n] \end{array} \right\}. \quad (6.2)$$

Lemma 6.2

$$E_{\text{xi does it}}^n \cap E_{\text{all pieces ok}}^n \subseteq E_{\text{reconst, Alg}^n(\tau, \cdot)}^m \quad (6.3)$$

Proof of Lemma 6.2. When the event $E_{\text{xi does it}}^n$ holds, then the set $\text{SolutionPieces}^n(\text{Input})$ is not empty. Thus $\text{Alg}^n(\text{Input})$ is the lexicographically smallest element of $\text{SolutionPieces}^n(\text{Input})$. When the event $E_{\text{all pieces ok}}^n$ also holds, then $\xi[[-2^n, 2^n] \preceq \text{Alg}^n(\text{Input}) \preceq \xi[[-9 \cdot 2^n, 9 \cdot 2^n]$. ■

Here is the main theorem of this section; it states that the events $E_{\text{xi does it}}^n$ and $E_{\text{all pieces ok}}^n$ occur very probably whenever the stopping times τ fulfill their task specified by $E_{\text{stop}, \tau}^m$:

Theorem 6.3 *For some constant $c_6 > 0$, $c_5 > 0$:*

$$P \left[E_{\text{stop}, \tau}^m \setminus (E_{\text{xi does it}}^n \cap E_{\text{all pieces ok}}^n) \right] \leq c_5 e^{-c_6 n}. \quad (6.4)$$

This theorem is proven the following three subsections. We split the proof into a purely combinatoric part and a probabilistic part. The combinatoric part (subsection 6.1 below for $E_{\text{xi does it}}^n$ and subsection 6.2 below for $E_{\text{all pieces ok}}^n$) shows that whenever some more “basic” events (named B_{\dots}^n below, where “...” stands for a varying label) and $E_{\text{stop},\tau}^m$ occur, then the events $E_{\text{xi does it}}^n$ and $E_{\text{all pieces ok}}^n$ occur, too. In the probabilistic part (subsection 6.3 below) we show that these basic events B_{\dots}^n are highly probable, at least when $E_{\text{stop},\tau}^m$ occurs.

The **Proof of Theorem 3.12** is an immediate consequence of Lemma 6.2 and Theorem 6.3.

6.1 Combinatorics concerning $E_{\text{xi does it}}^n$

In this subsection, we show that a piece of ξ centered at the origin passes all the tests specified by the Filter_i , provided some basic events B_{\dots}^n (specified below) hold.

Definition 6.4 For $n \in \mathbb{N}$ we define the following events:

$$B_{\text{sig rl}}^n := \left\{ \begin{array}{l} \text{For every right ladder path } \pi \in [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]^{[0, c_1 n/2[} \text{ and for} \\ \text{every admissible piece of path } \pi' \in \text{AdPath}(2 \cdot l2^{2n}, c_1 n/2): \\ \text{If } \xi \circ \pi = \xi \circ \pi', \text{ then } \pi(c_1 n/2 - 1) \geq \pi'(c_1 n/2 - 1). \end{array} \right\}, \quad (6.5)$$

$$B_{\text{sig rr}}^n := \left\{ \begin{array}{l} \text{For every right ladder path } \pi \in [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]^{[0, c_1 n/2[} \text{ and for} \\ \text{every admissible piece of path } \pi' \in \text{AdPath}(2 \cdot l2^{2n}, c_1 n/2): \\ \text{If } \xi \circ \pi = \xi \circ \pi', \text{ then } \pi(0) \leq \pi'(0). \end{array} \right\}. \quad (6.6)$$

Let $B_{\text{sig ll}}^n$ and $B_{\text{sig lr}}^n$ be defined just as $B_{\text{sig rl}}^n$ and $B_{\text{sig rr}}^n$ with “right ladder path” replaced by “left ladder path” and with “ \leq ” and “ \geq ” exchanged in (6.5) and (6.6). We set

$$B_{\text{signals}}^n := B_{\text{sig rl}}^n \cap B_{\text{sig rr}}^n \cap B_{\text{sig ll}}^n \cap B_{\text{sig lr}}^n, \quad (6.7)$$

$$E_{\text{signals II}}^n := \left\{ \begin{array}{l} \text{For every ladder path } \pi \in [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]^{[0, c_1 n[} \text{ and for every} \\ \text{admissible piece of path } \pi' \in \text{AdPath}(2 \cdot l2^{2n}, c_1 n): \\ \text{If } \xi \circ \pi = \xi \circ \pi', \text{ then } \pi(c_1 n/2) = \pi'(c_1 n/2). \end{array} \right\}. \quad (6.8)$$

Lemma 6.5 $B_{\text{signals}}^n \subseteq E_{\text{signals II}}^n$.

Proof of Lemma 6.5. Assume that the event B_{signals}^n occurs. Let $\pi \in [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]^{[0, c_1 n[}$ be a right ladder path and $\pi' \in \text{AdPath}(2 \cdot l2^{2n}, c_1 n)$. Assume that $\xi \circ \pi = \xi \circ \pi'$ holds. Looking at the first half of π and π' only (with the first points $(0, \pi(0))$, $(0, \pi'(0))$ dropped), we see $\pi(c_1 n/2) \geq \pi'(c_1 n/2)$, since $B_{\text{sig rl}}^n$ holds. Similarly, looking at the second half of π and π' only, we infer $\pi(c_1 n/2) \leq \pi'(c_1 n/2)$, since $B_{\text{sig rr}}^n$ holds. Therefore $\pi(c_1 n/2)$ and $\pi'(c_1 n/2)$ coincide. The case of left ladder paths is treated similarly. This shows that $E_{\text{signals II}}^n$ holds. ■

Definition 6.6 By definition, the event $B_{\text{all paths}}^n$ occurs if and only if the following holds: every admissible piece of path $R \in [-12 \cdot 2^n, 12 \cdot 2^n]^{[0, 3c_1 n[}$ occurs in the random walk S with start at most 2^{2n} time steps after some stopping time $\tau(k)$, $k < 2^{\alpha n}$. More formally:

$$B_{\text{all paths}}^n := \left\{ \begin{array}{l} \forall R \in \text{AdPaths}(12 \cdot 2^n, 3c_1 n) \exists k \in [0, 2^{\alpha n}[\exists j \in [0, 2^{2n}] : \\ \text{TimeShift}^{\tau(k)+j}(R) \subseteq S \end{array} \right\}. \quad (6.9)$$

The following auxiliary lemma helps us to show below that the true scenery ξ passes the test Filter_1 . Roughly speaking, it tells us that sufficiently many ladder words occur in the puzzle. This is important, since playing our puzzle game would lead to a failure when pieces were missing.

Lemma 6.7 *Assume that the event $B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop}, \tau}^m$ holds. Let $I \subseteq [-6 \cdot 2^n, 6 \cdot 2^n]$ be a right (or left) ladder interval with $|I| = 3c_1n$, and let $w_1, w_2, w_3 \in \mathcal{C}^{c_1n}$ with $(\xi[I]_{\rightarrow} = w_1w_2w_3$ (or $(\xi[I]_{\leftarrow} = w_1w_2w_3$ in the case of a left ladder interval). Then $(w_1, w_2, w_3) \in \text{Puzzle}_1^n(\text{Input})$.*

Proof of Lemma 6.7. Assume that I is a right ladder interval; the case of left ladder intervals can be treated in the same way by exchanging “left” and “right”. Let $I = I_1 \cup I_2 \cup I_3$, where I_1, I_2 , and I_3 denote the left, middle, and right third of I , respectively; thus $(\xi[I_i]_{\rightarrow} = w_i$, $i = 1, 2, 3$. Since the event $B_{\text{all paths}}^n$ holds, the straight path which steps through the elements of I from the left to the right in $3c_1n$ steps is realized at least once by the random walk $(S(t))_{t \geq 0}$ within time 2^{2n} of a stopping time $\tau(k)$, $k < 2^{\alpha n}$. Observing ξ along such a straight path generates the word $w_1w_2w_3$. Thus

$$(w_1, w_2, w_3) \in \text{PrePuzzle}^n(\text{Input}). \quad (6.10)$$

Let w'_2 be such that $(w_1, w'_2, w_3) \in \text{PrePuzzle}^n(\text{Input})$. In order to prove the claim $(w_1, w_2, w_3) \in \text{Puzzle}_1^n(\text{Input})$ it remains to show: $w_2 = w'_2$. When the event $E_{\text{stop}, \tau}^m$ holds, the stopping times of $\tau(k)$, $k < 2^{\alpha n}$, all stop the random walk $(S(t))_{t \geq 0}$ somewhere in the interval $[-2^n, 2^n]$. Within time 2^{2n} the random walk moves at most a distance $l2^{2n}$. Because of $w_1w'_2w_3 \in \text{PrePuzzle}^n(\text{Input})$, the word $w_1w'_2w_3$ occurs somewhere in the observations at most 2^{2n} time steps after a stopping time $\tau(k)$, $k < 2^{\alpha n}$. Within time 2^{2n} after a stopping time, the random walk cannot be further away from the origin than $l2^{2n} + 2^n \leq 2 \cdot l2^{2n}$, since the event $E_{\text{stop}, \tau}^m$ holds. Thus there exists an admissible piece of path $R' : [0, 3c_1n[\rightarrow [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]$ such that $\xi \circ R' = w_1w'_2w_3$. Let $R : [0, 3c_1n[\rightarrow I \subseteq [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]$ denote the right ladder path which passes through I from the left to the right. We know $\xi \circ R'[0, c_1n[= \xi \circ R'[0, c_1n[= w_1$ and $(\xi \circ R'[2c_1n, 3c_1n]_{\rightarrow} = (\xi \circ R[2c_1n, 3c_1n]_{\rightarrow} = w_3$. Furthermore, the event $E_{\text{signals II}}^n \supseteq B_{\text{signals}}^n$ holds; see Lemma 6.5. Abbreviating $x := c_1n/2$ and $y := 5c_1n/2$, this implies $R'(x) = R(x)$ and $R'(y) = R(y)$. But $R[[x, y]$ is a right ladder path; thus $R'[[x, y]$ must be the same right ladder path, since only right ladder paths can travel equally fast to the right as R does. Hence $w_2 = (\xi \circ R[[c_1n, 2c_1n]_{\rightarrow} = (\xi \circ R'[[c_1n, 2c_1n]_{\rightarrow} = w'_2$. This finishes the proof of Lemma 6.7. ■

Corollary 6.8 *If the event $B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop}, \tau}^m$ holds, then $\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_1^n(\text{Input})$.*

Proof of Corollary 6.8. Assume that $B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop}, \tau}^m$ holds, and let $I_2 \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$, $|I_2| = c_1n$, be a right ladder interval. Set $I_1 := I_2 - c_1nl_{\rightarrow}$ and $I_3 := I_2 + c_1nl_{\rightarrow}$; these are right ladder intervals adjacent to the left and to the right of I_2 , respectively. Thus $I := I_1 \cup I_2 \cup I_3$ is a right ladder interval, $|I| = 3c_1n$. Since $n \geq n_0$ and n_0 is large enough, we obtain $I \subseteq [-6 \cdot 2^n, 6 \cdot 2^n]$. We set $w_i := (\xi[I_i]_{\rightarrow}$, $i = 1, 2, 3$. We have $(w_1, w_2, w_3) \in \text{Puzzle}_1^n(\text{Input})$ by Lemma 6.7; thus $w_2 \in \text{Puzzle}_{\text{II}}^n(\text{Input})$. This finishes the proof of Corollary 6.8. ■

The following definitions are analogous to the definition of Filter_2^n and Filter_3^n , with the “reconstructed candidate” w replaced by the true scenery ξ , and with the domain $[-5 \cdot 2^n, 5 \cdot 2^n]$ replaced by the larger domain $[-9 \cdot 2^n, 9 \cdot 2^n]$. We insert the corresponding statements for left ladder intervals, too; this turns out to be useful only in the next subsection.

Definition 6.9

$$E_{\text{neighbor I}}^n := \left. \begin{array}{l} \text{For all right ladder intervals } I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n]: \text{ if } I \triangleright_n J, \text{ then} \\ ((\xi \uparrow I)_{\rightarrow}, (\xi \uparrow J)_{\rightarrow}) \in \text{Neighbors}^n(\tau, \eta). \\ \text{For all left ladder intervals } I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n]: \text{ if } I \triangleleft_n J, \text{ then} \\ ((\xi \uparrow I)_{\leftarrow}, (\xi \uparrow J)_{\leftarrow}) \in \text{Neighbors}^n(\tau, \eta). \end{array} \right\}, \quad (6.11)$$

$$E_{\text{neighbor II}}^n := \left. \begin{array}{l} \text{For all right ladder intervals } I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n], |I| = |J| = c_1 n: \\ \text{if } ((\xi \uparrow I)_{\rightarrow}, (\xi \uparrow J)_{\rightarrow}) \in \text{Neighbors}^n(\tau, \eta), \text{ then there is } q \in \mathbb{N} \text{ such that} \\ I \triangleright_n J + ql_{\rightarrow}. \\ \text{For all left ladder intervals } I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n], |I| = |J| = c_1 n: \text{ if} \\ ((\xi \uparrow I)_{\leftarrow}, (\xi \uparrow J)_{\leftarrow}) \in \text{Neighbors}^n(\tau, \eta), \text{ then there is } q \in \mathbb{N} \text{ such that } I \triangleleft_n J - ql_{\leftarrow}. \end{array} \right\}. \quad (6.12)$$

Lemma 6.10 *If the event $B_{\text{all paths}}^n$ holds, then the event $E_{\text{neighbor I}}^n$ holds too, and consequently $\xi \uparrow [-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_2^n(\text{Input})$.*

Proof of Lemma 6.10. Assume that the event $B_{\text{all paths}}^n$ holds. We treat only the case of right ladder intervals; the case of left ladder intervals can be treated in the same way by exchanging right with left, \rightarrow with \leftarrow , and \triangleright_n with \triangleleft_n .

Let $I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n]$ be right ladder intervals such that $I \triangleright_n J$. We need to prove $((\xi \uparrow I)_{\rightarrow}, (\xi \uparrow J)_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$. Let $i_l := \min I$, $i_r := \max I$, $j_l := \min J$, and $j_r := \max J$. Since $I \triangleright_n J$, there exists an admissible piece of path consisting of $h+1 = l|\mathcal{M}|+1$ points starting in i_r and ending in j_l . Since $I \triangleright_n J$ we have $|I|, |J| = c_1 n$. Thus there exists an admissible piece of path $R : [0, 2c_1 n + h - 1[\rightarrow [i_l, j_r]$ starting at i_l and ending in j_r ; furthermore we can require that $R \uparrow [0, c_1 n[$ and $R \uparrow (c_1 n + h - 1 + [0, c_1 n[$ are right ladder paths. Set $w_1 = (\xi \uparrow I)_{\rightarrow}$ and $w_2 = (\xi \uparrow J)_{\rightarrow}$; then $\xi \circ R = w_1 w w_2$ where $w \in \mathcal{C}^{h-1}$. Since $n \geq n_0$ holds and n_0 is large enough, we have $h \leq c_1 n$. Thus the piece of path R has length shorter than or equal to $3c_1 n$. The range $\text{rng}(R)$ of R fulfills $\text{rng}(R) \subseteq [-10 \cdot 2^n, 10 \cdot 2^n]$, and since $B_{\text{all paths}}^n$ holds, the random walk $(S(t))_{t \geq 0}$ “follows the path” R at least once within time 2^{2n} after a stopping time of τ . In other words, there exists $k \in [0, 2^{2n}[$ and $j \in [0, 2^{2n} - 2c_1 n - h + 1]$ such that for all $i \in [0, 2c_1 n + h - 1[$ we have $S(\tau(k) + j + i) = R(i)$. Thus we get $\xi \circ S \uparrow (\tau(k) + j + [0, 2c_1 n + h - 1]) \equiv w_1 w w_2$. This implies that $(w_1, w_2) \in \text{Neighbors}^n(\text{Input})$ and thus $((\xi \uparrow I)_{\rightarrow}, (\xi \uparrow J)_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$. ■

The following elementary number theoretic lemma serves to replace admissible pieces of path with more than h steps by admissible pieces of path with h steps, up to a sequence of maximal steps in one direction:

Lemma 6.11 *Let $s = (s_j)_{j=1, \dots, K} \in \mathcal{M}^K$, $K \in \mathbb{N}$. Then there is $(r_j)_{j=1, \dots, h} \in \mathcal{M}^h$ with*

$$\sum_{j=1}^h r_j + (K - h)l_{\rightarrow} - \sum_{j=1}^K s_j \in l_{\rightarrow} \mathbb{N}. \quad (6.13)$$

Similarly, there is $(r'_j)_{j=1, \dots, h} \in \mathcal{M}^h$ with

$$\sum_{j=1}^h r'_j - (K - h)l_{\leftarrow} - \sum_{j=1}^K s_j \in -l_{\leftarrow} \mathbb{N}. \quad (6.14)$$

Proof. In order to treat (6.13) and (6.14) simultaneously, let l_{\leftrightarrow} denote either l_{\rightarrow} or $-l_{\leftarrow}$. For $a \in \mathcal{M}$ let n_a denote the number of $j = 1, \dots, K$ such that $s_j = a$. Let $n'_a \in [0, |l_{\leftrightarrow}| \lfloor \cap(n_a + l_{\leftrightarrow})\mathbb{Z} \rfloor$ denote the remainder of n_a modulo l_{\leftrightarrow} . Then $\sum_{a \in \mathcal{M}} n'_a \leq h$. Choose any list $(r_j)_{j=1, \dots, h} \in \mathcal{M}^h$ having n'_a entries a for every $a \in \mathcal{M} \setminus \{l_{\leftrightarrow}\}$ and $h - \sum_{a \in \mathcal{M} \setminus \{l_{\leftrightarrow}\}} n'_a$ entries l_{\leftrightarrow} . Set

$$q := \frac{1}{l_{\leftrightarrow}} \sum_{a \in \mathcal{M}} (n_a - n'_a)(l_{\leftrightarrow} - a) \in \mathbb{N}; \quad (6.15)$$

note $(l_{\leftrightarrow} - a)/l_{\leftrightarrow} \geq 0$ and $n_a - n'_a \in |l_{\leftrightarrow}|\mathbb{N}$. Then

$$\begin{aligned} \sum_{j=1}^K (l_{\leftrightarrow} - s_j) &= \sum_{a \in \mathcal{M}} n_a (l_{\leftrightarrow} - a) \\ &= ql_{\leftrightarrow} + \sum_{a \in \mathcal{M}} n'_a (l_{\leftrightarrow} - a) = ql_{\leftrightarrow} + \sum_{j=1}^h (l_{\leftrightarrow} - r_j), \end{aligned} \quad (6.16)$$

which implies the claim (6.13) or (6.14), respectively. ■

Lemma 6.12 *If the event $E_{\text{signals II}}^n \cap E_{\text{stop}, \tau}^m$ holds, then the event $E_{\text{neighbor II}}^n$ holds, too, and consequently $\xi \llbracket [-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_3^n(\text{Input})$.*

Proof. Assume that the events $E_{\text{signals II}}^n$ and $E_{\text{stop}, \tau}^m$ hold. We treat here the case of right ladder intervals:

Let $I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n]$ be right ladder intervals with $|I| = |J| = c_1 n$, and assume $((\xi \lceil I \rceil_{\rightarrow}, (\xi \lceil J \rceil_{\rightarrow}) \in \text{Neighbors}^n(\text{Input}))$. We need to show $I \triangleright_n J + ql_{\rightarrow}$ for some $q \in \mathbb{N}$.

Using Definition 5.7 of Neighbors^n and the abbreviations $w_1 := (\xi \lceil I \rceil_{\rightarrow}$ and $w_2 := (\xi \lceil J \rceil_{\rightarrow}$, we see: There is an admissible piece of path $R : [0, 2c_1 n + h - 1]_{\rightarrow} \mathbb{Z}$ with the following properties:

- R is realized by the random walk S in during some time interval $D \subseteq \tau(k) + [0, 2^{2n}]$, $|D| = 2c_1 n + h - 1$, for some $k \in [0, 2^{2n}]$. This means: R equals $S \lceil D$ when time-shifted back to the origin.
- Observing the scenery ξ along R produces $w_1 w w_2$ for some $w \in \mathcal{C}^{h-1}$; i.e.: $\xi \circ R = w_1 w w_2$.

We know $|\tau(k)| \leq 2^n$ since the event $E_{\text{stop}, \tau}^m$ holds; thus R takes all its values in $[-(2^n + l2^{2n}), 2^n + l2^{2n}] \subseteq [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]$, since the random walk cannot travel faster than distance l per step. We examine the first $c_1 n$ steps of R : $(\xi \circ R \lceil [0, c_1 n] \rceil_{\rightarrow} = w_1 = (\xi \lceil I \rceil_{\rightarrow}$ implies $R(c_1 n/2) = \min I + c_1 n l_{\rightarrow}/2$, since the event $E_{\text{signals II}}^n$ holds; note that $x := \min I + c_1 n l_{\rightarrow}/2$ is the point in the middle of a right ladder path walking through I . The same argument applies to the last $c_1 n$ steps of R : $(\xi \circ R \lceil (c_1 n + h - 1 + [0, c_1 n]) \rceil_{\rightarrow} = w_2 = (\xi \lceil J \rceil_{\rightarrow}$ implies $R(3c_1 n/2 + h - 1) = \min J + c_1 n l_{\rightarrow}/2 =: y$; y is the point in the middle of J . The path R travels from x to y in $K := c_1 n + h - 1 \geq h$ steps, using some step sizes $(s_j)_{j=1, \dots, K} \in \mathcal{M}^K$. As a consequence of (6.13) in Lemma 6.11, there is $(r_j)_{j=1, \dots, h} \in \mathcal{M}^h$ with $\sum_{j=1}^h r_j + (K - h)l_{\rightarrow} - \sum_{j=1}^K s_j = ql_{\rightarrow}$ for some $q \in \mathbb{N}$. Since $\max I - x = (c_1 n/2 - 1)l_{\rightarrow}$ and $y - \min J = c_1 n l_{\rightarrow}/2$, we obtain $\min J - \max I = y - x - (c_1 n - 1)l_{\rightarrow} = \sum_{j=1}^K s_j - (c_1 n - 1)l_{\rightarrow} = \sum_{j=1}^h r_j - ql_{\rightarrow}$. This means $I \triangleright_n (J + ql_{\rightarrow})$, as we wanted to show.

Summarizing, this implies $\xi \llbracket [-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_3^n(\text{Input})$ and the first statement in the definition of $E_{\text{neighbor II}}^n$, which treats right ladder intervals.

The proof for left ladder intervals can be treated analogously. Altogether, we see that the event $E_{\text{neighbor II}}^n$ is valid. ■

Definition 6.13 We define the event

$$B_{\text{seed I}}^n := \left\{ \begin{array}{l} \text{For every modulo class } Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z} \text{ there exists } k \in [0, 2^{\alpha n}[\text{ such} \\ \text{that } S(\tau(k) + h) \in Z, S[(\tau(k) + h + [0, 3c_1nl_{\leftarrow}]]) \text{ is a right ladder} \\ \text{path, and } S[(\tau(k) + h + 3c_1nl_{\leftarrow} + [0, 3c_1nl_{\rightarrow}])] \text{ is a left ladder path.} \end{array} \right\}. \quad (6.17)$$

Lemma 6.14 If the events $B_{\text{all paths}}^n$, B_{signals}^n , $B_{\text{seed I}}^n$ and $E_{\text{stop}, \tau}^m$ hold, then $\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_4^n(\text{Input})$.

Proof of Lemma 6.14. Assume that the event $B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap B_{\text{seed I}}^n \cap E_{\text{stop}, \tau}^m$ holds. Let $Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$. Since $B_{\text{seed I}}^n$ holds, there exists a $k \in [0, 2^{\alpha n}[$ such that $S(\tau(k) + h) \in Z$, $R_1 := S[(\tau(k) + h + [0, 3c_1nl_{\leftarrow}]])$ is a right ladder path, and $R_2 := S[(\tau(k) + h + 3c_1nl_{\leftarrow} + [0, 3c_1nl_{\rightarrow}])]$ is a left ladder path. Since $E_{\text{stop}, \tau}^m$ holds, we know $S(\tau(k)) \in [-2^n, 2^n]$. Thus the random walk S cannot leave the interval $[-2 \cdot 2^n, 2 \cdot 2^n]$ during the time interval $\tau(k) + [h + 3c_1nl_{\leftarrow} + 3c_1nl_{\rightarrow}]$, since $(h + 3c_1nl_{\leftarrow} + 3c_1nl_{\rightarrow})l < 2^n$, and the random walk cannot travel faster than l per step. Thus R_1 and R_2 take all their values in $[-2 \cdot 2^n, 2 \cdot 2^n]$. Note that the right ladder path R_1 and the left ladder path R_2 traverse precisely the same interval, R_1 using step size l_{\rightarrow} to the right, and R_2 with step size $-l_{\leftarrow}$ back. The same is true when we restrict R_1 and R_2 to the smaller time intervals $[t_1, t'_1] := \tau(k) + h + c_1nl_{\leftarrow} + [0, c_2nl_{\leftarrow}]$ and $[t_2, t'_2] := \tau(k) + h + 3c_1nl_{\leftarrow} + 2c_1nl_{\rightarrow} + [-c_2nl_{\rightarrow}, 0]$, respectively: We have $S(t_1) = S(t'_2) =: a$, and $S(t'_1) = S(t_2) =: b$, and $S[[t_1, t'_1]]$ is a right ladder path: it traverses $[a, b]$ from the left to the right, while on $S[[t_2, t'_2]]$ it is a left ladder path; it traverses $[a, b]$ in opposite direction. In particular, reading only every l_{\leftarrow} th letter in $\chi[[t_1, t'_1]]$ and only every l_{\rightarrow} th letter in $\chi[[t_2, t'_2]]$ yield the same word, only in reversed direction:

$$(\chi[[t_1, t'_1] \cap (t_1 + l_{\leftarrow}\mathbb{Z})])_{\rightarrow} = (\xi[[a, b] \cap (a + l_{\rightarrow}l_{\leftarrow}\mathbb{Z})])_{\rightarrow} = (\chi[[t_2, t'_2] \cap (t_1 + l_{\leftarrow}\mathbb{Z})])_{\leftarrow}. \quad (6.18)$$

We consider the words $u_1u_2u_3 := \chi[(t_1 - c_1n + [0, 3c_1n])]$ and $v_1v_2v_3 := \chi[(t_2 - c_1n + [0, 3c_1n])]$ with $u_i, v_i \in \mathcal{C}^{c_1n}$; note that $t_1 - c_1n + [0, 3c_1n] \subseteq \text{domain}(R_1)$ and $t_2 - c_1n + [0, 3c_1n] \subseteq \text{domain}(R_2)$. We get $(u_1, u_2, u_3), (v_1, v_2, v_3) \in \text{Puzzle}_1^n(\text{Input})$ by Lemma 6.7. Hence we obtain $(w_1, w_2, w_3) \in \text{Seed}_1^n(\text{Input})$ by Definition (5.4), since the words $u_1u_2u_3$ and $v_1v_2v_3$ occur in the observations sufficiently close to a stopping time $\tau(k)$; more specifically: $t_1 - c_1n, t_2 - c_1n \in \tau(k) + [0, 7c_1nl]$. Consequently $u_2, v_2 \in \text{Seed}_{\text{II}}^n(\text{Input})$ by Definition (5.5). Finally we observe

$$(u_2[[0, c_2nl_{\leftarrow}] \cap l_{\leftarrow}\mathbb{Z}])_{\rightarrow} = (\xi[[a, b] \cap (a + l_{\rightarrow}l_{\leftarrow}\mathbb{Z})])_{\rightarrow} = (v_2[[0, c_2nl_{\rightarrow}] \cap l_{\rightarrow}\mathbb{Z}])_{\leftarrow} \quad (6.19)$$

by (6.18). Thus we have shown $u_2 \in \text{Seed}_{\text{III}}^n(\text{Input})$, see (5.6). Since $u_2 = \xi \circ S[(t_1 + [0, c_1n])]$, and since $S[(t_1 + [0, c_1n])]$ is a right ladder path with values in $Z \cap [-2 \cdot 2^n, 2 \cdot 2^n]$, this implies $\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_4^n(\text{Input})$. ■

Definition 6.15 For $n \in \mathbb{N}$, we define the following event:

$$B_{\text{unique fit}}^n := \left\{ \begin{array}{l} \text{For every } i, j \in \{1, \dots, l^2\}, \text{ every } i\text{-spaced interval } I \subseteq [-11 \cdot \\ 2^n, 11 \cdot 2^n], \text{ and every } j\text{-spaced interval } J \subseteq [-11 \cdot 2^n, 11 \cdot 2^n] \text{ with} \\ |I| = |J| \geq c_2n \text{ holds } (\xi[I]_{\leftarrow} \neq (\xi[J]_{\rightarrow}), \text{ and if } I \neq J, \text{ then} \\ (\xi[I]_{\rightarrow} \neq (\xi[J]_{\leftarrow}). \end{array} \right\}, \quad (6.20)$$

Lemma 6.16 If the event $B_{\text{unique fit}}^n$ holds, then $\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_5^n(\text{Input})$.

Proof. Using $c_2 \leq c_1$ (see subsection 2.1), this follows immediately from Definition 6.15 of the event $B_{\text{unique fit}}^n$, and of Definition 5.4 of Filter_5^n . ■

Theorem 6.17 $B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap B_{\text{seed I}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{stop},\tau}^m \subseteq E_{\text{xi does it}}^n$

Proof. We collect the statements of Lemmas/Corollary 6.5, 6.8, 6.10, 6.12, 6.14, and 6.16 in the following list:

$$\begin{array}{ll}
B_{\text{signals}}^n & \subseteq E_{\text{signals II}}^n, \\
B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop},\tau}^m & \subseteq \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_1^n(\text{Input})]\}, \\
B_{\text{all paths}}^n & \subseteq \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_2^n(\text{Input})]\}, \\
E_{\text{signals II}}^n \cap E_{\text{stop},\tau}^m & \subseteq \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_3^n(\text{Input})]\}, \\
B_{\text{all paths}}^n \cap B_{\text{signals}}^n \cap B_{\text{seed I}}^n \cap E_{\text{stop},\tau}^m & \subseteq \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_4^n(\text{Input})]\}, \\
B_{\text{unique fit}}^n & \subseteq \{\xi[[-5 \cdot 2^n, 5 \cdot 2^n] \in \text{Filter}_5^n(\text{Input})]\}.
\end{array}$$

The theorem is an immediate consequence these statements, using (6.1) and (5.9). ■

6.2 Combinatorics concerning $E_{\text{all pieces ok}}^n$

In this subsection, we show that a piece w that passes all the Filter_i occurs in the true scenery ξ near the origin, provided some “basic” events B_{\dots}^n hold.

Definition 6.18 *We define the events*

$$\begin{aligned}
B_{\text{recogn straight}}^n &:= & (6.21) \\
&\left\{ \begin{array}{l} \text{For every } R \in \text{AdPaths}(11 \cdot 2^n, c_1 n) \text{ with } R(c_1 n - 1) - R(0) \notin \{(c_1 n - 1)l_{\rightarrow}, -(c_1 n - 1)l_{\leftarrow}\} \\ \text{there is } \bar{R} \in \text{AdPaths}(12 \cdot 2^n, c_1 n) \text{ such that } R(0) = \bar{R}(0), R(c_1 n - 1) = \bar{R}(c_1 n - 1), \text{ and} \\ \xi \circ R \neq \xi \circ \bar{R}. \end{array} \right\},
\end{aligned}$$

$$E_{\text{only ladder}}^n := \left\{ \begin{array}{l} \text{For all } (w_1, w_2, w_3) \in \text{Puzzle}_1^n(\text{Input}) \text{ and every admissible piece of} \\ \text{path } R : [0, 3c_1 n[\rightarrow [-11 \cdot 2^n, 11 \cdot 2^n] \text{ with } \xi \circ R = w_1 w_2 w_3 \text{ holds:} \\ w_2 \text{ is a ladder word of } \xi[[-11 \cdot 2^n, 11 \cdot 2^n]. \end{array} \right\}. \quad (6.22)$$

Lemma 6.19 *We have*

$$B_{\text{all paths}}^n \cap B_{\text{recogn straight}}^n \subseteq E_{\text{only ladder}}^n. \quad (6.23)$$

Proof of Lemma 6.19. Assume that the event $B_{\text{all paths}}^n \cap B_{\text{recogn straight}}^n$ holds. Let $w_1 w_2 w_3 \in \text{Puzzle}_1^n(\text{Input})$, and let $R : [0, 3c_1 n[\rightarrow [-11 \cdot 2^n, 11 \cdot 2^n]$ be an admissible piece of path with $\xi \circ R = w_1 w_2 w_3$. We prove by contradiction that the event $E_{\text{only ladder}}^n$ holds: Assume w_2 is not a ladder word of $\xi[[-11 \cdot 2^n, 11 \cdot 2^n]$. Since $B_{\text{recogn straight}}^n$ holds, there exists an admissible piece of path $\bar{R} : [c_1 n, 2c_1 n[\rightarrow [-11 \cdot 2^n, 11 \cdot 2^n]$ such that $R(c_1 n) = \bar{R}(c_1 n)$ and $R(2c_1 n - 1) = \bar{R}(2c_1 n - 1)$, but $w_2 \neq (\xi \circ \bar{R})_{\rightarrow} =: w'_2$. Let $\check{R} : [0, 3c_1 n[\rightarrow [-11 \cdot 2^n, 11 \cdot 2^n]$ be the admissible piece of path which on $[c_1 n, 2c_1 n[$ is equal to \bar{R} and otherwise is equal to R . We have $\xi \circ \check{R} = w_1 w'_2 w_3$. Since $B_{\text{all paths}}^n$ holds, too, this implies that the random walk S follows the path of \check{R} within time 2^{2n} from a stopping time of $\tau(k)$, $k < 2^{\alpha n}$. The same is valid for R , maybe with a different stopping time $\tau(k')$. In other words: $w_1 w'_2 w_3 \in \text{PrePuzzle}^n(\text{Input})$ and $w_1 w_2 w_3 \in \text{PrePuzzle}^n(\text{Input})$. This implies the contradiction $w_1 w_2 w_3 \notin \text{Puzzle}_1^n(\text{Input})$; thus we have proved Lemma 6.19. ■

Definition 6.20 *We define the events*

$$B_{\text{outside out}}^n := \left\{ \begin{array}{l} \text{For every admissible piece of path} \\ R \in ([-2 \cdot l2^{2n}, 2 \cdot l2^{2n}] \setminus [-10 \cdot 2^n, 10 \cdot 2^n])^{[0, c_1 n/2]}: \xi \circ R \text{ is not strongly equivalent} \\ \text{to any ladder word of length } c_1 n/2 \text{ of } \xi[[-9 \cdot 2^n, 9 \cdot 2^n]. \end{array} \right\}, \quad (6.24)$$

$$E_{\text{mod class}}^n := \left\{ \begin{array}{l} \text{For all } w \in \text{Filter}_1^n(\text{Input}) \text{ and for all right ladder intervals } I \subseteq [-2 \cdot 2^n, 2 \cdot 2^n], |I| = c_1 n: \\ \text{If there is a right ladder interval } J_r \subseteq [-2 \cdot 2^n, 2 \cdot 2^n] \text{ with } w \upharpoonright I \equiv \xi \upharpoonright J_r, \text{ then} \\ \xi \upharpoonright ([-2^n, 2^n] \cap (J_r + l_{\rightarrow} \mathbb{Z})) \sqsubseteq w \upharpoonright (I + l_{\rightarrow} \mathbb{Z}) \sqsubseteq \xi \upharpoonright ([-9 \cdot 2^n, 9 \cdot 2^n] \cap (J_r + l_{\rightarrow} \mathbb{Z})), \text{ and} \\ \text{if } l_{\rightarrow} = l_{\leftarrow} \text{ and if there is a (left) ladder interval } J_l \subseteq [-2 \cdot 2^n, 2 \cdot 2^n] \text{ with } (w \upharpoonright I)^{\leftrightarrow} \equiv \xi \upharpoonright J_l, \\ \text{then } \xi \upharpoonright ([-2^n, 2^n] \cap (J_l + l_{\mathbb{Z}})) \sqsubseteq (w \upharpoonright (I + l_{\mathbb{Z}}))^{\leftrightarrow} \sqsubseteq \xi \upharpoonright ([-9 \cdot 2^n, 9 \cdot 2^n] \cap (J_l + l_{\mathbb{Z}})). \end{array} \right\}. \quad (6.25)$$

Informally speaking, the meaning of the event $E_{\text{mod class}}^n$ is the following: If a “reconstructed” piece of scenery w contains a correct “seed piece” $w \upharpoonright I$ over a sufficiently long ladder word, then the whole modulo class generated by I is reconstructed correctly. The reconstruction may generate the wrong orientation, but this is only allowed if left ladder intervals and right ladder intervals coincide, and if already the “seed piece” $w \upharpoonright I$ is reversed compared with the true scenery ξ .

The next lemma formalizes the intuitive idea of “playing a puzzle game”: We start with a seed word as reconstructed piece; then we append successively pieces of our puzzle that match to an ending of the growing reconstructed piece. This procedure continues until the reconstructed piece is large enough.

Lemma 6.21 *We have*

$$B_{\text{outside out}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{only ladder}}^n \cap E_{\text{stop}, \tau}^m \subseteq E_{\text{mod class}}^n \quad (6.26)$$

Proof of Lemma 6.21. Assume that the events on the left hand side of (6.26) hold. We claim that then $E_{\text{mod class}}^n$ holds, too. To prove this claim, let $w \in \text{Filter}_1^n(\text{Input})$, and let $I \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$, $|I| = c_1 n$ be a right ladder interval. Assume that $J \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$ is a ladder interval. We assume one of the following two cases:

- A) J is a right ladder interval, and $w \upharpoonright I \equiv \xi \upharpoonright J$;
- B) $l_{\rightarrow} = l_{\leftarrow}$ and $(w \upharpoonright I)^{\leftrightarrow} \equiv \xi \upharpoonright J$.

We treat both cases simultaneously as far as possible; in order to unify the notation, let \cdot^{\sim} denote the reversion operation \cdot^{\leftrightarrow} in case B and the identity operation in case A. We set $Z := J + l_{\rightarrow} \mathbb{Z} \in \mathbb{Z}/l_{\rightarrow} \mathbb{Z}$; then it remains to show:

$$\xi \upharpoonright ([-2^n, 2^n] \cap Z) \sqsubseteq (w \upharpoonright (I + l_{\rightarrow} \mathbb{Z}))^{\sim} \sqsubseteq \xi \upharpoonright ([-9 \cdot 2^n, 9 \cdot 2^n] \cap Z). \quad (6.27)$$

To prove the right hand side of (6.27), we prove by induction over all right ladder intervals I' with $I \subseteq I' \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$:

$$(w \upharpoonright I')^{\sim} \sqsubseteq \xi \upharpoonright ([-9 \cdot 2^n, 9 \cdot 2^n] \cap Z). \quad (6.28)$$

Once we have proven this, the right hand side of (6.27) follows from the special case $I' = [-5 \cdot 2^n, 5 \cdot 2^n] \cap (I + l_{\rightarrow} \mathbb{Z})$.

The induction starts with $I = I'$: in this case (6.28) holds since our assumption A) or B), respectively, implies $(w[I] \sim \sqsubseteq \xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$. For the induction step, assume that (6.28) holds for some I' . We enlarge I' by a single new point: let $I'' = I' \cup \{i\} \subseteq [-5 \cdot 2^n, 5 \cdot 2^n] \cap (I + l_{\rightarrow} \mathbb{Z})$ be a right ladder interval, $i \notin I'$. Let $I_i \subseteq I''$ be a right ladder interval with $|I_i| = c_1 n$ and $i \in I_i$. Using $w \in \text{Filter}_1^n(\text{Input})$ we see $w_2 := (w[I_i]_{\rightarrow} \in \text{Puzzle}_{\text{II}}^n(\text{Input})$. Hence there are $w_1, w_3 \in \mathcal{C}^{c_1 n}$ such that $(w_1, w_2, w_3) \in \text{Puzzle}_1^n(\text{Input}) \subseteq \text{PrePuzzle}^n(\text{Input})$. Thus $w_1 w_2 w_3$ occurs in the observation χ at most 2^{2n} time steps after a stopping time $\tau(k)$, $k < 2^{\alpha n}$; say $w_1 w_2 w_3$ is read there in χ while the random walk follows an admissible piece of path $R : [0, 3c_1 n] \rightarrow \mathbb{Z}$; (we shifted the time domain of R back to the origin). Since the event $E_{\text{stop}, \tau}^m$ holds, we have $|S(\tau(k))| \leq 2^n$. Within time 2^{2n} the random walk cannot travel farther than distance $l2^{2n}$; thus R has all its values in $[-(2^n + l2^{2n}), 2^n + l2^{2n}] \subseteq [-2 \cdot l2^{2n}, 2 \cdot l2^{2n}]$. Consider the ladder interval $I'_i := I_i \setminus \{i\} = I_i \cap I'$, $|I'_i| = c_1 n - 1 \geq c_1 n/2$: the induction hypothesis (6.28) implies $(w[I'_i] \sim \sqsubseteq \xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$; say $(w[I'_i] \sim \equiv \xi[D')$ for some right ladder interval $D' \subseteq [-9 \cdot 2^n, 9 \cdot 2^n] \cap Z$. Furthermore, $w'_2 := (w[I'_i]_{\rightarrow}$ is a subword of $w_2 = (w[I_i]_{\rightarrow}$ and thus also a subword of $\xi \circ R$. Hence we see, using that the event $B_{\text{outside out}}^n$ holds: R cannot take all of its values outside $[-10 \cdot 2^n, 10 \cdot 2^n]$; thus it has all its values in $[-10 \cdot 2^n - 3c_1 n l, 10 \cdot 2^n + 3c_1 n l] \subseteq [-11 \cdot 2^n, 11 \cdot 2^n]$. Since the event $E_{\text{only ladder}}^n$ holds, $w_2 = (w[I_i]_{\rightarrow}$ is a ladder word of $\xi[[-11 \cdot 2^n, 11 \cdot 2^n]$; say $w_2 = (\xi[D]_{\rightarrow}$ for some right ladder interval $D \subseteq [-11 \cdot 2^n, 11 \cdot 2^n]$ (we call this “case A₁”), or $w_2 = (\xi[D]_{\leftarrow}$ for some left ladder interval $D \subseteq [-11 \cdot 2^n, 11 \cdot 2^n]$ (call this “case B₁”). Thus w'_2 occurs as a (possibly reversed) ladder word

- as a subword of $(\xi[D]_{\rightarrow}$ in case A₁, or as a subword of $(\xi[D]_{\leftarrow}$ in case B₁;
- as $w'_2 = (\xi[D']_{\rightarrow}$ in case A, or as $w'_2 = (\xi[D']_{\leftarrow}$ in case B.

Since the event $B_{\text{unique fit}}^n$ holds, this implies $D' \subseteq D$, and furthermore the reading directions have to coincide: If case A holds, then case A₁ occurs, and if case B holds, then case B₁ occurs. Let $T : \mathbb{Z} \rightarrow \mathbb{Z}$ denote the translation (case A) or reflection (case B) that transports $w[I_i]$ to $\xi[D]$. Then T transports $w[I'_i]$ to $\xi[D']$, and thus – using once more that $B_{\text{unique fit}}^n$ holds – T is also the map that transports $w[I']$ to a subpiece of $\xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$ according to the induction hypothesis (6.28). Hence T transports $w[(I_i \cup I')] = w[I'']$ to an equivalent subpiece of $\xi[[-11 \cdot 2^n, 11 \cdot 2^n]$. To see that $T[w[I'']]$ is already a subpiece of $\xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$, we proceed as follows: T maps the nonempty seed interval $I \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$ to $J \subseteq [-2 \cdot 2^n, 2 \cdot 2^n] \cap Z$; thus it has the form $T(z) = \pm z + a$ with $|a| \leq 4 \cdot 2^n$. Consequently T maps the domain $[-5 \cdot 2^n, 5 \cdot 2^n]$ of w to a subset of $[-9 \cdot 2^n, 9 \cdot 2^n]$. This shows $(w[I''] \sim \sqsubseteq \xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$, which finishes our induction step and also the proof of the right hand side of the claim (6.27).

To prove the left hand side of (6.27), we observe that T^{-1} maps $[-2^n, 2^n]$ to a subset of $[-5 \cdot 2^n, 5 \cdot 2^n]$. Since T maps I to J , it maps the modulo class $I + l_{\rightarrow} \mathbb{Z}$ to $Z = J + l_{\rightarrow} \mathbb{Z}$; thus T^{-1} maps $[-2^n, 2^n] \cap Z$ to a subset of $(I + l_{\rightarrow} \mathbb{Z}) \cap [-5 \cdot 2^n, 5 \cdot 2^n] = (I + l_{\rightarrow} \mathbb{Z}) \cap \text{domain}(w)$. Since T^{-1} maps a subpiece of $\xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap Z)$ to $w[(I + l_{\rightarrow} \mathbb{Z})$, this implies the left hand side of the claim (6.27). This finishes the proof of Lemma 6.21. ■

Definition 6.22 *We define the event*

$$E_{\text{seed II}}^n := \left\{ \begin{array}{l} \text{Every } u \in \text{Seed}_{\text{II}}^n(\text{Input}) \text{ is a left or right ladder word of } \xi[[-2 \cdot \cdot \\ 2^n, 2 \cdot 2^n]. \text{ If } l_{\rightarrow} \neq l_{\leftarrow}, \text{ then every } u \in \text{Seed}_{\text{III}}^n(\text{Input}) \text{ is a right} \\ \text{ladder word of } \xi[[-2 \cdot 2^n, 2 \cdot 2^n]. \end{array} \right\}. \quad (6.29)$$

Lemma 6.23 *We have*

$$B_{\text{unique fit}}^n \cap B_{\text{signals}}^n \cap B_{\text{all paths}}^n \cap B_{\text{recogn straight}}^n \cap E_{\text{stop}, \tau}^m \subseteq E_{\text{seed II}}^n. \quad (6.30)$$

Proof of Lemma 6.23. Assume that the events on the left hand side of (6.30) hold. In order to show that the $E_{\text{seed II}}^n$ holds, let $w_2 \in \text{Seed}_{\text{II}}^n(\text{Input})$. We need to show that w_2 is a ladder word of $\xi[-2 \cdot 2^n, 2 \cdot 2^n]$. Using (5.5), we take $w_1, w_3 \in \mathcal{C}^{c_1 n}$ with $(w_1, w_2, w_3) \in \text{Seed}_{\text{I}}^n(\text{Input})$; thus $w_1 w_2 w_3 \equiv \eta[(\tau(k) + j + [0, 3c_1 n])]$ for some $k < 2^{\alpha n}$ and $j \in [0, 7c_1 n]$. Since $E_{\text{stop}, \tau}^m$ holds, we have $|S(\tau(k))| \leq 2^n$. Using $2^n + 7c_1 n l^2 + 3c_1 l n \leq 2 \cdot 2^n - c_1 n l$, we see that the random walk S is located inside the interval $[-2 \cdot 2^n + c_1 n l, 2 \cdot 2^n - c_1 n l]$ during the time interval $\tau(k) + 7c_1 n l + [0, 3c_1 n]$. The word $w_1 w_2 w_3$ is read along an admissible piece of path, say $R \in \text{AdPath}(2 \cdot 2^n - c_1 n l, 3c_1 n)$ with $\xi \circ R = w_1 w_2 w_3$; (the time interval is shifted back to the origin). The event $E_{\text{only ladder}}^n$ holds by Lemma 6.19, and we have $(w_1, w_2, w_3) \in \text{Puzzle}_{\text{I}}^n(\text{Input})$; hence w_2 is a ladder word of $\xi[-11 \cdot 2^n, 11 \cdot 2^n]$; say $w_2 = \xi \circ \pi$ for a ladder path $\pi : [0, c_1 n[\rightarrow [-11 \cdot 2^n, 11 \cdot 2^n]$. Let $\pi' = R[[c_1 n, 2c_1 n[$ be the middle piece of R , along which one observes $(\xi \circ \pi')_{\rightarrow} = w_2 = \xi \circ \pi$. Since the event $E_{\text{signals II}}^n$ holds by Lemma 6.5, we get $\pi'((3/2)c_1 n) = \pi(c_1 n/2)$; thus π takes least one value in $[-(2 \cdot 2^n - c_1 n l), 2 \cdot 2^n - c_1 n l]$; therefore all the values of π are in $[-2 \cdot 2^n, 2 \cdot 2^n]$. Thus w_2 is a ladder word of $\xi[-2 \cdot 2^n, 2 \cdot 2^n]$.

For the rest of the proof we assume $l_{\rightarrow} \neq l_{\leftarrow}$ and let $u \in \text{Seed}_{\text{III}}^n(\text{Input})$. It remains to show: u is a right ladder word of $\xi[-2 \cdot 2^n, 2 \cdot 2^n]$. Using Definition (5.6) of $\text{Seed}_{\text{III}}^n$, we choose $v \in \text{Seed}_{\text{II}}^n(\text{Input})$ with $(u[(l_{\leftarrow} \mathbb{Z} \cap [0, c_2 n l_{\leftarrow}]))_{\rightarrow} = (v[(l_{\rightarrow} \mathbb{Z} \cap [0, c_2 n l_{\rightarrow}]))_{\leftarrow}$. From the first part of the proof we get: u and v are ladder words of $\xi[-2 \cdot 2^n, 2 \cdot 2^n]$, since $u, v \in \text{Seed}_{\text{II}}^n(\text{Input})$. We distinguish three cases:

1. u is a right ladder word;
2. u and v are left ladder words;
3. u is a left ladder word and v is a right ladder word.

We need to show that case 1. holds; thus we prove that the cases 2. and 3. lead to a contradiction:

In case 2., let $u = (\xi[I]_{\leftarrow}$ and $v = (\xi[J]_{\leftarrow}$ for some left ladder intervals $I, J \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$, $|I| = |J| = c_1 n$. We get $(u[(l_{\leftarrow} \mathbb{Z} \cap [0, c_2 n l_{\leftarrow}]))_{\rightarrow} = (\xi[I']_{\leftarrow}$ for some l_{\leftarrow}^2 -spaced interval $I' \subseteq I$, $|I'| = c_2 n + 1$. Similarly, $(v[(l_{\rightarrow} \mathbb{Z} \cap [0, c_2 n l_{\rightarrow}]))_{\leftarrow} = (\xi[J']_{\rightarrow}$ for some $l_{\rightarrow} l_{\leftarrow}$ -spaced interval $J' \subseteq J$, $|J'| = c_2 n + 1$. Thus $(\xi[I']_{\leftarrow} = (\xi[J']_{\rightarrow}$, which is incompatible with the event $B_{\text{unique fit}}^n$.

In case 3., let $u = (\xi[I]_{\leftarrow}$ for some left ladder interval $I \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$ and $v = (\xi[J]_{\rightarrow}$ for some right ladder interval $J \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$, $|I| = |J| = c_1 n$. We get again $(u[(l_{\leftarrow} \mathbb{Z} \cap [0, c_2 n l_{\leftarrow}]))_{\rightarrow} = (\xi[I']_{\leftarrow}$ for some l_{\leftarrow}^2 -spaced interval $I' \subseteq I$, $|I'| = c_2 n + 1$. This time we have $(v[(l_{\rightarrow} \mathbb{Z} \cap [0, c_2 n l_{\rightarrow}]))_{\leftarrow} = (\xi[J']_{\leftarrow}$ for some l_{\rightarrow}^2 -spaced interval $J' \subseteq J$, $|J'| = c_2 n + 1$. Since $l_{\leftarrow}^2 \neq l_{\rightarrow}^2$, we have $I' \neq J'$. We obtain $(\xi[I']_{\leftarrow} = (\xi[J']_{\leftarrow}$, which is incompatible with the event $B_{\text{unique fit}}^n$.

Thus cases 2. and 3. cannot occur. Summarizing, we have proven that the event $E_{\text{seed II}}^n$ holds. \blacksquare

Definition 6.24 *If $l_{\rightarrow} = l_{\leftarrow}$, we define the event*

$$E_{\text{dist}}^n := \left\{ \begin{array}{l} \text{For all ladder intervals } I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n], |I| = |J| = c_1 n: \text{ if at least one} \\ \text{of } ((\xi[I]_{\rightarrow}, (\xi[J]_{\rightarrow})), ((\xi[I]_{\rightarrow}, (\xi[J]_{\leftarrow})), ((\xi[I]_{\leftarrow}, (\xi[J]_{\rightarrow})), \text{ or } ((\xi[I]_{\leftarrow}, (\xi[J]_{\leftarrow})) \text{ is in} \\ \text{Neighbors}^n(\text{Input}), \text{ then distance}(I, J) \leq 3 \cdot l c_1 n. \end{array} \right\} \quad (6.31)$$

In the case $l_{\rightarrow} \neq l_{\leftarrow}$, we set E_{dist}^n to be the sure event.

Lemma 6.25 $B_{\text{signals}}^n \cap E_{\text{stop},\tau}^m \subseteq E_{\text{dist}}^n$

Proof of Lemma 6.25. Assume that the event $B_{\text{signals}}^n \cap E_{\text{stop},\tau}^m$ holds, and that $l_{\rightarrow} = l_{\leftarrow} = l$. Let $I, J \subseteq [-9 \cdot 2^n, 9 \cdot 2^n]$, $|I| = |J| = c_1 n$ be right ladder intervals, and assume that there is a (w_1, w_2) among $((\xi \upharpoonright I)_{\rightarrow}, (\xi \upharpoonright J)_{\rightarrow}), ((\xi \upharpoonright I)_{\rightarrow}, (\xi \upharpoonright J)_{\leftarrow}), ((\xi \upharpoonright I)_{\leftarrow}, (\xi \upharpoonright J)_{\rightarrow}),$ or $((\xi \upharpoonright I)_{\leftarrow}, (\xi \upharpoonright J)_{\leftarrow})$ with $(w_1, w_2) \in \text{Neighbors}^n(\text{Input})$. By definition (5.7), some word $w_1 w w_2$ with $w \in \mathcal{C}^{h-1}$ occurs in the observations χ at most 2^{2n} time steps after a stopping time $\tau(k)$, $k < 2^{\alpha n}$. Since $E_{\text{stop},\tau}^m$ holds, the random walk remains in the interval $[-2 \cdot l 2^{2n}, 2 \cdot l 2^{2n}]$ during that time interval; say the random walk follows an admissible piece of path $R : [0, 2c_1 n + h - 1 \rightarrow [-2 \cdot l 2^{2n}, 2 \cdot l 2^{2n}]$ while producing the observations $\xi \circ R = w_1 w w_2$; (we shifted the time domain back to the origin). R consists of the three pieces $\pi'_1 = R \upharpoonright [0, c_1 n]$, $\pi' = R \upharpoonright [c_1 n + [0, h - 1]$, and $\pi'_2 = R \upharpoonright [c_1 n + h - 1 + [0, c_1 n]$ with $\xi \circ \pi'_1 = w_1$, $(\xi \circ \pi')_{\rightarrow} = w$, and $(\xi \circ \pi'_2)_{\rightarrow} = w_2$. Let $x_1 := c_1 n / 2$ and $x_2 := (3/2)c_1 n + h - 1$ be the points in the middle of the domain of π'_1 and π'_2 , respectively. Then

$$|\pi'_1(x_1) - \pi'_2(x_2)| \leq (c_1 n + h - 1)l, \quad (6.32)$$

since the path R cannot travel faster than l per step. The event $E_{\text{signals II}}^n$ holds by Lemma 6.5. Let $\pi_1 : [0, c_1 n \rightarrow I$ and $\pi_2 : c_1 n + h - 1 + [0, c_1 n \rightarrow J$ be ladder paths with range I and J , respectively; we choose these paths to be left or right ladder paths according to whether the reading direction is “ \leftarrow ” or “ \rightarrow ”. Hence, using $\xi \circ \pi_1 = w_1 = \xi \circ \pi'_1$ and $(\xi \circ \pi_2)_{\rightarrow} = w_2 = (\xi \circ \pi'_2)_{\rightarrow}$, we obtain $\pi'_1(x_1) = \pi_1(x_1)$ and $\pi'_2(x_2) = \pi_2(x_2)$. Consequently (6.32) implies

$$\text{distance}(I, J) \leq |\pi_1(x_1) - \pi_2(x_2)| \leq 3 \cdot l c_1 n. \quad (6.33)$$

Summarizing, we have shown that the event E_{dist}^n holds. ■

The following event $E_{\text{mod } \gamma \text{ ok}}^n$ compares modulo classes (modulo some γ) in “reconstructed” pieces w with modulo classes in the “true” scenery ξ . Roughly speaking, it states that all modulo classes are reconstructed correctly, and either all of them are reconstructed in the correct orientation (“case A”), or all of them are reversed (“case B”). Even more, reversion is only allowed for symmetric maximal jumps of the random walk. Our goal is to show that this event holds for $\gamma = 1$ (at least if the basic events B_{\dots} hold), but as intermediate steps, other values of γ are relevant, too.

Definition 6.26 For all divisors $\gamma \geq 1$ of l_{\rightarrow} , we define the event

$$E_{\text{mod } \gamma \text{ ok}}^n := \left. \begin{array}{l} \text{For all } w \in \text{SolutionPieces}^n(\text{Input}) \text{ there is a bijection } \iota_{\gamma} : \mathbb{Z}/\gamma\mathbb{Z} \rightarrow \mathbb{Z}/\gamma\mathbb{Z} \text{ such} \\ \text{that (at least) one of the following two cases holds:} \\ \text{A) } \forall Z \in \mathbb{Z}/\gamma\mathbb{Z}: \xi \upharpoonright [(-2^n, 2^n) \cap \iota_{\gamma}(Z)] \sqsubseteq w \upharpoonright Z \sqsubseteq \xi \upharpoonright [(-9 \cdot 2^n, 9 \cdot 2^n) \cap \iota_{\gamma}(Z)] \\ \text{B) } l_{\rightarrow} = l_{\leftarrow} \text{ and} \\ \quad \forall Z \in \mathbb{Z}/\gamma\mathbb{Z}: \xi \upharpoonright [(-2^n, 2^n) \cap \iota_{\gamma}(Z)] \sqsubseteq (w \upharpoonright Z)^{\leftrightarrow} \sqsubseteq \xi \upharpoonright [(-9 \cdot 2^n, 9 \cdot 2^n) \cap \iota_{\gamma}(Z)] \end{array} \right\}. \quad (6.34)$$

Lemma 6.27 For $\gamma = l_{\rightarrow}$, we have $E_{\text{seed II}}^n \cap E_{\text{mod class}}^n \cap E_{\text{dist}}^n \cap B_{\text{unique fit}}^n \subseteq E_{\text{mod } l_{\rightarrow} \text{ ok}}^n$.

Proof of Lemma 6.27. Assume that the event $E_{\text{seed II}}^n \cap E_{\text{mod class}}^n \cap E_{\text{dist}}^n \cap B_{\text{unique fit}}^n$ holds. Let $w \in \text{SolutionPieces}^n(\text{Input})$. Let $Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$. In order to define $\iota(Z) = \iota_{l_{\rightarrow}}(Z)$, we proceed as follows: Since $w \in \text{Filter}_4(\text{Input})$, there exists a right ladder interval $I \subseteq Z \cap [-2 \cdot 2^n, 2 \cdot 2^n]$ such that $(w[I]_{\rightarrow} \in \text{Seed}_{\text{III}}^n(\tau, \eta))$. We choose such an I . Then $(w[I]_{\rightarrow})$ is a left or right ladder word of $\xi[[-2 \cdot 2^n, 2 \cdot 2^n]]$, since the event $E_{\text{seed II}}^n$ holds. More specifically: for some right ladder interval $J \subseteq [-2 \cdot 2^n, 2 \cdot 2^n]$, at least one of the following two cases holds true:

$$\begin{aligned} \text{Case A}(Z): \quad & w[I] \equiv \xi[J], \\ \text{Case B}(Z): \quad & l_{\rightarrow} = l_{\leftarrow} \text{ and } (w[I]^{\leftarrow}) \equiv \xi[J]. \end{aligned}$$

We define $\iota(Z) := J + l_{\rightarrow}\mathbb{Z} \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$. Since the event $E_{\text{mod class}}^n$ holds, we get

$$\begin{aligned} \text{for Case A}(Z): \quad & \xi[([-2^n, 2^n] \cap \iota(Z))] \subseteq w[Z] \subseteq \xi[([-9 \cdot 2^n, 9 \cdot 2^n] \cap \iota(Z))], \\ \text{for Case B}(Z): \quad & \xi[([-2^n, 2^n] \cap \iota(Z))] \subseteq (w[Z]^{\leftarrow}) \subseteq \xi[([-9 \cdot 2^n, 9 \cdot 2^n] \cap \iota(Z))]. \end{aligned}$$

We claim that one of the following two cases occurs:

$$\begin{aligned} \text{Case A:} \quad & \text{For all modulo classes } Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z} \text{ holds Case A}(Z); \\ \text{Case B:} \quad & \text{For all modulo classes } Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z} \text{ holds Case B}(Z). \end{aligned}$$

This is obvious for $l_{\rightarrow} \neq l_{\leftarrow}$, since then Case B(Z) cannot occur. To prove the claim for $l_{\rightarrow} = l_{\leftarrow}$, we proceed as follows: For $Z \in \mathbb{Z}/l\mathbb{Z}$, let $T_Z : \mathbb{Z} \rightarrow \mathbb{Z}$ denote a translation (Case A(Z)) or reflection (Case B(Z)) which transports $w[Z]$ to a subpiece of $\xi[[-9 \cdot 2^n, 9 \cdot 2^n] \cap \iota(Z)]$. Let $Z, W \in \mathbb{Z}/l\mathbb{Z}$. We choose two right ladder intervals $I_1 \subseteq Z \cap [4 \cdot 2^n, 5 \cdot 2^n]$, $I_2 \subseteq W \cap [4 \cdot 2^n, 5 \cdot 2^n]$, $|I_1| = |I_2| = c_1 n$, with $I_1 \triangleright_n I_2$; such intervals exist, since $\text{supp } \mu^{*h}$ meets every modulo class (modulo l) and since $n \geq n_0$ is large enough. We abbreviate $I'_1 := T_Z[I_1]$ and $I'_2 := T_W[I_2]$. Since $w \in \text{Filter}_2(\text{Input})$ one has $((w[I_1]_{\rightarrow}, (w[I_2]_{\rightarrow})) \in \text{Neighbors}^n(\text{Input}))$. Let X_Z denote the symbol “ \rightarrow ” in the Case A(Z) and “ \leftarrow ” in the Case B(Z). Then $((w[I_1]_{\rightarrow}, (w[I_2]_{\rightarrow})) = ((\xi[I'_1]_{X_Z}, (\xi[I'_2]_{X_W}))$. Since the event E_{dist}^n holds, this implies $\text{distance}(I'_1, I'_2) \leq 3 \cdot l c_1 n$. However, T_Z maps $[-5 \cdot 2^n, 5 \cdot 2^n]$ to $[-9 \cdot 2^n - l, 9 \cdot 2^n + l]$; (the extra summand l arises since T_Z was specified only by its action on a modulo class). Thus it maps $I_1, I_2 \subseteq [4 \cdot 2^n, 5 \cdot 2^n]$ to a subset of $[4 \cdot 2^n - l, 9 \cdot 2^n + l]$ in the Case A(Z), and to a subset of $[-9 \cdot 2^n - l, -4 \cdot 2^n + l]$ in the Case B(Z). The same statement holds with Z replaced by W . The intervals $[4 \cdot 2^n - l, 9 \cdot 2^n + l]$ and $[-9 \cdot 2^n - l, -4 \cdot 2^n + l]$ are farther apart than $3 \cdot l c_1 n \geq \text{distance}(I'_1, I'_2)$; thus either both T_Z and T_W must be translations, or both must be reflections. Summarizing, we have shown so far that Case A holds or Case B holds.

It only remains to show that $\iota : \mathbb{Z}/l_{\rightarrow}\mathbb{Z} \rightarrow \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$ is bijective. Since $\mathbb{Z}/l_{\rightarrow}\mathbb{Z}$ is finite, it suffices to show that ι is injective: Let $Z, W \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$ with $\iota(Z) = \iota(W)$. Using the above maps T_Z, T_W again, we know

$$T_Z[Z \cap \text{domain}(w)] = T_Z[Z \cap [-5 \cdot 2^n, 5 \cdot 2^n]] \subseteq \iota(Z) \cap [-9 \cdot 2^n, 9 \cdot 2^n], \quad (6.35)$$

$$T_W[W \cap \text{domain}(w)] = T_W[W \cap [-5 \cdot 2^n, 5 \cdot 2^n]] \subseteq \iota(W) \cap [-9 \cdot 2^n, 9 \cdot 2^n]. \quad (6.36)$$

The sets on the right hand of (6.35) and (6.36) coincide; thus $T_Z[Z \cap [-5 \cdot 2^n, 5 \cdot 2^n]]$ and $T_W[W \cap [-5 \cdot 2^n, 5 \cdot 2^n]]$ overlap at least in $K \cap \iota(Z)$ for some interval K of length 2^n . We choose any right ladder interval $D \subseteq K \cap \iota(Z)$ with $|D| = c_1 n$ and set $D_1 := T_Z^{-1}[D]$ and $D_2 := T_W^{-1}[D]$. Then

$$\begin{aligned} \text{Case A:} \quad & (w[D_1]_{\rightarrow}) = (\xi[D]_{\rightarrow}) = (w[D_2]_{\rightarrow}), \\ \text{Case B:} \quad & (w[D_1]_{\rightarrow}) = (\xi[D]_{\leftarrow}) = (w[D_2]_{\rightarrow}); \end{aligned}$$

thus $w \in \text{Filter}_5(\text{Input})$ implies $D_1 = D_2$; hence $Z = D_1 + l_{\rightarrow}\mathbb{Z} = D_2 + l_{\rightarrow}\mathbb{Z} = W$. This shows that ι is indeed injective. ■

The next lemma contains a “step down” procedure in order to arrange correctly larger and larger modulo classes in a reconstructed piece of scenery w . Here is a rough idea for the rather complex construction:

Suppose we have already correctly reconstructed large pieces of the scenery ξ restricted to modulo classes (mod γ , say) up to a translation (and possibly a global reflection for all classes). Our task is to identify the relative translation between different modulo classes.

We start with a “reference” ladder word; it occurs over both, a ladder interval I in the reconstructed “candidate” scenery w , and a ladder interval J in the “true” scenery ξ (possibly reflected). Then we look for the rightmost “neighboring” ladder words that occur *not* in the same modulo class as the reference word, both in the candidate scenery and in the true scenery; we use here the “estimated” neighborhood relation “Neighbors”. Taking the rightmost “neighboring” words as our new starting point, we repeat this construction until we are sure after γ steps to re-enter the modulo class that we started with; say we arrive at ladder intervals I_γ and J_γ , respectively. In this way we obtain two “chains” (I_i) and (J_i) of neighboring ladder intervals; (J_i) belongs to the “true” scenery, and (I_i) belongs to the “reconstructed candidate” w .

Using the Definition of the tests “Filter_{2/3}”, and of the events $E_{\text{neighbor I/II}}^n$, we know that the “estimated” and the “geometrical” neighborhood relations coincide at least when taking only rightmost neighbors as above; this holds for both, the “reconstructed” piece w and for the “true” scenery ξ . The distance between I_γ and I equals the distance between J_γ and J , since this distance is not affected by a relative translation between different modulo classes; recall that I_γ and I belong to the same class modulo γ , and so do J_γ and J . Having identified the starting point and the end point of our two chains of intervals, there also no ambiguity left for the relative position of the intervals in between in the chain; but then we have successfully reconstructed the larger modulo class spanned by the whole chain (I_i) .

This construction is repeated recursively until we have correctly reconstructed the whole piece of scenery.

We describe the procedure formally:

Lemma 6.28 *Assume that the events $B_{\text{unique fit}}^n$, $E_{\text{neighbor I}}^n$, and $E_{\text{neighbor II}}^n$ hold true. Let $\gamma > 1$ be a divisor of l_{\rightarrow} , and assume that the event $E_{\text{mod } \gamma \text{ ok}}^n$ is valid. Then there is a divisor γ' of l_{\rightarrow} , with $1 \leq \gamma' < \gamma$ such that the event $E_{\text{mod } \gamma' \text{ ok}}^n$ is valid, too.*

Proof. Let γ be as in the hypothesis of the lemma. Every modulo class $Z \in \mathbb{Z}/\gamma\mathbb{Z}$ is a union of modulo classes $Z' \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$. Furthermore, every such modulo class $Z' \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$ has a nonempty intersection with $\text{supp } \mu^{*h}$. (One can see this as follows: Since 1 is the greatest common divisor of the elements of $\text{supp } \mu$, every integer can be written in the form $\beta l_{\rightarrow} + \sum_{j=1}^K s_j$ with $\beta \in \mathbb{Z}$, $K \in \mathbb{N}$, and $s_j \in \text{supp } \mu$ for $1 \leq j \leq K$. By Lemma 6.11 it suffices to take $K = h$; thus we get $\mathbb{Z} = \text{supp } \mu^{*h} + l_{\rightarrow}\mathbb{Z}$, which is equivalent to the above claim.)

Since we assume $\gamma > 1$, the set difference $\mathbb{Z} \setminus \gamma\mathbb{Z}$ contains at least one $Z \in \mathbb{Z}/\gamma\mathbb{Z}$ as a subset; thus $\mathbb{Z} \setminus \gamma\mathbb{Z}$ has at least one element in common with $\text{supp } \mu^{*h}$. Let $M_{\rightarrow} := \max[(\mathbb{Z} \setminus \gamma\mathbb{Z}) \cap \text{supp } \mu^{*h}]$ and $M_{\leftarrow} := -\min[(\mathbb{Z} \setminus \gamma\mathbb{Z}) \cap \text{supp } \mu^{*h}]$. Define γ' to be the greatest common divisor of γ and M_{\rightarrow} ; thus $\gamma' < \gamma$ since $M_{\rightarrow} \notin \gamma\mathbb{Z}$.

Let $w \in \text{SolutionPieces}^n(\text{Input})$. According to Definition (6.34) of $E_{\text{mod } \gamma \text{ ok}}^n$ we have to distinguish two cases A and B; however, we treat both cases simultaneously as far as possible. We

set

$$\zeta := \begin{cases} \xi \llbracket [-9 \cdot 2^n, 9 \cdot 2^n] & \text{in case A of (6.34),} \\ (\xi \llbracket [-9 \cdot 2^n, 9 \cdot 2^n])^{\leftrightarrow} & \text{in case B of (6.34).} \end{cases} \quad (6.37)$$

For $Z \in \mathbb{Z}/\gamma\mathbb{Z}$ we set $\tilde{\iota}_\gamma(Z) := \pm \iota_\gamma(\pm Z)$ with “+” in case A and “-” in case B; here the bijection $\iota_\gamma : \mathbb{Z}/\gamma\mathbb{Z} \rightarrow \mathbb{Z}/\gamma\mathbb{Z}$ is taken from Definition (6.34) of the event $E_{\text{mod } \gamma \text{ ok}}^n$. The introduction of $\tilde{\iota}_\gamma$ takes care of the inversion of modulo classes in ζ in case B. Since the event $E_{\text{mod } \gamma \text{ ok}}^n$ is valid, we have for all $Z \in \mathbb{Z}/\gamma\mathbb{Z}$:

$$\zeta \llbracket (\tilde{\iota}_\gamma(Z) \cap [-2^n, 2^n]) \sqsubseteq w \llbracket Z \sqsubseteq \zeta \llbracket \tilde{\iota}_\gamma(Z). \quad (6.38)$$

For $Z \in \mathbb{Z}/\gamma\mathbb{Z}$, let $T_Z : \mathbb{Z} \rightarrow \mathbb{Z}$ denote the translation which transports $w \llbracket Z$ to some $T_Z[w \llbracket Z] \subseteq \zeta \llbracket \tilde{\iota}_\gamma(Z)$; in particular $T_Z[Z] = \tilde{\iota}_\gamma(Z)$. T_Z is uniquely determined, since the event $B_{\text{unique fit}}^n$ holds. Of course, T_Z also depends on γ , but we suppress this in the notation, since γ is considered fixed for the moment. For $W \in \mathbb{Z}/\gamma\mathbb{Z}$, we set $\tilde{T}_W := (T_{\tilde{\iota}_\gamma^{-1}(W)})^{-1}$; thus $\tilde{T}_W[W] = \tilde{\iota}_\gamma^{-1}(W)$. For later use, we note

$$\zeta \llbracket (\tilde{\iota}_\gamma(Z) \cap [-2^n, 2^n]) \subseteq T_Z[w \llbracket Z] \subseteq \zeta \llbracket \tilde{\iota}_\gamma(Z). \quad (6.39)$$

We define

$$\zeta' := \bigcup_{Z \in \mathbb{Z}/\gamma\mathbb{Z}} T_Z[w \llbracket Z] \subseteq \zeta. \quad (6.40)$$

Note that $[-2^n, 2^n] \subseteq \text{domain}(\zeta')$. For (nonempty) ladder intervals I and J , we abbreviate $T_I := T_{I+\gamma\mathbb{Z}}$ and $\tilde{T}_J := \tilde{T}_{J+\gamma\mathbb{Z}}$.

Let the following data be given: $u \in \{w, \zeta'\}$, a right ladder interval I contained in the domain of u with $|I| = c_1 n$, and $k \in [0, \gamma]$. We define $\text{Seq}(I, u, k)$ to denote the set of all (I_0, \dots, I_k) with the following properties:

1. $I_0 = I$;
2. I_0, \dots, I_k are right ladder intervals contained in the domain of u with $|I_j| = c_1 n$, $0 \leq j \leq k$.
3. For all $j \in [0, k[$: $I_j + \gamma\mathbb{Z} \neq I_{j+1} + \gamma\mathbb{Z}$.
4. For all $j \in [0, k[$: $((u \llbracket I_j)_{\rightarrow}, (u \llbracket I_{j+1})_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$.

Of course $\text{Seq}(I, u, k)$ also depends on γ , Input , and n , but these parameters are considered fixed for the moment.

Let $\text{MaxSeq}(I, u, k)$ denote the set of all $(I_j)_{j=0, \dots, k} \in \text{Seq}(I, u, k)$ for which $\min I_k - \min I_0$ is maximal.

Given a modulo class $Z \in \mathbb{Z}/\gamma\mathbb{Z}$, we take a fixed right ladder interval $J \subseteq \tilde{\iota}_\gamma^{-1}(Z) \cap [0, (c_1 n + 1)l_{\rightarrow}] \subseteq \tilde{\iota}_\gamma^{-1}(Z) \cap \text{domain}(\zeta')$, $|J| = c_1 n$. Furthermore, we set $I := \tilde{T}_J J \subseteq Z \cap \text{domain}(w)$.

J serves as a “reference” interval in the “true” (only possibly reflected) piece of scenery ζ' , while I serves as a “reference” interval in the “reconstructed” piece of scenery w .

We prove by induction over k :

- $\text{MaxSeq}(I, w, k)$ contains a unique element $(I_j)_{j=0, \dots, k}$, namely

$$I_j = M_{\rightarrow} j + (c_1 n - 1)l_{\rightarrow} j + I. \quad (6.41)$$

- $\text{MaxSeq}(J, \zeta', k)$ contains a unique element $(J_j)_{j=0, \dots, k}$, too, namely

$$J_j = Mj + (c_1 n - 1)l_{\rightarrow} j + J, \quad (6.42)$$

where $M = M_{\rightarrow}$ in case A and $M = M_{\leftarrow}$ in case B.

This is obvious for $k = 0$. Here is the induction step $k - 1 \mapsto k$:

If $(I_j)_{j=0, \dots, k}$, $(J_j)_{j=0, \dots, k}$ are given by (6.41) and (6.42), then

$$(I_j)_{j=0, \dots, k} \in \text{Seq}(I, w, k) \quad \text{and} \quad (J_j)_{j=0, \dots, k} \in \text{Seq}(J, \zeta', k). \quad (6.43)$$

To see this, we check the conditions 1.–4. in the definition of Seq :

1. This is obvious.
2. The only nontrivial claims are $J_j \subseteq \text{domain}(\zeta')$ and $I_j \subseteq \text{domain}(w)$, $0 \leq j \leq k$. To prove the first claim, we observe $|\min J - \min J_j| \leq (M + c_1 n l)k \leq 2c_1 n l \gamma \leq 2c_1 n l^2$; thus we obtain for all $i \in J_j$: $|i| \leq 2c_1 n l^2 + (c_1 n + 1)l_{\rightarrow} \leq 2^n$; hence $J_j \subseteq [-2^n, 2^n] \subseteq \text{domain}(\zeta')$. To prove the second claim, we observe that $J + [-2^n/2, 2^n/2] \subseteq [-2^n, 2^n] = \text{domain}(\zeta')$ (recall $n \geq n_0$, and n_0 is large enough). We apply the translation \tilde{T}_J to $J + ([-2^n/2, 2^n/2] \cap \gamma\mathbb{Z})$ to obtain $I + ([-2^n/2, 2^n/2] \cap \gamma\mathbb{Z}) = \tilde{T}_J[J + ([-2^n/2, 2^n/2] \cap \gamma\mathbb{Z})] \subseteq \text{domain}(w) = [-5 \cdot 2^n, 5 \cdot 2^n]$. This implies $I + [-2^n/2 + \gamma, 2^n/2 - \gamma] \subseteq [-5 \cdot 2^n, 5 \cdot 2^n]$, since $[-5 \cdot 2^n, 5 \cdot 2^n]$ is an interval; consequently $I_j \subseteq I + [-2^n/2 + \gamma, 2^n/2 - \gamma] \subseteq \text{domain}(w)$, which proves the second claim.
3. This is a consequence of $\min I_{j+1} - \max I_j = M_{\rightarrow} \notin \gamma\mathbb{Z}$ and $\min J_{j+1} - \max J_j = M \notin \gamma\mathbb{Z}$.
4. Because of $\min I_{j+1} - \max I_j = M_{\rightarrow} \in \text{supp } \mu^{*h}$ we get $I_j \triangleright_n I_{j+1}$; thus the fact $w \in \text{Filter}_2^n(\text{Input})$ implies $((w \upharpoonright I_j)_{\rightarrow}, (w \upharpoonright I_{j+1})_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$; see Definition 5.4. Similarly $\min J_{j+1} - \max J_j = M_{\rightarrow} \in \text{supp } \mu^{*h}$ in case A and $\min J_{j+1} - \max J_j = M_{\leftarrow} \in -\text{supp } \mu^{*h}$ in case B. Hence we get $J_j \triangleright_n J_{j+1}$ in case A and $-J_j \triangleleft_n -J_{j+1}$ in case B; this implies $((\zeta' \upharpoonright J_j)_{\rightarrow}, (\zeta' \upharpoonright J_{j+1})_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$ in both cases, since the event $E_{\text{neighbor I}}^n$ holds; see Definition 6.9.

Thus the conditions 1.–4. are indeed valid.

To check the defining property of MaxSeq , consider another sequence

$$(I'_j)_{j=0, \dots, k} \in \text{Seq}(I, w, k) \quad \text{and} \quad (J'_j)_{j=0, \dots, k} \in \text{Seq}(J, \zeta', k). \quad (6.44)$$

Using our induction hypotheses

$$\text{MaxSeq}(I, w, k - 1) = \{(I_j)_{j=0, \dots, k-1}\}, \quad (6.45)$$

$$\text{MaxSeq}(J, \zeta', k - 1) = \{(J_j)_{j=0, \dots, k-1}\} \quad (6.46)$$

and

$$(I'_j)_{j=0, \dots, k-1} \in \text{Seq}(I, w, k - 1), \quad (J'_j)_{j=0, \dots, k-1} \in \text{Seq}(J, \zeta', k - 1), \quad (6.47)$$

we know

$$\min I_{k-1} - \min I_0 \geq \min I'_{k-1} - \min I'_0, \quad (6.48)$$

$$\min J_{k-1} - \min J_0 \geq \min J'_{k-1} - \min J'_0, \quad (6.49)$$

with equality only if $(I'_j)_{j=0,\dots,k-1} = (I_j)_{j=0,\dots,k-1}$ or $(J'_j)_{j=0,\dots,k-1} = (J_j)_{j=0,\dots,k-1}$.

We treat first case of the I 's: Using $((w \uparrow I'_{k-1})_{\rightarrow}, (w \uparrow I'_k)_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$ and $w \in \text{Filter}_3^n(\text{Input})$ we get $I'_{k-1} \triangleright_n I'_k + al_{\rightarrow}$ for some $a \in \mathbb{N}$; thus

$$\min I'_k - \max I'_{k-1} \leq \min I'_k + al_{\rightarrow} - \max I'_{k-1} \leq M_{\rightarrow} \quad (6.50)$$

by the maximality of M_{\rightarrow} and $I_k + \gamma\mathbb{Z} \neq I_{k-1} + \gamma\mathbb{Z}$; (see condition 3. in the definition of Seq , and recall $l_{\rightarrow} \in \gamma\mathbb{Z}$). Hence

$$\begin{aligned} \min I'_k - \min I'_0 &= (\min I'_k - \max I'_{k-1}) + (c_1 n - 1)l_{\rightarrow} + (\min I'_{k-1} - \min I'_0) \\ &\leq M_{\rightarrow} + (c_1 n - 1)l_{\rightarrow} + (\min I_{k-1} - \min I_0) = \min I_k - \min I_0. \end{aligned} \quad (6.51)$$

This proves

$$(I_j)_{j=0,\dots,k} \in \text{MaxSeq}(I, w, k). \quad (6.52)$$

Furthermore, using our induction hypothesis, equality in (6.51) can hold only if $(I'_j)_{j=0,\dots,k-1} \in \text{MaxSeq}(I, w, k-1)$ and $\min I'_k - \max I'_{k-1} = M_{\rightarrow}$, which is equivalent to $(I'_j)_{j=0,\dots,k} = (I_j)_{j=0,\dots,k}$.

We treat $(J_j)_{j=0,\dots,k}$ similarly: Since the event $E_{\text{neighbor II}}^n$ holds, $((\zeta' \uparrow J'_{k-1})_{\rightarrow}, (\zeta' \uparrow J'_k)_{\rightarrow}) \in \text{Neighbors}^n(\text{Input})$ implies

$$J'_{k-1} \triangleright_n J'_k + al_{\rightarrow} \quad \text{in case A,} \quad (6.53)$$

$$-J'_{k-1} \triangleleft_n -J'_k - al_{\leftarrow} \quad \text{in case B} \quad (6.54)$$

for some $a \in \mathbb{N}$; see Definition (6.12). This implies in both cases A and B, analogously to (6.50):

$$\min J'_k - \max J'_{k-1} \leq \min J'_k + al_{\rightarrow} - \max J'_{k-1} \leq M \quad (6.55)$$

by the maximality of M ; recall that $M = M_{\rightarrow}$ in case A and $M = M_{\leftarrow}$ in case B, and that $l_{\rightarrow} = l_{\leftarrow} \in \gamma\mathbb{Z}$ holds in case B; furthermore recall that J'_k and J'_{k-1} belong to different classes modulo γ . We repeat arguments similar to (6.51):

$$\begin{aligned} \min J'_k - \min J'_0 &= (\min J'_k - \max J'_{k-1}) + (c_1 n - 1)l_{\rightarrow} + (\min J'_{k-1} - \min J'_0) \\ &\leq M + (c_1 n - 1)l_{\rightarrow} + (\min J_{k-1} - \min J_0) = \min J_k - \min J_0, \end{aligned} \quad (6.56)$$

with equality only if $(J'_j)_{j=0,\dots,k} \in \text{MaxSeq}(J, \zeta', k-1)$ and $\min J'_k - \max J'_{k-1} = M$. This proves in analogy to (6.52):

$$\text{MaxSeq}(J, \zeta', k) = \{(J_j)_{j=0,\dots,k}\}. \quad (6.57)$$

Since \tilde{l}_{γ} is bijective, the facts $T_{I_j} I_j \subseteq \text{domain}(\zeta')$, $(\zeta' \uparrow T_{I_j} I_j)_{\rightarrow} = (w \uparrow I_j)_{\rightarrow}$, and $(I_j)_j \in \text{Seq}(I, w, k)$ imply

$$(T_{I_j} I_j)_j \in \text{Seq}(J, \zeta', k). \quad (6.58)$$

Similarly, $\tilde{T}_{J_j} J_j \subseteq \text{domain}(w)$, $(w \uparrow \tilde{T}_{J_j} J_j)_{\rightarrow} = (\zeta' \uparrow J_j)_{\rightarrow}$, and $(J_j)_j \in \text{Seq}(J, \zeta', k)$ imply

$$(\tilde{T}_{J_j} J_j)_j \in \text{Seq}(I, w, k). \quad (6.59)$$

Now we set $k = \gamma$. Observe that $I_\gamma + \gamma\mathbb{Z} = I_0 + \gamma\mathbb{Z}$ and $J_\gamma + \gamma\mathbb{Z} = J_0 + \gamma\mathbb{Z}$; hence $T_{I_0} = T_{I_\gamma}$ and $\tilde{T}_{J_0} = \tilde{T}_{J_\gamma}$. Thus, using (6.52), (6.57), (6.58), (6.59), and the defining property of MaxSeq , we obtain

$$\begin{aligned} \min I_\gamma - \min I_0 &= \min T_{I_\gamma} I_\gamma - \min T_{I_0} I_0 \\ &\leq \min J_\gamma - \min J_0 \\ &= \min \tilde{T}_{J_\gamma} J_\gamma - \min \tilde{T}_{J_0} J_0 \\ &\leq \min I_\gamma - \min I_0. \end{aligned} \tag{6.60}$$

Since the first and last term in (6.60) are identical, equality holds everywhere in (6.60). Hence, using (6.57), (6.58), and the defining property of MaxSeq again, we see

$$(T_{I_j} I_j)_j \in \text{MaxSeq}(J, \zeta', \gamma) \tag{6.61}$$

and thus $(T_{I_j} I_j)_j = (J_j)_j$, since $\text{MaxSeq}(J, \zeta', \gamma)$ is a singleton. Furthermore the facts (6.41), (6.42), $\gamma \neq 0$, and $T_{I_0} = T_{I_\gamma}$ imply $M_\rightarrow = M$, since

$$0 = (\min I_\gamma - \min I_0) - (\min J_\gamma - \min J_0) = M_\rightarrow \gamma - M\gamma. \tag{6.62}$$

A side remark: consequently case B cannot occur whenever $M_\rightarrow \neq M_-$. Using (6.41) and (6.42) again, we see that all translations T_{I_j} , $j = 0, \dots, \gamma$, coincide: $T_{I_j} = T_I$. We observe

$$(I_0 \cup \dots \cup I_\gamma) + \gamma\mathbb{Z} = I + \{jM_\rightarrow \mid j = 0, \dots, \gamma\} + \gamma\mathbb{Z} = I + \gamma'\mathbb{Z}; \tag{6.63}$$

recall that γ' was defined to be the greatest common divisor of M_\rightarrow and γ . Thus we have shown: the translations T_Z , $Z \in \mathbb{Z}/\gamma\mathbb{Z}$, depend only on the rougher modulo class $Z' = Z + \gamma'\mathbb{Z} \in \mathbb{Z}/\gamma'\mathbb{Z}$; hence $T_{Z+\gamma'\mathbb{Z}} := T_Z$ and $\iota_{\gamma'} : \mathbb{Z}/\gamma'\mathbb{Z} \rightarrow \mathbb{Z}/\gamma'\mathbb{Z}$, $\iota_{\gamma'}(Z') := \bigcup_{Z \subseteq Z', Z \in \mathbb{Z}/\gamma\mathbb{Z}} \iota_\gamma(Z)$ are well-defined. Since $\iota_\gamma : \mathbb{Z}/\gamma\mathbb{Z} \rightarrow \mathbb{Z}/\gamma\mathbb{Z}$ is a bijection, $\iota_{\gamma'}$ is a bijection, too. In analogy to $\tilde{\iota}_\gamma$, we introduce $\tilde{\iota}_{\gamma'}(Z') := \pm \iota_{\gamma'}(\pm Z')$, (“+” in case A, “−” in case B). As a consequence of (6.39), we obtain for all $Z' \in \mathbb{Z}/\gamma'\mathbb{Z}$:

$$\zeta[(\tilde{\iota}_{\gamma'}(Z') \cap [-2^n, 2^n]) \subseteq T_{Z'}[w[Z']] \subseteq \zeta[\tilde{\iota}_{\gamma'}(Z')]. \tag{6.64}$$

Hence the event $E_{\text{mod } \gamma' \text{ ok}}^n$ is valid. This finishes the proof of Lemma 6.28. ■

Lemma 6.29 $E_{\text{mod } l_\rightarrow \text{ ok}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{neighbor I}}^n \cap E_{\text{neighbor II}}^n \subseteq E_{\text{mod } 1 \text{ ok}}^n$.

Proof. Assume that $E_{\text{mod } l_\rightarrow \text{ ok}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{neighbor I}}^n \cap E_{\text{neighbor II}}^n$ holds. Let Γ denote the (random) set of all divisors $\gamma \geq 1$ of l_\rightarrow for which the event $E_{\text{mod } \gamma \text{ ok}}^n$ is valid. $\Gamma \neq \emptyset$, since $E_{\text{mod } l_\rightarrow \text{ ok}}^n$ holds. The smallest element of Γ cannot be bigger than 1 by Lemma 6.28; thus it must be equal to 1. This means that $E_{\text{mod } 1 \text{ ok}}^n$ holds. ■

Lemma 6.30 For $\gamma = 1$, we have $E_{\text{mod } 1 \text{ ok}}^n \subseteq E_{\text{all pieces ok}}^n$.

Proof. This is obvious, since there is only the trivial “modulo class” $Z = \iota_1(Z) = \mathbb{Z}$ remaining for $\gamma = 1$: In case A, one has $\xi[[-2^n, 2^n] \subseteq w \subseteq \xi[[-9 \cdot 2^n, 9 \cdot 2^n]]$, and in case B, one has $\xi[[-2^n, 2^n] \subseteq w^\leftrightarrow \subseteq \xi[[-9 \cdot 2^n, 9 \cdot 2^n]]$. ■

Theorem 6.31 $B_{\text{seed I}}^n \cap B_{\text{unique fit}}^n \cap B_{\text{all paths}}^n \cap B_{\text{outside out}}^n \cap B_{\text{recogn straight}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop},\tau}^m \subseteq E_{\text{all pieces ok}}^n$

Proof. We collect the results of Lemmas 6.5, 6.10, 6.12, 6.19, 6.21, 6.23, 6.25, 6.27, 6.29, and 6.30 in the following list:

$$\begin{array}{ll}
B_{\text{signals}}^n & \subseteq E_{\text{signals II}}^n, \\
B_{\text{all paths}}^n & \subseteq E_{\text{neighbor I}}^n, \\
E_{\text{signals II}}^n \cap E_{\text{stop},\tau}^m & \subseteq E_{\text{neighbor II}}^n, \\
B_{\text{all paths}}^n \cap B_{\text{recogn straight}}^n & \subseteq E_{\text{only ladder}}^n, \\
B_{\text{outside out}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{only ladder}}^n \cap E_{\text{stop},\tau}^m & \subseteq E_{\text{mod class}}^n, \\
B_{\text{unique fit}}^n \cap B_{\text{signals}}^n \cap B_{\text{all paths}}^n \cap B_{\text{recogn straight}}^n \cap E_{\text{stop},\tau}^m & \subseteq E_{\text{seed II}}^n, \\
B_{\text{signals}}^n \cap E_{\text{stop},\tau}^m & \subseteq E_{\text{dist}}^n, \\
E_{\text{seed II}}^n \cap E_{\text{mod class}}^n \cap E_{\text{dist}}^n \cap B_{\text{unique fit}}^n & \subseteq E_{\text{mod } l \rightarrow \text{ok}}^n, \\
E_{\text{mod } l \rightarrow \text{ok}}^n \cap B_{\text{unique fit}}^n \cap E_{\text{neighbor I}}^n \cap E_{\text{neighbor II}}^n & \subseteq E_{\text{mod I ok}}^n, \\
E_{\text{mod I ok}}^n & \subseteq E_{\text{all pieces ok}}^n.
\end{array}$$

The claim of the theorem is a simple combination of these inclusions. ■

6.3 Probabilistic estimates for basic events

In this subsection we show that the “basic events” B_{\dots} occur very probably. Together with the result of the previous subsections this shows that the partial reconstruction algorithms Alg^n yield with high probability a correctly reconstructed piece of scenery.

We start with an elementary auxiliary lemma:

Lemma 6.32 *Let $f : I_0 \rightarrow J$ be a finite injection without fixed points. Then there is $I' \subseteq I_0$ with $|I'| \geq |I_0|/3$ and $f[I'] \cap I' = \emptyset$.*

Proof. We construct recursively finite sequences (I_k) and (I'_k) , for $k - 1 < |I_0|/3$, of subsets of I_0 . The “loop invariants” of the recursion are: $f[I_k] \cap I'_k = \emptyset$, $f[I'_k] \cap I_k = \emptyset$, $f[I'_k] \cap I'_k = \emptyset$, $I_k \cap I'_k = \emptyset$, $|I'_k| = k$, and $|I_k| \geq |I_0| - 3k$.

The recursion starts with the given I_0 and with $I'_0 = \emptyset$. In the $(k + 1)$ st step, $k < |I_0|/3$, we choose any point $x \in I_k$, and define $I'_{k+1} := I'_k \cup \{x\}$. If $f^{-1}(x)$ exists, then we set $I_{k+1} := I_k \setminus \{x, f(x), f^{-1}(x)\}$; else we set $I_{k+1} := I_k \setminus \{x, f(x)\}$.

Note that the validity of the above “loop invariants” is indeed preserved by the recursion; the fact $f(x) \neq x$ is used for the third loop invariant.

Finally we set $I' := I'_k$ for $k := \min\{j \in \mathbb{N} \mid 3j \geq |I_0|\}$; then $I' \subseteq I_0$ is well-defined and fulfills the claims in Lemma 6.32. ■

Lemma 6.33 *There exists constants $c_{18}, c_{19} > 0$ not depending on n such that:*

$$P[(B_{\text{unique fit}}^n)^c] \leq c_{18}e^{-c_{19}n}. \quad (6.65)$$

Proof. Let $i, j \in \{1, \dots, l^2\}$, and let $I \subseteq [-11 \cdot 2^n, 11 \cdot 2^n]$ be a i -spaced interval, and $J \subseteq [-11 \cdot 2^n, 11 \cdot 2^n]$ be a j -spaced interval with $|I| = |J| \geq 1$. Let $f : I \rightarrow J$ be a monotonically increasing or decreasing bijection, but not the identity map; thus the case $I = J$ can only occur if f is decreasing.

We claim: For some constants $c_{12} > 0$ and $c_{13} > 2 \log 2 / c_2$ (not depending on i, j, I , or J) we have

$$P[\xi \circ f = \xi[I] \leq c_{12} e^{-c_{13}|I|}. \quad (6.66)$$

Note that $\xi \circ f = \xi[I]$ is equivalent to $(\xi[J])_{\rightarrow} = (\xi[I])_{\rightarrow}$ if f is increasing, and it is equivalent to $(\xi[J])_{\rightarrow} = (\xi[I])_{\leftarrow}$ if f is decreasing.

Before proving (6.66), let us show how it implies (6.65): There are at most l^2 choices for (i, j) , and given (i, j) , there are at most $(22 \cdot 2^n + 1)^2 \leq 500 \cdot 2^{2n}$ choices for (I, J) with $|I| = |J| = c_2 n$; finally there is one binary choice: f is increasing or decreasing. If $\xi \circ f \neq \xi[I]$ holds for all of these choices (with the trivial exception $I = J$ and $f = \text{id}$), then the event $B_{\text{unique fit}}^n$ is valid; note that it suffices to consider $|I| = |J| = c_2 n$ instead of $|I| = |J| \geq c_2 n$, since it suffices to consider subintervals of I, J consisting only of $c_2 n$ points. Hence (6.66) implies (6.65):

$$P[(B_{\text{unique fit}}^n)^c] \leq l^2 \cdot 500 \cdot 2^{2n} \cdot 2 \cdot c_{12} e^{-c_{13} c_2 n} = c_{18} e^{-c_{19} n}, \quad (6.67)$$

where $c_{18} := 1000l^2 c_{12}$ and $c_{19} := c_{13} c_2 - 2 \log 2 > 0$.

We prove (6.66) next: unless f is the identity map, it can have at most a single fixed point, since it is the restriction of some affine-linear map to the ladder interval I . Remove this fixed point from I , if it exists; call I_0 the set of all remaining points. By Lemma 6.32, there is $I' \subseteq I_0$ with $|I'| \geq |I_0|/3 \geq (|I| - 1)/3$ and $f[I'] \cap I' = \emptyset$. Hence $\xi[f[I']]$ and $\xi[I']$ are independent random pieces of scenery; thus

$$P[\xi \circ f = \xi[I] \leq P[\xi \circ f[I'] = \xi[I']] = |\mathcal{C}|^{-|I'|} \leq |\mathcal{C}|^{-(|I|-1)/3}, \quad (6.68)$$

thus (6.66) follows with $c_{13} := (\log |\mathcal{C}|)/3$ and $c_{12} := |\mathcal{C}|^{1/3}$. Note that $c_{13} c_2 - 2 \log 2 > 0$ since c_2 was required to be large enough; recall subsection 2.1. ■

Lemma 6.34 *There exist constants $c_{20}, c_{21} > 0$ not depending on n such that:*

$$P[(B_{\text{all paths}}^n)^c \cap E_{\text{stop}, \tau}^m] \leq c_{21} e^{-c_{20} n}. \quad (6.69)$$

Proof of Lemma 6.34. Let $k < 2^{\alpha n}$ and $R \in \text{AdPath}(12 \cdot 2^n, 3c_1 n)$. We set

$$B_R^{n,k} := \left\{ \exists j \in [0, 2^{2n}] : \text{TimeShift}^{\tau(k)+j}(R) \subseteq S \right\}, \quad (6.70)$$

$$E_{\text{stop}, \tau, k}^m := \left\{ \begin{array}{l} \tau_k(\chi) < 2^{12\alpha n_m}, |S(\tau_k(\chi))| \leq 2^{n_m}, \\ \tau_j(\chi) + 2 \cdot 2^{2n_m} \leq \tau_k(\chi) \text{ for } j < k \end{array} \right\}, \quad (6.71)$$

$$A_R^{n,k} := E_{\text{stop}, \tau, k}^m \setminus B_R^{n,k}. \quad (6.72)$$

Note that $B_{\text{all paths}}^n = \bigcap_{R \in \text{AdPath}(12 \cdot 2^n, 3c_1 n)} \bigcup_{k=0}^{2^{\alpha n} - 1} B_R^{n,k}$ and $E_{\text{stop}, \tau}^m \subseteq E_{\text{stop}, \tau, k}^m$ for $k \leq 2^{\alpha n}$, and thus

$$E_{\text{stop}, \tau}^m \setminus B_{\text{all paths}}^n \subseteq \bigcup_{R \in \text{AdPath}(12 \cdot 2^n, 3c_1 n)} \bigcap_{k=0}^{2^{\alpha n} - 1} A_R^{n,k}. \quad (6.73)$$

In the following, R runs over the set $\text{AdPath}(12 \cdot 2^n, 3c_1 n)$:

$$P \left[(B_{\text{all paths}}^n)^c \cap E_{\text{stop}, \tau}^m \right] \leq |\text{AdPath}(12 \cdot 2^n, 3c_1 n)| \max_R P \left[\bigcap_{k=0}^{2^{\alpha n} - 1} A_R^{n, k} \right], \quad (6.74)$$

$$|\text{AdPath}(12 \cdot 2^n, 3c_1 n)| \leq 25 \cdot 2^n |\mathcal{M}|^{3c_1 n}, \quad (6.75)$$

$$P \left[\bigcap_{k=0}^{2^{\alpha n} - 1} A_R^{n, k} \right] = \prod_{k=0}^{2^{\alpha n} - 1} P \left[A_R^{n, k} \mid \bigcap_{j < k} A_R^{n, j} \right], \quad (6.76)$$

$$P \left[A_R^{n, k} \mid \bigcap_{j < k} A_R^{n, j} \right] \leq P \left[(B_R^{n, k})^c \mid E_{\text{stop}, \tau, k}^m \cap \bigcap_{j < k} A_R^{n, j} \right]; \quad (6.77)$$

the last statement follows from the elementary fact $P[A \cap B | C] \leq P[A | B \cap C]$. Since $2^{2n} + 3c_1 n \leq 2 \cdot 2^{2n}$, we have $C_R^{n, k} := E_{\text{stop}, \tau, k}^m \cap \bigcap_{j < k} A_R^{n, j} \in \mathcal{F}_{\tau_k}$, i.e. one can decide whether the event $C_R^{n, k}$ holds by observing ξ and $S(0), \dots, S(\tau_k)$. Furthermore, if $C_R^{n, k}$ holds, then $|S(\tau_k(\chi))| \leq 2^n$, and as a consequence of the local Central Limit Theorem [4], Theorem 5.2 (page 132) we get: there is a constant $c_{22} > 0$ such that for all x, y with $|x| \leq 12 \cdot 2^n$ and $|y| \leq 2^n$: $P[y + S(j) = x \text{ for some } j \in [0, 2^{2n}]] \geq c_{22} 2^{-n}$; note that $y + S$ is a random walk starting in the point y . Note that we do not need the random walk to be aperiodic; it suffices that it can reach every integer, i.e. that the greatest common divisor of the elements of $|\mathcal{M}|$ is 1. Thus by the strong Markov property:

$$\inf_{|x| \leq 12 \cdot 2^n} P \left[S(\tau(k) + j) = x \text{ for some } j \in [0, 2^{2n}] \mid C_R^{n, k} \right] \geq c_{22} 2^{-n}. \quad (6.78)$$

Once it is in the starting point x , the probability that S follows an admissible path $R \in \text{AdPath}(12 \cdot 2^n, 3c_1 n)$ for the next $3c_1 n - 1$ steps is bounded from below by $\mu_{\min}^{3c_1 n}$. Here $\mu_{\min} := \min\{\mu(\{x\}) \mid x \in \mathcal{M}\}$ is the smallest positive probability for a jump. Therefore, using the strong Markov property again:

$$P \left[B_R^{n, k} \mid C_R^{n, k} \right] \geq c_{22} 2^{-n} \mu_{\min}^{3c_1 n}. \quad (6.79)$$

We combine (6.74)–(6.77) and (6.79) to obtain

$$\begin{aligned} P \left[E_{\text{stop}, \tau}^m \setminus B_{\text{all paths}}^n \right] &\leq 25 \cdot 2^n |\mathcal{M}|^{3c_1 n} (1 - c_{22} 2^{-n} \mu_{\min}^{3c_1 n})^{2^{\alpha n}} \\ &\leq 25 \cdot 2^n |\mathcal{M}|^{3c_1 n} \exp \left\{ -c_{22} 2^{-n} \mu_{\min}^{3c_1 n} 2^{\alpha n} \right\} \\ &\leq 25 \exp \left\{ n(\log 2 + 3c_1 \log |\mathcal{M}|) - c_{22} e^{n(\alpha \log 2 + 3c_1 \log \mu_{\min} - \log 2)} \right\}. \end{aligned} \quad (6.80)$$

Now $\alpha > 1 - 3c_1 \log_2 \mu_{\min}$ by our choice of α in subsection 2.1; thus the right hand side of the last inequality converges to 0 superexponentially fast as $n \rightarrow \infty$. Note that we may choose an upper bound $c_{21} e^{-c_{20} n}$ for the right hand side in 6.80, where neither c_{21} nor c_{20} depend on α or c_1 . This is true since $n \geq n_0$, and n_0 was chosen large enough, depending on c_1 and α ; recall subsection 2.1. This proves the lemma. ■

Lemma 6.35 *There exists a constant $c_{23} > 0$ not depending on n such that:*

$$P \left[(B_{\text{outside out}}^n)^c \right] \leq 160 e^{-c_{23} n}. \quad (6.81)$$

Proof of Lemma 6.35. The set $[-2 \cdot l2^{2n}, 2 \cdot l2^{2n}] \setminus [-10 \cdot 2^n, 10 \cdot 2^n]$ contains less than $4 \cdot l \cdot 2^{2n}$ points, and for every fixed starting point the number of admissible paths with $c_1 n/2$ points is equal to $|\mathcal{M}|^{c_1 n/2-1}$. Hence there are less than $4 \cdot l2^{2n} |\mathcal{M}|^{c_1 n/2}$ paths $R \in \text{AdPaths}(2 \cdot l2^{2n}, c_1 n/2)$ with $R(i) \notin [-10 \cdot 2^n, 10 \cdot 2^n]$ for all $i = 0, \dots, c_1 n/2$. On the other hand, there are less than $40 \cdot 2^n$ ladder words of length $c_1 n/2$ in $[-9 \cdot 2^n, 9 \cdot 2^n]$. The colors $\xi \circ R$ that a path in $R \in \text{AdPaths}(2^{2n}, c_1 n/2)$ with $R(i) \notin [-10 \cdot 2^n, 10 \cdot 2^n]$ for all $i = 0, \dots, c_1 n/2 - 1$ reads are independent of the colors inside $[-9 \cdot 2^n, 9 \cdot 2^n]$. Thus the probability that a given path $R \in \text{AdPaths}(2^{2n}, c_1 n/2)$ with $R(i) \notin [-10 \cdot 2^n, 10 \cdot 2^n]$ for all $i = 0, \dots, c_1 n/2 - 1$ reads the same colors as a fixed ladder word in $[-9 \cdot 2^n, 9 \cdot 2^n]$ is $|\mathcal{C}|^{-c_1 n/2}$. Thus

$$P[(B_{\text{outside out}}^n)^c] \leq 160l2^{3n} |\mathcal{M}|^{c_1 n/2} |\mathcal{C}|^{-c_1 n/2}. \quad (6.82)$$

Since $|\mathcal{M}| < |\mathcal{C}|$, the last expression becomes exponentially decreasing in n since $c_1 > 6/\log \frac{|\mathcal{C}|}{|\mathcal{M}|}$ since c_1 was chosen large enough; see subsection 2.1. This proves the lemma. ■

We prepare the treatment of the event $B_{\text{recognstraight}}^n$ by the following combinatoric lemma:

Lemma 6.36 *Let $c_{24} := 1/(2|\mathcal{M}|(l_{\rightarrow} + l_{\leftarrow}))$. There are two intervals $I_1, I_2 \subseteq [0, c_1 n[$ with $|I_1| = |I_2| \geq c_{24} c_1 n - 1$ such that the following statement is valid: For all $R \in \text{AdPaths}(11 \cdot 2^n, c_1 n)$ with $R(c_1 n - 1) - R(0) \notin \{(c_1 n - 1)l_{\rightarrow}, -(c_1 n - 1)l_{\leftarrow}\}$, there is $I \in \{I_1, I_2\}$ and an admissible path $\bar{R} \in \text{AdPaths}(12 \cdot 2^n, c_1 n)$ with the following properties:*

- $R(0) = \bar{R}(0), \quad R(c_1 n - 1) = \bar{R}(c_1 n - 1).$
- *At least one of the following holds:*
 1. *for all $(i, j) \in I \times I$ with $j < i$: $\bar{R}(i) \notin \{R(j), \bar{R}(j)\}$;*
 2. *for all $(i, j) \in I \times I$ with $i < j$: $\bar{R}(i) \notin \{R(j), \bar{R}(j)\}$.*

Proof. We define $k := \lfloor c_{24} c_1 n \rfloor$, $I' := [1, 2k] \subseteq [0, c_1 n[$, $I_1 := [1, k]$, and $I_2 := [k + 1, 2k]$. We observe $|I_1|, |I_2| \geq c_{24} c_1 n - 1$ and $I_1, I_2 \subseteq [0, c_1 n[$.

Let $R \in \text{AdPaths}(11 \cdot 2^n, c_1 n)$ be not a ladder path. We show first: There are $R', R'' \in \text{AdPaths}(12 \cdot 2^n, c_1 n)$ such that $R'(0) = R''(0) = R(0)$, $R'(c_1 n - 1) = R''(c_1 n - 1) = R(c_1 n - 1)$, $R'[I'$ and $R''[I'$ are ladder paths, and $R''[I' = r + R'[I'$ for some $r \neq 0$, i.e. $R''[I'$ is obtained from $R'[I'$ by a spatial translation.

To prove this claim, let $d = (d_i)_{i=1, \dots, c_1 n-1} \in \mathcal{M}^{c_1 n-1}$, $d_i := R(i) - R(i-1)$, be the jump sizes in R . Every other $\tilde{d} \in \mathcal{M}^{c_1 n-1}$ with $\sum_{i=1}^{c_1 n-1} \tilde{d}_i = \sum_{i=1}^{c_1 n-1} d_i$ gives rise to an admissible path $\tilde{R} \in \text{AdPaths}(12 \cdot 2^n, c_1 n)$, too, with $\tilde{R}(0) = R(0)$, $\tilde{R}(c_1 n - 1) = R(c_1 n - 1)$, and with jump sizes $\tilde{d}_i = \tilde{R}(i) - \tilde{R}(i-1)$; namely $\tilde{R}(i) := R(0) + \sum_{j=1}^i \tilde{d}_j$. Since \tilde{R} has its starting point and end point in $[-11 \cdot 2^n, 11 \cdot 2^n]$ and since $c_1 n l < 2^n$, the path \tilde{R} can indeed not leave the range $[-12 \cdot 2^n, 12 \cdot 2^n]$.

There are at most $|\mathcal{M}|$ possible values for d_i , but there are $c_1 n$ possible indices i ; thus at least one value $a \in \mathcal{M}$ occurs in the d_i at least $c_1 n/|\mathcal{M}|$ times. We choose $2k(a + l_{\leftarrow}) \geq 0$ indices i with $d_i = a$ and replace them by l_{\rightarrow} , and we choose $2k(l_{\rightarrow} - a) \geq 0$ different indices i with $d_i = a$ and replace them by $-l_{\leftarrow}$; note that $2k(a + l_{\leftarrow}) + 2k(l_{\rightarrow} - a) = 2k(l_{\rightarrow} + l_{\leftarrow}) \leq c_1 n/|\mathcal{M}|$. We end up with a new vector $\tilde{d} \in \mathcal{M}^{c_1 n-1}$ with $\sum_{i=1}^{c_1 n-1} \tilde{d}_i = \sum_{i=1}^{c_1 n-1} d_i$, since $2k(l_{\leftarrow} + l_{\rightarrow})a = 2k(a + l_{\leftarrow})l_{\rightarrow} + 2k(l_{\rightarrow} - a)(-l_{\leftarrow})$. \tilde{d} contains at least $2k$ entries with value l_{\rightarrow} , or it contains at least $2k$ entries with value $-l_{\leftarrow}$, since already the described replacement procedure has produced sufficiently many such entries. However, not all entries of \tilde{d} can equal l_{\rightarrow} ; similarly not all its

entries can equal $-l_{\leftarrow}$, since R is not a ladder path. We permute the entries of \tilde{d}_i in two different ways; the resulting vectors are called d' and d'' : First to obtain d' , permute the entries in \tilde{d} such that the first $2k$ permuted entries d'_i , $i = 1, \dots, 2k$ either all equal l_{\rightarrow} or all equal $-l_{\leftarrow}$; the order of the remaining entries is irrelevant. Second to obtain d'' , transpose the first entry d'_1 with a different entry $d'_i \neq d'_1$. Let R' and R'' be admissible pieces of paths with $R'(0) = R''(0) = R(0)$ and step sizes $d'_i = R'(i) - R'(i-1)$ and $d''_i = R''(i) - R''(i-1)$, respectively. Recall $I' = [1, 2k]$; then $R'[I']$ and $R''[I']$ are ladder paths, and $R''[I']$ is obtained from translating $R'[I']$ by $r := d''_1 - d'_1 \neq 0$. Thus our first claim holds.

$R'[I']$ is a right ladder path or a left ladder path. Without loss of generality, we assume that it is a right ladder path; the case of left ladder paths can be treated similarly by reversing directions in the arguments below. Furthermore, we assume without loss of generality $r > 0$; otherwise we exchange R' with R'' .

We are ready to prove the claim of the lemma; recall that k is a point in the middle of I' . There are two cases:

- If $R(k) > R'(k)$, then we take $I := I_1$ and $\bar{R} := R'$. Since $R'[I]$ is a right ladder path, it moves with maximal speed l_{\rightarrow} to the right. R cannot move faster than that to the right; thus $R(j) > R'(i)$ and $R'(j) > R'(i)$ for all $i, j \in I$ with $i < j$.
- If $R(k) \leq R'(k)$, then $R(k) < r + R'(k) = R''(k)$; this time we take $I := I_2$ and $\bar{R} := R''$. The same argument as above yields $R(j) < R''(i)$ and $R''(j) < R''(i)$ for all $i, j \in I$ with $j < i$.

This proves Lemma 6.36. ■

Lemma 6.37 *There exist positive constants c_{25} and c_{26} not depending on n such that:*

$$P[(B_{\text{recogn straight}}^n)^c] \leq c_{25}e^{-c_{26}n}. \quad (6.83)$$

Proof of Lemma 6.37. Given $R \in \text{AdPaths}(11 \cdot 2^n, c_1n)$ with $R(c_1n) - R(0) \notin \{(c_1n - 1)l_{\rightarrow}, -(c_1n - 1)l_{\leftarrow}\}$, we take $I = I(R) \subseteq [0, c_1n[$ and $\bar{R} \in \text{AdPaths}(12 \cdot 2^n, c_1n)$ as in Lemma 6.36. Without loss of generality assume that condition 1. in Lemma 6.36 is satisfied. We prove for all $I' \subseteq I$ by induction on $|I'|$:

$$P[(\xi \circ R)[I' = (\xi \circ \bar{R})[I']] = |\mathcal{C}|^{-|I'|}. \quad (6.84)$$

This is obvious for $I' = \emptyset$. For other I' , let $I'' := I' \setminus \max I'$. Then $\xi(\bar{R}(\max I'))$ is independent of $(\xi \circ R[I''], \xi \circ \bar{R}[I''])$, since they are generated by disjoint parts of the scenery. Thus

$$\begin{aligned} P[(\xi \circ R)[I' = (\xi \circ \bar{R})[I']] &= P[\xi(R(\max I')) = \xi(\bar{R}(\max I'))] \cdot P[(\xi \circ R)[I'' = (\xi \circ \bar{R})[I'']] \\ &= |\mathcal{C}|^{-1} |\mathcal{C}|^{-|I''|} = |\mathcal{C}|^{-|I'|}. \end{aligned} \quad (6.85)$$

By taking $I' = I$, we conclude $P[(\xi \circ R)[I = (\xi \circ \bar{R})[I]] \leq |\mathcal{C}|^{-|I|}$.

It does not suffice to multiply the last bound with the bound $23 \cdot 2^n |\mathcal{M}|^{c_1n} \geq |\text{AdPaths}(11 \cdot 2^n, c_1n)|$: the product may sometimes be bigger than 1.

To overcome this difficulty, we partition $\text{AdPaths}(12 \cdot 2^n, c_1n) \ni R$ into equivalence classes $[R]$: we put two paths into the same class if and only if they are mapped to the same value by the map $R \mapsto (R(0), R(c_1n - 1), I(R), R[I(R)])$; here $I(R) \in \{I_1, I_2\}$ is taken from Lemma

6.36. We bound the number of equivalence classes from above: For our purposes, a simple but rough bound suffices: There are at most $25 \cdot 2^n$ choices for each of $R(0)$, $R(c_1 n - 1)$, and $R(\min I(R))$, and there is a binary choice $I(R) \in \{I_1, I_2\}$; finally given $R(\min I(R))$, there are not more than $|\mathcal{M}|^k$ choices for $R[I(R)]$, where again $k = \lfloor c_{24} c_1 n \rfloor = |I(R)|$. Altogether the number of equivalence classes is bounded by $c_{27} 2^{3n} |\mathcal{M}|^k$, where $c_{27} := 2 \cdot 25^3$. We may choose a map $\text{AdPaths}(11 \cdot 2^n, c_1 n) \rightarrow \text{AdPaths}(12 \cdot 2^n, c_1 n)$, $R \mapsto \bar{R}$ such that \bar{R} depends only of the equivalence class $[R]$ and fulfills the claim in Lemma 6.36. We get

$$\begin{aligned} P[(B_{\text{recogn straight}}^n)^c] &\leq P[\exists R \in \text{AdPaths}(11 \cdot 2^n, c_1 n) : \xi \circ R[I(R)] = \xi \circ \bar{R}[I(R)]] \quad (6.86) \\ &\leq \sum_{[R]} P[\xi \circ R[I(R)] = \xi \circ \bar{R}[I(R)]] \\ &\leq c_{27} 2^{3n} (|\mathcal{M}|/|\mathcal{C}|)^k \leq c_{27} (|\mathcal{C}|/|\mathcal{M}|) \exp\{(3 \log 2 - c_{24} c_1 \log(|\mathcal{M}|/|\mathcal{C}|))n\}. \end{aligned}$$

We emphasize: the sum in the last but one expression runs over equivalence classes $[R]$, not over paths R ; the event $\{\xi \circ R[I(R)] = \xi \circ \bar{R}[I(R)]\}$ does not depend on the choice of $R \in [R]$. We have $c_{24} c_1 \log(|\mathcal{M}|/|\mathcal{C}|) - 3 \log 2 \geq 1$; recall from subsection 2.1 that c_1 is large enough. The estimate (6.86) proves the lemma with $c_{26} = 1$, $c_{25} = c_{27} |\mathcal{C}|/|\mathcal{M}|$. ■

Lemma 6.38 *There exist constants $c_{28} > 0$, $c_{29} > 0$ such that:*

$$P[(B_{\text{signals}}^n)^c] \leq c_{29} e^{-c_{28} n}. \quad (6.87)$$

Proof of Lemma 6.38. We show that

$$P[B_{\text{sig rr}}^n] \geq 1 - c_{30} e^{-c_{28} n} \quad (6.88)$$

for some constants $c_{30} > 0$ and $c_{28} > 0$. The proof for $B_{\text{sig rl}}^n$, $B_{\text{sig lr}}^n$, and $B_{\text{sig ll}}^n$ can be done analogously. Take a right ladder path $\pi \in [-2 \cdot l 2^{2n}, 2 \cdot l 2^{2n}]^{[0, c_1 n/2]}$ and an admissible piece of path $\pi' \in \text{AdPath}(2 \cdot l 2^{2n}, c_1 n/2)$ with $\pi(0) > \pi'(0)$. We show by induction over $j \in [0, c_1 n/2[$ with the abbreviation $I = [0, j + 1[$ and $I' = [0, j]$:

$$P[\xi \circ \pi[I] = \xi \circ \pi'[I']] = |\mathcal{C}|^{-j}. \quad (6.89)$$

Indeed, (6.89) is trivial for $j = 0$. For the step $j \mapsto j + 1$, we observe that $\pi(j)$ is right of all $\pi(i)$ and $\pi'(i)$, $i < j$, since π is a right ladder path and $\pi(0) > \pi'(0)$. Thus $\xi \circ \pi(j)$ is independent of the family $(\xi \circ \pi[I'], \xi \circ \pi'[I'])$. Therefore, using our induction hypothesis,

$$\begin{aligned} P[\xi \circ \pi[I] = \xi \circ \pi'[I]] &\quad (6.90) \\ &= P[\xi \circ \pi[I'] = \xi \circ \pi'[I']] \cdot P[\xi \circ \pi(j) = \xi \circ \pi'(j)] = |\mathcal{C}|^{-j-1}, \end{aligned}$$

For $j = c_1 n/2$ we obtain that

$$P[\xi \circ \pi[[0, c_1 n/2[= \xi \circ \pi'[[0, c_1 n/2]] \leq |\mathcal{C}|^{-c_1 n/2}. \quad (6.91)$$

There are no more than $4 \cdot l 2^{2n} + 1 \leq 5 \cdot l 2^{2n}$ such π and not more than $5 \cdot l 2^{2n} |\mathcal{M}|^{c_1 n/2}$ such π' . Therefore

$$P[(B_{\text{sig r}}^n)^c] \leq (5 \cdot l 2^{2n})^2 |\mathcal{M}|^{c_1 n/2} |\mathcal{C}|^{-c_1 n/2} \quad (6.92)$$

holds; consequently (6.88) is valid with $c_{29} = 25 \cdot l^2$ and $c_{28} := 1 \leq c_1 \log(|\mathcal{C}|/|\mathcal{M}|)/2 - 4 \log 2$. The last inequality holds, since $|\mathcal{C}| > |\mathcal{M}|$ and c_1 was chosen large enough; see subsection 2.1. ■

Lemma 6.39 *There exist constants $c_{31} > 0$ and $c_{32} > 0$ such that:*

$$P \left[(B_{\text{seed I}}^n)^c \cap E_{\text{stop}, \tau}^m \right] \leq c_{32} e^{-c_{31} n}. \quad (6.93)$$

Proof. We proceed similarly to the proof of Lemma 6.34. In the following, Z runs over all classes $Z \in \mathbb{Z}/l_{\rightarrow}\mathbb{Z}$. We set for all Z (compare with Definition (6.17) of $B_{\text{seed I}}^n$):

$$B_Z^{n,k} := \left\{ \begin{array}{l} S(\tau(k) + h) \in Z, S[(\tau(k) + h + [0, 3c_1 n l_{\leftarrow}]]) \text{ is a right ladder path,} \\ \text{and } S[(\tau(k) + h + 3c_1 n l_{\leftarrow} + [0, 3c_1 n l_{\rightarrow}]]) \text{ is a left ladder path.} \end{array} \right\}, \quad (6.94)$$

$$A_Z^{n,k} := E_{\text{stop}, \tau, k}^m \setminus B_Z^{n,k}. \quad (6.95)$$

where $E_{\text{stop}, \tau, k}^m$ is given by (6.71). Note that $B_{\text{seed I}}^n = \bigcap_Z \bigcup_{k=0}^{2^{\alpha n} - 1} B_Z^{n,k}$ and still $E_{\text{stop}, \tau}^m \subseteq E_{\text{stop}, \tau, k}^m$ for $k < 2^{\alpha n}$; thus

$$E_{\text{stop}, \tau}^m \setminus B_{\text{seed I}}^n \subseteq \bigcup_Z \bigcap_{k=0}^{2^{\alpha n} - 1} A_Z^{n,k}. \quad (6.96)$$

We obtain

$$P \left[E_{\text{stop}, \tau}^m \setminus B_{\text{seed I}}^n \right] \leq l_{\rightarrow} \max_Z P \left[\bigcap_{k=0}^{2^{\alpha n} - 1} A_Z^{n,k} \right] = l_{\rightarrow} \max_Z \prod_{k=0}^{2^{\alpha n} - 1} P \left[A_Z^{n,k} \middle| \bigcap_{j < k} A_Z^{n,j} \right], \quad (6.97)$$

$$P \left[A_Z^{n,k} \middle| \bigcap_{j < k} A_Z^{n,j} \right] \leq P \left[(B_Z^{n,k})^c \middle| E_{\text{stop}, \tau, k}^m \cap \bigcap_{j < k} A_Z^{n,j} \right]. \quad (6.98)$$

Since $h + 3c_1 n l_{\leftarrow} + 3c_1 n l_{\rightarrow} < 2 \cdot 2^{2n}$, we have $C_Z^{n,k} := E_{\text{stop}, \tau, k}^m \cap \bigcap_{j < k} A_Z^{n,j} \in \mathcal{F}_{\tau(k)}$. Using Lemma 6.11, we know $l_{\rightarrow}\mathbb{Z} + \text{supp } \mu^h = \mathbb{Z}$; hence $c_{33} := \inf_{x \in \mathbb{Z}} P[x + S(h) \in Z] > 0$; (note that the random walk $x + S$ starts in the point x). Moreover, given that $S(h) \in Z$, the probability to follow a right ladder path in Z in the subsequent $3c_1 n l_{\leftarrow}$ steps is $\mu(\{l_{\rightarrow}\})^{c_1 n l_{\leftarrow}}$, and the probability to follow then a left ladder path in the next $3c_1 n l_{\rightarrow}$ steps is $\mu(\{-l_{\leftarrow}\})^{c_1 n l_{\rightarrow}}$.

Thus by the strong Markov property:

$$P \left[B_Z^{n,k} \middle| C_Z^{n,k} \right] \geq c_{33} \mu(\{l_{\rightarrow}\})^{3c_1 n l_{\leftarrow}} \mu(\{-l_{\leftarrow}\})^{3c_1 n l_{\rightarrow}} = c_{33} e^{-c_{34} c_1 n}, \quad (6.99)$$

where $c_{34} := -3l_{\leftarrow} \log \mu(\{l_{\rightarrow}\}) - 3l_{\rightarrow} \log \mu(\{-l_{\leftarrow}\})$. We combine (6.97), (6.98) and (6.99) to obtain

$$P \left[E_{\text{stop}, \tau}^m \setminus B_{\text{seed I}}^n \right] \leq l_{\rightarrow} (1 - c_{33} e^{-c_{34} c_1 n})^{2^{\alpha n}} \leq l_{\rightarrow} \exp \left\{ -c_{33} e^{-c_{34} c_1 n} 2^{\alpha n} \right\}. \quad (6.100)$$

We have $c_{34} c_1 < \alpha \log 2$, since α was chosen large enough; see subsection 2.1. Thus the right hand side of the last inequality converges to 0 superexponentially fast as $n \rightarrow \infty$. This proves the lemma, since $n \geq n_0$ and n_0 was chosen large enough. ■

Finally we reap the results of this section:

Proof of Theorem 6.3. By Theorems 6.17 and 6.31 we have

$$E_{\text{xi does it}}^n \cap E_{\text{all pieces ok}}^n \supseteq \quad (6.101)$$

$$B_{\text{seed I}}^n \cap B_{\text{unique fit}}^n \cap B_{\text{all paths}}^n \cap B_{\text{outside out}}^n \cap B_{\text{recogn straight}}^n \cap B_{\text{signals}}^n \cap E_{\text{stop}, \tau}^m, \quad (6.102)$$

hence

$$\begin{aligned} E_{\text{stop},\tau}^m \setminus (E_{\text{xi does it}}^n \cap E_{\text{all pieces ok}}^n) &\subseteq (B_{\text{unique fit}}^n)^c \cup ((B_{\text{all paths}}^n)^c \cap E_{\text{stop},\tau}^m) \\ &\cup (B_{\text{outside out}}^n)^c \cup (B_{\text{recogn straight}}^n)^c \cup (B_{\text{signals}}^n)^c \cup ((B_{\text{seed 1}}^n)^c \cap E_{\text{stop},\tau}^m). \end{aligned} \quad (6.103)$$

Thus Theorem 6.3 follows from the main Lemmas 6.33, 6.34, 6.35, 6.37, 6.38, and 6.39 of this subsection. ■

7 How to find back: Correctness of the stopping times T_f

In this section, we prove Theorem 3.11.

Definition 7.1 Let $T = (T_k)_{k \in \mathbb{N}}$ be a sequence of \mathcal{G} -adapted stopping times. We define the events

$$E_{\text{no error},T}^m := \left\{ \forall k \geq 0 : \text{if } T_k(\chi) < 2^{12\alpha n_m}, \text{ then } |S(T_k(\chi))| \leq 2^{n_m} \right\}, \quad (7.1)$$

$$E_{\text{enough back}}^m := \left\{ \text{Up to time } 2^{12\alpha n_m}/8, S \text{ visits } 0 \text{ at least } \left. \vphantom{\left\{ \right.} \right. \right\} 2^{3\alpha n_m} \text{ times} \right\}. \quad (7.2)$$

We abbreviate

$$\Xi_{\text{reconst},f}^m := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P[E_{\text{reconst},f}^m \mid \xi] \geq \frac{1}{2} \right\}; \quad (7.3)$$

recall Definition (3.15) of the event $E_{\text{reconst},f}^m$.

Lemma 7.2 For some constant c_{35} and all $m \geq 0$:

$$1 - P[E_{\text{enough back}}^m] \leq c_{35} 2^{-\alpha n_m}. \quad (7.4)$$

Proof of Lemma 7.2. Let $(X_i)_{i \geq 1}$ denote the time difference between the $(i+1)$ st and the i -th visit of S at the origin. By recurrence, $(X_i)_{i \geq 1}$ is a.s. well defined, and by the strong Markov property it is i.i.d. with respect to P . Since S starts in the origin, X_1 is the first return time to the origin, and $\sum_{i=1}^j X_i$ is (a.s.) the time of the j -th visit at the origin. For the sake of this proof, we abbreviate: $x = 2^{12\alpha n_m}/8$ and $y = 2^{3\alpha n_m}$. Using

$$(E_{\text{enough back}}^m)^c = \left\{ \sum_{i=1}^y X_i \geq x \right\} \subseteq \left\{ \left(\sum_{i=1}^y X_i^{1/3} \right)^3 \geq x \right\} \quad (7.5)$$

and the Chebyshev-Markov inequality, we obtain the claim (7.4):

$$\begin{aligned} 1 - P[E_{\text{enough back}}^m] &\leq P \left[\sum_{i=1}^y X_i^{1/3} \geq x^{1/3} \right] \leq x^{-1/3} E \left[\sum_{i=1}^y X_i^{1/3} \right] \\ &= x^{-1/3} y E[X_1^{1/3}] = 2E[X_1^{1/3}] 2^{-\alpha n_m}. \end{aligned} \quad (7.6)$$

The fact $E[X_1^{1/3}] < \infty$ is proven in the appendix, Lemma A.1. ■

Definition 7.3 Let $v(k)$, $k \geq 0$, denote the $(k+1)$ st visit of S to the origin. We introduce a random set $\mathbb{T}'_f(\xi, \chi)$ and an event $E_{\text{when back recog}}^{m+1}$:

$$\mathbb{T}'_f(\xi, \chi) := \{t \in \mathbb{N} \mid \xi[[-2^{n_m}, 2^{n_m}] \preceq f(\theta^t(\chi)) \preceq \xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]\}, \quad (7.7)$$

$$E_{\text{when back recog}, f}^{m+1} := \left\{ \begin{array}{l} \text{For more than } 1/4 \text{ of the points } k \in [0, 2^{2\alpha n_{m+1}}[\text{ holds} \\ v(k2^{\alpha n_{m+1}}) \in \mathbb{T}'_f(\xi, \chi) \end{array} \right\}. \quad (7.8)$$

Lemma 7.4 If the event $E_{\text{reconst}, f}^m$ holds, then $\mathbb{T}_f(\chi) \supseteq \mathbb{T}'_f(\xi, \chi) \cap [0, 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m}[$.

Proof. We know $\xi[[-2^{n_m}, 2^{n_m}] \preceq f(\chi)$ by $E_{\text{reconst}, f}^m$. Let $t \in \mathbb{T}'_f(\xi, \chi)$, $t < 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m}$. Then we also have $\xi[[-2^{n_m}, 2^{n_m}] \preceq f(\theta^t \chi)$. Hence $t \in \mathbb{T}_f(\chi)$; to this end recall Definition (3.12) of the random set $\mathbb{T}_f(\chi)$. This implies the lemma. ■

Lemma 7.5 Assume that the events $E_{\text{no error}, T_f}^{m+1} \cap E_{\text{enough back}}^{m+1} \cap E_{\text{when back recog}, f}^{m+1}$ and $\mathbb{T}_f(\chi) \supseteq \mathbb{T}'_f(\xi, \chi) \cap [0, 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m}[$ hold. Then $E_{\text{stop}, T_f}^{m+1}$ holds, too.

Proof. Using $E_{\text{enough back}}^{m+1}$, we know

$$v(k2^{\alpha n_{m+1}}) \in [0, 2^{12\alpha n_{m+1}}/8] \subseteq [0, 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m}[\quad (7.9)$$

for all $k \in [0, 2^{2\alpha n_{m+1}}[$. Since the event $E_{\text{when back recog}, f}^{m+1}$ holds, we obtain $|\mathbb{T}_f(\chi)| \geq |\mathbb{T}'_f(\xi, \chi) \cap [0, 2^{12\alpha n_{m+1}} - 2 \cdot 2^{12\alpha n_m}[| \geq 2^{2\alpha n_{m+1}}/4$. By Definition (3.13) of the stopping times T_f , this yields $T_{f,k}(\chi) < 2^{12\alpha n_{m+1}}$ for all $k < (2^{2\alpha n_{m+1}}/4)/(2 \cdot 2^{2n_{m+1}}) = 2^{2(\alpha-1)n_{m+1}}/8$. The event $E_{\text{no error}, T_f}^{m+1}$ holds, and $2^{2(\alpha-1)n_{m+1}}/8 \geq 2^{\alpha n_{m+1}}$; recall that α and $n_{m+1} \geq n_0$ are large (see Section 2.1). Hence we obtain $|S(T_{f,k}(\chi))| \leq 2^{n_{m+1}}$ for all $k \in [0, 2^{\alpha n_{m+1}}[$; recall (7.1). Using Definition (3.13) again, we see that $T_{f,j}(\chi) + 2 \cdot 2^{2n_{m+1}} \leq T_{f,k}(\chi)$ is automatically fulfilled for $j < k$ whenever $T_{f,k}(\chi) < 2^{12\alpha n_{m+1}}$, which is the case at least for $k \in [0, 2^{\alpha n_{m+1}}[$. Summarizing, we have proven that the event $E_{\text{stop}, T_f}^{m+1}$ holds; recall its definition (3.14). ■

Lemma 7.6 $P \left[\left(E_{\text{when back recog}, f}^{m+1} \right)^c \cap \left\{ \xi \in \Xi_{\text{reconst}, f}^m \right\} \right] \leq 0.9^{2^{2\alpha n_{m+1}}}$.

Proof. We define Bernoulli random variables Y_k , $k \geq 0$, by $Y_k := 1$ if $v(k2^{\alpha n_{m+1}}) \in \mathbb{T}'_f(\xi, \chi)$, and $Y_k := 0$ else. Note that $v((k+1)2^{\alpha n_{m+1}}) - v(k2^{\alpha n_{m+1}}) \geq 2^{\alpha n_{m+1}} > 2 \cdot 2^{12\alpha n_m}$. Also note that $E_{\text{when back recog}, f}^{m+1} = \left\{ 2^{-2\alpha n_{m+1}} \sum_{k=0}^{2^{2\alpha n_{m+1}}-1} Y_k \geq 1/4 \right\}$. Because of the strong Markov property of the random walk $(S(k))_{k \geq 0}$ we have that conditioned under ξ the variables $(Y_k)_{k \geq 0}$ are i.i.d.; recall that $f(\chi)$ depends at most on $\chi[[-2^{12\alpha n_m}, 2^{12\alpha n_m}[$. If furthermore $\xi \in \Xi_{\text{reconst}, f}^m$ holds, then $E[Y_k \mid \xi] \geq 1/2$. Hence we obtain for these ξ , using the exponential Chebyshev inequality for the binomial variable $\sum_{k=0}^{2^{2\alpha n_{m+1}}-1} Y_k$:

$$\begin{aligned} P \left[\left(E_{\text{when back recog}, f}^{m+1} \right)^c \mid \xi \right] &= P \left[2^{-2\alpha n_{m+1}} \sum_{k=0}^{2^{2\alpha n_{m+1}}-1} Y_k \leq \frac{1}{4} \mid \xi \right] \\ &\leq E \left[e^{1/4 - Y_1} \mid \xi \right]^{2^{2\alpha n_{m+1}}} \leq \left(\frac{e^{1/4} + e^{-3/4}}{2} \right)^{2^{2\alpha n_{m+1}}} \leq 0.9^{2^{2\alpha n_{m+1}}}. \end{aligned} \quad (7.10)$$

This yields the claim of the lemma:

$$\begin{aligned} & P \left[\left(E_{\text{when back recog},f}^{m+1} \right)^c \cap \{ \xi \in \Xi_{\text{reconst},f}^m \} \right] \\ & \leq P \left[\left(E_{\text{when back recog},f}^{m+1} \right)^c \mid \xi \in \Xi_{\text{reconst},f}^m \right] \leq 0.9^{2^{2\alpha n_{m+1}}}. \end{aligned} \quad (7.11)$$

■

Lemma 7.7 $P \left[\left(E_{\text{no error},T_f}^{m+1} \right)^c \cap E_{\text{reconst},f}^m \right] \leq \frac{1}{3} e^{-n_{m+1}}.$

Proof. Let v_i denote the $(i+1)$ st time when the random walk S visits a point of $\mathbb{Z} \setminus [-2^{n_{m+1}} + 2l2^{12\alpha n_m}, 2^{n_{m+1}} - 2l2^{12\alpha n_m}]$. We set

$$E_{\text{wrong},i}^{m+1} := \{ \exists w \in \mathcal{C}^{2 \cdot 2^{n_m}} : w \preceq \xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]] \text{ and } w \preceq f(\theta^{v_i}(\chi)) \}. \quad (7.12)$$

If the event $E_{\text{wrong},i}^{m+1}$ occurs, then our procedure might fail to estimate correctly the location of the random walk: we might be misled to think that at time $v_i + 2 \cdot 2^{12\alpha n_m}$ we are close to the origin while we are not.

We claim that the following holds:

$$\left(E_{\text{no error},T_f}^{m+1} \right)^c \cap E_{\text{reconst},f}^m \subseteq \bigcup_{i=0}^{2^{12\alpha n_{m+1}} - 1} E_{\text{wrong},i}^{m+1}. \quad (7.13)$$

Indeed: If $\left(E_{\text{no error},T_f}^{m+1} \right)^c$ holds, then $|S(T_{f,k}(\chi))| > 2^{n_{m+1}}$ for some k with $T_{f,k} < 2^{12\alpha n_{m+1}}$ (see (7.1)); thus $|S(T_{f,k}(\chi) - 2 \cdot 2^{12\alpha n_m})| > 2^{n_{m+1}} - 2l2^{12\alpha n_m}$, since S cannot travel faster than speed l . This means $T_{f,k}(\chi) - 2 \cdot 2^{12\alpha n_m} = v_i$ for some $i < 2^{12\alpha n_{m+1}}$. Using Definition 3.8 of $T_{f,k}(\chi)$, this implies $v_i \in \mathbb{T}_f(\chi)$; hence there is $w \in \mathcal{C}^{2 \cdot 2^{n_m}}$ such that $w \preceq f(\chi)$ and $w \preceq f(\theta^{v_i}(\chi))$. Assuming that the event $E_{\text{reconst},f}^m$ holds, too, this implies $w \preceq f(\chi) \preceq \xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]$; see (3.15). This yields that $E_{\text{wrong},i}^{m+1}$ holds; recall (7.12). Summarizing, we have shown that (7.13) holds.

For all i , $f(\theta^{v_i}(\chi))$ depends only on $\chi[[v_i, v_i + 2 \cdot 2^{12\alpha n_m}[$, and S does not visit $[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]$ in this time interval $[v_i, v_i + 2 \cdot 2^{12\alpha n_m}[$, since the distance between $[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]$ and $\mathbb{Z} \setminus [-2^{n_{m+1}} + 2l2^{12\alpha n_m}, 2^{n_{m+1}} - 2l2^{12\alpha n_m}]$ is larger than $2l2^{12\alpha n_m}$, and since the random walk cannot travel faster than l steps per time unit.

Thus by the strong Markov property and by independence of S and ξ , we get: $\chi[[v_i, v_i + 2^{12\alpha n_m}]$ is independent of $\xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]$; therefore $f(\theta^{v_i}(\chi))$ is independent of $\xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]$, too.

The probability that a random word of length $2 \cdot 2^{n_m}$ which has i.i.d. letters with uniform distribution in \mathcal{C} is equal to a word which is independent of it is equal to $|\mathcal{C}|^{-2 \cdot 2^{n_m}}$. There are at most $37 \cdot 2^{n_m}$ words of a fixed length in $\xi[[-9 \cdot 2^{n_m}, 9 \cdot 2^{n_m}]]$ and also in $f(\theta^{v_k}(\chi))$, counting all reversed words, too. Thus there are at most $37^2 2^{2n_m}$ pairs of such words. It follows that

$$P \left[E_{\text{wrong},i}^{m+1} \right] \leq 37^2 2^{2n_m} |\mathcal{C}|^{-2 \cdot 2^{n_m}}. \quad (7.14)$$

Hence we get the claim of the lemma, using (7.13):

$$\begin{aligned} P \left[\left(E_{\text{no error},T_f}^{m+1} \right)^c \cap E_{\text{reconst},f}^m \right] & \leq \sum_{i=0}^{2^{12\alpha n_{m+1}} - 1} P \left[E_{\text{wrong},i}^{m+1} \right] \\ & \leq 2^{12\alpha n_{m+1}} \cdot 37^2 2^{2n_m} |\mathcal{C}|^{-2 \cdot 2^{n_m}} \leq \frac{1}{3} e^{-n_{m+1}}. \end{aligned} \quad (7.15)$$

For the last inequality, recall that $n_m \geq n_0$ is large enough, and note that $|\mathcal{C}|^{-2 \cdot 2^{n_m}}$ is the leading term of the last but one expression; also recall that $n_{m+1} = 2^{\lfloor \sqrt{n_m} \rfloor}$ is of a much smaller order than 2^{n_m} . ■

Proof of Theorem 3.11. By Lemmas 7.4 and 7.5, we know $E_{\text{no error}, T_f}^{m+1} \cap E_{\text{enough back}}^{m+1} \cap E_{\text{when back recog}, f}^{m+1} \cap E_{\text{reconst}, f}^m \subseteq E_{\text{stop}, T_f}^{m+1}$. Using some Boolean algebra, this implies

$$\begin{aligned} & (E_{\text{stop}, T_f}^{m+1})^c \cap E_{\text{reconst}, f}^m \cap \{\xi \in \Xi_{\text{reconst}, f}^m\} \\ & \subseteq \left(E_{\text{enough back}}^{m+1} \right)^c \cup \left((E_{\text{no error}, T_f}^{m+1})^c \cap E_{\text{reconst}, f}^m \right) \cup \left(\left(E_{\text{when back recog}, f}^{m+1} \right)^c \cap \{\xi \in \Xi_{\text{reconst}, f}^m\} \right). \end{aligned} \quad (7.16)$$

Consequently, using Definition (7.3) of $\Xi_{\text{reconst}, f}^m$ and Lemmas 7.2, 7.6, and 7.7:

$$\begin{aligned} & P \left[(E_{\text{stop}, T_f}^{m+1})^c \cap E_{\text{reconst}, f}^m \cap \left\{ P \left[E_{\text{reconst}, f}^m \mid \xi \right] \geq \frac{1}{2} \right\} \right] \\ & \leq P \left[\left(E_{\text{enough back}}^{m+1} \right)^c \right] + P \left[(E_{\text{no error}, T_f}^{m+1})^c \cap E_{\text{reconst}, f}^m \right] \\ & \quad + P \left[\left(E_{\text{when back recog}, f}^{m+1} \right)^c \cap \{\xi \in \Xi_{\text{reconst}, f}^m\} \right] \\ & \leq c_{35} 2^{-\alpha n_{m+1}} + \frac{1}{3} e^{-n_{m+1}} + 0.9 2^{2\alpha n_{m+1}} \leq e^{-n_{m+1}}; \end{aligned} \quad (7.17)$$

recall that α and $n_{m+1} \geq n_0$ are large (see Section 2.1). This proves Theorem 3.11. ■

8 Getting started: The first stopping times

In this section, we prove Theorem 3.10.

8.1 The stopping times T^0

We start with the definition of a sequence $T^0 = (T_k^0)_{k \geq 0}$ of \mathcal{G} -adapted stopping times with values in $[0, 2^{12\alpha n_0}]$. Roughly speaking, these times search for long blocks of 1's in the observation χ . Here is intuitive idea behind this construction: Since we conditioned on a large block of 1's to occur in the true scenery ξ close to the origin, observing a long block of 1's at a later time indicates with high probability that the random walk has returned close to the origin. This is true only up to a certain time horizon, since long blocks of 1's in the true scenery will occur far from the origin, as well.

Definition 8.1 *Let the random set $\mathbb{T}^0(\chi)$ be defined as follows:*

$$\mathbb{T}^0(\chi) := \{t \in [0, 2^{12\alpha n_0} - n_0^7] \mid \chi \upharpoonright [t, t + n_0^7] \text{ is constant } 1\}. \quad (8.1)$$

We arrange the elements of $\mathbb{T}^0(\chi)$ in increasing order: $t_0(0) < \dots < t_0(|\mathbb{T}^0(\chi)| - 1)$. We set

$$T_k^0(\chi) := \begin{cases} t_0(2 \cdot 2^{2n_0} k) + n_0^7 & \text{if } 2 \cdot 2^{2n_0} k < |\mathbb{T}^0(\chi)|, \\ 2^{12\alpha n_0} & \text{otherwise.} \end{cases} \quad (8.2)$$

Recall Definition (3.14):

$$E_{\text{stop}, T^0}^0 = \bigcap_{k=0}^{2\alpha n_0} \{T_k^0(\chi) < 2^{12\alpha n_0}, |S(T_k^0(\chi))| \leq 2^{n_0}, T_j^0(\chi) + 2 \cdot 2^{2n_0} \leq T_k^0(\chi) \text{ for } j < k\}. \quad (8.3)$$

Theorem 8.2 For some positive constants c_{36} and c_{37} holds $\tilde{P} \left[E_{\text{stop}, T^0}^0 \right] \geq 1 - c_{36} e^{-c_{37} n_0}$.

We prepare the proof of Theorem 8.2 by some Definitions and Lemmas. We use again the abbreviation $J_1 = [-2\ln_0^{20}, 2\ln_0^{20}]$.

Definition 8.3 (Analogue of Definition (7.2)) Let $I \subseteq J_1$ be an integer interval. We define

$$\tilde{E}_{\text{enough back}, I}^0 := \left\{ \text{Up to time } 2^{12\alpha n_0}/4, S \text{ visits the interval } I \text{ at least } 2^{3\alpha n_0} \text{ times} \right\}. \quad (8.4)$$

Lemma 8.4 For some constants $c_{38} > 0$ and $c_{39} > 0$, the following holds: If $I \subseteq J_1$, $|I| \geq l$, is an integer interval, then

$$\tilde{P}[\tilde{E}_{\text{enough back}, I}^0] \geq 1 - c_{38} e^{-c_{39} n_0}. \quad (8.5)$$

Proof. Let $T_I := \inf\{t \mid S(t) \in I\}$ be the entrance time of S into I . We show first: For some positive constants c_{40} and c_{41} (depending at most on the distribution μ of $S(1)$) we have:

$$\tilde{P}[T_I \geq 2^{12\alpha n_0}/8] \leq c_{40} e^{-c_{41} n_0}. \quad (8.6)$$

If $0 \in I$, this is trivial, since S starts in 0. Otherwise I contains only positive numbers, or it contains only negative numbers; without loss of generality we assume the first possibility. Let $z = \min I \in]0, 2\ln_0^{20}[$. Consider the interval $J :=]-2^{n_0}, z[\subseteq]-2^{n_0}, 2^{n_0}[$, and consider the exit time $H := \inf\{t \mid S(t) \notin J\}$ of J . Note that H is a.s. finite.

On the one hand, we know

$$\tilde{P}[H \geq t] \leq c_{42} e^{-c_{43} 2^{-2n_0} t} \quad (8.7)$$

for some constants $c_{42}, c_{43} > 0$ depending at most on the variance of $S(1)$, since in every time interval of size 2^{2n_0} the random walk has a positive probability to exit J , bounded away from 0. In particular, for $t = 2^{12\alpha n_0}/8$ the probability in (8.7) is superexponentially small in n_0 .

On the other hand, since S is a martingale and since S has jumps sizes bounded by l , we get $\tilde{P}[S(H) > 0] \geq 1 - (z + l)2^{-n_0}$. Furthermore, using again that S has jumps sizes bounded by l , we know the following: If $S(H) > 0$, then $S(H) \in I$ and $T_I = H$, since z is the leftmost point in I and $|I| \geq l$; the random walk cannot cross I without touching it.

Altogether, we have the following upper bound for the left hand side in (8.6):

$$\tilde{P}[H \geq 2^{12\alpha n_0}/8 \text{ or } S(H) < 0] \leq c_{40} e^{-c_{41} n_0}. \quad (8.8)$$

for some positive constants c_{40} and c_{41} .

Provided the random walk visits a point $x \in I$, the probability to visit this point again at least $2^{3\alpha n_m}$ times in the subsequent $2^{12\alpha n_m}/8$ time steps is at least $1 - c_{28} 2^{-\alpha n_0}$. This follows from Lemma 7.2, using the strong Markov property of the random walk; recall that the law of S with respect to P and with respect to \tilde{P} coincide. Combining this with (8.6) yields claim (8.5) of Lemma 8.4. ■

We remark: Lemma 8.4 holds not only for deterministic intervals I , but also for random ones, provided that I and S are independent. We use this below for the following specific choice of I , which depends on the scenery ξ , but not on S :

\tilde{P} -a.s. there is a (random) integer interval $J_0 \subseteq J_1 = [-2\ln_0^{20}, 2\ln_0^{20}]$ with $|J_0| \geq n_0^4$ such that $\xi \upharpoonright_{J_0}$ is constant 1; recall Definition 3.2. Just for definiteness we take the rightmost such J_0 . Let

$$I = I(\xi) := \left\{ z \in J_0 \mid \text{dist}(z, \mathbb{Z} \setminus J_0) > n_0^4/4 \right\}; \quad (8.9)$$

then I is \tilde{P} -a.s. well defined, and it is an integer interval containing $|I| \geq n_0^4/2 \geq l$ points.

Definition 8.5 (Modification of Definition 7.3) Let $w(k)$, $k \geq 0$, denote the $(k+1)$ st visit to the (random) set $I(\xi)$ by the random walk S . We introduce a random set $\mathbb{T}^{0'}$ and an event $\tilde{E}_{\text{when back recog}}^0$:

$$\mathbb{T}^{0'} := \{t \in \mathbb{N} \mid S(t) \in I(\xi) \text{ and } |S(j) - S(t)| \leq n_0^4/4 \text{ for } 0 \leq j - t \leq n_0^7\}, \quad (8.10)$$

$$\tilde{E}_{\text{when back recog}}^0 := \{\text{For more than } 1/4 \text{ of the points } k \in [0, 2^{2\alpha n_0}[\text{ holds } w(k2^{\alpha n_0}) \in \mathbb{T}^{0'}\}. \quad (8.11)$$

Lemma 8.6 (Modification of Lemma 7.6) $\tilde{P} \left[\left(\tilde{E}_{\text{when back recog}}^0 \right)^c \right] \leq 0.9^{2^{2\alpha n_0}}$.

Proof. We observe as in (4.16) by the submartingale inequality:

$$P[|S(j)| \leq n_0^4/4 \text{ for } 0 \leq j \leq n_0^7] \geq 1 - 4^2 n_0^{-8} E[S(n_0^7)^2] = 1 - \frac{16 \text{Var}[S(1)]}{n_0} \geq \frac{1}{2}, \quad (8.12)$$

since n_0 is large enough; see Subsection 2.1. Let Y_k denote the indicator function of the event $\{w(k2^{\alpha n_0}) \in \mathbb{T}^{0'}\}$; the Y_k are \tilde{P} -a.s. well defined. As a consequence of the strong Markov property, the Y_k , $k \in [0, 2^{2\alpha n_0}[$, are i.i.d. Bernoulli random variables; note that the stopping times $w(k2^{\alpha n_0})$, $k \in \mathbb{N}$, have at least the spacing $2^{\alpha n_0} > n_0^7$. Furthermore $\tilde{P}[Y_k = 1] \geq 1/2$, since this probability equals the left hand side in (8.12). The claim of the Lemma now follows by the same large deviation argument as in (7.10). ■

Lemma 8.7 (Analogue of Lemma 7.4) The inclusion $\mathbb{T}^0 \supseteq \mathbb{T}^{0'} \cap [0, 2^{12\alpha n_0} - n_0^7[$ holds \tilde{P} -almost surely.

Proof. Assume that the event BigBlock holds; this occurs \tilde{P} -almost surely. Then $I(\xi)$ is well defined. Let $t \in \mathbb{T}^{0'}$, $t < 2^{12\alpha n_0} - n_0^7$. Then $S(t) \in I(\xi)$, and during the subsequent n_0^7 steps, the random walk S cannot leave the interval J_0 , since it does not travel farther than $n_0^4/4$ (recall definition (8.10)), and since $\mathbb{Z} \setminus J_0$ is more distant than this from $I(\xi)$ (recall the definition of $I(\xi)$). Since $\xi \upharpoonright J_0$ is constant 1 by definition (3.4) of the event BigBlock, this implies that $S \upharpoonright [t, t + n_0^7]$ is constant 1; i.e. $t \in \mathbb{T}^0$. ■

Recall Definition (7.1):

$$E_{\text{no error}, T^0}^0 = \{\forall k \in \mathbb{N} : \text{If } T_k^0(\chi) < 2^{12\alpha n_0}, \text{ then } |S(T_k^0(\chi))| \leq 2^{n_0}\}. \quad (8.13)$$

Lemma 8.8 (Modification of Lemma 7.5) Assume that the events $E_{\text{no error}, T^0}^0 \cap \tilde{E}_{\text{enough back}, I(\xi)}^0 \cap \tilde{E}_{\text{when back recog}}^0$ and $\mathbb{T}^0(\chi) \supseteq \mathbb{T}^{0'} \cap [0, 2^{12\alpha n_0} - n_0^7[$ hold. Then E_{stop, T^0}^0 holds, too.

Proof. The proof is almost the same as for Lemma 7.5. Note that the small differences between the definitions of $\tilde{E}_{\text{enough back}, I(\xi)}^0$ and $E_{\text{enough back}}^{m+1}$ are not essential for the validity of the proof. ■

Definition 8.9 We define the event of sceneries

$$\Xi_{\text{no blocks}}^0 := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid \begin{array}{l} \text{For every (integer) interval } J \subseteq [-2l2^{12\alpha n_0}, 2l2^{12\alpha n_0}] \setminus J_1 \\ \text{with } |J| = n_0^2 \text{ holds: } \xi \upharpoonright J \text{ is not constant 1.} \end{array} \right\} \quad (8.14)$$

Lemma 8.10 For some positive constants c_{44}, c_{45} holds $\tilde{P}[\xi \in \Xi_{\text{no blocks}}^0] \geq 1 - c_{44}e^{-c_{45}n_0}$.

Proof. For every fixed interval $J \subseteq [-2 \cdot l2^{12\alpha n_0}, 2 \cdot l2^{12\alpha n_0}] \setminus J_1$ with $|J| = n_0^2$ we have

$$\tilde{P}[\xi[J \text{ is constant } 1]] = |\mathcal{C}|^{-n_0^2}, \quad (8.15)$$

which is superexponentially small in n_0 . Furthermore, there are less than $4l2^{12\alpha n_0}$ such intervals. Thus $\tilde{P}[\xi \notin \Xi_{\text{no blocks}}^0] \leq 4l2^{12\alpha n_0}|\mathcal{C}|^{-n_0^2}$, which is still superexponentially small in n_0 . This implies the lemma. Note that we may choose c_{44}, c_{45} independent α for n_0 large enough, even though $4l2^{12\alpha n_0}|\mathcal{C}|^{-n_0^2}$ does depend on α (see Subsection 2.1). ■

Lemma 8.11 For some constants $c_{46}, c_{47} > 0$ holds

$$\tilde{P}[(E_{\text{no error}, T^0}^0)^c] \leq c_{46}e^{-c_{47}n_0}. \quad (8.16)$$

Proof. Let \mathcal{X} be defined by

$$\mathcal{X} := \{x \in \mathbb{Z} | x + [-ln_0^7, ln_0^7] \subseteq [-2l2^{12\alpha n_0}, 2l2^{12\alpha n_0}] \setminus J_1\}. \quad (8.17)$$

As a consequence of Lemma 4.3 (with the parameters $k = n_0^7$ and $\kappa = n_0^2$) we know for every $\xi \in \mathcal{C}^{\mathbb{N}}$ such that $\xi[[-ln_0^7, ln_0^7]]$ contains no block of 1's of length n_0^2 :

$$P_\xi[\xi \circ S[[0, n_0^7] \text{ is constant } 1]] \leq e^{-c_7 n_0^3}. \quad (8.18)$$

Let $t \in \mathbb{N}$ and let $\xi \in \Xi_{\text{no blocks}}^0$. Using the Markov property of the random walk, (8.18) implies the following:

$$P_\xi[\xi \circ S[(t + [0, n_0^7]) \text{ is constant } 1 | S(t) \in \mathcal{X}]] \leq e^{-c_7 n_0^3}. \quad (8.19)$$

If $t < 2^{12\alpha n_0}$ and $|S(t)| > 2^{n_0}$ holds, then we know $S(t) \in \mathcal{X}$; note that $J_1 = [-2ln_0^{20}, 2ln_0^{20}]$ has a distance larger than ln_0^7 from $\mathbb{Z} \setminus [-2^{n_0}, 2^{n_0}]$, and recall that S cannot travel faster than with speed l , and that n_0 is large by Subsection 2.1.

Thus (8.19) implies

$$\begin{aligned} P_\xi[(E_{\text{no error}, T^0}^0)^c] & \leq P_\xi[\text{There is } t < 2^{12\alpha n_0} \text{ such that } |S(t)| > 2^{n_0} \text{ and } \xi \circ S[(t + [0, n_0^7]) \text{ is constant } 1]] \\ & \leq 2^{12\alpha n_0} e^{-c_7 n_0^3} \leq e^{-n_0}; \end{aligned} \quad (8.20)$$

for the last inequality recall that n_0 was chosen large enough, depending on α (see Subsection 2.1). Combining this with Lemma 8.10 yields for some positive constants c_{46}, c_{47} :

$$\tilde{P}[(E_{\text{no error}, T^0}^0)^c] \leq \tilde{P}[\xi \notin \Xi_{\text{no blocks}}^0] + \int_{\{\xi \in \Xi_{\text{no blocks}}^0\}} P_\xi[(E_{\text{no error}, T^0}^0)^c] d\tilde{P} \leq c_{46}e^{-c_{47}n_0}. \quad (8.21)$$

■

Proof of Theorem 8.2. From Lemmas 8.7 and 8.8 we know that

$$\tilde{P}[(E_{\text{stop}, T^0}^0)^c] \leq \tilde{P}[(E_{\text{no error}, T^0}^0)^c] + \tilde{P}[(\tilde{E}_{\text{enough back}, I(\xi)}^0)^c] + \tilde{P}[(\tilde{E}_{\text{when back recog}}^0)^c]. \quad (8.22)$$

Hence the claim of Theorem 8.2 is a consequence of Lemmas 8.4, 8.6, and 8.11. ■

8.2 The stopping times T^1

Unfortunately, the constructed stopping times T^0 are not good enough as arguments for the first reconstruction Algorithm “Algⁿ”: We cannot construct more than roughly $\exp(\text{const } n_0^4)$ reliable stopping times based on the way we build the T^0 stopping times; in fact we use much less than this number. Otherwise we cannot guarantee that they really stop the random walk with high probability close to the origin. However, the number $\exp(\text{const } n_0^4)$ is much too small to collect a sufficiently large puzzle for reconstructing at least the modified piece $\xi \upharpoonright J_1$ in the scenery using our reconstruction algorithm; to illustrate this fact, we remark that we have only roughly an upper bound $d\tilde{P}/dP \leq \exp(\text{const } n_0^{20})$; see Lemma 4.4. A modification of the parameters does not solve this problem; we need an essentially improved series of stopping times T^1 to get the reconstruction algorithm started.

Our construction of T^1 is partially parallel to the construction of the partial reconstruction algorithm Algⁿ, but it is also partially parallel to the construction of the stopping times T_f and T^0 : Roughly speaking, we collect a set of typical signals (“a puzzle”) at the very beginning and another one at a candidate time. Instead of matching the pieces together, we just compare the two puzzles: If the puzzles have a sufficiently high overlap, then they were generated with high probability at roughly the same location.

Fortunately, many constructions of the previous sections can be used again, up to small modifications: There are extra complications due to the presence of a modified domain J_1 . We keep the presentation as close as possible to the previous sections to show the parallelism. Here is the formal definition of the “new” puzzles and of T^1 :

Definition 8.12 *We set, using the abbreviation $\text{Input} := (T^0(\chi), \chi \upharpoonright [0, 2 \cdot 2^{12\alpha n_0}[$) again:*

$$\text{Puzzle}_{\text{III}}^{n_0}(\chi) := \tag{8.23}$$

$$\{(w_1, w_2, w_3) \in \text{Puzzle}_{\text{I}}^{n_0}(\text{Input}) \mid \exists k \in [0, 2^{\alpha n_0}[: w_1 w_2 w_3 \sqsubseteq \chi \upharpoonright (T_k^0(\chi) + [0, 2^{n_0}/l])\},$$

$$\text{Puzzle}_{\text{IV}}^{n_0}(\chi) := \{w_2 \in \mathcal{C}^{c_1 n_0} \mid \exists w_1, w_3 \in \mathcal{C}^{c_1 n_0} : (w_1, w_2, w_3) \in \text{Puzzle}_{\text{III}}^{n_0}(\chi)\}, \tag{8.24}$$

$$\mathbb{T}^1(\chi) := \left\{ t \in [0, 2^{12\alpha n_1} - 2 \cdot 2^{12\alpha n_0}[\mid \begin{array}{l} |\text{Puzzle}_{\text{IV}}^{n_0}(\chi) \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq 2^{n_0/3} \\ \text{and } |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0} \end{array} \right\}. \tag{8.25}$$

Finally we define another sequence $T^1 = (T_k^1)_{k \geq 0}$ of \mathcal{G} -adapted stopping times with values in $[0, 2^{12\alpha n_1}]$: Let $t_1(0) < \dots < t_1(|\mathbb{T}^1(\chi)| - 1)$ be the elements of $\mathbb{T}^1(\chi)$ arranged in increasing order. For $k \in \mathbb{N}$, we set

$$T_k^1(\chi) := \begin{cases} t_1(2 \cdot 2^{2n_1 k}) + 2 \cdot 2^{12\alpha n_0} & \text{if } 2 \cdot 2^{2n_1 k} < |\mathbb{T}^1(\chi)|, \\ 2^{12\alpha n_1} & \text{otherwise.} \end{cases} \tag{8.26}$$

Note that $T^0(\chi)$ only depends on $\chi \upharpoonright [0, 2^{12\alpha n_0}[$; thus $\text{Puzzle}_{\text{IV}}^{n_0}(\chi)$ only depends on $\chi \upharpoonright [0, 2 \cdot 2^{12\alpha n_0}[$, since $2^{n_0}/l \leq 2^{12\alpha n_0}$.

Definition 8.13 *Using the abbreviation $J_1 = [-2\ln_0^{20}, 2\ln_0^{20}]$ from Definition 3.2 again, we*

define the following random sets:

CorPaths :=

$$\left\{ R \in \mathbb{Z}^{[0, c_1 n_0[} \mid \begin{array}{l} R \text{ is an admissible piece of path, for every admissible piece of path} \\ R' : [0, c_1 n_0[\rightarrow \mathbb{Z} \text{ with } R'(0) = R(0) \text{ and } R'(c_1 n_0 - 1) = R(c_1 n_0 - 1) \\ \text{holds } \xi \circ R' = \xi \circ R, \text{ and there is such a path } R' \text{ which takes at} \\ \text{least one value in } J_1. \end{array} \right\}, \quad (8.27)$$

$$\text{Corrupted} := \{ \xi \circ R \in \mathcal{C}^{c_1 n_0} \mid R \in \text{CorPaths} \}, \quad (8.28)$$

$$\text{Center}_I := \{ w \in \mathcal{C}^{c_1 n_0} \mid w \text{ is a (left or right) ladder word of } \xi \uparrow ([-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}] \setminus J_1) \}, \quad (8.29)$$

$$\text{Center}_{II} := \text{Center}_I \cup \text{Corrupted}, \quad (8.30)$$

$$\text{Center}_{III} := \left\{ w \in \mathcal{C}^{c_1 n_0} \mid \begin{array}{l} w \text{ is a right ladder word of} \\ \xi \uparrow ([-2 \cdot 2^{n_0/2}, 2 \cdot 2^{n_0/2}] \setminus [-2^{n_0/2}, 2^{n_0/2}]) \end{array} \right\}. \quad (8.31)$$

Some of the definitions and lemmas below are only small modifications of previous definitions and lemmas, respectively. We underline the new pieces to show the differences.

Definition 8.14 (Modification of Definition 6.15) *We define:*

$$\tilde{B}_{\text{unique fit}}^{n_0} := \left\{ \begin{array}{l} \text{For every } i, j \in \{1, \dots, l^2\}, \text{ every } i\text{-spaced interval} \\ I \subseteq [-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}] \setminus J_1, \text{ and every } j\text{-spaced interval} \\ J \subseteq [-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}] \setminus J_1 \text{ with } |I| = |J| \geq c_2 n_0 \text{ holds} \\ \underline{(\xi \uparrow I)_\leftarrow \neq (\xi \uparrow J)_\rightarrow, \text{ and if } I \neq J, \text{ then } (\xi \uparrow I)_\rightarrow \neq (\xi \uparrow J)_\rightarrow.} \end{array} \right\}, \quad (8.32)$$

Lemma 8.15 (Modification of Lemma 6.33) *There exists constants $c_{18}, c_{19} > 0$ such that the following holds:*

$$\tilde{P} \left[(\tilde{B}_{\text{unique fit}}^{n_0})^c \right] \leq c_{18} e^{-c_{19} n_0}. \quad (8.33)$$

Proof. The proof of Lemma 6.33 remains literally true when we replace P by \tilde{P} , but additionally restrict I and J to be disjoint from J_1 , since the distributions of $\xi \uparrow (\mathbb{Z} \setminus J_1)$ with respect to \tilde{P} and with respect to P coincide; see Lemma 3.4. ■

Lemma 8.16 $|\text{Center}_I| \leq 46 \cdot 2^{n_0}$, $|\text{Corrupted}| \leq n_0^{41}$, and thus $|\text{Center}_{II}| \leq 50 \cdot 2^{n_0}$. *If the event $\tilde{B}_{\text{unique fit}}^{n_0}$ holds, then $|\text{Center}_{III}| \geq 2^{n_0/3}$.*

Proof. The first statement is obvious, since there are at most $23 \cdot 2^{n_0}$ choices for the leftmost point of a ladder interval in $[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$, and there is the binary choice “left” or “right”.

We show $|\text{Corrupted}| \leq n_0^{41}$ next: The number of pairs $(R(0), R(c_1 n_0 - 1)) \in \mathbb{Z}^2$ with $R \in \text{CorPaths}$ is bounded by $(|J_1| + c_1 n_0 l)^2 \leq n_0^{41}$; recall that n_0 was chosen to be large (see Subsection 2.1). Furthermore, every such pair gives rise to at most a single element of **Corrupted**, since different paths $R, R' \in \text{CorPaths}$ with the same starting point and the same end point generate the same word $\xi \circ R = \xi \circ R'$ by Definition (8.27). This shows $|\text{Corrupted}| \leq n_0^{41} \leq 4 \cdot 2^{n_0}$, since n_0 is large enough by Subsection 2.1. Using the definition of **Center_{II}**, we obtain $|\text{Center}_{II}| \leq 50 \cdot 2^{n_0}$.

Finally we show $|\text{Center}_{III}| \geq 2^{n_0/3}$. We observe that $[-2 \cdot 2^{n_0/2}, 2 \cdot 2^{n_0/2}] \setminus [-2^{n_0/2}, 2^{n_0/2}]$ is disjoint from J_1 . Assuming that $\tilde{B}_{\text{unique fit}}^{n_0}$ holds, this implies that all right ladder intervals

$I_1, I_2 \subseteq [-2 \cdot 2^{n_0/2}, 2 \cdot 2^{n_0/2}] \setminus [-2^{n_0/2}, 2^{n_0/2}]$, with $I_1 \neq I_2$ $|I_1| = |I_2| = c_1 n_0 \geq c_2 n_0$ generate pairwise different ladder words $(\xi[I_1]_{\rightarrow} \neq (\xi[I_2]_{\rightarrow})$. Since there are at least $2^{n_0/2} - c_1 n_0 \geq 2^{n_0/3}$ such ladder intervals (n_0 is large enough; see Subsection 2.1), there are at least as many ladder words $w \in \text{Center}_{\text{III}}$. ■

Lemma 8.17 *For every $x \in \mathbb{Z}$ with $|x| > 2 \cdot 2^{n_0} + 2l2^{12\alpha n_0}$ and for every $t \in [0, 2^{12\alpha n_1}[$ holds:*

$$\tilde{P} [S(t) = x, |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0}, \text{ and } |\text{Center}_{\text{II}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq 2^{n_0/3}] \leq \exp\{-2^{n_0/4}\}. \quad (8.34)$$

Proof. We set

$$\text{Outside}_{x,t} := \begin{cases} \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi) & \text{if } S(t) = x \text{ and } |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0}, \\ \emptyset & \text{else.} \end{cases} \quad (8.35)$$

The random set $\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)$ only depends on $\chi[[t, t + 2 \cdot 2^{12\alpha n_0}[$, and the random walk cannot travel a longer distance than $2l2^{12\alpha n_0}$ during the time interval $[t, t + 2 \cdot 2^{12\alpha n_0}[$. Given $S(t) = x$ and $|x| > 2 \cdot 2^{n_0} + 2l2^{12\alpha n_0}$, the random walk S cannot enter the interval $[-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}]$ during the time interval $[t, t + 2 \cdot 2^{12\alpha n_0}[$; thus $\text{Outside}_{x,t}$ depends only on S and $\xi[(\mathbb{Z} \setminus [-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}])]$. Hence, using Lemma 3.4 and $J_1 \subseteq [-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}]$, the random piece of scenery $\xi[[-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}]$ and the random set $\text{Outside}_{x,t}$ are independent with respect to \tilde{P} . Let \mathcal{I}_r denote the set of all right ladder intervals $I \subseteq ([-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}] \setminus J_1)$ with $|I| = c_1 n_0$. We define \mathcal{I}_l similarly with “right ladder intervals” replaced by “left ladder intervals”. We partition \mathcal{I}_r into $c_1 n_0 l_{\rightarrow}$ subsets, $\mathcal{I}'_r(1), \dots, \mathcal{I}'_r(c_1 n_0 l_{\rightarrow})$:

$$\mathcal{I}'_r(k) := \{I \in \mathcal{I}_r \mid \min I \in k + c_1 n_0 l_{\rightarrow} \mathbb{Z}\} \quad (8.36)$$

Let $k \in [1, c_1 n_0 l_{\rightarrow}]$ be fixed. Note that the cardinality $N := |\mathcal{I}'_r(k)|$ fulfills the bounds

$$\frac{2^{n_0}}{c_1 n_0 l_{\rightarrow}} \leq \frac{4 \cdot 2^{n_0}}{c_1 n_0 l_{\rightarrow}} - |J_1| - 2 \leq N \leq \frac{4 \cdot 2^{n_0}}{c_1 n_0 l_{\rightarrow}}. \quad (8.37)$$

Furthermore, the elements of $\mathcal{I}'_r(k)$ are pairwise disjoint; thus the family $(\xi[I])_{I \in \mathcal{I}'_r(k)}$ is i.i.d. and independent of $\text{Outside}_{x,t}$ (with respect to \tilde{P}). For $I \in \mathcal{I}_r$, we set $X_I^r := 1$ for $(\xi[I]_{\rightarrow} \in \text{Outside}_{x,t})$, and $X_I^r := 0$ otherwise. Similarly for $J \in \mathcal{I}_l$, let X_J^l denote the indicator function of the event $\{(\xi[J]_{\leftarrow} \in \text{Outside}_{x,t})\}$. Then, conditioned on a given value of $\text{Outside}_{x,t}$, the Bernoulli random variables X_I^r , $I \in \mathcal{I}'_r(k)$, are i.i.d. with respect to $\tilde{P}[\cdot \mid \text{Outside}_{x,t}]$. Furthermore we have, using $|\text{Outside}_{x,t}| \leq 50 \cdot 2^{n_0}$:

$$\tilde{P}[X_I^r = 1 \mid \text{Outside}_{x,t}] \leq |\text{Outside}_{x,t}| |\mathcal{C}|^{-|I|} \leq 50e^{(\log 2 - c_1 \log |\mathcal{C}|)n_0} =: p. \quad (8.38)$$

We set $Y_k^r := \sum_{I \in \mathcal{I}'_r(k)} X_I^r$. Consequently this random variable is stochastically dominated by a Binomial(N, p)–distributed random variable; note that Y_k^r is binomially distributed with respect to the conditioned measure $\tilde{P}[\cdot \mid \text{Outside}_{x,t}]$. A rough but simple large deviation estimate suffices for our purposes: Using the exponential Chebyshev inequality, we have for $a > 0$ and $\sigma := \log(a/p) > 0$:

$$\begin{aligned} \tilde{P}[Y_k^r \geq Na] &\leq E[e^{\sigma Y_k^r - Na}] \leq (pe^{\sigma(1-a)} + (1-p)e^{-\sigma a})^N = ((1+a-p)p^a a^{-a})^N \\ &\leq (e^a p^a a^{-a})^N = \exp\{Na(1 - \log(a/p))\} \end{aligned} \quad (8.39)$$

In particular, we obtain for the choice $a = N^{-1}2^{n_0/3}/(4c_1n_0l_{\rightarrow}) \geq 2^{-2n_0/3}/16$ (where we have used (8.37)), using (8.38): $\sigma = \log(a/p) \geq (c_1 \log |\mathcal{C}| - \frac{5}{3} \log 2)n_0 - \log 800 \geq c_1(\log |\mathcal{C}|)n_0/2 + 1$; the last inequality holds by our choice of c_1 and n_0 (see Subsection 2.1). Hence we obtain:

$$\begin{aligned} \tilde{P} \left[\sum_{I \in \mathcal{I}_r} X_I^r \geq \frac{2^{n_0/3}}{4} \right] &\leq \sum_{k=1}^{c_1 n_0 l_{\rightarrow}} \tilde{P} [Y_k^r \geq Na] \leq c_1 n_0 l_{\rightarrow} \exp\{Na(1 - \log(a/p))\} \\ &\leq c_1 n_0 l_{\rightarrow} \exp \left\{ -\frac{\log |\mathcal{C}|}{8l_{\rightarrow}} 2^{n_0/3} \right\} \leq \frac{1}{2} \exp\{-2^{n_0/4}\}. \end{aligned} \quad (8.40)$$

The same argument works for left ladder intervals, too:

$$\tilde{P} \left[\sum_{J \in \mathcal{I}_l} X_J^l \geq \frac{2^{n_0/3}}{4} \right] \leq \frac{1}{2} \exp\{-2^{n_0/4}\}. \quad (8.41)$$

Combining (8.41), (8.40), and $|\text{Corrupted}| \leq n_0^{41} \leq 2^{n_0/3}/2$ (see Lemma 8.16), we obtain

$$\begin{aligned} \tilde{P} \left[|\text{Center}_I \cap \text{Outside}_{x,t}| \geq 2^{n_0/3} \right] &\leq \tilde{P} \left[|\text{Center}_{II} \cap \text{Outside}_{x,t}| \geq \frac{2^{n_0/3}}{2} \right] \\ &\leq \tilde{P} \left[\sum_{I \in \mathcal{I}_r} X_I^r \geq \frac{2^{n_0/3}}{4} \right] + \tilde{P} \left[\sum_{J \in \mathcal{I}_l} X_J^l \geq \frac{2^{n_0/3}}{4} \right] \leq \exp\{-2^{n_0/4}\}. \end{aligned} \quad (8.42)$$

The claim (8.34) is an immediate consequence of this bound. ■

Due to the presence of the “modified” part $\xi[J_1]$, we define the following modification of $B_{\text{recogn straight}}^{n_0}$, which is a little weaker than the original version:

Definition 8.18 (Modification of Definition 6.18)

$$\begin{aligned} \tilde{B}_{\text{recogn straight}}^{n_0} &:= \quad (8.43) \\ &\left\{ \text{For every } R \in \text{AdPaths}(11 \cdot 2^{n_0}, c_1 n_0) \text{ with } R(c_1 n_0 - 1) - R(0) \notin \{(c_1 n_0 - 1)l_{\rightarrow}, (c_1 n_0 - 1)l_{\leftarrow}\} \text{ there is } \bar{R} \in \text{AdPaths}(12 \cdot 2^{n_0}, c_1 n_0) \text{ such that } R(0) = \bar{R}(0), R(c_1 n_0 - 1) = \bar{R}(c_1 n_0 - 1), \text{ and } \underline{\bar{R}} \text{ takes at least one value in } J_1, \text{ or } \xi \circ R \neq \xi \circ \bar{R}. \right\}, \end{aligned}$$

$$\tilde{E}_{\text{only ladder}}^{n_0} := \left\{ \text{For all } (w_1, w_2, w_3) \in \text{Puzzle}_1^{n_0}(\text{Input}) \text{ and every admissible piece of path } R : [0, 3c_1 n_0] \rightarrow [-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}] \text{ with } \xi \circ R = w_1 w_2 w_3 \text{ holds: } \right. \\ \left. w_2 \text{ is a ladder word of } \xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}], \text{ or } w_2 \in \underline{\text{Corrupted}}. \right\}. \quad (8.44)$$

Lemma 8.19 (Modification of Lemma 6.37) *There exist positive constants c_{25} and c_{26} not depending on n_0 such that:*

$$\tilde{P} \left[(\tilde{B}_{\text{recogn straight}}^{n_0})^c \right] \leq c_{25} e^{-c_{26} n_0}. \quad (8.45)$$

Proof. The proof of Lemma 6.37 requires only a small modification: Given $R \in \text{AdPaths}(11 \cdot 2^{n_0}, c_1 n_0)$, there are two cases: Either some $\bar{R} \in \text{AdPaths}(12 \cdot 2^{n_0}, c_1 n_0)$ with $\bar{R}(0) = R(0)$ and $\bar{R}(c_1 n_0 - 1) = R(c_1 n_0 - 1)$ touches J_1 (“case 1”), or no such \bar{R} touches J_1 (“case 2”).

- In the first case, the underlined new condition in definition (8.43) of $\tilde{B}_{\text{recogn straight}}^{n_0}$ is certainly satisfied.
- In the second case, we may proceed further just as in the proof of Lemma 6.37, with P replaced by \tilde{P} : formula (6.84) remains true in this case, since neither $R[I(R)]$ nor $\bar{R}[I(R)]$ touches J_1 . Recall that $\xi[(\mathbb{Z} \setminus J_1)]$ has the same distribution with respect to P as with \tilde{P} .

The rest of the proof of Lemma 6.37 still remains true when we replace P by \tilde{P} but remove all paths R from $\text{AdPaths}(11 \cdot 2^{n_0}, c_1 n_0)$ and $\text{AdPaths}(12 \cdot 2^{n_0}, c_1 n_0)$ that belong to the first case. ■

Lemma 8.20 (Modification of Lemma 6.19) *We have*

$$B_{\text{all paths}}^{n_0} \cap \tilde{B}_{\text{recogn straight}}^{n_0} \subseteq \tilde{E}_{\text{only ladder}}^{n_0}. \quad (8.46)$$

Proof. We describe the modifications required in the proof of Lemma 6.19: Assume the event $B_{\text{all paths}}^{n_0} \cap \tilde{B}_{\text{recogn straight}}^{n_0}$ holds, and let $w_1 w_2 w_3 \in \text{Puzzle}_I^{n_0}(\text{Input})$, and $R \in \text{AdPaths}(11 \cdot 2^{n_0}, 3c_1 n_0)$, $\xi \circ R = w_1 w_2 w_3$ as in the proof of Lemma 6.19. Again, we prove by contradiction that $\tilde{E}_{\text{only ladder}}^{n_0}$ holds: Assume that w_2 is not a ladder word of $\xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$ and $w_2 \notin \text{Corrupted}$. We distinguish two cases: Either the middle piece $R[[c_1 n_0, 2c_1 n_0[$ of R belongs to CorPaths when being time-shifted back to the origin (“case 1”), or it does not (“case 2”).

- In case 1, $w_2 = (\xi \circ R[[c_1 n_0, 2c_1 n_0])_{\rightarrow} \in \text{Corrupted}$ by Definition (8.28), which contradicts our assumption.
- In case 2, using Definition (8.27), there is an admissible piece of path $R' : [c_1 n_0, 2c_1 n_0[\rightarrow \mathbb{Z}$ with $R'(c_1 n_0) = R(c_1 n_0)$ and $R'(2c_1 n_0 - 1) = R(2c_1 n_0 - 1)$ such that $w'_2 := (\xi \circ R')_{\rightarrow} \neq (\xi \circ R[[c_1 n_0, 2c_1 n_0])_{\rightarrow}$ (“case 2.1”), or all admissible paths $R' : [c_1 n_0, 2c_1 n_0[\rightarrow \mathbb{Z}$ with $R'(c_1 n_0) = R(c_1 n_0)$ and $R'(2c_1 n_0 - 1) = R(2c_1 n_0 - 1)$ do not touch J_1 and fulfill $\xi \circ R' = \xi \circ R[[c_1 n_0, 2c_1 n_0[$ (“case 2.2”).
 - In case 2.1 we proceed just as in the proof of Lemma 6.19; this yields the contradiction $w_1 w_2 w_3 \notin \text{Puzzle}_I^{n_0}(\text{Input})$.
 - In case 2.2, we use that $R[[c_1 n_0, 2c_1 n_0[$ is not a ladder path, since w_2 is not a ladder word of $\xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$. Using Definition (8.43), this case is contradictory, too, since $\tilde{B}_{\text{recogn straight}}^{n_0}$ holds.

Thus all cases lead to a contradiction; this proves the Lemma. ■

Lemma 8.21 *If $\tilde{E}_{\text{only ladder}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds, then $\text{Puzzle}_{\text{IV}}^{n_0}(\chi) \subseteq \text{Center}_{\text{II}}$.*

Proof. Assume that $\tilde{E}_{\text{only ladder}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds, and let $w_2 \in \text{Puzzle}_{\text{IV}}^{n_0}(\chi)$. Take $w_1, w_3 \in \mathcal{C}^{c_1 n_0}$ with $(w_1, w_2, w_3) \in \text{Puzzle}_{\text{III}}^{n_0}(\chi)$ by (8.24). Then by (8.23), $(w_1, w_2, w_3) \in \text{Puzzle}_I^{n_0}(\text{Input})$, and $w_1 w_2 w_3$ occurs in the observations χ at most $2^{n_0}/l$ time steps after some stopping time $T_k^0(\chi)$, $0 \leq k < 2^{\alpha n_0}$. Since E_{stop, T^0}^0 holds, we have $|S(T_k^0)| \leq 2^{n_0}$; thus $w_1 w_2 w_3$ is read in χ while the random walk follows some admissible piece of path R with values in $[-2 \cdot 2^{n_0}, 2 \cdot 2^{n_0}] \subseteq [-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$. Since $\tilde{E}_{\text{only ladder}}^{n_0}$ holds, this implies: w_2 is a ladder word of $\xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$, or $w_2 \in \text{Corrupted}$. In the next argument, we use the following fact: If π is a ladder path and $\bar{\pi}$ is an admissible piece of path with the same length, starting point, and end point as π , then $\bar{\pi} = \pi$. Using this fact and the Definitions (8.27) and (8.28), we see: if w_2 is a ladder word of $\xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}]$, but not of $\xi[[-11 \cdot 2^{n_0}, 11 \cdot 2^{n_0}] \setminus J_1]$, then $w_2 \in \text{Corrupted}$, too. Thus we obtain $w_2 \in \text{Center}_{\text{II}}$. This proves the lemma. ■

Lemma 8.22 (Modification of Lemma 6.34) *There exist constants $c_{20}, c_{21} > 0$ such that:*

$$\tilde{P} \left[(B_{\text{all paths}}^{n_0})^c \cap E_{\text{stop}, T^0}^0 \right] \leq c_{21} e^{-c_{20} n_0}. \quad (8.47)$$

Proof. Again, the proof of Lemma 6.34 remains literally true when we replace P by \tilde{P} and τ by T^0 ; in particular note that the event $B_{\text{all paths}}^{n_0}$ depends only on the random walk S , but not on the scenery ξ , and the strong Markov property for S still holds with respect to \tilde{P} . ■

Recall Definition (7.1): $E_{\text{no error}, T^1}^1 = \{\forall k \in \mathbb{N} : \text{If } T_k^1(\chi) < 2^{12\alpha n_1}, \text{ then } |S(T_k^1(\chi))| \leq 2^{n_1}\}$.

Lemma 8.23 *For some constants $c_{48}, c_{49} > 0$ holds $\tilde{P} \left[E_{\text{no error}, T^1}^1 \right] \geq 1 - c_{48} e^{-c_{49} n_0}$.*

Proof. Using Definition 8.12 of T^1 and Lemmas 8.20, 8.21, and 8.17, we obtain (see also the explanations below):

$$\begin{aligned} & \tilde{P} \left[E_{\text{no error}, T^1}^1 \right] \quad (8.48) \\ & \geq \tilde{P} \left[\text{For all } t \in [0, 2^{12\alpha n_1}[\text{ holds: if } |\text{Puzzle}_{\text{IV}}^{n_0}(\chi) \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq 2^{n_0/3} \text{ and } \right. \\ & \quad \left. |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0}, \text{ then } |S(t + 2^{12\alpha n_0})| \leq 2^{n_1} \right] \\ & \geq \tilde{P}[\text{Puzzle}_{\text{IV}}^{n_0}(\chi) \subseteq \text{Center}_{\text{II}}] - \sum_{t=0}^{2^{12\alpha n_1}-1} \tilde{P} \left[\begin{array}{l} |\text{Center}_{\text{II}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq 2^{n_0/3}, \\ |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0}, \text{ and} \\ 2 \cdot 2^{n_0} + 2l2^{12\alpha n_0} < |S(t)| \leq l2^{12\alpha n_1} \end{array} \right] \\ & \geq \tilde{P}[B_{\text{all paths}}^{n_0} \cap \tilde{B}_{\text{recogn straight}}^{n_0} \cap E_{\text{stop}, T^0}^0] - 2^{12\alpha n_1} \cdot 2l2^{12\alpha n_1} \cdot \exp\{-2^{n_0/4}\} \\ & \geq \tilde{P}[E_{\text{stop}, T^0}^0] - \tilde{P}[(B_{\text{all paths}}^{n_0})^c \cap E_{\text{stop}, T^0}^0] - \tilde{P}[(\tilde{B}_{\text{recogn straight}}^{n_0})^c] - 2l \exp\{24(\log 2)\alpha n_1 - 2^{n_0/4}\} \\ & \geq 1 - c_{48} e^{-c_{49} n_0} \end{aligned}$$

for some constants $c_{48}, c_{49} > 0$. For the second inequality in (8.48), note that the random walk S cannot travel farther than $l2^{12\alpha n_1}$ within time $2^{12\alpha n_1}$; thus $|S(t + 2^{12\alpha n_0})| \leq 2^{n_1}$ or $2 \cdot 2^{n_0} + 2l2^{12\alpha n_0} < |S(t)| \leq l2^{12\alpha n_1}$ holds for all $t \in [0, 2^{12\alpha n_1}[$. In the last step of (8.48) we used Theorem 8.2, Lemmas 8.19 and 8.22, and the fact $n_1 = 2^{\lfloor \sqrt{n_0} \rfloor}$ (recall Definition 3.6); especially $\exp\{24(\log 2)\alpha n_1 - 2^{n_0/4}\}$ is superexponentially small in n_0 . The constants c_{48} and c_{49} need not depend on α , since n_0 was chosen large and α -dependent (see Subsection 2.1). ■

Definition 8.24 *We define the event*

$$B_{\text{all paths II}}^{n_0} := \left\{ \begin{array}{l} \forall R \in \text{AdPaths}(3 \cdot 2^{n_0/2}, 3c_1 n_0) \exists k \in [0, 2^{\alpha n_0}[\exists j \in [0, 2^{n_0}/l] : \\ \text{TimeShift}^{T_k^0(\chi)+j}(R) \subseteq S \end{array} \right\}. \quad (8.49)$$

Lemma 8.25 (Yet another modification of Lemma 6.34) *There exist constants $c_{50}, c_{51} > 0$ not depending on n_0 such that:*

$$\tilde{P} \left[(B_{\text{all paths II}}^{n_0})^c \cap E_{\text{stop}, T^0}^0 \right] \leq c_{50} e^{-c_{51} n_0}. \quad (8.50)$$

Proof. The proof is almost the same as the proof of Lemma 6.34; we only explain the differences. This time, some of the parameters in the proof of Lemma 6.34 must be changed: We replace (6.70) by $B_R^{n,k} := \left\{ \exists j \in [0, 2^{n_0}/l] : \text{TimeShift}^{\tau(k)+j}(R) \right\}$ and $\text{AdPath}(12 \cdot 2^n, 3c_1 n)$ by $\text{AdPaths}(3 \cdot$

$2^{n_0/2}, 3c_1n_0$); then the estimate (6.75) is replaced by $|\text{AdPaths}(3 \cdot 2^{n_0/2}, 3c_1n_0)| \leq 7 \cdot 2^{n_0/2} |\mathcal{M}|^{3c_1n_0}$. Then (6.78) changes to

$$\inf_{|x| \leq 3 \cdot 2^{n_0/2}} \tilde{P} \left[S(\tau(k) + j) = x \text{ for some } j \in [0, 2^{n_0}/l] \mid C_R^{n_0, k} \right] \geq c_{52} 2^{-n_0/2} \quad (8.51)$$

for some constant $c_{52} > 0$. Hence the right hand side of (6.79) gets replaced by $c_{52} 2^{-n_0/2} \mu_{\min}^{3c_1n_0}$. We end up with the following modified version of (6.80):

$$\tilde{P} \left[E_{\text{stop}, T^0}^0 \setminus B_{\text{all paths II}}^{n_0} \right] \leq 7 \exp \left\{ n_0 \left(\frac{\log 2}{2} + 3c_1 \log |\mathcal{M}| \right) - c_{52} e^{n_0(\alpha \log 2 + 3c_1 \log \mu_{\min} - (\log 2)/2)} \right\}, \quad (8.52)$$

which still converges superexponentially fast to 0 as $n_0 \rightarrow \infty$. This proves the Lemma. ■

Definition 8.26 (Modification of Definition 6.4)

$$\tilde{B}_{\text{signals}}^{n_0} := \left\{ \begin{array}{l} \text{For every right ladder path } \pi \in ([-2l2^{2n_0}, 2l2^{2n_0}] \setminus J_1)^{[0, c_1n_0/2[} \text{ and} \\ \text{for every admissible piece of path } \pi' \in \text{AdPath}(2l2^{2n_0}, c_1n_0/2): \\ \text{If } \xi \circ \pi = \xi \circ \pi', \text{ then } \pi(0) \leq \pi'(0) \text{ and } \pi(c_1n_0/2 - 1) \geq \pi'(c_1n_0/2 - 1). \end{array} \right\}, \quad (8.53)$$

$$\tilde{E}_{\text{signals II}}^{n_0} := \left\{ \begin{array}{l} \text{For every ladder path } \pi \in ([-2l2^{2n_0}, 2l2^{2n_0}] \setminus J_1)^{[0, c_1n_0[} \text{ and for} \\ \text{every admissible piece of path } \pi' \in \text{AdPath}(2l2^{2n_0}, c_1n_0): \\ \text{If } \xi \circ \pi = \xi \circ \pi', \text{ then } \pi(c_1n_0/2) = \pi'(c_1n_0/2). \end{array} \right\}. \quad (8.54)$$

Note that π' in the last two definitions may well have some of its values in J_1 .

Lemma 8.27 (Modification of Lemma 6.38) *There exist constants $c_{28} > 0, c_{29} > 0$ not depending on n_0 such that:*

$$\tilde{P} \left[(\tilde{B}_{\text{signals}}^{n_0})^c \right] \leq c_{29} e^{-c_{28}n_0}. \quad (8.55)$$

Proof. The proof of Lemma 6.38 remains literally true when we consider paths

$$\pi \in ([-2l2^{2n_0}, 2l2^{2n_0}] \setminus J_1)^{[0, c_1n_0[} \quad (8.56)$$

only, but replace P by \tilde{P} . Note that in the induction step in proof of Lemma 6.38 $\xi \circ \pi(j)$ is independent of the family $(\xi \circ \pi[I', \xi \circ \pi'[I')$ with respect to \tilde{P} , too, even if π' touches the “corrupted” domain J_1 ; see (6.90) and a few lines before this formula. This is true because π does not touch J_1 , and $\xi \upharpoonright J_1$ is independent of $\xi \upharpoonright (\mathbb{Z} \setminus J_1)$ by Lemma 3.4. Thus formula (6.90) remains true when P is replaced by \tilde{P} . ■

Lemma 8.28 (Modification of Lemma 6.5) $\tilde{B}_{\text{signals}}^{n_0} \subseteq \tilde{E}_{\text{signals II}}^{n_0}$.

Proof. When we consider paths π only which do not touch J_1 , the proof of Lemma 6.5 remains literally true in this modified case, too. ■

Lemma 8.29 (Modification of Lemma 6.7)

Assume that the event $B_{\text{all paths}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds. Let $I \subseteq [-6 \cdot 2^{n_0}, 6 \cdot 2^{n_0}] \setminus J_1$ be a right ladder interval with $|I| = 3c_1n_0$, and let $w_1, w_2, w_3 \in \mathcal{C}^{c_1n_0}$ with $(\xi \upharpoonright I)_{\rightarrow} = w_1 w_2 w_3$. Then $(w_1, w_2, w_3) \in \text{Puzzle}_1^{n_0}(\text{Input})$.

Proof. The proof of Lemma 6.7 remains literally true when we consider only intervals I that are disjoint from J_1 , $n = n_0$, and replace $B_{\text{signals}}^{n_0}$ and $E_{\text{signals II}}^{n_0}$ by their modified versions; note that the ladder path R in the proof of Lemma 6.7 does not touch J_1 , but we need not assume this for R' . ■

Lemma 8.30 *If the event $B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds, then $\text{Center}_{\text{III}} \subseteq \text{Puzzle}_{\text{IV}}^{n_0}(\chi)$.*

Proof. Assume that $B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds, and let $w_2 \in \text{Center}_{\text{III}}$. Then $w_2 = (\xi[I]_{\rightarrow}$ for some right ladder interval $I \subseteq [-2 \cdot 2^{n_0/2}, 2 \cdot 2^{n_0/2}] \setminus [-2^{n_0/2}, 2^{n_0/2}]$, $|I| = c_1 n_0$. We take the larger right ladder interval $I' \supseteq I$, $|I'| = 3c_1 n_0$, with $c_1 n_0$ extra points to the left of I and another $c_1 n_0$ extra points to the right of I ; then $I' \subseteq [-3 \cdot 2^{n_0/2}, 3 \cdot 2^{n_0/2}] \setminus J_1 \subseteq [-6 \cdot 2^{n_0/2}, 6 \cdot 2^{n_0/2}] \setminus J_1$; note that $\text{dist}(J_1, \mathbb{Z} \setminus [-2^{n_0/2}, 2^{n_0/2}]) > c_1 n_0 l$ and $\text{dist}([-2 \cdot 2^{n_0/2}, 2 \cdot 2^{n_0/2}], \mathbb{Z} \setminus [-3 \cdot 2^{n_0/2}, 3 \cdot 2^{n_0/2}]) > c_1 n_0 l$; recall that n_0 is chosen large enough (Subsection 2.1). Then $(\xi[I']_{\rightarrow} = w_1 w_2 w_3$ for some $w_1, w_3 \in \mathcal{C}^{c_1 n_0}$, and Lemma 8.29 implies $(w_1, w_2, w_3) \in \text{Puzzle}_{\text{I}}^{n_0}(\text{Input})$. Let R denote the (unique) right ladder path $R : [0, 3c_1]_{\rightarrow} I'$. Since $B_{\text{all paths II}}^{n_0}$ holds, the random walk S follows R (time-shifted) at most $2^{n_0}/l$ time steps after some stopping time $T_k^0(\chi)$, $k \in [0, 2^{\alpha n_0}[$. Then $\xi \circ R = w_1 w_2 w_3$; thus $(w_1, w_2, w_3) \in \text{Puzzle}_{\text{III}}^{n_0}(\chi)$ by Definition (8.23); hence $w_2 \in \text{Puzzle}_{\text{IV}}^{n_0}(\chi)$ by Definition (8.24). This proves the lemma. ■

Definition 8.31 *We set*

$$E_{\text{center}}^1 := \{ |\text{Center}_{\text{III}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\chi)| \geq 2^{n_0/3} \text{ and } |\text{Puzzle}_{\text{IV}}^{n_0}(\chi)| \leq 50 \cdot 2^{n_0} \}, \quad (8.57)$$

$$\Xi_{\text{center}}^1 := \left\{ \xi \in \mathcal{C}^{\mathbb{Z}} \mid P[E_{\text{center}}^1 \mid \xi] \geq \frac{1}{2} \right\}. \quad (8.58)$$

The sets E_{center}^1 and Ξ_{center}^1 play an analogous role for the stopping times T^1 as $E_{\text{reconst}, f}^m$ and $\Xi_{\text{reconst}, f}^m$ play for the “higher level” stopping times in Section 7.

Lemma 8.32 *For some positive constants c_{53} and c_{54} holds $\tilde{P}[\xi \in \Xi_{\text{center}}^1] \geq 1 - c_{53} e^{-c_{54} n_0}$.*

Proof. If the events $B_{\text{all paths}}^{n_0}$, $B_{\text{all paths II}}^{n_0}$, $\tilde{B}_{\text{signals}}^{n_0}$, E_{stop, T^0}^0 , $\tilde{E}_{\text{only ladder}}^{n_0}$, and $\tilde{B}_{\text{unique fit}}^{n_0}$ hold, then we have $|\text{Center}_{\text{III}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\chi)| = |\text{Center}_{\text{III}}| \geq 2^{n_0/3}$ and $|\text{Puzzle}_{\text{IV}}^{n_0}(\chi)| \leq |\text{Center}_{\text{II}}| \leq 50 \cdot 2^{n_0}$ by Lemmas 8.21, 8.30, and 8.16. By Lemma 8.20, we can replace $\tilde{E}_{\text{only ladder}}^{n_0}$ in the above list of events by $\tilde{B}_{\text{recogn straight}}^{n_0}$. Thus we have

$$\begin{aligned} \tilde{P}[E_{\text{center}}^1] &\geq \tilde{P} \left[B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0 \cap \tilde{B}_{\text{recogn straight}}^{n_0} \cap \tilde{B}_{\text{unique fit}}^{n_0} \right] \\ &\geq 1 - c_{55} e^{-c_{54} n_0} \end{aligned} \quad (8.59)$$

for some positive constants c_{55} and c_{54} by Theorem 8.2 and Lemmas 8.22, 8.25, 8.19, 8.27, and 8.15.

Hence we obtain the following:

$$\frac{1}{2} \tilde{P}[\xi \notin \Xi_{\text{center}}^1] = \frac{1}{2} \tilde{P} \left[P[(E_{\text{center}}^1)^c \mid \xi] > \frac{1}{2} \right] \leq \tilde{P}[(E_{\text{center}}^1)^c] \leq c_{55} e^{-c_{54} n_0}; \quad (8.60)$$

recall that $\tilde{P}[\cdot \mid \xi]$ and $P[\cdot \mid \xi]$ coincide. ■

Definition 8.33 (Yet another modification of Definition 7.3) Let $v(k)$ denote again the $(k+1)$ st visit of S to the origin. We define

$$\mathbb{T}^{1'}(\xi, \chi) := \left\{ t \in \mathbb{N} \mid \begin{array}{l} |\text{Center}_{\text{III}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq 2^{n_0/3} \\ \text{and } |\text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \leq 50 \cdot 2^{n_0} \end{array} \right\}, \quad (8.61)$$

$$E_{\text{when back recog}}^1 := \left\{ \begin{array}{l} \text{For more than } 1/4 \text{ of the points } k \in [0, 2^{2\alpha n_1}[\\ \text{holds } v(k2^{\alpha n_1}) \in \mathbb{T}^{1'}(\xi, \chi) \end{array} \right\}. \quad (8.62)$$

Lemma 8.34 (Yet another analogue to Lemma 7.4) If the event $B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0$ holds, then $\mathbb{T}^1(\chi) \supseteq \mathbb{T}^{1'}(\xi, \chi) \cap [0, 2^{12\alpha n_1} - 2 \cdot 2^{12\alpha n_0}[$.

Proof. Assuming that the event $B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}}^0$ holds, we know $\text{Center}_{\text{III}} \subseteq \text{Puzzle}_{\text{IV}}^{n_0}(\chi)$ by Lemma 8.30; thus $|\text{Puzzle}_{\text{IV}}^{n_0}(\chi) \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)| \geq |\text{Center}_{\text{III}} \cap \text{Puzzle}_{\text{IV}}^{n_0}(\theta^t \chi)|$ for all t . This implies the claim $\mathbb{T}^1(\chi) \supseteq \mathbb{T}^{1'}(\xi, \chi) \cap [0, 2^{12\alpha n_1} - 2 \cdot 2^{12\alpha n_0}[$ of the lemma; recall Definition (8.25) of $\mathbb{T}^1(\chi)$. ■

Lemma 8.35 (Yet another modification of Lemma 7.5) Assume that the events $E_{\text{no error}, T^1}^1 \cap E_{\text{enough back}}^1 \cap E_{\text{when back recog}}^1$ and $\mathbb{T}^1(\chi) \supseteq \mathbb{T}^{1'}(\xi, \chi) \cap [0, 2^{12\alpha n_1} - 2 \cdot 2^{12\alpha n_0}[$ hold. Then E_{stop, T^1}^1 holds, too.

Proof. Replacing $\mathbb{T}_f^{(\prime)}$ by $\mathbb{T}^{1(\prime)}$ and T_f by T^1 , the proof of Lemma 7.5 remains literally true. ■

Lemma 8.36 (Yet another modification of Lemma 7.6) We have the bound

$$\tilde{P} \left[(E_{\text{when back recog}}^1)^c \cap \{ \xi \in \Xi_{\text{center}}^1 \} \right] \leq 0.9^{2^{2\alpha n_1}}. \quad (8.63)$$

Proof. The proof of Lemma 7.6 remains literally true when one replaces $E_{\text{when back recog}, f}^{m+1}$ by $E_{\text{when back recog}}^1$, $\Xi_{\text{reconst}, f}^m$ by Ξ_{center}^1 , $m+1$ by 1, and P by \tilde{P} ; recall $P[\cdot|\xi] = \tilde{P}[\cdot|\xi]$. ■

Proof of Theorem 3.10. By Lemmas 8.34 and 8.35 we know

$$E_{\text{no error}, T^1}^1 \cap E_{\text{enough back}}^1 \cap E_{\text{when back recog}}^1 \cap B_{\text{all paths}}^{n_0} \cap B_{\text{all paths II}}^{n_0} \cap \tilde{B}_{\text{signals}}^{n_0} \cap E_{\text{stop}, T^0}^0 \subseteq E_{\text{stop}, T^1}^1. \quad (8.64)$$

Since $E_{\text{enough back}}^1$ depends only on S but not on ξ , we have $\tilde{P}[(E_{\text{enough back}}^1)^c] = P[(E_{\text{enough back}}^1)^c]$. Thus, using Lemmas 8.23, 7.2, 8.36, 8.32, 8.22, 8.25, 8.27, and Theorem 8.2, we know

$$\begin{aligned} & \tilde{P}[(E_{\text{stop}, T^1}^1)^c] \\ & \leq \tilde{P}[(E_{\text{no error}, T^1}^1)^c] + \tilde{P}[(E_{\text{enough back}}^1)^c] + \tilde{P} \left[(E_{\text{when back recog}}^1)^c \cap \{ \xi \in \Xi_{\text{center}}^1 \} \right] \\ & \quad + \tilde{P} \left[\xi \notin \Xi_{\text{center}}^1 \right] + \tilde{P}[(B_{\text{all paths}}^{n_0})^c \cap E_{\text{stop}, T^0}^0] + \tilde{P}[(B_{\text{all paths II}}^{n_0})^c \cap E_{\text{stop}, T^0}^0] \\ & \quad + \tilde{P}[(\tilde{B}_{\text{signals}}^{n_0})^c] + \tilde{P}[(E_{\text{stop}}^0)^c] \\ & \leq c_{48}e^{-c_{49}n_0} + c_{35}2^{-\alpha n_1} + 0.9^{2^{2\alpha n_1}} + c_{53}e^{-c_{54}n_0} + c_{21}e^{-c_{20}n_0} \\ & \quad + c_{50}e^{-c_{51}n_0} + c_{29}e^{-c_{28}n_0} + c_{36}e^{-c_{37}n_0} \\ & \leq e^{-c_4 n_0}, \end{aligned} \quad (8.65)$$

since n_0 is chosen large enough (see Subsection 2.1) ■

A Appendix: The first return time to the origin

Lemma A.1 *Let X_1 be the first return time of the random walk S to the origin. Then $E[X_1^\beta] < \infty$ holds whenever $0 < \beta < \frac{1}{2}$.*

Proof. Let $k := l_\rightarrow + l_\leftarrow$, and note that $P[S_k = 0] > 0$; e.g. one may perform l_\leftarrow steps of size l_\rightarrow and then l_\rightarrow steps of size $-l_\leftarrow$. As a consequence of the local central limit theorem, there is a constant $c_{56} > 0$ such that $P[S_n = 0] \geq c_{56}n^{-1/2}$ holds for all $n \in k\mathbb{N}$, $n > 0$. (By choosing c_{56} small enough, this even holds for small values $n \in k\mathbb{N} \setminus \{0\}$.)

For $\lambda > 0$, let

$$L[\lambda] := \sum_{n=0}^{\infty} e^{-\lambda n} P[S_n = 0]. \quad (\text{A.1})$$

Then we have for $0 < \lambda \leq 1/k$ and $c_{57} := \frac{c_{56}}{e\sqrt{2k}}$:

$$L[\lambda] \geq \sum_{m=1}^{\lfloor (\lambda k)^{-1} \rfloor} e^{-\lambda mk} P[S_{mk} = 0] \geq \lfloor (\lambda k)^{-1} \rfloor \cdot e^{-1} \cdot c_{56} \lfloor (\lambda k)^{-1} \rfloor^{-1/2} \geq \frac{c_{57}}{\sqrt{\lambda}}. \quad (\text{A.2})$$

Let X_j denote the $(j+1)$ st visit of the random walk S to the origin; this is consistent with the above definition of X_1 . Then

$$L[\lambda] = \sum_{j=0}^{\infty} E[e^{-\lambda X_j}] = \sum_{j=0}^{\infty} E[e^{-\lambda X_1}]^j = \frac{1}{1 - E[e^{-\lambda X_1}]}. \quad (\text{A.3})$$

Using (A.2), this implies for $0 < \lambda \leq 1/k$:

$$1 - E[e^{-\lambda X_1}] = \frac{1}{L[\lambda]} \leq c_{57}^{-1} \sqrt{\lambda}. \quad (\text{A.4})$$

We observe the elementary fact

$$\int_0^{\infty} (1 - e^{-\lambda x}) \lambda^{-1-\beta} d\lambda = -\Gamma(-\beta) x^\beta, \quad (\text{A.5})$$

which holds since $\beta < 1$; here Γ denotes the Gamma-function. Note that $-\Gamma(-\beta) > 0$, since $0 < \beta < 1$.

By Fubini's theorem, this implies

$$\begin{aligned} E[X_1^\beta] &= \frac{-1}{\Gamma(-\beta)} \int_0^{\infty} (1 - E[e^{-\lambda X_1}]) \lambda^{-1-\beta} d\lambda \\ &\leq \frac{-1}{\Gamma(-\beta)} \left(\int_0^{1/k} (1 - E[e^{-\lambda X_1}]) \lambda^{-1-\beta} d\lambda + \int_{1/k}^{\infty} \lambda^{-1-\beta} d\lambda \right). \end{aligned} \quad (\text{A.6})$$

The second integral in the last bracket is finite, since $\beta > 0$. The first integral in the bracket is estimated as follows, using (A.4):

$$\int_0^{1/k} (1 - E[e^{-\lambda X_1}]) \lambda^{-1-\beta} d\lambda \leq c_{57}^{-1} \int_0^{1/k} \lambda^{-\frac{1}{2}-\beta} d\lambda < \infty; \quad (\text{A.7})$$

the last integral is finite since $\beta < \frac{1}{2}$. This proves the lemma. ■

References

- [1] Itai Benjamini and Harry Kesten. Distinguishing sceneries by observing the scenery along a random walk path. *J. Anal. Math.*, 69:97–135, 1996.
- [2] Frank den Hollander and Jeffrey E. Steif. Mixing properties of the generalized T, T^{-1} -process. *J. Anal. Math.*, 72:165–202, 1997.
- [3] W. Th. F. den Hollander. Mixing properties for random walk in random scenery. *Ann. Probab.*, 16(4):1788–1802, 1988.
- [4] Richard Durrett. *Probability: theory and examples*. Duxbury Press, Belmont, CA, second edition, 1996.
- [5] C. Douglas Howard. Detecting defects in periodic scenery by random walks on \mathbb{Z} . *Random Structures Algorithms*, 8(1):59–74, 1996.
- [6] C. Douglas Howard. Orthogonality of measures induced by random walks with scenery. *Combin. Probab. Comput.*, 5(3):247–256, 1996.
- [7] C. Douglas Howard. Distinguishing certain random sceneries on \mathbb{Z} via random walks. *Statist. Probab. Lett.*, 34(2):123–132, 1997.
- [8] Steven Arthur Kalikow. T, T^{-1} transformation is not loosely Bernoulli. *Ann. of Math. (2)*, 115(2):393–409, 1982.
- [9] M. Keane and W. Th. F. den Hollander. Ergodic properties of color records. *Phys. A*, 138(1-2):183–193, 1986.
- [10] Harry Kesten. Detecting a single defect in a scenery by observing the scenery along a random walk path. In *Itô's stochastic calculus and probability theory*, pages 171–183. Springer, Tokyo, 1996.
- [11] Harry Kesten. Distinguishing and reconstructing sceneries from observations along random walk paths. In *Microsurveys in discrete probability (Princeton, NJ, 1997)*, pages 75–83. Amer. Math. Soc., Providence, RI, 1998.
- [12] A. Lenstra and H. Matzinger. Reconstructing a 4-color scenery by observing it along a recurrent random walk path with unbounded jumps. In preparation, Eurandom, 2001.
- [13] Elon Lindenstrauss. Indistinguishable sceneries. *Random Structures Algorithms*, 14(1):71–86, 1999.
- [14] M. Löwe and H. Matzinger. Reconstruction of sceneries with correlated colors. Eurandom Report 99-032, Eurandom, 1999. Submitted to *Stochastic Processes and their Applications*.
- [15] M. Löwe and H. Matzinger. Scenery reconstruction in two dimensions with many colors. Eurandom Report 99-018, Eurandom, 1999. Submitted to *The Annals of Applied Probability*.
- [16] H. Matzinger. *Reconstructing a 2-color scenery by observing it along a simple random walk path with holding*. PhD-thesis, Cornell University, 1999.

- [17] H. Matzinger. Reconstructing a 2-color scenery by observing it along a simple random walk path. Eurandom Report 2000-003, Eurandom, 2000. Submitted to *The Annals of Applied Probability*.
- [18] H. Matzinger. Reconstructing a 2-color scenery in polynomial time by observing it along a simple random walk path with holding. Eurandom Report 2000-002, Eurandom, 2000. Submitted to *Probability Theory and Related Fields*.
- [19] Heinrich Matzinger. Reconstructing a three-color scenery by observing it along a simple random walk path. *Random Structures Algorithms*, 15(2):196–207, 1999.