

## Uncertainty in exposure to air pollution

**Citation for published version (APA):**

Pebesma, E., Helle, K. B., Stasch, C., Rasouli, S., Timmermans, H. J. P., Walker, S., & Denby, B. (2013). Uncertainty in exposure to air pollution. In *Proceedings of the General Assembly of the European Geosciences Union (EGU 2013), 7-12 April 2013, Vienna, Austria* (Geophysical research abstracts; Vol. 15).

**Document status and date:**

Published: 01/01/2013

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.



## Uncertainty in exposure to air pollution

Edzer Pebesma (1), Kristina Helle (1), Stasch Christoph (1), Soora Rasouli (2), Harry Timmermans (2), Sam-Erik Walker (3), and Bruce Denby (3)

(1) University of Münster, Institute for Geoinformatics, Germany (edzer.pebesma@uni-muenster.de), (2) Eindhoven University of Technology (TUE), The Netherlands, (3) Norwegian Institute for Air Research (NILU), Norway

To assess exposure to air pollution for a person or for a group of people, one needs to know where the person or group is as a function of time, and what the air pollution is at these times and locations. In this study we used the Albatross activity-based model to assess the whereabouts of people and the uncertainties in this, and a probabilistic air quality system based on TAPM/EPISODE to assess air quality probabilistically. The outcomes of the two models were combined to assess exposure to air pollution, and the errors in it. We used the area around Rotterdam (Netherlands) as a case study.

As the outcomes of both models come as Monte Carlo realizations, it was relatively easy to cancel one of the sources of uncertainty (movement of persons, air pollution) in order to identify their respective contributions, and also to compare evaluations for individuals with averages for a population of persons. As the output is probabilistic, and in addition spatially and temporally varying, the visual analysis of the complete results poses some challenges. This case study was one of the test cases in the UncertWeb project, which has built concepts and tools to realize the uncertainty-enabled model web. Some of the tools and protocols will be shown and evaluated in this presentation.

For the uncertainty of exposure, the uncertainty of air quality was more important than the uncertainty of peoples locations. This difference was stronger for PM10 than for NO<sub>2</sub>.

The workflow was implemented as generic Web services in UncertWeb that also allow for other inputs than the simulated activity schedules and air quality with other resolution. However, due to this flexibility, the Web services require standardized formats and the overlay algorithm is not optimized for the specific use case resulting in a data and processing overhead. Hence, we implemented the full analysis in parallel in R, for this specific case as the model web solution had difficulties with massive data.