

High-frequency AC breakdown in near-atmospheric pressure noble gasses

Citation for published version (APA):

Sobota, A., Kanters, J. H. M., Gendre, M. F., Manders, F., Dijk, van, J., Veldhuizen, van, E. M., & Haverlag, M. (2011). High-frequency AC breakdown in near-atmospheric pressure noble gasses. In *Proceedings of the 64th Annual Gaseous Electronics Conference 2011, 14-18 November 2011, Salk Lake City, USA* (pp. 3.00002-).

Document status and date:

Published: 01/01/2011

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Abstract Submitted
for the GEC11 Meeting of
The American Physical Society

Sorting Category: 2.12 (E)

High-frequency AC breakdown in near-atmospheric pressure noble gasses ANA SOBOTA, J.H.M. KANTERS, Eindhoven University of Technology, M.F. GENDRE, F. MANDERS, Philips Lighting, J. VAN DIJK, E.M. VAN VELDHUIZEN, Eindhoven University of Technology, M. HAVERLAG, Philips Lighting — AC breakdown has been extensively studied in two frequency ranges. At the low-frequency end breakdown happens at least once in every half voltage cycle. At the high-frequency end electron dynamics cannot follow the changes in the imposed electric field. The transitional range is not well known. Here we examine AC breakdown process in argon and xenon at near-atmospheric pressure in this transitional range. We used enclosed pin-pin geometry, with electrode gap length of 4 or 7 mm. The driving frequency was varied between 60 kHz and 1 MHz. Both experiments and numerical simulations were performed. We explain the observed timing and appearance of the discharge, the fact that the minimum breakdown voltage in this frequency range is significantly lower than in pulsed, DC or low-frequency AC discharges and the decreasing trend of minimum breakdown voltage with the increase of the driving frequency. The analysis of the experimental and numerical results show that this is a type of breakdown process that marks the transition from pulse-like and low-frequency AC, which features streamers and single polarity discharges, to a regime where ions start playing an important role in the process and can to a large degree determine the characteristics of the discharge.

Prefer Oral Session
 Prefer Poster Session

Ana Sobota
a.sobota@tue.nl
Eindhoven University of Technology

Date submitted: 31 May 2011

Electronic form version 1.4