

The effects of sustained inhomogeneous deformation on muscle cells

Citation for published version (APA):

Peeters, E. A. G., Bouten, C. V. C., Oomens, C. W. J., & Baaijens, F. P. T. (1999). *The effects of sustained inhomogeneous deformation on muscle cells*. Poster session presented at Mate Poster Award 1999 : 4th Annual Poster Contest.

Document status and date:

Published: 01/01/1999

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

The effects of sustained inhomogeneous deformation on muscle cells



E.A.G. Peeters, C.V.C. Bouten, C.W. Oomens,
F.P.T. Baaijens

Eindhoven University of Technology,
Faculty of Mechanical Engineering,
Section Materials Technology,
P.O. Box 513, 5600 MB Eindhoven, the Netherlands



Introduction

This project is embedded within the existing research line on soft tissue damage and adaptation, more specifically the project *Aetiology of Pressure sores*. Pressure sores are defined as localised areas of tissue degeneration in skin and underlying tissue resulting from prolonged mechanical load. It affects in particular disabled and debilitated patients and its occurrence is likely to increase due to the higher average age of the population [1].




fig 1. Areas which are most susceptible to pressure sores development

Objective

It is aimed to investigate the relationship between the degree of cell deformation, cell defence response and cell damage. The ultimate goal is to identify an objective marker for cell damage.

Methods

Numerical

A model of a single muscle cell under compression will be developed. It should incorporate the passive mechanical properties as well as metabolic phenomena and signaling cascades which relate the cellular deformation to cellular damage




fig 2. Compressive loading device

Experimental

In-vitro experiments on cultured muscle cells will be performed in order to determine:

- mechanical properties → input for the numerical model
- (patho)physiological effects.

A loading device capable of compressing populations of cells embedded in a gel-like matrix is being developed (figure 1).

Results

In preliminary experiments myoblasts (muscle precursor cells) were seeded into agarose gel and subsequently compressed up to a compressive strain of 20%. No statistically significant area and volume changes were observed. The cell deformation index (ratio between X and Y diameter) appeared to reduce from a mean of 0.98 to 0.66 [2].




fig 3. Deformation of myoblasts

Discussion

Until now, only general information about the deformation of compressed muscle cells is available. Future work should reveal what specific function the load bearing elements of the cell (the cell membrane and the cytoskeleton) and certain signaling molecules (proteins and ions) have in the complex cascades of mechano-transduction.

References:

- [1] BADER, D.L.: *Pressure sores: clinical practice and scientific approach* (London, MacMillan Scientific & Medical, 1992)
- [2] PEETERS, E.A.G.: *Deformation of myoblasts, mounted in agarose constructs* (Eindhoven, MT 98.007, 1998)