

Constituents of *Alpinia Breviligulata* Gagnep. from Vietnam

Citation for published version (APA):

Dung, N. X., Chinh, T. D., Rang, D. D., & Leclercq, P. A. (1994). Constituents of *Alpinia Breviligulata* Gagnep. from Vietnam. In *Abstracts of Eight Asian Symposium on Medicinal Plants, species and other natural products (ASOMPS VIII), 12-16 June 1994, Melaka, Malaysia* (pp. CP3A-)

Document status and date:

Published: 01/01/1994

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

CONSTITUENTS OF *ALPINIA BREVILIGULATA* GAGNEP. FROM VIETNAM

Nguyễn XUÂN DŨNG*, Trịnh ĐÌNH CHÍNH, Đỗ ĐÌNH RĂNG and Piet A. LECLERCQ*

Centre for Education and Development of Chromatography of Vietnam (EDC Vietnam),
3 Giải Phóng Street, Hai Bà District, 10 000 Hanoi, Vietnam. Fax : 84 4 291256

The Essential oils of the Root, Rhizome, Stem, Leaf, Flower, Fruit-peel and Seed of *Alpinia breviligulata* Gagnep. from Vietnam were analyzed by a combination of capillary GC, GC/MS, IR and NMR and found to contain :

* More than thirty components in the Root oil, in which 1,8- cineole (6.0%), borneol (5.5%), fenchyl acetate (8.8%), β -caryophyllene (7.5%), α -humulene (10.8%) and caryophyllene oxide (13.0%) are main compounds [1].

* More than thirty compounds in the Rhizome oil. The major components were β -pinene (10.0%), 1,8-cineole (5.9%), bornyl acetate (7.0%), β -caryophyllene (8.0%), α -humulene (7.9%) and caryophyllene oxide (10.5%) [1].

* More than forty components in the Stem and Leaf oils of which the major ones were found to be α -pinene (14.3%), caryophyllene oxide (27.5%), and γ -eudesmol (6.2%) in the Stem oil and α -pinene (17.7%) and caryophyllene oxide (23.1%) in the Leaf oil [2].

* More than thirty five components in the Flower oil of which α -pinene (12.7%), β -pinene (20.2%) and β -caryophyllene (14.0%) were found to be the main compounds. The components which provide the characteristic aroma are verbenone, bornyl acetate, hedycarol, (E)-nerolidol, guaiol and β -eudesmol [3].

* More than thirty components in the Fruit-peel oil. The major constituents were found to be β -pinene (21.2%), α -terpineol (7.0%), β -caryophyllene (5.7%), α -humulene (5.5%), and caryophyllene oxide (8.4%) [4].

* More than twenty compounds in the Seed oil of which the major ones were found to be geranyl acetate (8.8%), α -humulene (6.1%) and (E,E)-farnesol (65.0%).

These compounds and another components such as (Z)-nerolidol, (Z, E)-farnesol, (E,E)-farnesol, (E,E)-farnenyl acetate, citronellyl acetate, geraniol, octyl acetate, α -terpineol, linalool, hexyl acetate and other acetates are thought to be responsible for the pleasant floral odor of this oil [4]. This result shows that the Seed oil of *Alpinia breviligulata* Gagnep. is a valuable source for (E,E)-farnesol. The application of the *Alpinia breviligulata* Gagnep. in traditional medicine is also presented.

Nguyễn Xuân Dũng, Trịnh Đình Chính, Đỗ Đình Răng and Piet A. Leclercq (1993)

1) JEOR (accepted); 2) JEOR (accepted); 3) JEOR 5(5) 575-576; 4) JEOR (accepted)