

The influence of curvature on extracellular matrix components in 3D tissue engineered bone constructs cultured under static and perfused conditions

Citation for published version (APA):

Vetsch, J. R., Müller, R., & Hofmann, S. (2016). The influence of curvature on extracellular matrix components in 3D tissue engineered bone constructs cultured under static and perfused conditions. In *Towards Future Regenerative Therapies (TERMIS-EU Conference)* (pp. 166). (European Cells and Materials; Vol. 31, No. Supplement 1).

Document status and date:

Published: 01/01/2016

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

The influence of curvature on extracellular matrix components in 3D tissue engineered bone constructs cultured under static and perfused conditions

JR Vetsch¹, R Müller¹, S Hofmann^{1,2,3}

¹ [Institute for Biomechanics, ETH Zurich, Zurich, CH.](#) ² [Department of Biomedical Engineering and](#)
³ [Institute for Complex Molecular Systems, TU Eindhoven, Eindhoven, NL.](#)

INTRODUCTION: The adult skeleton constantly adapts its structure to changes in long-term mechanical loading. This continuous process of bone resorption and formation is called bone-remodelling [1]. The refilling of resorption sites *in vivo* has been shown to be most probably curvature driven [2]. *In vitro*, essential work on the influence of curvature on non-mineralized tissue in static cultures has been performed. Local curvature increased tissue growth formed by osteoblasts [3]. Cytoskeletal changes have been observed when comparing endothelial cells cultured on curved or flat surfaces. The changes were additionally enhanced when cells were exposed to perfusion [4]. The effects of curvature and perfusion on mineralized extracellular matrix (ECM) formation have, however, never been observed in 3D. Hence, the aim of this study was to investigate the effects of curvature on different ECM components in a 3D tissue engineered bone construct under static or perfused conditions.

METHODS: Porous silk fibroin scaffolds were produced by introducing three channels of different sizes using biopsy punches. The relative curvatures of the three channels were set to -50% in curvature (S channel, d=1mm) or +50% in curvature (L channel, d=3mm) with respect to the M channel (d=1.5mm). Scaffolds were seeded with 5×10^6 human mesenchymal stem cells each, supplied with osteogenic medium and cultured in either static or perfused bioreactors (N=10 per group). Micro-computed tomography (μ CT) was performed weekly to monitor mineralized ECM formation. Mineralized ECM was evaluated in three different volumes: (i) the 'full scaffold volume', (ii) the 'void volume' of the channel (90% of channel diameter), and (iii) the 'full volume' of the channel (channel diameter plus 1mm). Histology was performed to visualize cell nuclei and non-mineralized ECM stained with Haematoxylin&Eosin (H&E) and collagen stained with Sirius Red (SR).

RESULTS: μ CT monitoring showed curvature dependent ingrowth of mineralized ECM into the 'void volume' of the channels (Fig. 1). The coverage of the channels increased with curvature

and was additionally increased by perfusion (Fig. 1A-C, right). The morphology of mineralized ECM was more cortical-like in static samples (Fig. 1A-C, left) compared to the more trabecular-like structure in perfused samples (Fig. 1A-C, right).




Fig. 1: 3D mineralized ECM in (A) L, (B) M, and (C) S channels (top view). A-C left: static samples. A-C right: perfused samples. Scale bar: 1mm.




Fig. 2: H&E staining of: (A-C) static and (D-F) perfused samples. Scale bar: 1mm.

Histology images confirmed the results observed by μ CT. H&E staining showed less coverage of channels with cells and non-mineralized ECM when cultured under static (Fig. 2A-C) compared to perfused (Fig. 2D-F) conditions. SR staining showed the same patterns for collagen.

DISCUSSION & CONCLUSIONS: This study showed a clear dependence of different ECM components like mineralization or collagen on local curvature. Perfusion showed to enhance the effects of curvature. Additionally, the results observed suggest a possible application of the scaffold model to investigate circular critical size defects in trabecular or cortical bone *in vitro*.

REFERENCES: ¹S.C. Cowin & D.H. Hegedus (1976) *J Elast* **6**:313-26. ²H. Jinnai et al (2002) *Bone* **30**:191-4. ³M. Rumpler et al (2008) *J R Soc Interface* **5**:1173-80. ⁴M.D. Frame et al (2000) *Microcirculation* **7**:419-27.

ACKNOWLEDGEMENTS: This study has been funded by the EU Seventh Framework Programme (FP/2007-2013); Grant Agreement n. 262948.