

## Clustering spectrum of scale-free networks

**Citation for published version (APA):**

Stegehuis, C., van der Hofstad, R. W., Janssen, A. J. E. M., & van Leeuwen, J. S. H. (2017). Clustering spectrum of scale-free networks. *Physical Review E*, 96(4), Article 042309.  
<https://doi.org/10.1103/PhysRevE.96.042309>

**DOI:**

[10.1103/PhysRevE.96.042309](https://doi.org/10.1103/PhysRevE.96.042309)

**Document status and date:**

Published: 26/10/2017

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

## Clustering spectrum of scale-free networks

Clara Stegehuis, Remco van der Hofstad, A. J. E. M. Janssen, and Johan S. H. van Leeuwen

*Department of Mathematics and Computer Science, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands*

(Received 13 June 2017; published 26 October 2017)

Real-world networks often have power-law degrees and scale-free properties, such as ultrasmall distances and ultrafast information spreading. In this paper, we study a third universal property: three-point correlations that suppress the creation of triangles and signal the presence of hierarchy. We quantify this property in terms of  $\bar{c}(k)$ , the probability that two neighbors of a degree- $k$  node are neighbors themselves. We investigate how the clustering spectrum  $k \mapsto \bar{c}(k)$  scales with  $k$  in the hidden-variable model and show that  $\bar{c}(k)$  follows a *universal curve* that consists of three  $k$  ranges where  $\bar{c}(k)$  remains flat, starts declining, and eventually settles on a power-law  $\bar{c}(k) \sim k^{-\alpha}$  with  $\alpha$  depending on the power law of the degree distribution. We test these results against ten contemporary real-world networks and explain analytically why the universal curve properties only reveal themselves in large networks.

DOI: [10.1103/PhysRevE.96.042309](https://doi.org/10.1103/PhysRevE.96.042309)

### I. INTRODUCTION

Most real-world networks have power-law degrees so that the proportion of nodes having  $k$  neighbors scales as  $k^{-\tau}$  with exponent  $\tau$  between 2 and 3 [1–4]. Power-law degrees imply various intriguing scale-free network properties, such as ultrashort distances [5,6] and the absence of percolation thresholds when  $\tau < 3$  [7,8]. Empirical evidence has been matched by random graph null models that are able to explain mathematically why and how these properties arise. This paper deals with another fundamental property observed in many scale-free networks related to three-point correlations that suppress the creation of triangles and signal the presence of hierarchy. We quantify this property in terms of the *clustering spectrum*, the function  $k \mapsto \bar{c}(k)$  with  $\bar{c}(k)$  as the probability that two neighbors of a degree- $k$  node are neighbors themselves.

In *uncorrelated* networks the clustering spectrum  $\bar{c}(k)$  remains constant and independent of  $k$ . However, the majority of real-world networks have spectra that decay in  $k$  as first observed in technological networks including the Internet [9,10]. Figure 1 shows the same phenomenon for a social network: YouTube users as vertices and edges indicating friendships between them [11].

Close inspection suggests the following properties not only in Fig. 1, but also in the nine further networks in Fig. 10 in Appendix E. The right end of the spectrum appears to be of the power-law form  $k^{-\alpha}$ ; approximate values of  $\alpha$  give rise to the dashed lines; (ii) the power law is only approximate and kicks in for rather large values of  $k$ . In fact, the slope of  $\bar{c}(k)$  decreases with  $k$ ; (iii) there exists a transition point: the minimal degree as of which the slope starts to decline faster and settles on its limiting (large  $k$ ) value.

For scale-free networks a decaying  $\bar{c}(k)$  is taken as an indicator for the presence of modularity and hierarchy [10], architectures that can be viewed as collections of subgraphs with dense connections within themselves and sparser ones between them. The existence of clusters of dense interaction signals hierarchical or nearly decomposable structures. When the function  $\bar{c}(k)$  falls off with  $k$ , low-degree vertices have relatively high clustering coefficients, hence, creating small modules that are connected through triangles. In contrast, high-degree vertices have very low clustering coefficients

and therefore act as bridges between the different local modules. This also explains why  $\bar{c}(k)$  is not just a local property and, when viewed as a function of  $k$ , measures crucial mesoscopic network properties, such as modularity, clusters, and communities. The behavior of  $\bar{c}(k)$  also turns out to be a good predictor for the macroscopic behavior of the network. Randomizing real-world networks while preserving the shape of the  $\bar{c}(k)$  curve produces networks with very similar component sizes as well as similar hierarchical structures as the original network [16]. Furthermore, the shape of  $\bar{c}(k)$  strongly influences the behavior of networks under percolation [17]. This places the  $\bar{c}(k)$  curve among the most relevant indicators for structural correlations in network infrastructures.

In this paper, we obtain a precise characterization of clustering in the hidden-variable model, a tractable random graph null model. We start from an explicit form of the  $\bar{c}(k)$  curve for the hidden-variable model [18–20]. We obtain a detailed description of the  $\bar{c}(k)$  curve in the large-network limit that provides rigorous underpinning of the empirical observations (i)–(iii). We find that the decay rate in the hidden-variable model is significantly different from the exponent  $\bar{c}(k) \sim k^{-1}$  that has been found in a hierarchical graph model [10] as well as in the preferential attachment model [21] and a preferential attachment model with enhanced clustering [22]. Furthermore, we show that before the power-law decay of  $\bar{c}(k)$  kicks in,  $\bar{c}(k)$  first has a constant regime for small  $k$  and a logarithmic decay phase. This characterizes the entire clustering spectrum of the hidden-variable model.

This paper is structured as follows. Section II introduces the random graph model and its local clustering coefficient. Section III presents the main results for the clustering spectrum. Section IV explains the shape of the clustering spectrum in terms of an energy minimization argument, and Sec. V quantifies how fast the limiting clustering spectrum arises as a function of the network size. We conclude with a discussion in Sec. VI and present all mathematical derivations of the main results in the Appendices.

### II. HIDDEN VARIABLES

As a null model we employ the hidden-variable model [18,23–26]. Given  $N$  nodes, hidden-variable models are




FIG. 1.  $\bar{c}(k)$  for the YouTube social network.

defined as follows. Associate with each node a hidden-variable  $h$  drawn from a given probability distribution function,

$$\rho(h) = Ch^{-\tau} \tag{1}$$

for some constant  $C$ . Next join each pair of vertices independently according to a given probability  $p(h, h')$  with  $h$  and  $h'$  as the hidden variables associated with the two nodes. Many networks can be embedded in this hidden-variable framework, but particular attention goes to the case in which the hidden variables have themselves as the structure of the degrees of a real-world network. In that case the hidden-variable model puts soft constraints on the degrees, which typically is easier to analyze than hard constraints as in the configuration model [4,27–29]. Chung and Lu [30] introduced the hidden-variable model in the form

$$p(h, h') \sim \frac{hh'}{N\langle h \rangle}, \tag{2}$$

so that the expected degree of a node equals its hidden variable.

We now discuss the structural and natural cutoff because both will play a crucial role in the description of the clustering spectrum. The structural cutoff is defined as the largest possible upper bound on the degrees required to guarantee single edges, whereas the natural cutoff characterizes the maximal degree in a sample of  $N$  vertices. For scale-free networks with exponent  $\tau \in (2, 3]$  the structural cutoff scales as  $\sqrt{N}$ , whereas the natural cutoff scales as  $N^{1/(\tau-1)}$ , which gives rise to structural negative correlations and possibly other finite-size effects. If one wants to avoid such effects, then the maximal




FIG. 2. Clustering spectrum  $h \mapsto c(h)$  with three different ranges for  $h$ : the flat range, logarithmic decay, and the power-law decay.

value of the product  $hh'$  should never exceed  $N\langle h \rangle$ , which can be guaranteed by the assumption that the hidden degree  $h$  is smaller than the structural cutoff  $h_s = \sqrt{N\langle h \rangle}$ . Although this restricts  $p(h, h')$  in (2) within the interval  $[0, 1]$ , banning degrees larger than the structural cutoff strongly violates the reality of scale-free networks where degrees all the way up to the natural cutoff  $(N\langle h \rangle)^{1/(\tau-1)}$  need to be considered. We therefore work with (although many asymptotically equivalent choices are possible; see Ref. [31] and Appendix A)

$$p(h, h') = \min\left(1, \frac{hh'}{N\langle h \rangle}\right), \tag{3}$$

putting no further restrictions on the range of the hidden variables (and hence degrees).

In this paper, we will work with  $c(h)$ , the local clustering coefficient of a randomly chosen vertex with hidden-variable  $h$ . However, when studying local clustering in real-world data sets, we can only observe  $\bar{c}(k)$ , the local clustering coefficient of a vertex of degree  $k$ . In Appendix C we show that the approximation  $\bar{c}(k) \approx c(h)$  is highly accurate. We start from the explicit expression for  $c(h)$  [18], which measures the probability that two randomly chosen edges from  $h$  are neighbors, i.e.,

$$c(h) = \int_{h'} \int_{h''} p(h'|h)p(h', h'')p(h''|h)dh''dh', \tag{4}$$

with  $p(h'|h)$  as the conditional probability that a randomly chosen edge from an  $h$ -vertex is connected to an  $h'$  vertex and  $p(h, h')$  as in (3). The goal is now to characterize the  $c(h)$  curve [and hence the  $\bar{c}(k)$  curve].

### III. UNIVERSAL CLUSTERING SPECTRUM

The asymptotic evaluation of the double integral (4) in the large- $N$  regime reveals three different ranges, defined in terms of the scaling relation between the hidden-variable  $h$  and the network size  $N$ . The three ranges together span the entire clustering spectrum as shown in Fig. 2. The detailed calculations are deferred to Appendix A.

The first range pertains to the smallest-degree nodes, i.e., vertices with a hidden variable that does not exceed  $N^{\beta(\tau)}$  with  $\beta(\tau) = \frac{\tau-2}{\tau-1}$ . In this case we show that

$$c(h) \propto N^{2-\tau} \ln N, \quad h \leq N^{\beta(\tau)}. \tag{5}$$

In particular, here the local clustering does not depend on the degree and in fact corresponds with the large- $N$  behavior of the global clustering coefficient [31,32]. Note that the interval  $[0, \beta(\tau)]$  diminishes when  $\tau$  is close to 2, a possible explanation for why the flat range associated with Range I is hard to recognize in some of the real-world data sets.

Range II considers nodes with hidden variables (degrees) above the threshold  $N^{\beta(\tau)}$  but below the structural cutoff  $\sqrt{N}$ . These nodes start experiencing structural correlations, and close inspection of the integral (4) yields

$$c(h) \propto N^{2-\tau} \left[ 1 + \ln\left(\frac{\sqrt{N}}{h}\right) \right], \quad N^{\beta(\tau)} \leq h \leq \sqrt{N}. \tag{6}$$

This range shows relatively slow logarithmic decay in the clustering spectrum and clearly is visible in the ten data sets.

TABLE I. Data sets.  $N$  denotes the number of vertices,  $\tau$  denotes the exponent of the tail of the degree distribution estimated by the method proposed in Ref. [27] together with the goodness of fit criterion proposed in Ref. [27] (when the goodness of fit is at least 0.10, a power-law tail cannot be rejected), and  $\alpha$  denotes the exponent of  $c(k)$ .

	$N$	$\tau$	Goodness of fit	$\alpha$
Hudong	1.984.484	2.30	0.00	0.85
Baidu	2.141.300	2.29	0.00	0.80
Wordnet	146.005	2.47	0.00	1.01
Google web	875.713	2.73	0.00	1.03
AS-Skitter	1.696.415	2.35	0.06	1.12
TREC-WT10g	1.601.787	2.23	0.00	0.99
Wiki-talk	2.394.385	2.46	0.00	1.54
Catster and Dogster	623.766	2.13	0.00	1.20
Gowalla	196.591	2.65	0.80	1.24
YouTube	1.134.890	2.22	0.00	1.05

Range III considers hidden variables above the structural cutoff when the restrictive effect of degree-degree correlations becomes more evident. In this range we find that

$$c(h) \propto \frac{1}{N} \left( \frac{h}{N} \right)^{-2(3-\tau)}, \quad h \geq \sqrt{N}, \quad (7)$$

hence power-law decay with a power-law exponent  $\alpha = 2(3 - \tau)$ . Such power-law decay has been observed in many real-world networks [4,10,33–36] where most networks were found to have a power-law exponent close to one. The asymptotic relation (7) shows that the exponent  $\alpha$  decreases with  $\tau$  and takes values in the entire range of (0,2). Table I contains estimated values of  $\alpha$  for the ten data sets.

IV. ENERGY MINIMIZATION

We now explain why the clustering spectrum splits into three ranges using an argument that minimizes the energy needed to create triangles among nodes with specific hidden variables.

In all three ranges for  $h$ , there is one type of “most likely” triangle as shown in Fig. 3. This means that most triangles containing a vertex  $v$  with hidden-variable  $h$  are triangles with two other vertices  $v'$  and  $v''$  with hidden-variables  $h'$  and  $h''$  of specific sizes, depending on  $h$ . The probability that a triangle




FIG. 3. Orders of magnitude of the major contributions in the different  $h$  ranges. The highlighted edges are present with asymptotically positive probability. (a)  $h < \sqrt{N}$  and (b)  $h > \sqrt{N}$ .

is present among  $v$ ,  $v'$ , and  $v''$  can be written as

$$\min \left( 1, \frac{hh'}{N\langle h \rangle} \right) \min \left( 1, \frac{hh''}{N\langle h \rangle} \right) \min \left( 1, \frac{h'h''}{N\langle h \rangle} \right). \quad (8)$$

Although the probability that such a triangle exists among the three nodes thus increases with  $h'$  and  $h''$ , the number of such nodes decreases with  $h'$  and  $h''$  because vertices with higher  $h$  values are rarer. Therefore, the maximum contribution to  $c(h)$  results from a trade-off between large enough  $h', h''$  for a likeliness of the occurrence of the triangle and  $h', h''$  small enough to have enough copies. Thus, having  $h' > N\langle h \rangle/h$  is not optimal since then the probability that an edge exists between  $v$  and  $v'$  no longer increases with  $h'$ . This results in the bound,

$$h', h'' \leq \frac{N\langle h \rangle}{h}. \quad (9)$$

Similarly,  $h'h'' > N\langle h \rangle$  is also suboptimal since then further increasing  $h'$  and  $h''$  does not increase the probability of an edge between  $v'$  and  $v''$ . This gives as a second bound,

$$h'h'' \leq N\langle h \rangle. \quad (10)$$

In Ranges I and II,  $h < \sqrt{N\langle h \rangle}$  so that  $N\langle h \rangle/h > \sqrt{N\langle h \rangle}$ . In this situation we reach bound (10) before we reach bound (9). Therefore, the maximum contribution to  $c(h)$  comes from  $h'h'' \approx N$ , where also  $h', h'' < N\langle h \rangle/h$  because of bound (9). Here the probability that the edge between  $v'$  and  $v''$  exists is high, whereas the other two edges have a low probability to be present as shown in Fig. 3(a). Note that for  $h$  in Range I, bound (9) is superfluous since in this regime  $N\langle h \rangle/h > h_c$ , whereas the network does not contain vertices with hidden variables larger than  $h_c$ . This bound indicates the minimal values of  $h'$  such that an  $h$  vertex is guaranteed to be connected to an  $h'$  vertex. Thus, vertices in Range I are not even guaranteed to have connections to the highest-degree vertices, hence they are not affected by the single-edge constraints. Therefore the value of  $c(h)$  in Range I is independent of  $h$ .

In Range III,  $h > \sqrt{N\langle h \rangle}$  so that  $N\langle h \rangle/h < \sqrt{N\langle h \rangle}$ . Therefore, we reach bound (9) before we reach bound (10). Thus, we maximize the contribution to the number of triangles by choosing  $h', h'' \approx N\langle h \rangle/h$ . Then the probability that the edges from  $v$  to  $v'$  and from  $v$  to  $v''$  are present is high, whereas the probability that the edge between  $v'$  and  $v''$  exists is low as illustrated in Fig. 3(b).

V. CONVERGENCE RATE

We next ask how large networks should be, or become, before they reveal the features of the universal clustering spectrum. In other words, although the results in this paper are shown for the large- $N$  limit, for what finite  $N$  values can we expect to see the different ranges and clustering decay? To bring networks of different sizes  $N$  on a comparable footing, we consider

$$\sigma_N(t) = \frac{\ln[c(h)/c(h_c)]}{\ln(N\langle h \rangle)}, \quad h = (N\langle h \rangle)^t \quad (11)$$

for  $0 \leq t \leq \frac{1}{\tau-1}$ . The slope of  $\sigma_N(t)$  can be interpreted as a measure of the decay of  $c(h)$  at  $h = (N\langle h \rangle)^t$ , and all curves share the same right end of the spectrum; see Appendix B for




FIG. 4.  $\sigma_N(t)$  for  $N = 10^4$ ,  $10^6$ , and  $10^8$  together with the limiting function using  $\tau = 2.25$  for which  $\frac{1}{\tau-1} = 0.8$ .

more details. Figure 4 shows this rescaled clustering spectrum for synthetic networks generated with the hidden-variable model with  $\tau = 2.25$ . Already  $10^4$  vertices reveal the essential features of the spectrum: the decay and the three ranges. Increasing the network size further to  $10^5$  and  $10^6$  nodes shows that the spectrum settles on the limiting curve. Here we note that the real-world networks reported in Figs. 1 and 10 are also of order  $10^5$ – $10^6$  nodes, see Table I.

Figure 4 also brings to bear a potential pitfall when the goal is to obtain statistically accurate estimates for the slope of  $c(h)$ . Observe the extremely slow convergence to the limiting curve for  $N = \infty$ ; a well documented property of certain clustering measures [31,32,37,38]. In Appendix B we again use the integral expression (4) to characterize the limiting curve for  $N = \infty$  and the rate of convergence as function of  $N$ , and indeed extreme  $N$  values are required for statistically reliable slope estimates for, e.g.,  $t$  values of  $\frac{1}{2}$  and  $\frac{1}{\tau-1}$ ; this also is apparent from visual inspection of Fig. 4. Therefore, the estimates in Table I only serve as indicative values of  $\alpha$ . Finally, observe that Range II disappears in the limiting curve due to the rescaling in (11) but again only for extreme  $N$  values. Because this paper is about structure rather than statistical estimation, the slow convergence in fact provides additional support for the persistence of Range II in Figs. 1 and 10.

Table I also shows that the relation  $\alpha = -2(3 - \tau)$  is inaccurate for the real-world data sets, in turn, affecting the theoretical boundaries of the three regimes indicated in Fig. 10. One explanation for this inaccuracy is that the real-world networks might not follow pure power-law distributions as measured by the goodness of fit criterion in Table I and visualized in Appendix D. Furthermore, real-world networks usually are highly clustered and contain community structures, whereas the hidden-variable model is locally treelike. These modular structures may explain, for example, why the power-law decay of the hidden-variable model is less pronounced in the three social networks of Fig. 10. It is remarkable that, despite these differences between hidden-variable models and real-world networks, the global shape of the  $c(k)$  curve of the hidden-variable model is still visible in these heavy-tailed real-world networks.

## VI. DISCUSSION

The hidden-variable model gives rise to *single-edge* networks in which pairs of vertices can only be connected once.




FIG. 5.  $\bar{c}(k)$  for a hidden-variable model with connection probabilities (12) (the solid line) and an erased configuration model (the dashed line). The presented values of  $\bar{c}(k)$  are averages over  $10^4$  realizations of networks of size  $N = 10^5$ .

Hierarchical modularity and the decaying clustering spectrum have been contributed to this restriction that no two vertices have more than one edge connecting them [9,39–42]. The physical intuition is that the single-edge constraint leads to far fewer connections between high-degree vertices than anticipated based on randomly assigned edges. We have indeed confirmed this intuition not only through analytically revealing the universal clustering curve, but also by providing an alternative derivation of the three ranges based on energy minimization and structural correlations.

We now show that the clustering spectrum revealed by using the hidden-variable model also appears for a second widely studied null model. This second model cannot be the configuration model (CM), which preserves the degree distribution by making connections between vertices in the most random way possible [6,43]. Indeed, because of the random edge assignment, the CM has no degree correlations, leading in the case of scale-free networks with diverging second moment to uncorrelated networks with non-negligible fractions of self-loops (a vertex joined to itself) and multiple connections (two vertices connected by more than one edge). This picture changes dramatically when self-loops and multiple edges are avoided, a restriction mostly felt by the high-degree nodes, who can no longer establish multiple edges among each other.

We therefore consider the erased configuration model (ECM) that takes a sample from the CM and then erases all the self-loops and multiple edges. Although this removes some of the edges in the graph, thus violating the hard constraint, only a small proportion of the edges is removed so that the degree of vertex  $j$  in the ECM is still close to  $D_j$  [44, Chap. 7]. In the ECM, the probability that a vertex with degree  $D_i$  is connected to a vertex with degree  $D_j$  can be approximated by  $1 - e^{-D_i D_j / \langle D \rangle N}$  [45, Eq. (4.9)]. Therefore, we expect the ECM and the hidden-variable model to have similar properties (see, e.g., Ref. [31]) when we choose

$$p(h, h') = 1 - e^{-hh' / N \langle h \rangle} \approx \frac{hh'}{N \langle h \rangle}. \quad (12)$$

Figure 5 illustrates how both null models generate highly similar spectra, which provides additional support for the claim that the clustering spectrum is a universal property of simple scale-free networks. The ECM is more difficult

to deal with compared to the hidden-variable models since edges in the ECM are not independent. In particular, we expect that these dependencies vanish for the  $k \mapsto \bar{c}(k)$  curve. Establishing the universality of the  $k \mapsto \bar{c}(k)$  curve for other random graph null models, such as the ECM, networks with an underlying geometric space [46], or hierarchical configuration models [47], is a major research direction. The ECM and the hidden-variable model are both null models with soft constraints on the degrees. Putting hard constraints on the degrees with the CM has the nice property that simple graphs generated using this null model are uniform samples of all simple graphs with the same degree sequence. Dealing with such uniform samples is notoriously hard when the second moment of the degrees is diverging, for example, since the CM will yield many edges between high-degree vertices. This makes sampling uniform graphs difficult [48–50]. Thus, the joint requirement of hard degree and single-edge constraints as in the CM presents formidable technical challenges. Whether our results for the  $k \mapsto \bar{c}(k)$  curve for soft-constraint models also carry over to these uniform simple graphs is a challenging open problem.

In this paper we have investigated the presence of triangles in the hidden-variable model. We have shown that, by first conditioning on the node degree, there arises a unique most likely triangle with two other vertices of specific degrees. We not only have explained this insight heuristically, but also reflected it in the elaborate analysis of the double integral

for  $c(h)$  in Appendix A. As such, we have introduced an intuitive and tractable mathematical method for asymptotic triangle counting. It is likely that the method carries over to counting other motifs, such as squares or complete graphs of larger sizes. For any given motif and first conditioning on the node degree, we again expect to find specific configurations that are most likely. Further mathematical challenges need to be overcome though because we expect that the most likely configurations critically depend on the precise motif topologies and the associated energy minimization problems.

### ACKNOWLEDGMENTS

This work was supported by NWO TOP Grant No. 613.001.451 and by the NWO Gravitation Networks Grant No. 024.002.003. The work of R.v.d.H. was supported further by the NWO VICI Grant No. 639.033.806. The work of J.S.H.v.L. was supported further by an NWO TOP-GO Grant No. 613.001.012 and by an ERC Starting Grant.

### APPENDIX A: DERIVATION FOR THE THREE RANGES

In this Appendix, we compute  $c(h)$  in (4), and we show that  $c(h)$  can be approximated by (5), (6), or (7), depending on the value of  $h$ . Throughout the Appendix, we assume that  $p(h, h') = \min(1, hh'/h_s^2)$  and  $\rho(h) = Ch^{-\tau}$ . Then, the derivation of  $c(h)$  in Ref. [16] yields

$$c(h) = \frac{\int_1^{h_c} \int_1^{h_c} \rho(h') p(h, h') \rho(h'') p(h, h'') p(h', h'') dh'' dh'}{\left[ \int_1^{h_c} \rho(h') p(h, h') dh' \right]^2} = \frac{\int_1^{h_c} \int_1^{h_c} (h' h'')^{-\tau} \min\left(\frac{hh'}{h_s^2}, 1\right) \min\left(\frac{hh''}{h_s^2}, 1\right) \min\left(\frac{h'h''}{h_s^2}, 1\right) dh'' dh'}{\left[ \int_1^{h_c} (h')^{-\tau} \min\left(\frac{hh'}{h_s^2}, 1\right) dh' \right]^2}. \quad (\text{A1})$$

Computing  $c(h)$  also will allow us to compute

$$\sigma_N(t) = \frac{\ln[c(h)/c(h_{\text{ref}})]}{\ln(N\langle h \rangle)}, \quad h = (N\langle h \rangle)^t \quad (\text{A2})$$

for  $0 \leq t \leq \frac{1}{\tau-1}$ , where  $h_{\text{ref}} \in [0, h_c]$  is fixed. We are interested in computing the value of  $\sigma_N(t)$  for large values of  $N$ .

Adopting the standard choices [31],

$$h_s = \sqrt{N\langle h \rangle}, \quad h_c = (N\langle h \rangle)^{1/(\tau-1)}, \quad (\text{A3})$$

and setting  $h_{\min} = 1$  gives

$$\langle h \rangle = \frac{\tau - 1}{\tau - 2} \frac{1 - N^{2-\tau}}{1 - N^{1-\tau}}. \quad (\text{A4})$$

For ease of notation in the proofs below, we will use

$$a = h_s^{-1} = (N\langle h \rangle)^{-1/2}, \quad b = \frac{h_c}{h_s} = (N\langle h \rangle)^{[(3-\tau)/2(\tau-1)]}, \quad (\text{A5})$$

and

$$r(u) = \min(u, 1). \quad (\text{A6})$$

In this notation, (A1) can be written succinctly as

$$c(h) = \frac{\int_a^b \int_a^b (xy)^{-\tau} r(ahx)r(ahy)r(xy)dx dy}{\left[ \int_a^b x^{-\tau} r(ahx)dx \right]^2}. \quad (\text{A7})$$

Because of the four min operators in expression (A1), we have to consider various  $h$  ranges. We compute the value of  $c(h)$  in these three ranges one by one.

**Range I.**  $h < h_s^2/h_c$

We now show that, in this range,

$$c(h) \approx \frac{\tau - 2}{3 - \tau} h_s^{4-2\tau} \ln\left(\frac{h_c^2}{h_s^2}\right) \propto N^{2-\tau} \ln N, \quad (\text{A8})$$

which proves (5).

This range corresponds to  $h < 1/(ab)$  with  $a$  and  $b$  as in (A5). In this range,  $r(ahx) = ahx$  and  $r(ahy) = ah y$  for all  $x \in [a, b]$ . This yields for  $c(h)$ ,

$$c(h) = \frac{\int_a^b \int_a^b (xy)^{1-\tau} r(xy)dx dy}{\left[ \int_a^b x^{1-\tau} dx \right]^2}. \quad (\text{A9})$$

For the denominator we compute

$$\int_a^b x^{1-\tau} dx = \frac{a^{2-\tau} - b^{2-\tau}}{\tau - 2}. \quad (\text{A10})$$

Since  $a \ll b$ , this can be approximated as

$$\frac{a^{2-\tau} - b^{2-\tau}}{\tau - 2} \approx \frac{a^{2-\tau}}{\tau - 2}. \quad (\text{A11})$$

We can compute the numerator of (A9) as

$$\begin{aligned} \int_a^b \int_a^b (xy)^{1-\tau} r(xy)dx dy &= \int_a^{1/b} \int_a^b (xy)^{2-\tau} dx dy + \int_{1/b}^b \int_a^{1/x} (xy)^{2-\tau} dx dy + \int_{1/b}^b \int_{1/x}^b (xy)^{1-\tau} dx dy \\ &= \frac{(b^{\tau-3} - a^{3-\tau})(b^{3-\tau} - a^{3-\tau})}{(3-\tau)^2} + \frac{1}{3-\tau} \left( \ln(b^2) - \frac{a^{3-\tau}(b^{3-\tau} - b^{\tau-3})}{3-\tau} \right) \\ &\quad + \frac{1}{2-\tau} \left( \frac{b^{2-\tau}(b^{2-\tau} - b^{\tau-2})}{2-\tau} - \ln(b^2) \right) \\ &= \frac{\ln(b^2)}{(3-\tau)(\tau-2)} - \frac{1 - b^{4-2\tau}}{(\tau-2)^2} + \frac{1 - 2(ab)^{3-\tau} + a^{6-2\tau}}{(3-\tau)^2}. \end{aligned} \quad (\text{A12})$$

The first of these three terms dominates when

$$\frac{3-\tau}{\tau-1} \frac{\ln(N\langle h \rangle)}{(3-\tau)(\tau-2)} \gg \frac{1}{(\tau-2)^2}, \quad (\text{A13})$$

and

$$\frac{3-\tau}{\tau-1} \frac{\ln(N\langle h \rangle)}{(3-\tau)(\tau-2)} \gg \frac{1}{(3-\tau)^2}, \quad (\text{A14})$$

where we have used that  $b^2 = (N\langle h \rangle)^{(3-\tau)/(\tau-1)}$ . Thus, when  $\ln(N\langle h \rangle)$  is large compared to  $(\tau-1)/(\tau-2)$  and  $(\tau-1)(\tau-2)/(\tau-3)^2$ , we obtain

$$c(h) \approx \frac{\tau-2}{3-\tau} a^{2\tau-4} \ln(b^2) \propto N^{2-\tau} \ln(N), \quad (\text{A15})$$

which proves (A8).

**Range II.**  $h_s^2/h_c < h < h_s$

In this range, we show that

$$c(h) \approx h_s^{4-2\tau} \frac{\ln\left(\frac{h_s^2}{h^2}\right) + M}{(\tau-2)(3-\tau)} \propto N^{2-\tau} [\ln(N/h^2) + M] \quad (\text{A16})$$

for some positive constant  $M$ , which proves (6).

This range corresponds to  $(ab)^{-1} < h < a^{-1}$ . For these values of  $h$ , we have  $ahx, ah y = 1$  for  $x, y = (ah)^{-1} \in (1, b)$  and  $xy = 1$  for  $y = 1/x \in [a, b]$  when  $b^{-1} < x < b$ . Then for the denominator of (A7) we compute

$$\begin{aligned} \int_a^{1/(ah)} ahx^{1-\tau} dx + \int_{1/(ah)}^b x^{-\tau} dx &= \frac{1}{\tau-2} [a^{3-\tau} h - (ah)^{\tau-1}] + \frac{1}{\tau-1} [(ah)^{\tau-1} - b^{1-\tau}] \\ &= ah \left( \frac{a^{2-\tau}}{\tau-2} - \frac{(ah)^{\tau-2}}{(\tau-1)(\tau-2)} - \frac{b^{1-\tau}/(ah)}{\tau-1} \right). \end{aligned} \tag{A17}$$

Splitting up the integral in the numerator results in

$$\begin{aligned} \text{Num}(h) &= \int_a^b \int_a^b (xy)^{-\tau} r(ahx)r(ahy)r(xy) dx dy = \int_{1/(ah)}^b \int_{1/(ah)}^b (xy)^{-\tau} dy dx + 2ah \int_{1/(ah)}^b \int_{1/x}^{1/(ah)} (xy)^{-\tau} y dy dx \\ &+ 2ah \int_{1/(ah)}^b \int_a^{1/x} (xy)^{1-\tau} y dy dx + a^2 h^2 \int_{ah}^{1/(ah)} \int_a^{1/x} (xy)^{2-\tau} dy dx + a^2 h^2 \int_{ah}^{1/(ah)} \int_{1/x}^{1/(ah)} (xy)^{1-\tau} dy dx \\ &+ a^2 h^2 \int_a^{ah} \int_a^{1/(ah)} (xy)^{2-\tau} dy dx =: I_1 + I_2 + I_3 + I_4 + I_5 + I_6, \end{aligned} \tag{A18}$$

where the factors 2 arise by symmetry of the integrand in  $x$  and  $y$ . Computing these integrals yields

$$I_1 = a^2 h^2 \left( \frac{(ah)^{\tau-2} - a^{-1} b^{1-\tau} h^{-1}}{\tau-1} \right)^2, \tag{A19}$$

$$I_2 = 2a^2 h^2 \left( \frac{1 - 1/(abh)}{\tau-2} - \frac{(ah)^{2\tau-4}}{(\tau-1)(\tau-2)} [1 - (abh)^{1-\tau}] \right), \tag{A20}$$

$$I_3 = 2a^2 h^2 \left( \frac{1 - 1/(abh)}{3-\tau} - \frac{h^{\tau-3} [1 - (abh)^{2-\tau}]}{(3-\tau)(\tau-2)} \right), \tag{A21}$$

$$I_4 = a^2 h^2 \left( \frac{\ln[(ah)^{-2}]}{3-\tau} + \frac{(a^2 h)^{3-\tau} - h^{\tau-3}}{(3-\tau)^2} \right), \tag{A22}$$

$$I_5 = a^2 h^2 \left( \frac{\ln[(ah)^{-2}]}{\tau-2} - \frac{1 - (ah)^{2\tau-4}}{(\tau-2)^2} \right), \tag{A23}$$

$$I_6 = a^2 h^2 \left( \frac{1 - h^{\tau-3} + a^{6-2\tau} - (a^2 h)^{3-\tau}}{(3-\tau)^2} \right). \tag{A24}$$

We have  $ah < 1 < ahb$ , and so the leading behavior of  $\text{Num}(h)$  is determined by the terms involving  $\ln[(ah)^{-2}]$  in  $I_3$  and  $I_4$ , all other terms being bounded. Retaining only these dominant terms, we get

$$\text{Num}(h) = a^2 h^2 \frac{\ln[(ah)^{-2}]}{(\tau-2)(3-\tau)} [1 + o(1)] \tag{A25}$$

provided that  $ah \rightarrow 0$  as  $N \rightarrow \infty$ . In terms of the variable  $t$  in  $h = (N\langle h \rangle)^t$ , see (11) and (A2), this condition holds when we restrict to  $t \in [(\tau-2)/(\tau-1), \frac{1}{2} - \varepsilon]$  for any  $\varepsilon > 0$ . Furthermore, from (A17),

$$\left( \int_a^b x^{-\tau} r(ahx) dx \right)^2 = a^2 h^2 \left( \frac{a^{2-\tau}}{\tau-2} \right)^2 [1 + o(1)]. \tag{A26}$$

Hence, when  $ah \rightarrow 0$ , we have

$$c(h) = \frac{\tau-2}{3-\tau} a^{2\tau-4} \ln[(ah)^{-2}] [1 + o(1)] \propto N^{2-\tau} \ln(N/h^2). \tag{A27}$$

We compute  $c(h = 1/a)$  asymptotically by retaining only all constant terms between brackets in (A19)–(A24) since all other terms vanish or tend to 0 as  $N \rightarrow \infty$ . This gives

$$\text{Num}(h = 1/a) = a^2 h^2 \left( \frac{1}{(\tau-1)^2} + \frac{2}{\tau-2} - \frac{2}{(\tau-1)(\tau-2)} + \frac{2}{3-\tau} + \frac{1}{(3-\tau)^2} \right) [1 + o(1)] = Pa^2 h^2 [1 + o(1)], \tag{A28}$$

where  $P = \frac{1}{(\tau-1)^2} + \frac{1}{(3-\tau)^2} + \frac{2}{\tau-1} + \frac{2}{3-\tau}$ . Together with (A26), we find

$$c(h = 1/a) = P(\tau-2)^2 a^{2\tau-4} [1 + o(1)] \propto N^{2-\tau}. \tag{A29}$$

In Ref. [31], it has been shown that  $c(h)$  decreases in  $h$ , and then (A16) follows from (A27) and (A29).



**Range III.**  $h_s < h < h_c$

We now show that when  $h_s < h < h_c$ , then

$$c(h) \approx \frac{1}{(3 - \tau)^2} (h_s/h)^{6-2\tau} h_s^{4-2\tau} \propto N^{5-2\tau} h^{2\tau-6}, \tag{A30}$$

which proves (7).

This range corresponds to  $1/a < h < b/a$ . The denominator of (A7) remains the same as in the previous range and is given by (A17). Splitting up the integral in the numerator of (A7) now results in

$$\begin{aligned} \text{Num}(h) &= \int_a^b \int_a^b (xy)^{-\tau} r(ahx)r(ahy)r(xy) dx dy = \int_{1/(ah)}^{ah} \int_{1/x}^b (xy)^{-\tau} dy dx + \int_{ah}^b \int_{1/(ah)}^b (xy)^{-\tau} dy dx \\ &+ \int_{1/(ah)}^{ah} \int_{1/(ah)}^{1/x} (xy)^{1-\tau} dy dx + 2ah \int_{ah}^b \int_{1/x}^{1/(ah)} (xy)^{-\tau} y dy dx + 2ah \int_{1/(ah)}^{ah} \int_a^{1/(ah)} (xy)^{1-\tau} y dy dx \\ &+ 2ah \int_{ah}^b \int_a^{1/x} (xy)^{1-\tau} y dy dx + a^2 h^2 \int_a^{1/(ah)} \int_a^{1/(ah)} (xy)^{2-\tau} dy dx =: I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7. \end{aligned} \tag{A31}$$

Computing these integrals yields

$$I_1 = a^2 h^2 \left( (ah)^{-2} \frac{\ln(a^2 h^2)}{\tau - 1} + \frac{b^{1-\tau} [(ah)^{-\tau-1} - (ah)^{\tau-3}]}{(\tau - 1)^2} \right), \tag{A32}$$

$$I_2 = a^2 h^2 \left( \frac{(ah)^{-2} + b^{2-2\tau} (ah)^{-2}}{(\tau - 1)^2} - \frac{b^{1-\tau} [(ah)^{\tau-3} + (ah)^{-\tau-1}]}{(\tau - 1)^2} \right), \tag{A33}$$

$$I_3 = a^2 h^2 \left( -(ah)^{-2} \frac{\ln(a^2 h^2)}{\tau - 2} + \frac{(ah)^{2\tau-6} - (ah)^{-2}}{(\tau - 2)^2} \right), \tag{A34}$$

$$I_4 = 2a^2 h^{-2} \left( -\frac{(abh)^{-1}}{\tau - 2} + \frac{(ah)^{-2}}{\tau - 1} + \frac{b^{1-\tau} (ah)^{\tau-3}}{(\tau - 1)(\tau - 2)} \right), \tag{A35}$$

$$I_5 = 2a^2 h^2 \left( \frac{(ah)^{2\tau-6} + h^{1-\tau} a^{4-2\tau} - h^{\tau-3} - (ah)^{-2}}{(3 - \tau)(\tau - 2)} \right), \tag{A36}$$

$$I_6 = 2a^2 h^2 \left( \frac{(ab)^{2-\tau} h^{-1} - h^{1-\tau} a^{4-2\tau}}{(3 - \tau)(\tau - 2)} - \frac{(abh)^{-1} - (ah)^{-2}}{3 - \tau} \right), \tag{A37}$$

$$I_7 = a^2 h^2 \left( \frac{a^{6-2\tau} - 2h^{\tau-3} + (ah)^{2\tau-6}}{\tau - 3} \right). \tag{A38}$$

A careful inspection of the terms between brackets in (A32) and (A38) shows that the terms involving  $(ah)^{2\tau-6}$  are dominant when  $ah \rightarrow \infty$ . In terms of the variable  $t$  in  $h = (N\langle h \rangle)^t$ , see (11) and (A2), we have that  $ah \rightarrow \infty$  when we restrict to  $t \in [\frac{1}{2} + \varepsilon, 1/(\tau - 1)]$  for any  $\varepsilon > 0$ . When we retain only these dominant terms, we have, when  $ah \rightarrow \infty$ ,

$$\text{Num}(h) = a^2 h^2 (ah)^{2\tau-6} \left( \frac{1}{(\tau - 2)^2} + \frac{2}{(3 - \tau)(\tau - 2)} + \frac{1}{(3 - \tau)^2} \right) [1 + o(1)] = a^2 h^2 \frac{(ah)^{2\tau-6}}{(\tau - 2)^2 (3 - \tau)^2} [1 + o(1)]. \tag{A39}$$

Using (A26) again, we get, when  $ah \rightarrow \infty$ ,

$$c(h) = \frac{1}{(3 - \tau)^2} (ah)^{2\tau-6} a^{2\tau-4} [1 + o(1)] \propto N^{5-2\tau} h^{2\tau-6}. \tag{A40}$$

Furthermore,  $c(1/a)$  is given by (A29), whereas  $c(h)$  decreases in  $h$ . This gives (A30).

**Other connection probabilities**

In Ref. [31] we have presented a class of functions  $r(u) = uf(u)$ ,  $u \geq 0$  so that

$$p(h, h') = r(u) \quad \text{with } u = \frac{hh'}{h_s^2} \tag{A41}$$

has appropriate monotonicity properties. The maximal member  $r(u) = \min(u, 1)$  of this class yields  $p$  in (3) and is quite representative of the whole class whereas allowing explicit computation and asymptotic analysis of  $c(h)$  as in Ref. [31] and this paper. Figure 6 shows that other asymptotically equivalent choices, such as  $r(u) = u/(1 + u)$  and  $r(u) = 1 - e^{-u}$ , have comparable clustering spectra. A minor difference is that the choice  $r(u) = \min(1, u)$  for  $p$  in (3) forces  $c(h)$  to be constant on the range of  $h \leq N^{\beta(\tau)}$ , whereas the other two choices show a gentle decrease.




FIG. 6.  $c(h)$  for  $r(u) = \min(u, 1)$  (the solid line),  $r(u) = u/(1 + u)$  (the dashed line), and  $r(u) = 1 - e^{-u}$  (the dotted line), obtained by calculating (A7) numerically.

### Limiting form of $\sigma_N(t)$ and finite-size effects

We consider  $\sigma_N(t)$  as in (A2) with  $h_{\text{ref}} = 0$ . Using (A8), (A16), and (A30), it readily is seen that

$$\lim_{N \rightarrow \infty} \sigma_N(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ (3 - \tau)(1 - 2t), & \frac{1}{2} \leq t \leq \frac{1}{\tau - 1}. \end{cases} \quad (\text{A42})$$

Hence, some of the detailed information that is present in (A8), (A16), and (A30), disappears when taking the limit as in (A42). This is in particular so for the  $\ln N$  factor in (A8) and the logarithmic decaying factor  $\ln(N^2/h)$  in Region II.

Consider  $\sigma_N(t)$  of (A2) with  $h_{\text{ref}} = h_c$  as is performed in Fig. 4. It follows from the detailed forms of (A8) and (A30) that

$$\sigma_N(0) = \frac{\ln[c(0)/c(h_c)]}{\ln(N\langle h \rangle)} = \gamma + \frac{\ln(\beta y)}{y}, \quad (\text{A43})$$

where

$$\gamma = \frac{(3 - \tau)^2}{\tau - 1}, \quad \beta = (\tau - 2)\gamma, \quad y = \ln(N\langle h \rangle). \quad (\text{A44})$$

We have that  $\sigma_N(0) \rightarrow \gamma$  as  $N \rightarrow \infty$ , and the right-hand side of (A43) exceeds this limit  $\gamma$  from  $y = 1/\beta$  onwards with a maximum excess  $\beta/e$  for  $N\langle h \rangle$  as large as  $\exp(e/\beta)$ . This explains why the deviation of  $\sigma_N(0)$  from its limit value in Fig. 4 persists when  $\tau = 9/4$  (for which  $e^{e/\beta} = 3 \times 10^{10}$ ).

### APPENDIX B: EXACT AND ASYMPTOTIC RESULT FOR THE DECAY RATE OF $c(h)$ AT $h = h_c$ AND $h = h_s$

We let  $h_c = (N\langle h \rangle)^{1/(\tau-1)}$  where we assume that  $N$  is so large that  $h_c \leq N$ . This requires  $N$  to be on the order of  $(1/\varepsilon)^{1/\varepsilon}$ , where  $\varepsilon = \tau - 2$ . We again consider the function  $\sigma_N(t)$  of (11),

$$\sigma_N(t) = \frac{\ln[c(h)/c(h_{\text{ref}})]}{\ln(N\langle h \rangle)}, \quad h = (N\langle h \rangle)^t \quad (\text{B1})$$

for  $0 \leq t \leq \frac{1}{\tau-1}$  and  $h_{\text{ref}}$  is fixed so that

$$c(h) = c(h_{\text{ref}})(N\langle h \rangle)^{\sigma_N(t)}, \quad h = (N\langle h \rangle)^t. \quad (\text{B2})$$

When we fix  $t_0$  and linearize  $\sigma_N(t)$  around  $t_0$ , we get

$$\begin{aligned} c(h) &\approx c(h_{\text{ref}})(N\langle h \rangle)^{\sigma_N(t_0) + (t-t_0)\sigma'_N(t_0)} \\ &= c(h_0) \left( \frac{h}{h_0} \right)^{\sigma'_N(t_0)}, \end{aligned} \quad (\text{B3})$$

so that  $\sigma'_N(t) = \frac{d}{dt} \sigma_N(t)$  is a measure for the decay rate of  $c(h)$  at  $h = h_0 = (N\langle h \rangle)^{t_0}$ .

In this Appendix, we compute an exact expression for  $\sigma'_N(t)$  at  $t = \frac{1}{\tau-1}$ , we compute its limit as  $N \rightarrow \infty$  and discuss convergence speed, and we show that this limit is a lower bound for  $\sigma'_N(t)$ .

More precisely, we show the following result:

*Proposition 1.* Let  $a$  and  $b$  be as in (A5). Then

$$\sigma'_N \left( \frac{1}{\tau - 1} \right) = -2 \left( \frac{A + \frac{3-\tau}{\tau-2}C}{A + \frac{4-\tau}{\tau-2}C} - \frac{D}{E + D} \right), \quad (\text{B4})$$

where

$$A = \frac{1}{b^2} \left( \frac{-\ln(b^2)}{(\tau-1)(\tau-2)} - \frac{1 - b^{2(1-\tau)}}{(\tau-1)^2} + \frac{b^{2(\tau-2)} - 1}{(\tau-2)^2} \right), \quad (\text{B5})$$

$$C = \left( \frac{b^{\tau-3} - a^{3-\tau}}{3 - \tau} \right)^2, \quad (\text{B6})$$

$$D = \frac{1}{b} \frac{b^{\tau-1} - b^{1-\tau}}{\tau - 1}, \quad (\text{B7})$$

$$E = \frac{a^{2-\tau} - b^{\tau-2}}{\tau - 2}. \quad (\text{B8})$$

Furthermore,

$$\sigma'_N \left( \frac{1}{\tau - 1} \right) > \lim_{M \rightarrow \infty} \sigma'_M \left( \frac{1}{\tau - 1} \right) = -2(3 - \tau) \quad (\text{B9})$$

for all  $N$ .

The limiting value in (B9) is consistent with the limiting value of  $\sigma_N(t)$  that has been found in (A42). We assess this convergence result with plots. Although these indicate that the limits are reached only for very large  $N$ , especially when  $\tau$  is close to 2, it can also be seen that the limiting shape of  $\sigma_N(t)$  already shows up for considerably smaller  $N$ .

To start the proof of Proposition 1, note that in the  $a, b$  notation of (A5),

$$c(h) = \frac{K(h)}{J(h)}, \quad 0 \leq h \leq h_c, \quad (\text{B10})$$

where

$$K(h) = \int_a^b \int_a^b (xy)^{2-\tau} f(ahx) f(ahy) f(xy) dx dy, \quad (\text{B11})$$

$$J(h) = \left( \int_a^b x^{1-\tau} f(ahx) dx \right)^2, \quad (\text{B12})$$




FIG. 7.  $\sigma'_N(t)$  plotted against  $N$  for (a)  $t = \frac{1}{\tau-1}$  and (b)  $t = \frac{1}{2}$ . The dashed line gives the limiting value of  $\sigma'_N(t)$  as  $N \rightarrow \infty$ .

with  $f(u) = \min(1, u^{-1})$ . Note that  $r(u) = uf(u)$ , see (A6). We compute

$$\begin{aligned} \sigma'_N(t) &= \frac{d}{dt} \left( \frac{\ln\{c[(N\langle h \rangle)^t]/c(h_{\text{ref}})\}}{\ln(N\langle h \rangle)} \right) \\ &= (N\langle h \rangle)^t \ln(N\langle h \rangle) \frac{c'[(N\langle h \rangle)^t]}{c[(N\langle h \rangle)^t] \ln(N\langle h \rangle)} \\ &= h \frac{c'(h)}{c(h)}, \quad h = (N\langle h \rangle)^t, \end{aligned} \quad (\text{B13})$$

where the prime on  $c$  indicates differentiation with respect to  $h$ . With (B10) we get

$$\frac{c'(h)}{c(h)} = \frac{K'(h)}{K(h)} - \frac{J'(h)}{J(h)}, \quad (\text{B14})$$

and we have to evaluate  $K(h)$ ,  $K'(h)$ ,  $J(h)$ , and  $J'(h)$  at

$$h = h_c = b/a. \quad (\text{B15})$$

*Lemma 1.*

$$K(h_c) = A + \frac{4-\tau}{2-\tau}C, \quad K'(h_c) = \frac{-2a}{b} \left( A + \frac{3-\tau}{\tau-2}C \right), \quad (\text{B16})$$

$$J(h_c) = (D+E)^2, \quad J'(h_c) = -\frac{2a}{b}(D+E)D, \quad (\text{B17})$$

with  $A, C, D, E$  as in (B5)–(B8).

From Lemma 1, (B13), and (B15) we get (B4) in Proposition 1.

*Proof of Lemma 1.* Since  $h_c = b/a$ ,

$$K(h_c) = \int_a^b \int_a^b (xy)^{2-\tau} f(bx)f(by)f(xy) dx dy. \quad (\text{B18})$$

With  $f(u) = \min(1, u^{-1})$  we split up the integration range  $[a, b] \times [a, b]$  into the four regions  $[a, 1/b] \times [a, 1/b]$ ,  $[1/b, b] \times [1/b, b]$ ,  $[1/b, b] \times [a, 1/b]$ , and  $[a, 1/b] \times [1/b, b]$ , where we observe that  $a \leq 1/b \leq 1 \leq b$ . Here  $\times$

denotes the Cartesian product. We first get

$$\begin{aligned} &\int_a^{1/b} \int_a^{1/b} (xy)^{2-\tau} f(bx)f(by)f(xy) dx dy \\ &= \int_a^{1/b} \int_a^{1/b} (xy)^{2-\tau} dx dy \\ &= \left( \frac{b^{\tau-3} - a^{3-\tau}}{3-\tau} \right)^2 = C. \end{aligned} \quad (\text{B19})$$

Next,

$$\begin{aligned} &\int_{1/b}^b \int_{1/b}^b (xy)^{2-\tau} f(bx)f(by)f(xy) dx dy \\ &= \int_{1/b}^b \int_{1/b}^b (xy)^{2-\tau} \frac{1}{bx} \frac{1}{by} f(xy) dx dy \\ &= \frac{1}{b^2} \int_{1/b}^b \int_{1/b}^b (xy)^{1-\tau} f(xy) dx dy. \end{aligned} \quad (\text{B20})$$

The remaining double integral with  $\tau + 1$  instead of  $\tau$  has been evaluated in Ref. [31, Appendix C, (C3)] as

$$-\frac{\ln(b^2)}{(\tau-1)(\tau-2)} - \frac{1-b^{2(1-\tau)}}{(\tau-1)^2} + \frac{b^{2(\tau-2)-1}}{(\tau-2)^2} = b^2 A. \quad (\text{B21})$$

Finally, the two double integrals over  $[1/b, b] \times [a, 1/b]$  and  $[a, 1/b] \times [1/b, b]$  are by symmetry both equal to

$$\begin{aligned} &\int_{1/b}^b \int_a^{1/b} (xy)^{2-\tau} f(bx)f(by)f(xy) dx dy \\ &= \int_{1/b}^b \int_a^{1/b} (xy)^{2-\tau} \frac{1}{bx} dx dy \\ &= \frac{1}{b} \frac{b^{\tau-2} - b^{2-\tau} b^{\tau-3} - a^{3-\tau}}{\tau-2} = \frac{(b^{\tau-3} - a^{3-\tau})^2}{(\tau-2)(3-\tau)} \\ &= \frac{3-\tau}{\tau-2} C. \end{aligned} \quad (\text{B22})$$

Here we have used that, see (A5),

$$b^{1-\tau} = a^{3-\tau}. \quad (\text{B23})$$

Now the expression in (B16) for  $K(h_c)$  follows.

To evaluate  $K'(h_c)$ , we observe by symmetry that

$$K'(h) = 2 \int_a^b \int_a^b (xy)^{2-\tau} axf'(ahx)f(ahy)f(xy)dx dy. \quad (\text{B24})$$

At  $h = h_c$ , we have  $ah = b$ , and so

$$K'(h_c) = 2 \frac{a}{b} \int_a^b \int_a^b (xy)^{2-\tau} bxf'(bx)f(by)f(xy)dx dy. \quad (\text{B25})$$

Now  $uf'(u) = 0$  for  $0 \leq u \leq 1$  and  $uf'(u) = -f(u)$  for  $u \geq 1$ . Hence, splitting up the integration range into the four regions as earlier, we see that those over  $[a, 1/b] \times [a, 1/b]$  and  $[a, 1/b] \times [1/b, b]$  vanish while those over  $[1/b, b] \times [1/b, b]$  and  $[1/b, b] \times [a, 1/b]$  give rise to the same double integrals as in (B20) and (B22), respectively. This yields the expression in (B16) for  $K'(h_c)$ .

The evaluation of  $J(h_c)$  and  $J'(h_c)$  is straightforward from (B12) with  $ah = b$  and a splitting of the integration range  $[a, b]$  into  $[a, 1/b]$  and  $[1/b, b]$ . This yields (B17), and the proof of Lemma 1 is complete.

We now turn to the limiting behavior of  $\sigma'_N(\frac{1}{\tau-1})$  as  $N \rightarrow \infty$ . For this we write

$$0 < \frac{D}{D+E} = \frac{1 - b^{2(1-\tau)}}{\frac{\tau-1}{\tau-2}(ab)^{2-\tau} - \frac{1}{\tau-2} - \frac{1}{\tau-1}b^{2(1-\tau)}}, \quad (\text{B26})$$

in which

$$b^{2(1-\tau)} = (N\langle h \rangle)^{\tau-3} \rightarrow 0, \quad (\text{B27})$$

$$(ab)^{2-\tau} = (N\langle h \rangle)^{(\tau-2)^2/(\tau-1)} \rightarrow \infty, \quad (\text{B28})$$

as  $N \rightarrow \infty$ . Hence,  $D/(D+E) \rightarrow 0$  as  $N \rightarrow \infty$ . Furthermore, we write

$$C = \frac{b^{2(\tau-3)}}{(\tau-3)^2} [1 - (ab)^{3-\tau}]^2, \quad (\text{B29})$$

and

$$A = \frac{b^{2(\tau-3)}}{(\tau-2)^2} (1-F), \quad (\text{B30})$$

where

$$\begin{aligned} F &= b^{-2(\tau-2)} \left[ \frac{\tau-2}{\tau-1} \ln(b^2) + \left( \frac{\tau-2}{\tau-1} \right)^2 (1 - b^{2(1-\tau)}) + 1 \right] \\ &= \frac{1}{\tau-1} b^{-2(\tau-2)} \ln(b^{2(\tau-2)}) \left[ 1 + O\left( \frac{1}{\ln(b)} \right) \right]. \end{aligned} \quad (\text{B31})$$

Now, using (B23), we have

$$(ab)^{3-\tau} = b^{-2(\tau-2)} = (N\langle h \rangle)^{[(\tau-2)(3-\tau)]/(\tau-1)} \rightarrow 0, \quad (\text{B32})$$

as  $N \rightarrow \infty$ . Thus, we get

$$\lim_{N \rightarrow \infty} \frac{A + \frac{3-\tau}{2-\tau}C}{A + \frac{4-\tau}{2-\tau}C} = \frac{\frac{1}{(\tau-2)^2} + \frac{3-\tau}{\tau-2} \frac{1}{(3-\tau)^2}}{\frac{1}{(\tau-2)^2} + \frac{4-\tau}{\tau-2} \frac{1}{(3-\tau)^2}} = 3 - \tau, \quad (\text{B33})$$

and this yields (B9).

Note that  $D/(D+E)$  approaches 0 much slower than the limit in (B33) is reached when  $\tau$  is close to 2, compare (B28) and (B33). Thus, we can concentrate on  $D/(D+E)$ , and the




FIG. 8.  $\bar{c}(k)$  (the dashed line) and  $c(h)$  (the solid line) for  $N = 10^5$ , averaged over  $10^4$  realizations.

relative deviation of  $\sigma'_N(t)$  from  $-2(3-\tau)$  is approximately,

$$\begin{aligned} \frac{2D}{D+E} \frac{1}{2(3-\tau)} &\approx \frac{\tau-2}{3-\tau} \frac{1}{(ab)^{2-\tau} - 1} \\ &\approx \frac{\tau-2}{3-\tau} (N\langle h \rangle)^{-[(\tau-2)^2/(\tau-1)]}. \end{aligned} \quad (\text{B34})$$

We finally turn to the inequality in (B9) in Proposition 1. Obviously, we have

$$\sigma'_N\left(\frac{1}{\tau-1}\right) > -2 \frac{A + \frac{3-\tau}{\tau-2}C}{A + \frac{4-\tau}{\tau-2}C}. \quad (\text{B35})$$

We will show that

$$\frac{A + \frac{3-\tau}{\tau-2}C}{A + \frac{4-\tau}{\tau-2}C} \leq \frac{A_{\text{as}} + \frac{3-\tau}{\tau-2}C_{\text{as}}}{A_{\text{as}} + \frac{4-\tau}{\tau-2}C_{\text{as}}} = 3 - \tau, \quad (\text{B36})$$

where

$$A_{\text{as}} = \frac{b^{2(\tau-3)}}{(\tau-2)^2}, \quad C_{\text{as}} = \frac{b^{2(\tau-3)}}{(3-\tau)^2}, \quad (\text{B37})$$

the asymptotic form of  $A$  and  $C$  as  $N \rightarrow \infty$  obtained from (B30) and (B29) by deleting  $F$  and  $(ab)^{3-\tau}$ , respectively. The function,

$$x \in [0, \infty) \mapsto \frac{1 + \frac{3-\tau}{\tau-2}x}{1 + \frac{4-\tau}{\tau-2}x} \quad (\text{B38})$$

is decreasing in  $x \geq 0$ , and so it suffices to show that

$$\frac{C_{\text{as}}}{A_{\text{as}}} \leq \frac{C}{A}, \quad \text{i.e., that } \frac{C_{\text{as}}}{C} \leq \frac{A_{\text{as}}}{A}. \quad (\text{B39})$$

We have from (B29) that

$$\frac{C_{\text{as}}}{C} = \frac{1}{[1 - (ab)^{3-\tau}]^2}, \quad (\text{B40})$$

and from (B30) and (B31) that

$$\begin{aligned} \frac{A}{A_{\text{as}}} &= 1 - F = 1 - b^{-2(\tau-2)} - b^{-2(\tau-2)} \\ &\times \left[ \frac{\tau-2}{\tau-1} \ln(b^2) + \left( \frac{\tau-2}{\tau-1} \right)^2 (1 - b^{2(1-\tau)}) \right]. \end{aligned} \quad (\text{B41})$$

Using that  $(ab)^{3-\tau} = b^{-2(\tau-2)}$ , see (B32), we see that the inequality  $C_{as}/C \leq A_{as}/A$  in (B39) is equivalent to

$$(1 - b^{-2(\tau-2)})^2 \geq 1 - b^{-2(\tau-2)} - b^{-2(\tau-2)} \times \left[ \frac{\tau-2}{\tau-1} \ln(b^2) + \left( \frac{\tau-2}{\tau-1} \right)^2 (1 - b^{2(1-\tau)}) \right]. \quad (\text{B42})$$

Using that  $(1-u)^2 - (1-u) = -u(1-u)$  and dividing through by  $u = b^{-2(\tau-2)}$ , we see that (B42) is equivalent to

$$\frac{\tau-2}{\tau-1} \ln(b^2) + \left( \frac{\tau-2}{\tau-1} \right)^2 (1 - b^{2(1-\tau)}) \geq 1 - b^{-2(\tau-2)}. \quad (\text{B43})$$

With  $y = \ln(b^2) \geq 0$ , we write (B43) as

$$K(y) := \left( \frac{\tau-2}{\tau-1} \right)^2 (1 - e^{(1-\tau)y}) + \frac{\tau-2}{\tau-1} y - (1 - e^{(2-\tau)y}) \geq 0. \quad (\text{B44})$$

Taylor development of  $K(y)$  at  $y = 0$  yields

$$K(y) = 0y^0 + 0y^1 + 0y^2 + \frac{1}{6}(\tau-2)^2 y^3 + \dots \quad (\text{B45})$$

Furthermore,

$$K''(y) = (\tau-2)^2 e^{(1-\tau)y} (e^y - 1) > 0, \quad y > 0. \quad (\text{B46})$$

Therefore,  $K(0) = K'(0) = 0$ , whereas  $K''(y) > 0$  for  $y > 0$ . This gives  $K(y) > 0$  when  $y > 0$  as required.

Similar to Proposition 1, we can derive the following result for  $\sigma'_N(\frac{1}{2})$ :


*Proposition 2.*

$$\sigma'_N\left(\frac{1}{2}\right) = -2 \left[ \frac{G + H}{\left[1 + \left(\frac{\tau-1}{3-\tau}\right)^2\right]G + 2H} - \frac{I}{I + J} \right], \quad (\text{B47})$$

where

$$G = \left( \frac{1 - b^{1-\tau}}{\tau - 1} \right)^2, \quad (\text{B48})$$

$$I = \frac{1 - b^{1-\tau}}{\tau - 1}, \quad (\text{B49})$$



$$J = \frac{b^{(\tau-2)(\tau-1)/(3-\tau)} - 1}{\tau - 2}, \quad (\text{B50})$$

$$H = \frac{1 - 1/b - b^{1-\tau}(1 - b^{2-\tau})}{(\tau - 2)(3 - \tau)} - \frac{1 - b^{1-\tau}}{(\tau - 1)(\tau - 2)}. \quad (\text{B51})$$

Furthermore,

$$\sigma'_N\left(\frac{1}{2}\right) > \lim_{M \rightarrow \infty} \sigma'_M\left(\frac{1}{2}\right) = -1 + \frac{2(\tau - 2)}{3 - (\tau - 2)^2} \quad (\text{B52})$$

for all  $N$ .

Figure 7 shows the values of  $\sigma'_N(\frac{1}{2})$  and  $\sigma'_N(\frac{1}{\tau-1})$  for finite-size networks together with their limiting values. For example, when  $\tau = 2.25$ , Fig. 7(a) shows that  $N$  needs to be on the order of  $10^{16}$  for the slope to be close to its limiting value of  $-1.5$ . When, for example,  $N = 10^6$ , we see that the slope is much smaller: approximately  $-1.1$ . This makes a statistical estimation of the true underlying power-law exponent  $\alpha$  extremely challenging, especially for the relevant regime  $\tau$  close to 2 because enormous amounts of data should be available to get sufficient statistical accuracy. Most data sets, even the largest available networks used in this paper, are simply not large enough to have sufficiently many samples from the large-degree region to get a statistically accurate estimate of the power-law part. This also explains why based on smaller data sets it is common to assume that  $\alpha$  is roughly one [4,10,33–36]. Comparing Figs. 7(a) and 7(b) shows that the convergence to the limiting value is significantly faster at point  $t = \frac{1}{2}$  than at point  $t = \frac{1}{\tau-1}$ .

### APPENDIX C: FROM HIDDEN VARIABLES TO DEGREES

In this paper, we focus on computing  $c(h)$ , the local clustering coefficient of a randomly chosen vertex with hidden-variable  $h$ . However, when studying local clustering in real-world data sets, we can only observe  $\bar{c}(k)$ , the local clustering coefficient of a vertex of degree  $k$ . In this Appendix, we show that, for the hidden-variable model, the difference between these two methods of computing the clustering coefficient is




FIG. 9. The probability that the degree of a vertex exceeds  $x$  in (a) the largest five networks of Table I and (b) the smallest five networks in Table I.




FIG. 10.  $\bar{c}(k)$  for several information [red; (a)–(c)], technological [green; (d)–(f)], and social [blue; (g)–(i)] real-world networks. (a) Hudong encyclopedia [12], (b) Baidu encyclopedia [12], (c) WordNet [13], (d) TREC-WT10g web graph [14], (e) Google web graph [11], (f) Internet on the autonomous systems level [11], (g) Catster and Dogster social networks [15], (h) Gowalla social network [11], and (i) Wikipedia communication network [11]. The different shadings indicate the theoretical boundaries of the regimes as in Fig. 2 with  $N$  and  $\tau$  as in Table I.

small and asymptotically negligible. We consider

$$c(h) = \frac{\int_1^{h_c} \int_1^{h_c} (h'h'')^{2-\tau} p(h,h')p(h,h'')p(h',h'')dh'dh''}{\left(\int_1^{h_c} x^{1-\tau} p(h,h')dh'\right)^2}. \tag{C1}$$

We define  $\bar{c}(k)$  as the average clustering coefficient over all vertices of degree  $k$ . By Ref. [32], the probability that a vertex with hidden-variable  $h$  has degree  $k$  equals

$$g(k|h) = \frac{e^{-h} h^k}{k!}. \tag{C2}$$

Then, by Ref. [32],

$$\bar{c}(k) = \begin{cases} \frac{1}{P(k)} \int_1^{h_c} \rho(h)c(h)g(k|h)dh, & k \geq 2, \\ 0, & k < 2, \end{cases} \tag{C3}$$

where  $\bar{c}(k) = 0$  for  $k < 2$  because a vertex with a degree less than 2 cannot be part of a triangle. Here,

$$P(k) = \int_1^{h_c} g(k|h)\rho(h)dh \tag{C4}$$

is the probability that a randomly chosen vertex has degree  $k$ .

First we consider the case where  $h > N^{(\tau-2)/(\tau-1)}$ . The Chernoff bound gives for the tails of the Poisson distribution that

$$\mathbb{P}(\text{Poi}(\lambda) > x) \leq e^{-\lambda} \left(\frac{e\lambda}{x}\right)^x, \quad x > \lambda, \tag{C5}$$

$$\mathbb{P}(\text{Poi}(\lambda) < x) \leq e^{-\lambda} \left( \frac{e\lambda}{x} \right)^x, \quad x < \lambda. \quad (\text{C6})$$

Let  $k(h)$  be the degree of a node with hidden-variable  $h$ . Then, for any  $M > 1$ ,

$$\sum_{k=Mh}^{\infty} g(k|h) \leq \left( \frac{e^{M-1}}{M^M} \right)^h, \quad (\text{C7})$$

and for any  $\delta \in (0, 1)$ ,

$$\sum_{k=1}^{\delta h} g(k|h) \leq \left( \frac{e^{\delta-1}}{\delta^\delta} \right)^h. \quad (\text{C8})$$

Because  $e^{x-1}/x^x < 1$  for  $x \neq 1$ , (C7) and (C8) tend to zero as  $h \rightarrow \infty$ . Therefore, for  $h$  large,

$$k(h) = h[1 + o(1)], \quad (\text{C9})$$

with high probability. Therefore, when  $k$  is large,

$$\bar{c}(k) \approx c(k). \quad (\text{C10})$$

Thus,  $c(h)$  is very similar to  $\bar{c}(k)$ .

On the other hand, for  $h \ll h_s^2/h_c$ ,

$$\sum_{h_s^2/h_c}^{\infty} g(k|h) \leq e^{-h} \left( \frac{eh}{h_s^2/h_c} \right)^{h_s^2/h_c}, \quad (\text{C11})$$

which is small by the assumption on  $h$ . Thus,

$$P(k) \approx \int_1^{h_s^2/h_c} g(k|h)\rho(h)dh. \quad (\text{C12})$$

Furthermore,  $c(h) = c(0)$  in this regime of  $h$ . This results in

$$\bar{c}(k) \approx \frac{c(0) \int_1^{h_s^2/h_c} \rho(h)g(k|h)dh}{\int_1^{h_s^2/h_c} \rho(h)g(k|h)dh} = c(0). \quad (\text{C13})$$

Therefore,  $\bar{c}(h) \approx c(h)$  also when  $h$  is small. Figure 8 shows that indeed the difference between  $\bar{c}(k)$  and  $c(k)$  is small. When  $\tau$  approaches 2, the difference becomes larger. We see that, for small values of  $k$ ,  $\bar{c}(k)$  and  $c(k)$  are not very close. This is due to the fact that (C1) does not take into account that a vertex with hidden-variable  $h$  may have less than two neighbors so that its local clustering is zero. In Ref. [31] we show how to adjust (A7) to account for this.

#### APPENDIX D: DEGREE DISTRIBUTIONS

Figure 9 shows the degree distributions of all ten networks of Table I.

#### APPENDIX E: ADDITIONAL DATA SETS

Figure 10 presents the clustering spectrum of nine additional data sets.

- 
- [1] R. Albert, H. Jeong, and A.-L. Barabási, *Nature (London)* **401**, 130 (1999).
- [2] M. Faloutsos, P. Faloutsos, and C. Faloutsos, in *Proceedings of the Conference on Applications, Technologies, Architectures, and Protocols for Computer Communication, SIGCOMM '99, Cambridge, MA, 1999* (ACM, New York, 1999), Vol. 29, pp. 251–262.
- [3] H. Jeong, B. Tombor, R. Albert, Z. N. Oltvai, and A.-L. Barabási, *Nature (London)* **407**, 651 (2000).
- [4] A. Vázquez, R. Pastor-Satorras, and A. Vespignani, *Phys. Rev. E* **65**, 066130 (2002).
- [5] R. van der Hofstad, G. Hooghiemstra, and D. Znamenski, *Electron. J. Probab.* **12**, 703 (2007).
- [6] M. E. J. Newman, S. H. Strogatz, and D. J. Watts, *Phys. Rev. E* **64**, 026118 (2001).
- [7] S. Janson, *Electron. J. Probab.* **14**, 86 (2009).
- [8] R. Pastor-Satorras and A. Vespignani, *Phys. Rev. Lett.* **86**, 3200 (2001).
- [9] R. Pastor-Satorras, A. Vázquez, and A. Vespignani, *Phys. Rev. Lett.* **87**, 258701 (2001).
- [10] E. Ravasz and A.-L. Barabási, *Phys. Rev. E* **67**, 026112 (2003).
- [11] J. Leskovec and A. Krevl, SNAP Datasets: Stanford large network dataset collection, <http://snap.stanford.edu/data> (2014), date of access: 14/03/2017.
- [12] X. Niu, X. Sun, H. Wang, S. Rong, G. Qi, and Y. Yu, in *The Semantic Web—ISWC 2011: 10th International Semantic Web Conference, Bonn, Germany, 2011*, edited by L. Aroyo *et al.* (Springer, Heidelberg/New York, 2011), pp. 205–220.
- [13] G. Miller and C. Fellbaum, Wordnet: An electronic lexical database (1998).
- [14] P. Bailey, N. Craswell, and D. Hawking, *Inf. Process. Manage.* **39**, 853 (2003).
- [15] J. Kunegis, in *Proceedings of the 22nd International Conference on World Wide Web, WWW '13 Companion* (ACM, New York, 2013), pp. 1343–1350.
- [16] P. Colomer-de Simón, M. A. Serrano, M. G. Beiró, J. I. Alvarez-Hamelin, and M. Boguñá, *Sci. Rep.* **3**, 2517 (2013).
- [17] M. A. Serrano and M. Boguñá, *Phys. Rev. Lett.* **97**, 088701 (2006).
- [18] M. Boguñá and R. Pastor-Satorras, *Phys. Rev. E* **68**, 036112 (2003).
- [19] M. A. Serrano, M. Boguñá, R. Pastor-Satorras, and A. Vespignani, in *Large Scale Structure and Dynamics of Complex Networks: From Information Technology to Finance and Natural Science*, edited by G. Caldarelli and A. Vespignani (World Scientific, Singapore, 2007), p. 35.
- [20] S. N. Dorogovtsev, *Phys. Rev. E* **69**, 027104 (2004).
- [21] A. Krot and L. O. Prokhorenkova, in *Algorithms and Models for the Web Graph*, edited by D. F. Gleich, J. Komjáthy, and N. Litvak, Lecture Notes in Computer Science Vol. 9479 (Springer, Cham, 2015), pp. 15–28.
- [22] G. Szabó, M. Alava, and J. Kertész, *Phys. Rev. E* **67**, 056102 (2003).

- [23] J. Park and M. E. J. Newman, *Phys. Rev. E* **70**, 066117 (2004).
- [24] B. Bollobás, S. Janson, and O. Riordan, *Random Struct. Alg.* **31**, 3 (2007).
- [25] T. Britton, M. Deijfen, and A. Martin-Löf, *J. Stat. Phys.* **124**, 1377 (2006).
- [26] I. Norros and H. Reittu, *Adv. Appl. Probab.* **38**, 59 (2006).
- [27] A. Clauset, C. R. Shalizi, and M. E. J. Newman, *SIAM Rev.* **51**, 661 (2009).
- [28] M. E. J. Newman, *SIAM Rev.* **45**, 167 (2003).
- [29] S. Dhara, R. van der Hofstad, and J. S. H. van Leeuwen, *Electron. J. Probab.* **22**, 66 (2016).
- [30] F. Chung and L. Lu, *Proc. Natl. Acad. Sci. USA* **99**, 15879 (2002).
- [31] R. van der Hofstad, A. J. E. M. Janssen, J. S. H. van Leeuwen, and C. Stegehuis, *Phys. Rev. E* **95**, 022307 (2017).
- [32] P. Colomer-de-Simon and M. Boguñá, *Phys. Rev. E* **86**, 026120 (2012).
- [33] M. A. Serrano and M. Boguñá, *Phys. Rev. E* **74**, 056114 (2006).
- [34] M. Catanzaro, G. Caldarelli, and L. Pietronero, *Phys. Rev. E* **70**, 037101 (2004).
- [35] J. Leskovec, *Dynamics of Large Networks* (ProQuest, Ann Arbor, MI, 2008).
- [36] D. Krioukov, M. Kitsak, R. S. Sinkovits, D. Rideout, D. Meyer, and M. Boguñá, *Sci. Rep.* **2**, 793 (2012).
- [37] M. Boguñá, C. Castellano, and R. Pastor-Satorras, *Phys. Rev. E* **79**, 036110 (2009).
- [38] A. J. E. M. Janssen and J. S. H. van Leeuwen, *Europhys. Lett.* **112**, 68001 (2015).
- [39] S. Maslov, K. Sneppen, and A. Zaliznyak, *Phys. A* **333**, 529 (2004).
- [40] J. Park and M. E. J. Newman, *Phys. Rev. E* **68**, 026112 (2003).
- [41] M. E. J. Newman, *Phys. Rev. Lett.* **89**, 208701 (2002).
- [42] M. E. J. Newman, *Phys. Rev. E* **67**, 026126 (2003).
- [43] B. Bollobás, *Eur. J. Combin.* **1**, 311 (1980).
- [44] R. van der Hofstad, *Random Graphs and Complex Networks* (Cambridge University Press, Cambridge, UK, 2017), Vol. 1.
- [45] R. van der Hofstad, G. Hooghiemstra, and P. Van Mieghem, *Random Struct. Alg.* **27**, 76 (2005).
- [46] M. Á. Serrano, D. Krioukov, and M. Boguñá, *Phys. Rev. Lett.* **100**, 078701 (2008).
- [47] C. Stegehuis, R. van der Hofstad, and J. S. H. van Leeuwen, *Phys. Rev. E* **94**, 012302 (2016).
- [48] R. Milo, N. Kashtan, S. Itzkovitz, M. E. J. Newman, and U. Alon, [arXiv:cond-mat/0312028](https://arxiv.org/abs/cond-mat/0312028).
- [49] F. Viger and M. Latapy, *Lecture Notes in Computer Science* (Springer, Berlin/Heidelberg, 2005), pp. 440–449.
- [50] C. I. D. Genio, H. Kim, Z. Toroczkai, and K. E. Bassler, *PLoS ONE* **5**, e10012 (2010).