

A geometric characterization of the classical Lie algebras

Citation for published version (APA):

Cuypers, H., & Fleischmann, Y. (2017). A geometric characterization of the classical Lie algebras. *arXiv*, Article 1707.02084. <https://arxiv.org/abs/1707.02084>

Document status and date:

Published: 07/07/2017

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

A geometric characterization of the classical Lie algebras

Hans Cuypers*, Yael Fleischmann †
 Department of Mathematics and Computer Science
 Eindhoven University of Technology
 P.O. Box 513, 5600 MB Eindhoven
 The Netherlands
 email: f.g.m.t.cuypers@tue.nl

July 10, 2017

1 Introduction

An element $x \neq 0$ in a Lie algebra \mathfrak{g} over a field \mathbb{F} with Lie product $[\cdot, \cdot]$ is called a *extremal element* if we have

$$[x, [x, \mathfrak{g}]] \subseteq \mathbb{F}x.$$

(Or, in case the characteristic of \mathbb{F} is two, when $[x, [x, \mathfrak{g}]] = 0$ and some additional conditions are satisfied. See Section 3.)

Long root elements in classical Lie algebras are examples of pure extremal elements. Arjeh Cohen *et al.* [8] initiated the investigation of Lie algebras generated by extremal elements in order to provide a geometric characterization of the classical Lie algebras generated by their long root elements. He and Gabor Ivanyos studied the so-called *extremal geometry* with as points the 1-dimensional subspaces of \mathfrak{g} generated by extremal elements of \mathfrak{g} and as lines the 2-dimensional subspaces of \mathfrak{g} all whose nonzero vectors are extremal.

*corresponding author

†present address: Institut für Mathematik, Warburger Str. 100, 33098 Paderborn, yael.fleischmann@math.uni-paderborn.de

‡This work is (partly) financed by the Netherlands Organisation for Scientific Research (NWO), project 613.000.905.

The main result of [7] and [6], based on the work of Kasikova and Shult [12], states that, if \mathfrak{g} is of finite dimension, simple and generated by extremal elements, then its extremal geometry either contains no lines or is the root shadow space of a spherical building. This raises the question whether one can recover the algebra \mathfrak{g} from its extremal geometry.

In [9], Cuypers, Roberts and Shpectorov show that in case the building is of simply laced type, the Lie algebra \mathfrak{g} is indeed a classical Lie algebra of the same type as the building and its extremal elements are the long root elements.

In this paper we prove the following result.

Theorem 1.1. *Let \mathfrak{g} be a simple Lie algebra generated by its set of extremal elements. If the extremal geometry of \mathfrak{g} is the root shadow space of a spherical building of rank at least 3, then \mathfrak{g} is, up to isomorphism, uniquely determined by its extremal geometry.*

As we already noticed, the long root elements in (forms of) Chevalley Lie algebras are extremal (see Section 7). So, as a corollary of the above result we obtain for most of the simple classical Lie algebras that they are characterized by their extremal geometry. The classes of simple classical Lie algebras, which are generated by their long root elements, but escape the above result, are those of relative type A_1 , B_2 , C_n or G_2 . The extremal geometries of these Lie algebras of type A_1 , B_2 or C_n have extremal geometries with no lines.

Simple Lie algebras \mathfrak{g} over a field \mathbb{F} generated by extremal elements whose extremal geometry has no lines have been studied in [10]. There it has been shown that either one can extend the underlying field \mathbb{F} quadratically and obtain an extremal geometry with lines, or \mathfrak{g} is a symplectic Lie algebra. So, it only remains to study Lie algebras whose extremal geometry is a root shadow space of type G_2 .

We notice that finite dimensional simple Lie algebras defined over algebraically closed fields of characteristic ≥ 7 and generated by extremal elements have been shown to be classical by Premet [13]. In Theorem 1.1, however, no restriction is imposed on the field \mathbb{F} , nor on the dimension of the Lie algebra. Our results are independent of the classification of spherical buildings of rank ≥ 3 .

The present paper is organized as follows. In Section 2 we provide definitions of, and some results on the geometric concepts used in this paper. In particular, we describe point-line geometries and root shadow spaces, and

present some results on embeddings of point-line geometries in projective spaces.

Section 3 is devoted to extremal elements. We present various results, mainly from [7], needed for the proof of our main results. This includes the construction and classification of the extremal geometries of finite dimensional simple Lie algebras generated by their extremal elements.

In Section 4 we start with the proof of Theorem 1.1, by proving that the embedding of the extremal geometry of a Lie algebra \mathfrak{g} as in the hypothesis of Theorem 1.1 into the projective space $\mathbb{P}(\mathfrak{g})$ is determined by the isomorphism type of the geometry. This proof is continued in Section 5, where we show that, given the isomorphism type of the extremal geometry of \mathfrak{g} and its embedding into $\mathbb{P}(\mathfrak{g})$, the Lie product is fixed up to a scalar.

In Section 6 we finish the proof of Theorem 1.1 by combining all the results obtained.

The final Section 7 is devoted to examples. In this section we show that the long root elements in Chevalley Lie algebras are extremal.

Acknowledgment. Parts of this paper can be found in the second author's PhD-thesis [11], which was written under supervision of Arjeh Cohen and the first author. We thank Arjeh Cohen, and also Kieran Roberts and Sergey Shpectorov, for many inspiring discussions on the topic.

2 Root shadow spaces and polarized embeddings

In this section we discuss the geometric results needed to prove our main result, Theorem 1.1.

A *point-line geometry* $(\mathcal{P}, \mathcal{L})$ consists of a set \mathcal{P} of *points* and a set \mathcal{L} of *lines*, being subsets of the point set \mathcal{P} of size at least 2. A point-line geometry is called a *partial linear space* if any two points are on at most one common line.

Let $(\mathcal{P}, \mathcal{L})$ be a point-line geometry. A subset X of \mathcal{P} is called a *subspace* of the point-line geometry $(\mathcal{P}, \mathcal{L})$ if every line meeting X in at least 2 points is fully contained in X . We often identify a subspace X with the point-line geometry $(X, \{l \in \mathcal{L} \mid l \subseteq X\})$.

The subset X is called a *geometric hyperplane*, or just *hyperplane*, if it meets every line in one point, or contains the line. Notice that a hyperplane is a subspace.

The collinearity graph of $(\mathcal{P}, \mathcal{L})$ is the graph with the points as vertices, and two points adjacent if and only if they are on a common line.

The distance between two points of $(\mathcal{P}, \mathcal{L})$ is the distance in the collinearity graph, and the diameter of $(\mathcal{P}, \mathcal{L})$ is the diameter of its collinearity graph. If $p \in \mathcal{P}$ is a point and d the diameter of $(\mathcal{P}, \mathcal{L})$, then p^\perp denotes the set of all points $q \in \mathcal{P}$ which are at distance $< d$ from p .

A *projective embedding* $\epsilon : \Gamma \rightarrow \mathbb{P}$ of a point-line geometry $\Gamma = (\mathcal{P}, \mathcal{L})$ into a projective space \mathbb{P} is an injective map ϵ from \mathcal{P} into the point set of the projective space \mathbb{P} such that the image of \mathcal{P} spans \mathbb{P} and the image of any line in \mathcal{L} comprises all projective points of a projective line in \mathbb{P} . Note that this induces an injection from \mathcal{L} into the line set of $\mathbb{P}(V)$. Moreover, if Γ can be embedded into a projective space, then Γ is a partial linear space.

Two embeddings $\epsilon : \Gamma \rightarrow \mathbb{P}$ and $\epsilon' : \Gamma \rightarrow \mathbb{P}'$ are called *isomorphic*, notation $\epsilon \cong \epsilon'$, if there is an isomorphism $\phi : \mathbb{P} \rightarrow \mathbb{P}'$ with $\epsilon' = \phi \circ \epsilon$.

In this paper we only consider embeddings into projective spaces which are obtained from vector spaces over a field.

Let V be a vector space and $\epsilon : \Gamma \rightarrow \mathbb{P}(V)$ be an embedding. Suppose $t : V \rightarrow W$ is a surjective semilinear transformation, with the property that $K := \ker(t)$ intersects the span $\langle \epsilon(p), \epsilon(q) \rangle$ for any pair p, q of points in \mathcal{P} trivially. Then ϵ can be carried onwards to the cosets of K , and we obtain an embedding $\epsilon' : \Gamma \rightarrow \mathbb{P}(W)$. We call ϵ' the *morphic image* of ϵ , or we say that ϵ' is *derived* from ϵ or ϵ *covers* ϵ' . In particular, $\epsilon'(p) := t(\epsilon(p))$ is a 1-space in W for all $p \in \mathcal{P}$. We write $\epsilon' = \epsilon/K$.

If all embeddings ϵ' of Γ can be obtained as morphic images from a fixed embedding ϵ , then this ϵ is called *absolute* or *absolutely universal*.

Let $\epsilon : \Gamma \rightarrow \mathbb{P}(V)$ be an arbitrary projective embedding of Γ into the projective space $\mathbb{P}(V)$ for some vector space V . We call ϵ *polarized* if and only if $\epsilon(p^\perp)$ is contained in a proper hyperplane of \mathbb{P} for all $p \in \mathcal{P}$.

The *radical* R_ϵ of a polarized embedding ϵ is the intersection

$$R_\epsilon := \bigcap_{p \in \mathcal{P}} \langle \epsilon(p^\perp) \rangle.$$

Here $\langle \epsilon(p^\perp) \rangle$ denotes the subspace of V generated by $\epsilon(p^\perp)$.

Lemma 2.1. *Let $\psi : \Gamma \rightarrow \mathbb{P}$ be a projective embedding covering a polarized embedding ϕ . Then ψ is polarized.*

Moreover, the kernel of the projection of ψ to ϕ is contained in the radical of ψ .

Proof. The first statement is trivial.

Now suppose, ϕ is an embedding of Γ into the projective space \mathbb{P}' . If the kernel K of the projection τ of ψ to ϕ is not contained in the radical of ψ ,

then there is an element $x \in \mathcal{P}$ such that $\langle \psi(x^\perp) \rangle$ does not contain K . But that implies that the image under τ of the hyperplane $\langle \psi(x^\perp) \rangle$ of \mathbb{P} is the full space $\mathbb{P}(W)$. This contradicts that ϕ is polarized. \diamond

Proposition 2.2. *Let ψ be a cover of a polarized embedding ϕ of Γ . If the radical of ϕ is trivial, then ϕ is isomorphic to ψ modulo its radical R_ψ .*

Proof. The projection τ of ψ onto ϕ maps the radical of ψ into the radical of ϕ . However, since the radical of ϕ is trivial, we find the kernel of τ to be the radical R_ψ . \diamond

Theorem 2.3. *Suppose Γ admits an absolutely universal embedding and a polarized embedding ϕ with trivial radical.*

Then any polarized embedding ψ of Γ covers ϕ .

Proof. Let ϵ be the absolutely universal embedding of Γ . By Lemma 2.1, ϵ is polarized and both ψ and ϕ are isomorphic to the quotient of ϵ by a subspace K_ψ and K_ϕ , respectively, of its radical R_ϵ .

Since the radical of ϕ is trivial, we find K_ϕ to be equal to R_ϵ . But this implies that $K_\psi \subseteq K_\phi$ and ψ clearly covers ϕ . \diamond

Theorem 2.4. *Suppose $(\mathcal{P}, \mathcal{L})$ is a point-line geometry admitting an absolutely universal embedding ϵ . If ϕ is a polarized embedding of $(\mathcal{P}, \mathcal{L})$, then*

$$\phi/R_\phi \cong \psi/R_\epsilon.$$

In particular, if ϕ has trivial radical, it is unique up to isomorphism.

Proof. Lemma 2.1 shows that $\phi \cong \epsilon/R$ for some $R \subseteq R_\epsilon$. The radical of ϵ/R is $R_\epsilon/R \cong R_\phi$, and we get

$$\phi/R_\phi \cong (\epsilon/R)/R_\phi \cong (\epsilon/R)/(R_\epsilon/R) \cong \epsilon/R_\epsilon.$$

\diamond

We notice that similar results have been obtained by Blok [1].

The point-line geometries we want to consider in this paper are the so-called root shadow spaces of spherical buildings. We view a building as chamber system and use the notation of [19].

Let Δ be an irreducible spherical building of type X_n and with corresponding root system Φ . (In the case that the building is of type BC_n , we only consider the type B_n .) Fix the index set $I = \{1, \dots, n\}$ of Δ .

For any subset J of I , we define a J -shadow in Δ to be an $(I \setminus J)$ -residue. If $j \in J$, then a j -line is the set of all J -shadows containing chambers from a given j -panel. Let \mathcal{P} be the set of J -shadows, called *points* and \mathcal{L} be the set of j -lines for $j \in J$. We say that the point-line geometry $(\mathcal{P}, \mathcal{L})$ is the J -shadow space (or *shadow space of type $X_{n,J}$*). If J is the set $\{j\}$, then we write $X_{n,j}$ instead of $X_{n,J}$.

Of importance to us are the *root shadow spaces*. These are the shadow spaces of type $X_{n,J}$, (or $X_{n,j}$, when $J = \{j\}$) where J is a so-called root set, which can be defined as follows. Fix a base $B = \{\alpha_1, \dots, \alpha_n\}$ for Φ and denote the longest root with respect to B by α . Then the root set J is the subset I consisting of all elements $i \in I$ with $(\alpha, \alpha_i) \neq 0$. The root set is $J = \{1, n\}$ for A_n , $J = \{2\}$ for $(BC)_n$, D_n , E_6 and G_2 , $J = \{1\}$ for E_7 and F_4 , and $J = \{8\}$ for E_8 . (We use the Bourbaki labelling of the Dynkin diagrams as in [19].)

We provide some more concrete descriptions of some examples of root shadow spaces.

Example 2.5. Let \mathbb{P} be a projective geometry of projective dimension at least 2 and suppose \mathbb{H} is a set of hyperplanes of \mathbb{P} forming a subspace of the dual of \mathbb{P} such that the intersection of all hyperplanes in \mathbb{H} is empty. The set P is the set of incident point-hyperplane pairs (p, H) where $p \in \mathbb{P}$ and $H \in \mathbb{H}$. As lines in L we take all subsets of P consisting of all $(p, H) \in P$ with p running over the points of a fixed line ℓ of \mathbb{P} and H a fixed hyperplane in \mathbb{H} containing ℓ , or, dually, p being a fixed point inside a fixed hyperline K which is the intersection of two elements from \mathbb{H} and H running over all hyperplanes containing K .

If the dimension of \mathbb{P} is $n < \infty$, then (P, L) is a root shadow space of type $A_{n, \{1, n\}}$.

Example 2.6. Let V be a vector space over a field \mathbb{F} of characteristic $\neq 2$, and Q a nondegenerate quadratic form on V with associated symmetric bilinear form f . Assume that the Witt index of Q is $n \geq 3$.

Let P be the set of all isotropic 2-spaces of V and L the set of all subsets of P consisting of all isotropic 2-spaces on a fixed isotropic 1-space and contained in a fixed isotropic 3-space. Then (P, L) is a root shadow space of type $BC_{n,2}$ (or $D_{n+1,2}$ in case Q is split).

More generally, if Π is a nondegenerate polar space of rank n at least 3, then let P be the set of all lines of Π and L the set of all pencils of lines on a point and inside a singular plane. Then, for finite n , the space (P, L) is a root shadow space of type $BC_{n,2}$ or $D_{n+1,2}$.

We recall some well known properties of root shadow spaces and their collinearity graph.

Proposition 2.7. [4] *Let $\Gamma = (\mathcal{P}, \mathcal{L})$ be a root shadow space of a spherical building of rank at least 2. Then the following holds:*

- (a) Γ is a partial linear space.
- (b) The diameter of the collinearity graph is ≤ 3 .
- (c) Let p be a point. The set p^\perp is a hyperplane of Γ .

We close this section with the following result of A. Kasikova and E. Shult and an application of Theorem 2.4:

Theorem 2.8 ([12]). *Let $\Gamma = (\mathcal{E}, \mathcal{F})$ be an embeddable root shadow space of type $\text{BC}_{n,2}$ ($n \geq 3$), $\text{D}_{n,2}$ ($n \geq 4$), $\text{E}_{6,2}$, $\text{E}_{7,1}$, $\text{E}_{8,8}$ or $\text{F}_{4,1}$. Then Γ admits an absolutely universal embedding.*

Proof. Kasikova and Shult prove the existence for each of these cases in [12]. The case $\text{BC}_{n,2}$ for $n \geq 4$ can be found in [12, 4.8] and $\text{BC}_{3,2}$ in 4.7, $\text{D}_{n,2}$ for $n \geq 5$ is covered in 4.5 and the special case $\text{D}_{4,2}$ is treated in 4.1, $\text{E}_{6,2}$, $\text{E}_{7,1}$ and $\text{E}_{8,8}$ in 4.11, and $\text{F}_{4,1}$ in 4.9. \diamond

Corollary 2.9. *Let $\Gamma = (\mathcal{E}, \mathcal{F})$ be a root shadow space of type $\text{BC}_{n,2}$ ($n \geq 3$), $\text{D}_{n,2}$ ($n \geq 4$), $\text{E}_{6,2}$, $\text{E}_{7,1}$, $\text{E}_{8,8}$ or $\text{F}_{4,1}$ admitting a polarized embedding. Then Γ admits, up to isomorphism, a unique polarized embedding with trivial radical.*

3 Extremal elements

In this section we provide some definitions and collect some results on extremal elements, mainly from [7].

Definition 3.1. Let \mathfrak{g} be a Lie algebra over the field \mathbb{F} . A nonzero element $x \in \mathfrak{g}$ is called *extremal* if there is a map $g_x : \mathfrak{g} \rightarrow \mathbb{F}$, called the *extremal form* on x , such that

$$[x, [x, y]] = 2g_x(y)x \tag{3.1}$$

and moreover

$$[[x, y], [x, z]] = g_x([y, z])x + g_x(z)[x, y] - g_x(y)[x, z] \tag{3.2}$$

and

$$[x, [y, [x, z]]] = g_x([y, z])x - g_x(z)[x, y] - g_x(y)[x, z] \quad (3.3)$$

for every $y, z \in \mathfrak{g}$.

The last two identities are called the *Premet identities*. If the characteristic of \mathbb{F} is not 2, then the Premet identities follow from Equation 3.1. See [7].

As a consequence, $[x, [x, \mathfrak{g}]] \subseteq \mathbb{F}x$ for an extremal $x \in \mathfrak{g}$. We call $x \in \mathfrak{g}$ a *sandwich* if $[x, [x, y]] = 0$ and $[x, [y, [x, z]]] = 0$ for every $y, z \in \mathfrak{g}$. So, a sandwich is an element x for which the extremal form g_x can be chosen to be identically zero. We introduce the convention that g_x is *identically zero whenever x is a sandwich in \mathfrak{g}* . An extremal element is called *pure* if it is not a sandwich.

We denote the set of extremal elements of a Lie algebra by $E(\mathfrak{g})$ or, if \mathfrak{g} is clear from the context, by E . Accordingly, we denote the set $\{\mathbb{F}x | x \in E(\mathfrak{g})\}$ of *extremal points* in the projective space on \mathfrak{g} by $\mathcal{E}(\mathfrak{g})$ or \mathcal{E} .

Example 3.2. Let V be a vector space over a field \mathbb{F} with dual space V^* . Suppose W^* is a subspace of V^* annihilating V .

On $V \otimes W^*$ we can define a Lie bracket by linear extension of the following product for pure tensors $v \otimes \phi$ and $w \otimes \psi$:

$$[v \otimes \phi, w \otimes \psi] = (v \otimes \psi)\phi(w) - (w \otimes \phi)\psi(v).$$

A pure tensor $v \otimes \phi$ is called *singular* if $\phi(v) = 0$. Let g be the \mathbb{F} -bilinear form on $V \otimes W^*$ defined by

$$g(v \otimes \phi, w \otimes \psi) = -\psi(v)\phi(w)$$

for $v \otimes \phi, w \otimes \psi \in V \otimes W^*$. Then for all singular pure tensors $v \otimes \phi$ and tensors $w \otimes \psi$ we have

$$\begin{aligned} [v \otimes \phi, [v \otimes \phi, w \otimes \psi]] &= [v \otimes \phi, \phi(w)v \otimes \psi - \psi(v)w \otimes \phi] \\ &= -\psi(v)\phi(w)v \otimes \phi - \psi(v)\phi(w)v \otimes \phi \\ &= -2\psi(v)\phi(w)v \otimes \phi \\ &= 2g(v \otimes \phi, w \otimes \psi)v \otimes \phi. \end{aligned}$$

In characteristic $\neq 2$ this implies that the singular pure tensors are extremal elements in $V \otimes W^*$. This also holds true in characteristic 2. (It is straightforward, but somewhat tedious, to check that the Premet identities also hold.)

An element $v \otimes \phi \in V \otimes V^*$ acts linearly on V by

$$v \otimes \phi(w) = \phi(w)v$$

for all $w \in V$. This provides an isomorphism between $V \otimes V^*$ and the finitary general linear Lie algebra $\mathfrak{fgl}(V)$. (A linear map is finitary if its kernel has finite codimension.) The singular pure tensors in $V \otimes V^*$ (or $V \otimes W^*$) generate (subalgebras of) $\mathfrak{fsl}(V)$, the finitary special Lie algebra, or $\mathfrak{sl}(V)$ in case V has finite dimension.

Example 3.3. Let V be a vector space over the field \mathbb{F} of characteristic $\neq 2$ and $Q : V \rightarrow \mathbb{F}$ a nondegenerate quadratic form with associated bilinear form b . For $v \in V$ we denote by b_v the linear form $w \in V \mapsto b(v, w)$.

Then, for $v, w \in V$ consider the following element in $V \otimes V^*$:

$$s_{v,w} = v \otimes b_w - w \otimes b_v.$$

If v and w are linearly independent and $Q(v) = Q(w) = b(v, w) = 0$ then we call $s_{v,w}$ a *Siegel element*. Now let $s_{v,w}$ be a Siegel element, then for all $x, y \in V$ we have

$$\begin{aligned} b(s_{v,w}(x), y) &= b(b(w, x)v - b(v, x)w, y) \\ &= b(w, x)b(v, y) - b(v, x)b(w, y) \\ &= b(x, b(y, v)w - b(y, w)v) \\ &= -b(x, s_{v,w}(y)). \end{aligned}$$

The Siegel elements generate (and even linearly span) the orthogonal Lie algebra $\mathfrak{so}(V, b) = \{s \in \mathfrak{fsl}(V) \mid b(s(x), y) = -b(s(y), x)\}$.

A straightforward computation yields

$$[s_{v,w}, [s_{v,w}, s_{x,y}]] = (b(w, x)b(w, y) - b(v, x)b(v, y))s_{v,w},$$

which shows that $s_{v,w}$ is extremal.

For $x \in E$ and $\lambda \in \mathbb{F}$ we define the map $\exp(x, \lambda) : \mathfrak{g} \rightarrow \mathfrak{g}$ by

$$\exp(x, \lambda)y = y + \lambda[x, y] + \lambda^2 g_x(y)x$$

for all $y \in \mathfrak{g}$.

Proposition 3.4. [7, Lemma 15] *Let $x \in E$ be a pure and $\lambda \in \mathbb{F}$. Then $\exp(x, \lambda)$ is an automorphism of \mathfrak{g} .*

Let $x \in E$ be pure. By $\text{Exp}(x)$ we denote the set $\{\exp(x, \lambda) \mid \lambda \in \mathbb{F}\}$. Since, for $\lambda, \mu \in \mathbb{F}$ we have $\exp(x, \lambda)\exp(x, \mu) = \exp(x, \lambda + \mu)$, we find that $\text{Exp}(x)$ is a subgroup of $\text{Aut}(\mathfrak{g})$ isomorphic to the additive group of \mathbb{F} .

Clearly, $\text{Exp}(x) = \text{Exp}(\lambda x)$ for $\lambda \in \mathbb{F}^*$. Therefore we can define $\text{Exp}(\langle x \rangle)$ to be equal to $\text{Exp}(x)$.

Proposition 3.5. [7, Lemma 24-27] *For $x, y \in E$ we have one of the following:*

- (a) $\mathbb{F}x = \mathbb{F}y$;
- (b) $[x, y] = 0$ and $\lambda x + \mu y \in E \cup \{0\}$ for all $\lambda, \mu \in \mathbb{F}$;
- (c) $[x, y] = 0$ and $\lambda x + \mu y \in E$ only if $\lambda = 0$ or $\mu = 0$;
- (d) $z := [x, y] \in E$, and x, z and y, z are as in case (b);
- (e) $\langle x, y \rangle$ is isomorphic to $\mathfrak{sl}_2(\mathbb{F})$.

Moreover, $g_x(y) \neq 0$ if and only if $\langle x, y \rangle$ is isomorphic to $\mathfrak{sl}_2(\mathbb{F})$.

In each of the cases described in the above proposition we find that the subalgebra generated by two extremal elements x and y is linearly spanned by all its extremal elements. Indeed, this is trivial in the cases that x and y commute. If x and y do not commute, then $\exp(x, 1)y = y + [x, y] + g_x(y)y$ is extremal and spans together with x and y the subalgebra $\langle x, y \rangle$.

Moreover, inside \mathfrak{sl}_2 we can see that $g_x(y) = g_y(x)$. This leads to the the following.

Proposition 3.6. [7, Proposition 20] *Let \mathfrak{g} be generated by $E(\mathfrak{g})$. Then \mathfrak{g} is linearly spanned by E and there is a unique bilinear symmetric form $g : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{F}$ such that the linear form g_x coincides with $y \mapsto g(x, y)$ for each $x \in E$. Moreover, this form is associative in the sense that $g(x, [y, z]) = g([x, y], z)$ for all $x, y, z \in \mathfrak{g}$.*

The form g is called the *extremal form* on \mathfrak{g} . As the form g is associative, its radical $\text{rad}(g) := \{u \in \mathfrak{g} \mid g_u(z) = 0 \forall z \in \mathfrak{g}\}$ is an ideal in \mathfrak{g} .

Notice that the extremal form f from [8] satisfies $f = 2g$.

Definition 3.7. For $x, y \in E$ extremal elements we define

$$(x, y) \in \begin{cases} E_{-2}, & \iff \mathbb{F}x = \mathbb{F}y, \\ E_{-1}, & \iff [x, y] = 0, (x, y) \notin E_{-2} \text{ and } \mathbb{F}x + \mathbb{F}y \subseteq E \cup \{0\}, \\ E_0, & \iff [x, y] = 0 \text{ and } (x, y) \notin E_{-2} \cup E_{-1}, \\ E_1, & \iff [x, y] \neq 0 \text{ and } g(x, y) = 0, \\ E_2, & \iff g(x, y) \neq 0. \end{cases}$$

For the corresponding extremal points $\langle x \rangle, \langle y \rangle$, we define

$$(\langle x \rangle, \langle y \rangle) \in \mathcal{E}_i \iff (x, y) \in E_i.$$

Let $x \in E$. Then $y \in E_i(x)$ denotes that $(x, y) \in E_i$. By $E_{\leq i}(x)$ we denote the set $\bigcup_{-2 \leq j \leq i} E_j(x)$. Similarly, if $x \in \mathcal{E}$, then $\mathcal{E}_i(x)$ consists of all y with $(x, y) \in \mathcal{E}_i$, and $\mathcal{E}_{\leq i}(x)$ denotes $\bigcup_{-2 \leq j \leq i} \mathcal{E}_j(x)$.

Proposition 3.8. *If $x, y \in E$ with $g_x(y) \neq 0$, then $\langle x \rangle$ and $\langle y \rangle$ are in the same $\text{Aut}(\mathfrak{g})$ -orbit. In particular, if the graph on $(\mathcal{E}, \mathcal{E}_2)$ is connected, then the set of all extremal points is a single $\text{Aut}(\mathfrak{g})$ -orbit.*

Proof. Let $x, y \in E$ with $g_x(y) = g_y(x) \neq 0$. Then, after replacing y by a scalar multiple, we can assume $g_x(y) = g_y(x) = 1$. But then $\exp(x, 1)y = y + [x, y] + x = \exp(y, 1)x$. So, $\exp(y, -1)\exp(x, 1)$ maps y to x . The proposition follows immediately. \diamond

Definition 3.9. Let \mathcal{E} be the set of extremal points of the Lie algebra \mathfrak{g} and let \mathcal{F} be the set of projective lines $\mathbb{F}x + \mathbb{F}y$ for $(x, y) \in \mathcal{E}_{-1}$. Hereby, we identify a 2-space with the set of 1-spaces it contains. Then the point-line space $(\mathcal{E}, \mathcal{F})$ is called the *extremal geometry* of \mathfrak{g} . We denote it by $\Gamma(\mathfrak{g})$, or in case it is clear what \mathfrak{g} is, by Γ .

The *rank* of $\Gamma(\mathfrak{g})$ is the maximal dimension (as a linear subspace of \mathfrak{g}) of a subspace X of $\Gamma(\mathfrak{g})$ in which any two points are collinear.

Example 3.10. Let V be a vector space over a field \mathbb{F} and W^* a subspace of V^* annihilating V . Let \mathcal{E} be the set of extremal points of $\langle v \otimes \phi \rangle$ where $v \in V$ and $\phi \in W^*$ with $\phi(v) = 0$. Then each extremal point $\langle v \otimes \phi \rangle$ corresponds to an incident point-hyperplane pair $(\langle v \rangle, \ker(\phi))$ of $\mathbb{P}(V)$. The extremal geometry with point set \mathcal{E} is isomorphic to the geometry described in Example 2.5. In particular, if V has dimension $n + 1 < \infty$, then the extremal geometry is a root shadow space of type $A_{n, \{1, n\}}$.

The extremal geometry is embedded into a subspace the projective space $\mathbb{P}(V \otimes W^*)$.

Example 3.11. Consider a vector space V over a field \mathbb{F} of characteristic $\neq 2$, and equipped with a nondegenerate quadratic form Q of Witt index at least 2.

Then we can consider the extremal geometry whose point set is the set of extremal points generated by Siegel elements. To each such extremal

point corresponds a unique isotropic line on (V, Q) . The extremal geometry is isomorphic to the geometry described in Example 2.6. Thus, if the Witt index of Q is finite and greater than 2, then the extremal geometry is a root shadow space of type $BC_{n,2}$ or $D_{n+1,2}$.

Again the extremal geometry is embedded into the subspace of $\mathbb{P}(V \otimes V^*)$ generated by the extremal points.

Proposition 3.12. [7] *Let $x, y \in \mathcal{E}$. Then we have the following:*

- (a) $(x, y) \in \mathcal{E}_{-2} \Leftrightarrow x = y$;
- (b) $(x, y) \in \mathcal{E}_{-1} \Leftrightarrow x$ and y are distinct but collinear in $\Gamma(\mathfrak{g})$;
- (c) $(x, y) \in \mathcal{E}_0 \Leftrightarrow x$ and y are at distance 2 and have ≥ 2 common neighbors in $\Gamma(\mathfrak{g})$;
- (d) $(x, y) \in \mathcal{E}_1 \Leftrightarrow x$ and y are at distance 2 and have a unique common neighbor in $\Gamma(\mathfrak{g})$, the point $[x, y]$;
- (e) $(x, y) \in \mathcal{E}_2 \Leftrightarrow x$ and y are at distance 3 in $\Gamma(\mathfrak{g}) \Leftrightarrow \langle x, y \rangle \cong \mathfrak{sl}_2(\mathbb{F})$.

The following lemma characterizes collinearity.

Lemma 3.13. [7, Lemma 27] *Let $x, y \in E$ be linearly independent. Then $(x, y) \in E_{-1} \Leftrightarrow$ there are $\lambda, \mu \in \mathbb{F}^*$ with $\lambda x + \mu y \in E$.*

We will use the following fundamental result of Cohen and Ivanyos (see [6] and [7]) in the next section.

Theorem 3.14. [7, Theorem 28] *Suppose that the extremal geometry Γ of a Lie algebra \mathfrak{g} , generated by its set of extremal elements and equipped with a nondegenerate extremal form g , has finite rank. Then a connected component of $\Gamma(\mathfrak{g})$ is isomorphic to a root shadow space of type $A_{n, \{1, n\}}$ ($n \geq 2$), $BC_{n,2}$ ($n \geq 3$), $D_{n,2}$ ($n \geq 4$), $E_{6,2}$, $E_{7,1}$, $E_{8,8}$, $F_{4,1}$ or $G_{2,2}$ or consists of a single point.*

Furthermore \mathfrak{g} is a direct sum of the Lie algebras generated by the connected components of Γ . Moreover, in each such Lie subalgebra the extremal points form a single orbit under the automorphism group of the Lie subalgebra.

Proof. The result directly follows from Theorem 28 of [7], except for the last sentence. But this follows from the fact that the subgraph $(\mathcal{E}, \mathcal{E}_2)$ induced on the connected components of Γ is connected, cf. Lemma 5 of [7], and Proposition 3.8. \diamond

Note that the labeling of the Coxeter diagrams follows [3].

4 The embedding

We fix the properties that we assume for Lie algebras in this section. If not mentioned otherwise, any Lie algebra in the remainder of this section is supposed to fulfill the conditions of Setting 4.1.

Setting 4.1. By \mathfrak{g} we denote a Lie algebra generated by its set E of extremal elements and with nondegenerate extremal form g . By $\Gamma = (\mathcal{E}, \mathcal{F})$, we denote the extremal geometry of \mathfrak{g} . We assume Γ to be connected and not a single point. So, in particular we assume that $\mathcal{E}_{-1} \neq \emptyset$.

Considering \mathfrak{g} as a vector space, and its projective geometry $\mathbb{P}(\mathfrak{g})$, the *natural* projective embedding of the extremal geometry $\Gamma = (\mathcal{E}, \mathcal{F})$ into $\mathbb{P}(\mathfrak{g})$ is defined to be the injection

$$\phi : \mathcal{E} \rightarrow \mathbb{P}(\mathfrak{g}), \text{ with } \phi(x) = x \text{ for } x \in \mathcal{E}.$$

By definition of lines in \mathcal{F} , the image under ϕ of a line $l \in \mathcal{F}$ is the full set $\phi(l)$ of points of some projective line in $\mathbb{P}(\mathfrak{g})$. Moreover, as the extremal points in \mathcal{E} linearly span \mathfrak{g} (see 3.6), the set $\phi(\mathcal{E})$ spans $\mathbb{P}(\mathfrak{g})$. So, ϕ is indeed a projective embedding.

Lemma 4.2. *The embedding ϕ is polarized.*

Proof. For each $x \in \mathcal{E}$ we find $\phi(\mathcal{E}_{\leq 1}(x))$ to be contained in the hyperplane $\{y \in \mathfrak{g} \mid g(x, y) = 0\}$. See Proposition 3.12. \diamond

Theorem 4.3. *Suppose \mathfrak{g}_1 and \mathfrak{g}_2 are two Lie algebras as in Setting 4.1, each of them generated by its set of extremal elements and equipped with a nondegenerate extremal form. Assume their corresponding extremal geometries Γ_1 and Γ_2 are isomorphic to each other and admit an absolutely universal embedding. Then their natural embeddings are isomorphic.*

Proof. We can apply the results of Section 2 and find by 2.4 that the natural embeddings ϕ_1 and ϕ_2 are isomorphic, provided their radicals are trivial.

Since, for $i = 1, 2$, the radical R_i of embedding ϕ_i is the intersection of all the subspaces $\langle \mathcal{E}_{\leq 1}(x) \rangle$ where x runs through the set of extremal points of \mathfrak{g}_i , we find these radicals to be contained in the radical of the extremal form g_i of \mathfrak{g}_i .

As the radicals of the forms g_1 and g_2 are trivial by assumptions, the radicals of the embeddings are also trivial. \diamond

The above has the following consequences.

Corollary 4.4. *Let \mathfrak{g}_1 and \mathfrak{g}_2 be Lie algebras as in 4.1. Assume the corresponding extremal geometries Γ_1 and Γ_2 are isomorphic to each other and to a connected root shadow space of type $BC_{n,2}$ ($n \geq 3$), $D_{n,2}$ ($n \geq 4$), $E_{6,2}$, $E_{7,1}$, $E_{8,8}$, or $F_{4,1}$. Then their natural embeddings are isomorphic.*

Proof. By Corollary 2.9 we find that Γ_i , with $i = 1, 2$, admits an absolutely universal embedding. So Theorem 4.3 applies. \diamond

Remark 4.5. For root shadow spaces of type $A_{n,\{1,n\}}$ and of type $G_{2,2}$ it is not known whether they admit an absolutely universal embedding.

The results of Völklein [18] imply that the natural embeddings of the extremal geometries of type $A_{n,\{1,n\}}$ and of type $G_{2,2}$ of the Chevalley Lie algebras of type A_n and G_2 do have a universal cover (see Section 7).

Blok and Pasini [2] obtain some partial results on embeddings of the geometries of type $A_{n,\{1,n\}}$ under some extra conditions on the underlying field. Van Maldeghem and Thas [14] show that the natural embedding of finite dual Cayley hexagons in the Chevalley Lie algebra is, up to isomorphism, the unique embedding of the hexagon in dimension ≥ 14 .

In the next section we will prove that given an embedding of the extremal geometry of a Lie algebra there is, up to a scalar multiple, at most one Lie bracket corresponding to it. This implies that the Lie structures \mathfrak{g}_1 and \mathfrak{g}_2 , in the cases considered in 4.4 are isomorphic.

5 Uniqueness of the Lie product

Let \mathfrak{g} be a Lie algebra generated by its set of extremal elements E with respect to a nondegenerate extremal form g . As before let $\Gamma = (\mathcal{E}, \mathcal{F})$ be the extremal geometry of \mathfrak{g} . In the previous section we have seen that the natural embedding of Γ into the projective space $\mathbb{P}(\mathfrak{g})$ is uniquely determined (up to isomorphism), if Γ admits an absolutely universal embedding. Our goal is to prove that not only the embedding of the extremal geometry is uniquely determined, but that also the Lie product is determined up to scalar multiples.

In this section we assume the following.

Setting 5.1. Let \mathfrak{g} be a Lie algebra as in 4.1, with Γ naturally embedded into the projective space $\mathbb{P}(\mathfrak{g})$.

Let $[\cdot, \cdot]$ denote the Lie product on \mathfrak{g} . We consider a second Lie product $[\cdot, \cdot]_1$ defining a Lie algebra \mathfrak{g}_1 on the vector space underlying \mathfrak{g} with extremal form g_1 and also Γ as extremal geometry.

We want to show that $[\cdot, \cdot]_1 = \lambda[\cdot, \cdot]$ for some fixed $\lambda \in \mathbb{F}^*$. Notice that the relations \mathcal{E}_i with $-2 \leq i \leq 2$ are determined by Γ (see 3.12). So elements $x, y \in \mathcal{E}$ are in relation \mathcal{E}_i in \mathfrak{g} if and only if they are in relation \mathcal{E}_i in \mathfrak{g}_1 .

Lemma 5.2. *Let $(x, y) \in E_{\leq 1}$, then there is a $\lambda \in \mathbb{F}^*$ such that $[x, y]_1 = \lambda[x, y]$.*

Proof. If $(x, y) \in E_{\leq 0}$, then $[x, y]_1 = 0 = [x, y]$.

If $(x, y) \in E_1$, then, by Proposition 3.12, both $[x, y]$ and $[x, y]_1$ span the unique point in \mathcal{E} collinear to both $\langle x \rangle$ and $\langle y \rangle$. So indeed, there is a $\lambda \in \mathbb{F}^*$ with $[x, y]_1 = \lambda[x, y]$. \diamond

Now we concentrate on the subalgebra of \mathfrak{g} generated by a pair of points in \mathcal{E}_2 . Such a subalgebra is isomorphic to $\mathfrak{sl}_2(\mathbb{F})$.

Lemma 5.3. *Let $(x, y) \in E_2$ be a generating a subalgebra \mathfrak{h} of \mathfrak{g} . Then there is a $\lambda \in \mathbb{F}^*$ such that for all $v, w \in \mathfrak{h}$ we have $[v, w]_1 = \lambda[v, w]$.*

Proof. Without loss of generality suppose that $g(x, y) = 1$. Inside both \mathfrak{g} and \mathfrak{g}_1 the elements x and y generate a subalgebra \mathfrak{h} and \mathfrak{h}_1 , respectively, isomorphic to \mathfrak{sl}_2 .

We first prove that \mathfrak{h} and \mathfrak{h}_1 are equal as linear subspaces of \mathfrak{g} .

Inside $\Gamma(\mathfrak{g})$ we take two distinct lines l_1 and l_2 on $\langle x \rangle$ with $[l_1, l_2] = \langle x \rangle$. Notice that such lines exist. For $i = 1, 2$, fix a point $\langle x_i \rangle$ on l_i which is at distance 2 from $\langle y \rangle$. Let $y_i := [y, x_i]$. Then for each point $\langle z_1 \rangle$ on the line through $\langle x_1 \rangle$ and $\langle y_1 \rangle$, there is a point $\langle z_2 \rangle$ on the line through $\langle x_2 \rangle$ and $\langle y_2 \rangle$ which is in relation \mathcal{E}_1 with $\langle z_1 \rangle$. This follows from the observation that the group $\langle \text{Exp}(x), \text{Exp}(y) \rangle$ leaves the lines $\langle x_1, y_1 \rangle$ and $\langle x_2, y_2 \rangle$ invariant and is transitive on the points of these lines.

We claim that both in \mathfrak{g}_1 and \mathfrak{g}_2 the unique common neighbour $\langle z \rangle = \langle [z_1, z_2] \rangle$ of $\langle z_1 \rangle$ and $\langle z_2 \rangle$ is inside the subalgebra generated by x and y .

Indeed, within $\text{Aut}(\mathfrak{g})$ we find that the elements of $\text{Exp}(\langle x \rangle)$ fix, for $i = 1, 2$, the point $\langle x_i \rangle$ as well as the line spanned by x_i and y_i . Moreover, $\text{Exp}(\langle x \rangle)$ acts transitively on the points of this line different from $\langle x_i \rangle$. Thus there is an element $e \in \text{Exp}(\langle x \rangle)$ that maps $\langle y_1 \rangle$ to $\langle z_1 \rangle$. As e leaves the line spanned by x_2 and y_2 invariant, it maps $\langle y_2 \rangle$ to the unique point on this line which is at distance 2 from $\langle z_1 \rangle$, the point $\langle z_2 \rangle$. But then $\langle y \rangle$ is mapped to $\langle z \rangle$ by the element e , as $\langle z \rangle$ is the unique common neighbor of $\langle z_1 \rangle$ and $\langle z_2 \rangle$. This clearly implies that $\langle z \rangle$ is inside the subalgebra \mathfrak{h} of \mathfrak{g} generated by x and y . In particular, x, y and z linearly span the subalgebra \mathfrak{h} of \mathfrak{g} generated by x and y .




Figure 1: Configuration of points

But similarly, these three elements are also contained in the subalgebra \mathfrak{h}_1 of \mathfrak{g}_1 generated by x and y and span this subalgebra. So, the subalgebras \mathfrak{h} and \mathfrak{h}_1 have to coincide as linear subspaces. The above proves more. Indeed, it shows that the point $\langle z \rangle$ is in the $\text{Exp}(\langle x \rangle)$ -orbit of $\langle y \rangle$ both with respect to $[\cdot, \cdot]$ and to $[\cdot, \cdot]_1$. Actually, these two orbits have to be equal.

In \mathfrak{g} this orbit, together with $\langle x \rangle$, consists of all 1-spaces spanned by elements

$$ax + by + c[x, y],$$

where $a, b, c \in \mathbb{F}$ satisfy $ab = c^2$.

Now, suppose

$$[x, y]_1 = \alpha x + \beta y + \gamma[x, y],$$

for some fixed $\alpha, \beta, \gamma \in \mathbb{F}$. Note that $[x, y]_1 \neq 0 \neq [x, y]$, since $(x, y) \in E_2$ with respect to both Lie products by the general assumptions in 5.1. Note moreover that $\gamma \neq 0$, since otherwise 3.13 leads to a contradiction.

The images of y under elements from $\text{Exp}(\langle x \rangle)$, but now with respect to $[\cdot, \cdot]_1$, are of the form

$$y + \lambda[x, y]_1 + \lambda^2 g_1(x, y)x = (\lambda^2 g_1(x, y) + \lambda\alpha)x + (1 + \lambda\beta)y + \lambda^2 \gamma^2 [x, y]$$

where $\lambda \in \mathbb{F}$. These elements are also extremal in \mathfrak{g} and hence satisfy the equation

$$(1 + \lambda\beta)(\lambda^2 g_1(x, y) + \lambda\alpha) = \lambda^2 \gamma^2.$$

This implies that the cubic equation

$$(1 + \beta X)(g_1(x, y)X^2 + \alpha X) = \gamma^2 X^2$$

has $|\mathbb{F}|$ zeros. So, if $|\mathbb{F}| > 3$, this means

$$\alpha = \beta = 0, \text{ and } \gamma^2 = g_1(x, y),$$

and we deduce

$$[x, y]_1 = \gamma[x, y].$$

But this implies that $[\cdot, \cdot]_1$ equals $\gamma[\cdot, \cdot]$ and hence is a scalar multiple of $[\cdot, \cdot]$.

In case that $|\mathbb{F}| = 2$, the above equation for $\lambda = 1$ reads as follows:

$$(1 + \alpha)(1 + \beta) = \gamma^2.$$

Now if $\alpha = 1$ or $\beta = 1$, it follows that $\gamma^2 = 0$ and so $\gamma = 0$, which is a contradiction. So also here, $\alpha = \beta = 0$ must hold.

In case that $|\mathbb{F}| = 3$, then setting λ to be ± 1 implies that we have the following two equations for α and β :

$$\begin{aligned} (1 + \beta)(g_1(x, y) + \alpha) &= 1 \\ (1 - \beta)(g_1(x, y) - \alpha) &= 1. \end{aligned}$$

But then $\alpha = \beta = 0$ and $g_1(x, y) = 1$. So, we conclude that the Lie product is unique up to scalar multiples. \diamond

Lemma 5.4. *Let $x \in E$ and $l \in \mathcal{F}$. Suppose $y_1, y_2 \in E$ span l . Then we can find an element $\lambda \in \mathbb{F}^*$ with $[x, y_i]_1 = \lambda[x, y_i]$ for both $i = 1$ and $i = 2$.*

Proof. Under the given conditions, Lemma 5.2 and Lemma 5.3 imply that there exist λ_i for $i = 1, 2$ with $[x, y_i]_1 = \lambda_i[x, y_i]$. If $[x, y_1] = 0$ or $[x, y_2] = 0$, then clearly we can take λ_1 and λ_2 to be equal. So assume $[x, y_1] \neq 0 \neq [x, y_2]$ and let $y_3 := -(y_1 + y_2)$ such that there exists λ_3 with $[x, y_3]_1 = \lambda_3[x, y_3]$.

Suppose $\lambda_1 \neq \lambda_2$. We find

$$0 = [x, y_1 + y_2 + y_3] = [x, y_1] + [x, y_2] + [x, y_3]$$

and

$$\begin{aligned} 0 &= [x, y_1 + y_2 + y_3]_1 = [x, y_1]_1 + [x, y_2]_1 + [x, y_3]_1 \\ &= \lambda_1[x, y_1] + \lambda_2[x, y_2] + \lambda_3[x, y_3]. \end{aligned}$$

But this implies that $(\lambda_1 - \lambda_2)[x, y_1] + (\lambda_3 - \lambda_2)[x, y_3] = 0$ and hence

$$\begin{aligned} 0 &= [x, (\lambda_1 - \lambda_2)y_1 + (\lambda_3 - \lambda_2)y_3] \\ &= [x, (\lambda_1 - \lambda_2)y_1 + (\lambda_3 - \lambda_2)(-y_1 - y_2)] \\ &= [x, (\lambda_1 - \lambda_3)y_1 + (\lambda_2 - \lambda_3)y_2]. \end{aligned}$$

With the definition $z := (\lambda_1 - \lambda_3)y_1 + (\lambda_2 - \lambda_3)y_2 \neq 0$, we have $[x, z] = 0$ and we find $z \in \mathcal{E}_{\leq 0}(x)$ and hence $[x, z]_1 = 0$ by Lemma 5.2. Since $[x, y_1] \neq 0 \neq [x, y_2]$, the element z is not a multiple of y_1 or y_2 , and hence there are $\mu_1, \mu_2 \in \mathbb{F}^*$ with $y_2 = \mu_1 y_1 + \mu_2 z$. Now we find

$$\begin{aligned} [x, y_2]_1 &= [x, \mu_1 y_1 + \mu_2 z]_1 = [x, \mu_1 y_1]_1 + [x, \mu_2 z]_1 \\ &= \lambda_1 \mu_1 [x, y_1] + 0 = \lambda_1 \mu_1 [x, y_1] \end{aligned}$$

and

$$[x, y_2] = [x, \mu_1 y_1 + \mu_2 z] = \mu_1 [x, y_1] + 0 = \mu_1 [x, y_1].$$

So, $[x, y_2]_1 = \lambda_1 [x, y_2]$, contradicting λ_1 to be different from λ_2 . \diamond

We are now ready to prove the main objective of this section.

Theorem 5.5. *Let \mathfrak{g} be a Lie algebra generated by its set of extremal elements E , equipped with the Lie product denoted by $[\cdot, \cdot]$ and a nondegenerate extremal form g .*

Assume that there is a second Lie product $[\cdot, \cdot]_1$ defined on the underlying vector space, with corresponding nondegenerate extremal form g_1 , giving rise to the same extremal geometry Γ . Then, there is a $\lambda \in \mathbb{F}^$ with $[x, y]_1 = \lambda [x, y]$ for all $x, y \in \mathfrak{g}$.*

Proof. Fix a pair $(x, y) \in E_2$. Then, by Lemma 5.3, there is a $\lambda \in \mathbb{F}^*$ with $[x, y]_1 = \lambda [x, y]$. We prove that this element λ is the one we are looking for. We begin with the proof that for all $z \in E$ we have $[x, z]_1 = \lambda [x, z]$.

Suppose $z \in E$ is different from y . If $[x, z] = 0$, then $z \in E_{\leq 0}(y)$, hence also $[x, z]_1 = 0$, and $[x, z]_1 = \lambda [x, z]$.

If $z \in E_2(x)$, then, as Γ has diameter 3, we can find elements z_1 and z_2 in $E_{\geq 1}(x)$ such that $\langle y, z_1 \rangle$, $\langle z_1, z_2 \rangle$, and $\langle z_2, z \rangle$ are in \mathcal{F} . (Notice that we allow these subspaces to be equal to each other.) Now we can apply the above Lemma 5.4 to each of these lines and eventually find that as $[x, z_1] \neq 0 \neq [x, z_2]$ that $[x, z]_1 = \lambda [x, z]$.

Finally consider the case where $z \in E_1(x)$. Notice that $[x, z] \in E$. As g is nondegenerate, we can find an element $u \in E_2([x, z])$. This u is in $E_1(z'')$ for some $z'' \in \langle [x, z], z \rangle \setminus \langle [x, z] \rangle$. As $\text{Exp}(\langle x \rangle)$ is transitive on the points of $\langle [x, z], z \rangle$ distinct from $\langle [x, z] \rangle$, we can assume that $u \in E_1(z)$. Now take z' to be $[u, z]$. Then $z' \in E_1([x, z])$. If $z' \in E_{\leq 0}(x)$, then $[[x, z], z'] = -[[z, z'], x] - [[z', x], z] = 0$, which is a contradiction with $z' \in E_1([x, z])$. So, $z' \in E_1(x)$, but then, as $z + z' \in E$, we have $[x, z + z'] = [x, z] + [x, z'] \in E$ and, by 3.13, $z' \in E_{-1}([x, z])$. Again a contradiction. Hence, $z' \in E_2(x)$.

By the above we have $[x, z']_1 = \lambda[x, z']$ and Lemma 5.4 implies now that $[x, z]_1 = \lambda[x, z]$.

Since we started with a fixed $x \in E$, it remains to show that the scalar factor λ is independent of x . Clearly, multiplying x with a nonzero scalar, does not change the value of λ .

Suppose $x_1 \in E$ with $(x, x_1) \in E_{-1}$. Then let $z \in E_2(x)$. Then $[z, x]_1 = \lambda[z, x]$. If $z \in E_2(x_1)$ then applying the above to z instead of x we find $[z, x_1]_1 = \lambda[z, x_1]$. So also $[x_1, z]_1 = \lambda[x_1, z]$. If $z \notin E_2(x_1)$, then let $x_2 = x + x_1$. The element x_2 is extremal and $z \in E_2(x_2)$, so, by similar arguments as used above, we have $[x_2, z]_1 = \lambda[x_2, z]$. Now pick an element $z_1 \in E_2(x_1)$. Then $z_1 \in E_2(x)$ and $[x, z_1]_1 = \lambda[x, z_1]$, or $z_1 \in E_2(x_2)$ and $[x_2, z_1]_1 = \lambda[x_2, z_1]$. As above we get $[x_1, z_1]_1 = \lambda[x_1, z_1]$ and λ is also the scalar for x_1 .

Now connectedness of $\Gamma(\mathfrak{g})$ implies that the scalar is the same for all extremal elements in E .

But then λ is the same for all pairs (x, y) of extremal elements in $E \times E$. Since E generates \mathfrak{g} , we find for all $x, y \in \mathfrak{g}$ that $[x, y]_1 = \lambda[x, y]$. \diamond

6 Conclusions

Combining the results of the previous two sections, we finally can characterize the Lie algebras under consideration by their extremal geometry.

Theorem 6.1. *Let \mathfrak{g} be a Lie algebra generated by its set E of extremal elements with respect to the extremal form g with trivial radical. If the natural embedding of the extremal geometry $\Gamma(\mathfrak{g})$ into $\mathbb{P}(\mathfrak{g})$ admits an absolutely universal cover, then \mathfrak{g} is uniquely determined (up to isomorphism) by $\Gamma(\mathfrak{g})$.*

Proof. We combine the previous results. So let \mathfrak{g}_1 be a second Lie algebra with isomorphic extremal geometry $\Gamma(\mathfrak{g}_1) \cong \Gamma(\mathfrak{g})$. By 4.3, the projective embeddings of $\Gamma(\mathfrak{g})$ and $\Gamma(\mathfrak{g}_1)$ are isomorphic and therefore \mathfrak{g} and \mathfrak{g}_1 are not only isomorphic vector spaces, but they also have the same Lie structure, as a consequence of Theorem 5.5. \diamond

Corollary 6.2. *Let $\mathfrak{g}_1, \mathfrak{g}_2$ be two Lie algebras as in 4.1 with extremal geometries $\Gamma(\mathfrak{g}_1) \cong \Gamma(\mathfrak{g}_2)$ isomorphic to a root shadow space of type $BC_{n,2}$ ($n \geq 3$), $D_{n,2}$ ($n \geq 4$), $E_{6,2}$, $E_{7,1}$, $E_{8,8}$, or $F_{4,1}$. Then $\mathfrak{g}_1 \cong \mathfrak{g}_2$.*

Proof. As a consequence of the main result 2.8 in [12], the root shadow space of \mathfrak{g}_1 (and \mathfrak{g}_2) of one of the given types has an absolutely universal cover. So Theorem 6.1 applies. \diamond

The above results provide a proof of Theorem 1.1. Indeed, suppose \mathfrak{g} is a simple Lie algebra generated by its set E of extremal elements. Then the extremal form g is nondegenerate. By the assumptions of Theorem 1.1 and Theorem 3.14 we find the extremal geometry $\Gamma(\mathfrak{g})$ to be a root shadow space of type $A_{n,\{1,n\}}$ ($n \geq 3$), $BC_{n,2}$ ($n \geq 3$), $D_{n,2}$ ($n \geq 4$), $E_{6,2}$, $E_{7,1}$, $E_{8,8}$, or $F_{4,1}$. Now we can apply the above corollary to find that \mathfrak{g} is uniquely determined by its extremal geometry, unless the extremal geometry is of type $A_{n,\{1,n\}}$ ($n \geq 3$). In that case we refer to [9].

7 Extremal elements in classical Lie algebras

In this final section we investigate extremal elements in classical Lie algebras. Although most of the results are well known, we present them now within the theory of extremal elements.

Let Φ be an irreducible root system with dot product (\cdot, \cdot) and Φ^+ the set of positive roots. Then consider

$$\mathfrak{ch}_\Phi = \bigoplus_{\alpha \in \Phi^+} \mathbb{Z}x_\alpha + \mathbb{Z}x_{-\alpha} + \mathbb{Z}h_\alpha$$

where for $\alpha \in \Phi^+$ the elements x_α , $x_{-\alpha}$ and h_α form a basis. Let $h_{-\alpha} = -h_\alpha$. Then on \mathfrak{ch}_Φ we define a bilinear product

$$[\cdot, \cdot] : \mathfrak{ch}_\Phi \times \mathfrak{ch}_\Phi \rightarrow \mathfrak{ch}_\Phi$$

by the following rules:

$$\begin{aligned} [h_\alpha, h_\beta] &= 0 \\ [h_\alpha, x_\beta] &= 2 \frac{(\beta, \alpha)}{(\alpha, \alpha)} x_\beta \\ [x_\alpha, x_\beta] &= \begin{cases} N_{\alpha, \beta} x_{\alpha+\beta} & \text{if } \alpha + \beta \in \Phi, \\ h_\alpha & \text{if } \beta = -\alpha, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where $\alpha, \beta \in \Phi$. The numbers $N_{\alpha, \beta}$ are integral *structure constants* chosen to be $\pm(p_{\alpha, \beta} + 1)$, where $p_{\alpha, \beta}$ is the biggest number such that $-p_{\alpha, \beta}\alpha + \beta$ is a root.

For a suitable choice of signs this product defines a Lie algebra called the *integral Chevalley Lie algebra*, see [5, 15]. The formal basis elements x_α , $\alpha \in \Phi$, and h_α with α positive, form a *Chevalley basis* of \mathfrak{ch}_Φ . A Lie

algebra $\mathfrak{ch}_\Phi(\mathbb{F}) := \mathfrak{ch}_\Phi \otimes \mathbb{F}$ obtained by tensoring with a field \mathbb{F} will be called a *Chevalley Lie algebra*.

In this section, we investigate extremal elements in a Chevalley Lie algebra $\mathfrak{ch}_\Phi(\mathbb{F})$ over a field \mathbb{F} . To avoid several technical difficulties, we restrict our attention to the case that the characteristic of \mathbb{F} is different from 2. (Notice that most results remain true in the characteristic 2 case.)

Example 7.1. Let V be a vector space over the field \mathbb{F} of dimension $n + 1$. If e_1, \dots, e_{n+1} is a basis of V with dual basis $\phi_1, \dots, \phi_{n+1}$, then the elements $x_{\epsilon_i - \epsilon_j} = e_i \otimes \phi_j$, with $i \neq j$ and $h_{\epsilon_i - \epsilon_j} = e_i \otimes \phi_i - e_j \otimes \phi_j$ form a Chevalley basis of type $A_n = \{\epsilon_i - \epsilon_j \mid 1 \leq i \neq j \leq n + 1\}$ for the Lie algebra $\mathfrak{sl}(V)$ of Example 3.2.

Suppose V has dimension $n + 1 = 2m$, the characteristic of \mathbb{F} is not 2, and b is a symmetric bilinear form on V satisfying $b(e_i, e_j) = 0$ unless $\{i, j\} = \{2k - 1, 2k\}$ for some k with $1 \leq k \leq m$, in which case $b(e_i, e_j) = 1$. Denote by b_i the linear form b_{e_i} as in 3.3. Then the Siegel elements $x_{\epsilon_i - \epsilon_j} = e_{2i-1} \otimes b_{2j-1} - e_{2j} \otimes b_{2i}$, $x_{\epsilon_i + \epsilon_j} = e_{2i-1} \otimes b_{2j} - e_{2j-1} \otimes b_{2i}$ and $x_{-\epsilon_i - \epsilon_j} = e_{2i} \otimes b_{2j-1} - e_{2j} \otimes b_{2i-1}$ together with the elements $h_\alpha = [x_\alpha, x_{-\alpha}]$ form a Chevalley basis of type $D_m = \{\epsilon_i \pm \epsilon_j \mid 1 \leq i, j \leq m, i \neq j\}$ in $\mathfrak{so}(V, b)$.

A *long (short) root element* in $\mathfrak{ch}_\Phi(\mathbb{F})$ is an element in the $\text{Aut}(\mathfrak{ch}_\Phi(\mathbb{F}))$ -orbit of an element x_α where α is a long (short) root in Φ .

We notice that, since x_α is ad-nilpotent for both α being short or long, we find $\exp(x_\alpha, \lambda)$, where $\lambda \in \mathbb{F}$, to be an automorphism of $\mathfrak{ch}_\Phi(\mathbb{F})$. Here $\exp(x_\alpha, \lambda) = \exp(\lambda x_\alpha)$ in the language of [5].

Proposition 7.2. *The long root elements in $\mathfrak{ch}_\Phi(\mathbb{F})$ are extremal.*

Proof. Let α be a long root in Φ . As the characteristic of \mathbb{F} is different from 2, it suffices to prove that $[x_\alpha, [x_\alpha, y]] \in \mathbb{F}x_\alpha$ for all y in the Chevalley basis.

If $y = x_\beta$ for any $\beta \in \Phi$ different from $-\alpha$, then $[x_\alpha, [x_\alpha, y]] = [x_\alpha, [x_\alpha, x_\beta]] = 0$ since at least one of $\alpha + \beta$ and $2\alpha + \beta$ is not in Φ .

If $y = x_{-\alpha}$, then $[x_\alpha, [x_\alpha, y]] = [x_\alpha, [x_\alpha, x_{-\alpha}]] = [x_\alpha, h_\alpha] = -2x_\alpha$.

Finally for $y = h_\beta$ we have $[x_\alpha, [x_\alpha, y]] = [x_\alpha, -2\frac{(\alpha, \beta)}{(\beta, \beta)}x_\alpha] = 0$. \diamond

Lemma 7.3. *The long root elements linearly span $\mathfrak{ch}_\Phi(\mathbb{F})$.*

Proof. First we show that every x_β , where β is a short root in Φ , is a linear combination of long root elements.

If Φ is of type B_n or C_n and β is a short root, then we can find a long root α such that $2\beta + \alpha$ is long. Indeed, if Φ is of type B_n then,

without loss of generality we can assume $\beta = \epsilon_1$ and take $\alpha = \epsilon_2 - \epsilon_1$ and if Φ is of type C_n then β can be assumed to be $-\epsilon_1 + \epsilon_2$ and we can take $\alpha = 2\epsilon_1$. But then $\exp(x_{-\alpha-\beta}, 1)(x_{\alpha+2\beta}) = x_{\alpha+2\beta} - x_\beta - x_{-\alpha}$, so $x_\beta = -\exp(x_{-\alpha-\beta}, 1)(x_{\alpha+2\beta}) + x_{\alpha+2\beta} + x_{-\alpha}$, a linear combination of three long root elements.

If Φ is of type F_4 , every short root is inside a B_2 subsystem and we can apply the above.

So it remains to consider the G_2 -case and β a short root. Then we can pick a long root α such that $\alpha, \alpha + 3\beta$ and $2\alpha + 3\beta$ are long. Notice that $\alpha + \beta$ is short. But then $\exp(x_{-\alpha-\beta}, 1)(x_{2\alpha+3\beta}) + \exp(x_{-\alpha-\beta}, 1)(x_{2\alpha+3\beta}) = 2x_\beta + 2x_{2\alpha+3\beta}$.

So $x_\beta = \frac{1}{2}(\exp(x_{-\alpha-\beta}, 1)(x_{2\alpha+3\beta}) + \exp(x_{-\alpha-\beta}, 1)(x_{2\alpha+3\beta}) - 2x_{2\alpha+3\beta})$, and hence a linear combination of three long root elements.

Now consider the elements h_α , where $\alpha \in \Phi^+$. We have

$$\exp(x_\alpha, 1)(x_{-\alpha}) = x_{-\alpha} + h_\alpha - x_\alpha,$$

from which we deduce that h_α is a linear combination of short or long root elements and hence also a linear combination of long root elements. \diamond

Let g be the extremal form on $\mathfrak{ch}_\Phi(\mathbb{F})$, and denote by \mathcal{E} the set of extremal points of $\mathfrak{ch}_\Phi(\mathbb{F})$.

Lemma 7.4. *The extremal points $\langle x_\alpha \rangle$, with $\alpha \in \Phi$ a long root, are contained in a single connected component of the graph $(\mathcal{E}, \mathcal{E}_2)$.*

Proof. We denote adjacency in $(\mathcal{E}, \mathcal{E}_2)$ by \sim .

Let α, β be distinct long roots and $\langle x_\alpha \rangle$ and $\langle x_\beta \rangle$ be the corresponding extremal points. Obviously, if $\alpha = -\beta$, the points are adjacent in the $(\mathcal{E}, \mathcal{E}_2)$.

If $\alpha \neq -\beta$ and $(\alpha, \beta) \neq 0$, we can assume, (up to replacing β by $-\beta$) that $\langle \alpha, \beta \rangle = -1$ and $\alpha + \beta$ is also a root. This implies that $[x_\alpha, x_\beta] = N_{\alpha, \beta} x_{\alpha+\beta}$ with $N_{\alpha, \beta} \neq 0$. By 3.5, it follows that $x_\alpha + x_{\alpha+\beta}$ is extremal.

As $g(x_{-\alpha}, x_\alpha + x_{\alpha+\beta}) = g(x_{-\alpha}, x_\alpha) + g(x_{-\alpha}, x_{\alpha+\beta}) = 1$, we find $\langle x_\alpha \rangle \sim \langle x_{-\alpha} \rangle \sim \langle x_\alpha + x_{\alpha+\beta} \rangle$. Moreover, as $g(x_{-\alpha-\beta}, x_\alpha + x_{\alpha+\beta}) = 1$ we can also find a path from $\langle x_{\alpha+\beta} \rangle$ to $\langle x_\alpha + x_{\alpha+\beta} \rangle$ and hence from $\langle x_\alpha \rangle$ to $\langle x_{\alpha+\beta} \rangle$. Similarly, there is a path from $\langle x_\beta \rangle$ to $\langle x_{\alpha+\beta} \rangle$ and $\langle x_\alpha \rangle$ and $\langle x_\beta \rangle$ are in the same connected component.

It remains to consider the case where $(\alpha, \beta) = 0$, so the roots α and β are orthogonal to each other. If Φ is not of type C_n , we can find a long root γ such that $(\alpha, \gamma) \neq 0$ and $(\beta, \gamma) \neq 0$ and then apply the above, to conclude that both $\langle x_\alpha \rangle$ and $\langle x_\beta \rangle$ are in the connected component of $(\mathcal{E}, \mathcal{E}_2)$ containing $\langle x_\gamma \rangle$.

If Φ is of type C_n , the long roots are $\pm 2\varepsilon_i$ and the short roots are $\pm(\varepsilon_i \pm \varepsilon_j)$, where $1 < i \leq j < n$. Note that in this case, we have the root lengths $\sqrt{2}$ and 2. We will show that for $i \neq j$ the elements $\langle x_{-2\varepsilon_i} \rangle$ and $\langle x_{2\varepsilon_j} \rangle$ are in the same $(\mathcal{E}, \mathcal{E}_2)$ -component.

Define $y = \exp(x_{-\varepsilon_i - \varepsilon_j}, 1)(x_{2\varepsilon_i}) = x_{2\varepsilon_i} - x_{\varepsilon_i - \varepsilon_j} - x_{-2\varepsilon_j}$, which is a long root element since $x_{2\varepsilon_i}$ is one.

Bilinearity of g gives $g(x_{-2\varepsilon_i}, y) = g(x_{-2\varepsilon_i}, x_{2\varepsilon_i}) - g(x_{-2\varepsilon_i}, x_{\varepsilon_i - \varepsilon_j}) - g(x_{-2\varepsilon_i}, x_{-2\varepsilon_j}) = 1 + 0 + 0 \neq 0$. Moreover $g(x_{2\varepsilon_j}, y) = g(x_{2\varepsilon_j}, x_{2\varepsilon_i}) - g(x_{2\varepsilon_j}, x_{\varepsilon_i - \varepsilon_j}) - g(x_{2\varepsilon_j}, x_{-2\varepsilon_j}) = 0 + 0 + 1 \neq 0$. So we have a path $\langle x_{-2\varepsilon_i} \rangle \sim \langle y \rangle \sim \langle x_{2\varepsilon_j} \rangle$ in $(\mathcal{E}, \mathcal{E}_2)$. \diamond

As we have seen before, the radical of the extremal form g is an ideal in the Lie algebra $\mathfrak{ch}_\Phi(\mathbb{F})$. By $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ we denote the quotient Lie algebra $\mathfrak{ch}_\Phi(\mathbb{F})/\text{rad}(g)$. For $x \in \mathfrak{ch}_\Phi(\mathbb{F})$ we denote by \bar{x} the element $x + \text{rad}(g) \in \overline{\mathfrak{ch}_\Phi(\mathbb{F})}$.

The elements \bar{x} in $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ where x is a long root element not in $\text{rad}(g)$, are also called long root elements. The set of extremal points of $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is denoted by $\bar{\mathcal{E}}$.

Proposition 7.5. *The Lie algebra $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is simple. It is generated by its extremal elements.*

These extremal elements are all scalar multiples of long root elements.

Proof. Assume that $x \in \overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is extremal, but not in the same connected component of $(\mathcal{E}, \mathcal{E}_2)$ as some element $\bar{x}_\alpha \in \overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ with $\alpha \in \Phi$ long. Since $\mathfrak{ch}_\Phi(\mathbb{F})$ is generated by its long root elements, that are in one connected component by 7.4, it follows by 3.14 that $[x, \overline{\mathfrak{ch}_\Phi(\mathbb{F})}] = 0$ and $x \in Z(\overline{\mathfrak{ch}_\Phi(\mathbb{F})}) = \{0\}$. A contradiction. So $(\mathcal{E}, \mathcal{E}_2)$ and $\Gamma(\overline{\mathfrak{ch}_\Phi(\mathbb{F})})$ are connected.

But this implies that all extremal points are in one single orbit under the automorphism group of $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$, see 3.14, and hence all extremal elements are scalar multiples of long root elements.

Now assume \mathfrak{i} is a nonzero ideal of $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$. Let $0 \neq i \in \mathfrak{i}$. Then, as $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is spanned by the images of long root elements, there is an extremal x with $\bar{g}(x, i) \neq 0$, where \bar{g} is the extremal form on $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$. But then $\bar{g}(x, i)x = [x, [x, i]] \in \mathfrak{i}$.

If $\langle y \rangle$ is in relation \mathcal{E}_2 with $\langle x \rangle$, then inside $\langle x, y \rangle \cong \mathfrak{sl}_2$ we find that $g(y, x)y = [y, [y, x]]$ is in \mathfrak{i} . By connectedness of $(\mathcal{E}, \mathcal{E}_2)$ we find all extremal elements to be in \mathfrak{i} and hence $\mathfrak{i} = \overline{\mathfrak{ch}_\Phi(\mathbb{F})}$. This shows that $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is simple. \diamond

If Φ is an irreducible root system of type X_n , $n \geq 2$, then the extremal geometry of the Lie algebra $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is a root shadow space of type $X_{n,J}$ for some set J , except when $X_n = A_2$ and the characteristic of \mathbb{F} is 3. This is explained by the following. The radical of g is usually equal to the center of \mathfrak{g} . (See [11] for details.) The only exception is the case where $X_n = G_2$ and the characteristic of \mathbb{F} is equal to 3. In this exceptional case, the short roots are contained in the radical of g . This radical is 7-dimensional and $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ is isomorphic to $\overline{\mathfrak{ch}_{\Phi'}(\mathbb{F})}$, where Φ' is a root system of type A_2 . So, if $X_n = A_2$ and the characteristic of \mathbb{F} is 3, then the extremal geometry is a root shadow space of type $G_{2,\{1,2\}}$. (We notice that in characteristic 2 more exceptions occur.)

The extremal geometry of $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$, when Φ is of type A_1 or C_n , does not contain lines.

Let \mathfrak{g} be a simple Lie algebra over a subfield \mathbb{K} of \mathbb{F} such that $\mathfrak{g} \otimes_{\mathbb{K}} \mathbb{F}$ is isomorphic to $\hat{\mathfrak{g}}$. Then we call \mathfrak{g} a *form* of $\hat{\mathfrak{g}}$.

Several forms of the Lie algebras $\overline{\mathfrak{ch}_\Phi(\mathbb{F})}$ do contain long root elements. Forms of Chevalley Lie algebras of type A, B, C and D, give rise to various orthogonal and unitary Lie algebras generated by their long root elements. The extremal geometries are then root shadow spaces of type $BC_{n,2}$ or have no lines. In case Φ is of type D_4 one also obtains examples of extremal geometries which are of type $G_{2,2}$. Forms of Chevalley Lie algebras of type E lead to extremal geometries without lines, as well as root shadow spaces of type $A_{2,\{1,2\}}$, $G_{2,2}$, and $F_{4,1}$. We obtain examples of Lie algebras generated by their extremal elements having an extremal geometry isomorphic to the root shadow space of a building of algebraic type. See for example [16, 17, 19].

References

- [1] R. J. Blok, *Highest weight modules and polarized embeddings of shadow spaces*, J. Algebraic Combin. **34** (2011), no. 1, 67–113.
- [2] R. J. Blok and A. Pasini, *On absolutely universal embeddings*, Discrete Math. **267** (2003), no. 1-3, 45–62. Combinatorics 2000 (Gaeta).
- [3] N. Bourbaki, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*, Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968 (French).
- [4] F. Buekenhout and A. M. Cohen, *Diagram geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 57, Springer, Heidelberg, 2013. Related to classical groups and buildings.

- [5] Roger W. Carter, *Simple groups of Lie type*, John Wiley & Sons, London-New York-Sydney, 1972. Pure and Applied Mathematics, Vol. 28.
- [6] A. M. Cohen and G. Ivanyos, *Root filtration spaces from Lie algebras and abstract root groups*, J. Algebra **300** (2006), no. 2, 433–454.
- [7] ———, *Root shadow spaces*, European J. Combin. **28** (2007), no. 5, 1419–1441.
- [8] A. M. Cohen, A. Steinbach, R. Ushirobira, and D. Wales, *Lie algebras generated by extremal elements*, J. Algebra **236** (2001), no. 1, 122–154.
- [9] H. Cuypers, K. Roberts, and S. Shpectorov, *Recovering the Lie algebra from its extremal geometry*, J. Algebra **441** (2015), 196–215.
- [10] H. Cuypers and Y. Fleischmann, *A geometric characterization of the symplectic Lie algebra* (2017).
- [11] Y. Fleischmann, *A Geometric Approach to Classical Lie Algebras*, PhD thesis, Technische Universiteit Eindhoven, 2015.
- [12] A. Kasikova and E. Shult, *Absolute embeddings of point-line geometries*, J. Algebra **238** (2001), no. 1, 265–291.
- [13] A. A. Premet, *Inner ideals in modular Lie algebras*, Vestsi Akad. Navuk BSSR Ser. Fiz.-Mat. Navuk **5** (1986), 11–15, 123 (Russian, with English summary).
- [14] J. A. Thas and H. Van Maldeghem, *Full embeddings of the finite dual split Cayley hexagons*, Combinatorica **24** (2004), no. 4, 681–698.
- [15] J. Tits, *Sur les constantes de structure et le théorème d’existence des algèbres de Lie semi-simples*, Inst. Hautes Études Sci. Publ. Math. **31** (1966), 21–58.
- [16] Jacques Tits, *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Mathematics, Vol. 386, Springer-Verlag, Berlin-New York, 1974.
- [17] J. Tits and R. M. Weiss, *Moufang polygons*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2002.
- [18] H. Völklein, *On the geometry of the adjoint representation of a Chevalley group*, J. Algebra **127** (1989), no. 1, 139–154.
- [19] R. M. Weiss, *The structure of spherical buildings*, Princeton University Press, Princeton, NJ, 2003.