

# The predictability of visual outcome of epiretinal membrane peeling by pre-operative OCT images

**Citation for published version (APA):**

Takkenberg, D. E. J., Berendschot, T. T. J. M., & Goezinne, F. (2018). The predictability of visual outcome of epiretinal membrane peeling by pre-operative OCT images. *Acta Ophthalmologica*, 96(1), 9-9.

**Document status and date:**

Published: 01/03/2018

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

average flare at 1 month was higher than in healthy controls ( $p < 0.001$ ). Long term flare levels after DMEK (Group II) measured 9.6 ( $\pm 4.2$ ) ph/ms and was higher in eyes associated with allograft rejection ( $n = 6$ ) vs those without rejection (16.7 ( $\pm 7.8$ ) vs 9.3 ( $\pm 3.8$ ) ph/ms, respectively,  $p < 0.001$ ). All rejection eyes had flare values above 10 ph/ms.

**Conclusions:** Aqueous flare after DMEK quickly decreased within the first postoperative month indicating a fast recovery of the blood-aqueous barrier. Long-term flare levels was higher in eyes associated with rejection, suggesting persistent subclinical inflammation. A flare level above 10 ph/ms may be used as a threshold for identifying eyes associated with or at risk of allograft rejection after DMEK.

**Financial disclosures:** Drs Baydoun and Dapena are consultants for DORC International/ Dutch Ophthalmic USA. Dr. Melles is a consultant for DORC International/ Dutch Ophthalmic USA and SurgiCube International. All other authors have no potential interest to disclose.

### iOCT assisted DMEK: towards more efficient and safe surgery

M.B. Muijzer, C. van Luijk, N. Soeters, R.P.L. Wisse

*Utrecht Cornea Research Group, Department of Ophthalmology, University Medical Center Utrecht, Utrecht, The Netherlands*

**Introduction:** Intra-operative optical coherence tomography (iOCT) is a novel and promising imaging method in ocular surgery, where conventional OCT technology is integrated in the ophthalmic microscope (Zeiss Lumera 700 RESCAN). Proper imaging of the thin graft in Descemet Membrane Endothelial Keratoplasty (DMEK) surgery is a particular advantage, and this study reports on our experiences of iOCT in DMEK surgery.

**Patient and method:** DMEK surgery was conducted by one surgeon in a standardized manner and cases were retrospectively clustered in three equally sized groups: the first underwent DMEK surgery without iOCT, the second group with iOCT first experiences, and the third group with standardized iOCT use. iOCT was used in particular for assessment of graft orientation, interface and apposition. The surgeons first 10 DMEK cases were excluded from analysis.

**Results:** During May 2016 and November 2017 48 cases with 3 month follow-up were included for analysis. Surgery time decreased significantly between different groups (ANOVA;  $p < 0.001$ ) with mean surgery times of 68 min  $\pm$  16, 58 min  $\pm$  12, resp. 46 min  $\pm$  14. iOCT enabled the surgeon to refrain from post-insertion pressurizing of the globe. Corrected distance visual acuity appeared comparable (ANOVA;  $p > 0.171$ ). The incidence of adverse events remained equal. 3/33.3%, 5/25% resp. 3/15.8% graft detachments needed rebubbling in the three treatment groups (Fisher exact;  $p = 0.809$ )

**Conclusion:** iOCT enables the surgeon to better assess graft orientation, interface and adherence during corneal transplant surgery. A better visualization potentially leads to a shorter surgery time without compromising the safety of the procedure.

### Combined phacovitrectomy or sequential cataract surgery in macula-off retinal detachment

A. Hajjaj<sup>1</sup>, S. Manning<sup>1,2</sup>, R. Wubbels<sup>2</sup>, K. van Overdam<sup>1,2</sup>

<sup>1</sup>Department of Vitreoretinal Surgery, The Rotterdam Eye Hospital, Rotterdam, The Netherlands, <sup>2</sup>The Rotterdam Ophthalmic Institute, Rotterdam, The Netherlands

**Purpose:** Compare refractive outcomes between cataract surgery combined with pars plana vitrectomy (PPV) and sequential cataract surgery after PPV, in macula-off rhegmatogenous retinal detachment (RRD).

**Methods:** Multi-surgeon, retrospective case series. Consecutive cases of PPV for macula-off RRD, in 12 months, with combined (combined

group) or sequential (sequential group) phacoemulsification cataract surgery.

**Results:** There were 73 eyes in the combined and 77 eyes in the sequential group. Axial length (AL) measurement was not possible in 26/73 eyes in the combined group. Contralateral eye biometry was used in 17 eyes and contralateral AL in 9 eyes. Four eyes with silicone oil in situ at follow-up (3 in the combined and 1 in the sequential group), were excluded from refractive outcome analysis. The combined group compared as follows to the sequential group: prediction error:  $-0.50 \pm 0.91$  vs  $-0.32 \pm 0.68$  D ( $p = 0.21$ ); absolute biometry prediction error:  $0.74 \pm 0.72$  D vs  $0.58 \pm 0.47$  D ( $p = 0.33$ ); refractive outcome within  $\pm 0.5$  D of target: 27/70 (39%) vs 40/76 (53%); within  $\pm 1$  D of target: 55/70 (79%) vs 64/76 (84%); outside 2 D of target: 7/70 (10%) vs 2/76 (2.6%). In cases where AL measurement was possible in the combined group (45/70), refractive outcome within  $\pm 0.5$  D, within  $\pm 1$  D and outside 2D of target, were 17/45 (38%), 37/45 (82%) and 2/45 (4.4%).

**Conclusion:** Cataract surgery combined with PPV had worse refractive outcome and more refractive surprises, than PPV and sequential cataract surgery in macula-off RRD.

**Proprietary interest declaration:** None of the authors have any proprietary interests in relation to this study.

### The predictability of visual outcome of epiretinal membrane peeling by pre-operative OCT images

D.E.J. Takkenberg, T.T.J.M. Berendschot, F. Goezinne

*University Eye Clinic Maastricht, Maastricht, The Netherlands*

**Purpose:** To determine the predictability of the visual acuity (VA) outcome of inner limiting membrane (ILM) peeling for epiretinal membrane (ERM), as estimated by vitreoretinal surgeons based on pre-operative OCT scans.

**Patients and methods:** All patients who underwent pars plana vitrectomy (PPV) for ERM peeling in the University Eye Clinic Maastricht between 01-01-2015 and 01-01-2016 were included in this retrospective study. Pre-operative OCT images were collected and were anonymously reviewed by 3 vitreoretinal surgeons and a fellow vitreoretinal surgery, after which pre- and postoperative VA was estimated by them and compared to the real postoperative VA.

**Results:** 136 eyes underwent surgery, 12 eyes were excluded: 7 due to incomplete VA-data, 5 because of retinal detachment within 3 months after surgery. The mean age of the remaining 124 patients, 69 men and 55 women, was  $72.2 \pm 6.8$  years. The mean post-operative LogMAR VA,  $0.27 \pm 0.27$ , was significantly better than the pre-operative VA:  $0.42 \pm 0.24$  ( $p < 0.001$ ). We found a significant correlation between the predicted postoperative VA and the actual post-operative VA in 3 of the 4 vitreoretinal surgeons, however rather low (range 0.17–0.44). Moreover, we found significant differences in the estimates of the post-operative VA between the different surgeons and in comparison with the actual post-operative VA ( $p < 0.001$ ).

**Conclusions:** Generally, results from PPV ILM peeling are good. In this study, vitreoretinal surgeons could not reliably predict post-operative VA by reviewing pre-operative OCT images.

### Posterior vitreous detachment, its characteristics and the risk of developing retinal tears: a systematic review

O. Gishiti<sup>1</sup>, J. Verhoeks<sup>1</sup>, R.A.J. van den Nieuwenhof<sup>2</sup>, K. van Overdam<sup>1</sup>

<sup>1</sup>The Rotterdam Eye Hospital, Rotterdam, The Netherlands,

<sup>2</sup>Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands

**Introduction:** Flashes and floaters are the main symptoms of posterior vitreous detachment, which is shown to be associated with increased risks of developing retinal tears, retinal detachment or vitreous