

Cell-free microcompartmentalised transcription-translation for the prototyping of synthetic communication networks

Citation for published version (APA):

Dubuc, E., Pieters, P. A., van der Linden, A. J., van Hest, J. C. M., Huck, W. T. S., & de Greef, T. F. A. (2019). Cell-free microcompartmentalised transcription-translation for the prototyping of synthetic communication networks. *Current Opinion in Biotechnology*, 58, 72-80. <https://doi.org/10.1016/j.copbio.2018.10.006>

Document license:

CC BY

DOI:

[10.1016/j.copbio.2018.10.006](https://doi.org/10.1016/j.copbio.2018.10.006)

Document status and date:

Published: 01/08/2019

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.



ELSEVIER



Cell-free microcompartmentalised transcription–translation for the prototyping of synthetic communication networks

Emilien Dubuc^{1,2,3}, Pascal A Pieters^{1,2,3}, Ardjan J van der Linden^{1,2,3}, Jan CM van Hest⁴, Wilhelm TS Huck⁵ and Tom FA de Greef^{1,2,3,6}

Recent efforts in synthetic biology have shown the possibility of engineering distributed functions in populations of living cells, which requires the development of highly orthogonal, genetically encoded communication pathways. Cell-free transcription–translation (TXTL) reactions encapsulated in microcompartments enable prototyping of molecular communication channels and their integration into engineered genetic circuits by mimicking critical cell features, such as gene expression, cell size, and cell individuality within a community. In this review, we discuss the uses of cell-free transcription–translation reactions for the development of synthetic genetic circuits, with a special focus on the use of microcompartments supporting this reaction. We highlight several studies where molecular communication between non-living microcompartments and living cells have been successfully engineered.

Addresses

¹Laboratory of Chemical Biology, Department of Biomedical Engineering, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands

²Institute for Complex Molecular Systems, Department of Biomedical Engineering, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands

³Computational Biology Group, Department of Biomedical Engineering, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands

⁴Department of Biomedical Engineering & Department of Chemical Engineering and Chemistry, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

⁵Department of Physical Organic Chemistry, Institute for Molecules and Materials, Radboud University, Nijmegen 6525 HP, The Netherlands

⁶Institute for Molecules and Materials, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

Corresponding author: de Greef, Tom FA (t.f.a.d.greef@tue.nl)

Current Opinion in Biotechnology 2019, 58C:72–80

This review comes from a themed issue on **Nanobiotechnology**

Edited by **Giovanni Maglia** and **Wesley R Browne**

<https://doi.org/10.1016/j.copbio.2018.10.006>

0958-1669/© 0001 Elsevier Ltd. All rights reserved.

Introduction

Synthetic biology traditionally utilises genetic circuits in order to implement novel functions in living cells [1,2]. The fundamental approach of synthetic biology is based on the use of building blocks assembled in functional modules, culminating in genetic circuits that can display complex spatiotemporal behaviour such as oscillations, synchronisation and combinatorial control of endogenous signalling pathways [3–5,6^{••}]. However, progressing from module-based to system-focussed synthetic biology remains an ongoing challenge, partly due to our limited knowledge of complex function integration into living systems. For instance, to facilitate the engineering of synthetic cellular consortia capable of displaying collective behaviour, a broader understanding of the mechanisms by which cells communicate in their natural environment is required.

Simplified models of biological systems could help identify key molecular parameters and as such have the potential to uncover generalisable concepts. Proteins and other components of the transcription and translation machinery can be extracted from cells as a lysate or purified and reconstituted (PURE) in order to perform gene expression *in vitro* with a good control over the biochemical composition. Here, compartmentalised cell-free transcription–translation (TXTL) reactions constitute ideal model systems for deducing the rules of network composition, developing new communication pathways between cellular mimics and living cells [7], and creating genetic devices for the implementation of synthetic communication between cells of different species [8].

In this review, we intend to first summarise the use of TXTL reactions for prototyping and implementation of novel genetic networks. We will discuss how microfluidic technologies and semipermeable microcapsules provide essential tools for cell-free synthetic biology in a context where size, composition, and individuality of the reaction are of particular importance. Finally, we will highlight important contributions of encapsulated TXTL to the development of synthetic communication paths.

Cell-free reactions for applied synthetic biology: circuit testing, behaviour prediction and forward engineering


As the complexity of synthetic gene networks increases, more resources are shared between synthetic circuits and their host. As a result, cells experience a decrease in fitness, which limits their growth and the efficiency of synthetic circuits [9**]. Furthermore, implementing synthetic circuits within cellular hosts exposes these networks to interactions with endogenous pathways, negatively influencing the behaviour of both host and synthetic circuitry. Cell-free synthetic biology has emerged as a powerful tool for testing new genetic networks in a controlled biochemical context, as well as for predicting cellular responses to such networks [10*,11**]. Here we summarise most recent studies using cell-free TXTL for prototyping novel genetic modules and circuits

and facilitating their integration into cellular hosts (Figure 1a).

Characterisation of novel regulatory parts

Development of complex genetic circuits requires the use of well-characterised regulatory modules. Cell-free TXTL allows the rapid and thorough testing of new genetic parts thus enabling such characterisation (Figure 1b upper panel). DNA and RNA sequence-specific regulation is receiving particular interest as it yields novel regulators with high degree of orthogonality. Researchers recently used a cell-free synthetic biology approach for the testing of novel regulatory elements with further *in vivo* applications, leveraging the strong potential of riboregulation [12*,13*,14,15], and dCas9-based repression [16].

Figure 1



Cell-free reactions for characterising, testing, and optimising complex circuits.

Cell-free TXTL reactions can be used in combination with mathematical modelling to test complex circuits, and identify optimal conditions for the implementation of novel biological functions *in vivo* (a). Cell-free reactions are used to characterise the behaviour of novel isolated circuit parts [19**] (upper panel (b)) and test the behaviour of circuits combining novel parts (lower panel (b)). Various versions of similar networks can rapidly be tested in TXTL, which can be used to reveal retroactive interactions between modules (upper panel (c)), as well as unexpected effects when combining components into a single construct (lower panel (c)). Cell burden occurs when a synthetic circuit excessively mobilises resources that are also necessary for endogenous circuits (left panel (d)). A capacity monitor reports possible cell burden during the implementation of novel circuits using TXTL and allows the design of networks generating minimal burden [21] (right panel (d)). TXTL allows the testing of various biochemical conditions, such as cofactor, salt, and enzyme concentrations, in order to optimise a TXTL-based reaction [30] (e).

Testing and forward engineering of synthetic gene networks with complex behaviour

After characterising regulatory building blocks, novel genetic circuits can be designed and assembled (Figure 1b lower panel). *In vitro* characterisation is often done under batch conditions, but these conditions do not ensure a constant supply of substrate, as well as the removal of by-products, and the renewal of information, which are prerequisites for the implementation of higher-order regulatory behaviours. In contrast, flow reactors enable the testing of regulatory modules and the implementation of complex networks in conditions mimicking cellular homeostasis [17^{••},18^{••}], as well as the forward engineering of such networks into bacterial hosts [19^{••}]. In addition, TXTL in flow reactors constitutes an effective method for approaching synthetic biology from control theory perspectives, where compositional context, cell heterogeneity and division become controlled parameters instead of poorly defined variables [20]. Finally, we anticipate flow reactors could serve as platform for studying retroactivity (Figure 1c upper panel) between modules or networks sharing common resources, and test different compositions of a network (Figure 1c lower panel).

Resource burden

Synthetic gene networks mobilise resources essential to the cell, which are often found in limited supply (Figure 1d left panel). In order to rapidly estimate the resource cost of integrating novel synthetic circuits, Borkowski *et al.* coupled a TXTL batch system to a capacitor circuit, which facilitated circuit transposition into cells [21] (Figure 1d right panel). Coupled to a regulatory dCas9-based negative feedback, this circuit could provide direct supervision over the resource consumption of synthetic circuits [22]. By leveraging the possibility to tune parameters critical to the TXTL reaction, such as DNA instructions and DNA concentration, multiple other systems have also been used to screen and identify resource competition [19^{••},17^{••},23].

Tuneability, context, and composition

One major advantage of cell-free TXTL reactions is the possibility to adjust the parameters influencing the performance of a synthetic circuit, whereas *in vivo* methods rely predominantly on host resources and characteristics. Egbert and Klavins established that, in living cells, the performance of a non-endogenous genetic circuit is highly context-dependent, as a network can yield its designed function in one host strain but fails to achieve this function in another strain [24]. The choice of a bacterial host could be guided by identifying *in vitro* which parameters are essential for the desired behaviour and selecting a strain matching these context requirements. To further optimise host context, many regulatory elements critical for protein synthesis can be added or expressed *in situ*, such as MazF ribonuclease [25], GamS [26^{••}] and Chi6 [27] DNase inhibitors, GreA/B transcription elongation

factors [28], and ClpXP protease [29]. Karim *et al.* composed a mixture of lysates enriched in specific enzymes or with enzymes expressed *in situ* in order to identify major directions for improving the yield of *n*-butanol synthesis [30] (Figure 1e). Furthermore, Yeung *et al.* leveraged the tuneability of TXTL systems in order to show that supercoiling is one of the main factors explaining differences in expression levels within an intergenic context [31].

Alternatives to *Escherichia coli*

The majority of cell-free synthetic biology work is based on cell extracts from the *E. coli* bacterium. However, other prokaryotes have been used to provide novel TXTL systems, and could become new model organisms due to their interesting features, such as growth traits and facilitated genetic manipulation [23,32–35]. Furthermore, additional eukaryotic TXTL systems were recently developed, opening methods for *in vitro* studies with a closer focus on human-centred applications [36–38].

Engineered compartments to study physicochemical properties of living systems

Cells display a high intrinsic concentration of biomolecules, a reduced volume, and an amphiphilic interface between their internal volume and external environment. Engineering synthetic compartments of controlled size and composition for the conduction of TXTL reaction enables the study of gene expression at the relevant scale. We distinguish three experimental microcompartments that have been employed to perform TXTL or prototype synthetic genetic networks based on cell-free TXTL, first, PDMS-based compartments, second, lipid-based compartments, and third, coacervates, also known as complex aqueous two-phase systems (ATPS).

PDMS-based microcompartments


Complex network implementation

In contrast with methods using free-floating DNA gene templates, Bar-Ziv *et al.* proposed a method for performing TXTL reaction on a chip in which the gene template DNA was attached to a functionalised surface, first under batch conditions using wheat-germ extract [39], and subsequently in microfluidic flow reactors using *E. coli* cell extract [18^{••}]. The latter, alongside the work of Niederholtmeyer *et al.* [17^{••}] (Figure 2a), was amongst the first examples of long-term TXTL, and demonstrated the implementation of advanced synthetic gene networks such as oscillators. Although challenging to fabricate and use, the devices developed by Niederholtmeyer *et al.* are a major technological breakthrough that will facilitate the development of numerous synthetic gene circuits in the future.

Reaction-diffusion gene networks

Gene expression-pattern formation plays an essential role in morphogenesis and is a relevant study-case for cell-free

Figure 2



Microfluidic technologies for studying gene expression at the cellular scale.

Continuous flow reactors allow the implementation of out-of-equilibrium gene networks in TXTL. Valves precisely control addition and mixing of fresh TXTL reagents, enabling the implementation of complex networks such as oscillators [19**] (a). Control over geometry and diffusion allows the study of gene expression propagation and pattern formation in compartmentalised flow reactors using DNA brushes [41] (b). Microdroplets generated on a microchip allow the screening of a vast range of conditions influencing network behaviour in TXTL, such as DNA template concentrations, shown here for the implementation of an incoherent feedforward loop [44] (c). Liposomes encapsulating TXTL can be obtained by double-emulsion techniques ((d), top-left panel). Liposomes are used for isolating genetic cascades [75] ((d), bottom-left panel) and creating units containing self-replicating genetic information [53**] ((d), right panel). Upon increase of ionic strength, TXTL mixture can phase separate, resulting in a highly active microcompartment [47] ((e) left panel). DNA-functionalised hydrogels are used to create membraneless compartments capable of gene expression ((e) right panel).

synthetic biology. In 2005, Isalan *et al.* implemented a coarse mimic of a *Drosophila melanogaster* morphogenic gene network [40]. Three genes were coupled to paramagnetic beads at defined locations inside a cm-long batch chamber containing wheat germ lysate-based TXTL, yielding the emergence of a reaction-diffusion network allowing the formation of gene expression patterns. This work highlighted important criteria for the implementation of pattern-forming gene networks, first, control over the sublocation of network components, second, control over resource competition and depletion, and third, control over protein degradation. The previously described microfluidic system developed by the

Bar-Ziv group provided a platform within which each of these critical criteria can be controlled (Figure 2b). Tayar *et al.* leveraged this technical improvement to study the emergence of expression patterns from out-of-equilibrium gene networks, first using a bistable switch [41], and later a series of oscillators coupled in space [42**]. In a follow-up study, Pardatscher *et al.* developed a lithography technique in order to functionalise a surface with up to three distinct DNA strands on a chip supporting both lysate-based and PURE-based TXTL reaction [43]. This technique will allow the study of complex interactions between multiple, spatially resolved, genes and the formation of more complex spatial patterns.

Lipid-based microcompartments

The cell membrane constitutes an amphiphilic interface between the interior of the cell and its environment. This interface can be mimicked by encapsulating TXTL reactions inside single (water-in-oil) or double (water-in-oil-in-water) emulsions in order to study large number of these reactions at a cellular scale.

High-throughput screening

Large numbers of TXTL microdroplets in oil with a controlled dispersity are easy to generate, store, and remain stable over hours, which makes them particularly interesting for screening large numbers of parameters influencing TXTL reactions (Figure 2c). This feature was elegantly employed by Hori *et al.*, who optimised a cell-free genetic circuit based on an incoherent feedforward topology using fluorescently barcoded droplets in which lysate-based TXTL reactions can take place [44].

Confinement

The reduced size of microdroplets is particularly interesting to study physical effects such as confinement. Guan *et al.* encapsulated *Xenopus* egg extract in microdroplets of various size. The authors showed that the extract could undergo several mitotic oscillations, and described the influence of the size of the compartments on the period [45]. Sakamoto *et al.* investigated the influence of surface-to-volume ratio of microcompartments on the efficiency of lysate-based TXTL, suggesting a deleterious interaction between membrane lipids and translation machinery [46].

Molecular crowding

Macromolecular crowding effects, which emerge in highly concentrated media such as the cytosol, can be artificially induced in TXTL mixtures using crowding additives inside microdroplets. Molecular crowding has been shown to influence the spatial segregation of biomolecules and kinetics of cell-free TXTL reactions [47,48]. In addition, Norred *et al.* showed that noise in transcription reaction in systems displaying reduced diffusion resulted in gene expression bursts, which are likely to occur in cells but are not observable in bulk PURE-based TXTL reactions [49].

Semi-permeable liposomes

Given that droplets are intrinsically closed systems, they cannot be utilised for long-term protein expression, nor as analogues of semi-permeable lipid bilayers for mimicking the cell membrane. Noireaux *et al.* first implemented lysate-based TXTL reaction in a liposome permeated by haemolysin pores, which enabled the exchange of nutrients and by-products with a feeding solution, resulting in long-term gene expression [50]. Later, Garamella *et al.* tested various transcriptional cascades inside liposomes [26**] (Figure 2d left panel). Majumder *et al.* implemented a lysate-based TXTL liposome sensitive to osmotic changes

via expression of the MscL calcium channel [51]. Moreover, lysate encapsulating liposomes were recently used by Krinsky *et al.* to serve as containers for the production and intratumoural delivery of a toxin protein [52]. Finally, Van Nies *et al.* recently described the first liposomes capable of isothermally replicating DNA using self-encoded proteins in PURE system, constituting a major step towards the construction of a true synthetic cell capable of full replication [53**] (Figure 2d right panel).

Membraneless compartments

Under critical ionic concentrations, biomolecules displaying a high electrical charge or high multivalency can undergo complex liquid-liquid phase separation, which yields their partition into membraneless compartments known as coacervates (Figure 2e left panel). Coacervates obtained from TXTL mixture are therefore highly enriched in TXTL machinery. There is growing evidence that phase separation plays a major role in the regulation of protein activity, spatial segregation of nucleic acids, and functional organisation of the cell [54–56]. Phase separation also gives rise to the emergence of partitioning, confinement, and crowding effects influencing cell-free gene expression [47,57,58]. TXTL was first combined with hydrogels by Park *et al.*, who used a micropad of DNA hydrogel incubated in TXTL mixture [59]. The DNA hydrogel, consisting of genes coding for a reporter protein, was not permeable to TXTL machinery, so protein expression occurred at the surface of the gel but not inside. Thiele *et al.* used hyaluronic acid gel beads in which DNA template was covalently attached, and incubated the gel beads in presence of TXTL mixture inside a microdroplet [60]. In contrast to the gel used by Park *et al.*, gel beads used in this study were porous enough to allow TXTL to occur inside the gel beads. The authors showed that transcription and translation reactions were confined inside the gel as the mRNA remained trapped inside the beads. Finally Zhou *et al.* produced hydrogel particles containing DNA template, ribosomes, and His-tagged TXTL proteins from PURE system. All biomolecules remained trapped inside the hydrogel particles, so TXTL could occur over several days by continuous supply of feeding buffer (Figure 2d left panel) [61].

TXTL and communication

Interactions between gut microbiota and the human body recently gained in interest as studies highlighted the impact of microorganisms on the metabolism [62], immune responses [63], and the recurrence of cancer [64]. The development of new biomolecular tools inspired by natural quorum sensing systems is therefore of particular relevance in order to understand multicellular communication [8] and to engineer synthetic communication devices [6**,65]. Furthermore, to avoid the difficulty of engineering organisms with increasingly complex genetic circuits, the development of orthogonal communication channels that enable distributed

functions within a bacterial community has recently received particular attention [8,66]. Nevertheless, the prototyping of synthetic and highly orthogonal molecular communication channels that mediate collective behaviour in populations of living cells remains challenging. Compartmentalised TXTL systems can play an important role in the development of such pathways, as they can mimic communication between cells whilst remaining modular and easy to control.

Quorum sensor characterisation

Quorum sensors are naturally occurring genetic communication systems found in many bacterial species in a variety of chemical messenger-transcription factor-operator systems. Quorum sensors are essential building blocks for engineering molecular communication channels in synthetic biology [6,8]. Halleran and Murray recently characterised a series of quorum sensors using lysate-based TXTL, prior to their implementation *in vivo*, enabling them to accurately predict crosstalk between different systems [67]. In addition, Wen *et al.* used lysate-based TXTL reactions to identify the presence of quorum sensing signals in patients suffering from lung infection by *Pseudomonas aeruginosa*, demonstrating the therapeutic relevance of quorum sensing detection and characterisation [68].


Artificial cell communication

Combination of microfluidic technologies with TXTL allows the generation of cell mimics capable of communication processes based on gene instructions. Booth *et al.* engineered a tissue mimic by 3D-printing TXTL-containing droplets into organised layers [69]. The PURE-based TXTL reactions produced α -haemolysin upon light-activation in order to create communication channels between individual droplets in a programmable manner. In a related approach, Findlay *et al.* programmed specific communication between two droplets by implementing the active transport of a signalling molecule through LacY transporter proteins produced *in situ* using PURE-based TXTL reaction [70]. Alternatively, communication was implemented between emitter TXTL-liposomes and receiver-proteinosomes [71]. By establishing communication between mammalian and bacterial synthetic cell models (synells), Adamala *et al.* provided the first example of diffusive one-way communication between fully synthetic cells of different organisms (Figure 3a) [72]. Such synells comprised liposomes containing either HeLa or *E. coli* extract, as well as quorum sensing DNA circuits. However, to engineer true collective behaviour in populations of synells requires further development of this technology to allow bidirectional exchange of information.

Natural-artificial cell communication

Schwarz-Schilling *et al.* achieved communication between two distinct droplet populations—encapsulating

Figure 3



TXTL in liposomes for prototyping novel communication channels. Through the exchange of chemical signalling molecules, liposomes containing TXTL mixture and quorum sensing circuits can be used as artificial cells to establish communication with other artificial cells (a) or with living cells (b).

either TXTL reaction mixture or bacteria [73]. Each population was alternatively equipped with either a receiver circuit or an emitter circuit, so droplets could exchange chemical signals further processed by TXTL or bacteria. Later, Lentini *et al.* implemented communication between lysate-based or PURE-based synthetic cells and various bacteria by combining multiple quorum sensing devices from various species [74] (Figure 3b). In doing so, the authors were able to establish two-way communication between synthetic cells and *Vibrio fischeri* bacteria. Furthermore, authors produced a synthetic cell capable of interfacing communications between various bacterial species which were previously unable to communicate. The synells were programmed to receive a chemical signal from one bacterium (*V. fischeri*), which induced the production of an enzyme disrupting an ongoing communication between an emitter bacterium (*P. aeruginosa*) and a receiver bacterium (engineered *E. coli*), a phenomenon also known as quorum quenching.

Conclusion and outlook

Cell-free TXTL reactions are becoming an essential tool to prototype novel genetic circuits and can facilitate their integration into living hosts. In combination with microfluidic technologies and the development of novel semi-permeable microcapsules, TXTL reactions constitute a unique platform for the study of gene expression and coupling to other cellular processes in a biochemically relevant microenvironment.

It has been established that size and confinement have a considerable impact on TXTL reactions [45,46]. For this reason, more attention should be paid to implementing TXTL reactions in liposomes of defined size and dispersity. In contrast to flow reactors, complex networks have not yet been implemented in liposomes. Possible reasons are the passive diffusion of nutrients, which could be limiting for the TXTL reaction, as well as an absence of active removal of the information. Potential improvements could come from active import and export of substrate and by-products, implementation of a sustained metabolism, all integrated with the enzymatic mRNA and protein degradation methods described in this review. Another remaining challenge is to discover and characterise novel quorum sensors, allowing communication between artificial cells and natural human cells.

Current efforts are exploring possibilities to establish communication between artificial and natural systems, working towards therapeutic applications wherein information exchange between living cells are mediated or altered. We anticipate that micro-compartmentalised TXTL—by accelerating the characterisation of novel genetic circuits and orthogonal molecular communication channels—will have a compelling contribution to the field of synthetic biology.

Conflict of interest statement

Nothing declared.

Acknowledgements

This work was supported by the European Research Council, ERC (project n. 677313 BioCircuit) an NWO-VIDI grant from the Netherlands Organization for Scientific Research (NWO, 723.016.003), funding from the Ministry of Education, Culture and Science (Gravity programs, 024.001.035 & 024.003.013)

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Khalil AS, Collins JJ: **Synthetic biology: applications come of age.** *Nat Rev Genet* 2010, **11**:367-379.
 2. Collins J: **Synthetic biology: bits and pieces come to life.** *Nature* 2012, **483**.
 3. Stricker J, Cookson S, Bennett MR, Mather WH, Tsimring LS, Hasty J: **A fast, robust and tunable synthetic gene oscillator.** *Nature* 2008, **456**:516-519.
 4. Danino T, Mondragón-Palomino O, Tsimring L, Hasty J: **A synchronized quorum of genetic clocks.** *Nature* 2010, **463**:326-330.
 5. Nielsen AA, Voigt CA: **Multi-input CRISPR/Cas genetic circuits that interface host regulatory networks.** *Mol Syst Biol* 2014, **10**:763.
 6. Din MO, Danino T, Prindle A, Skalak M, Selimkhanov J, Allen K, Julio E, Atolia E, Tsimring LS, Bhatia SN *et al.*: **Synchronized cycles of bacterial lysis for in vivo delivery.** *Nature* 2016, **536**:81-85.
- Salmonella* cells are engineered to periodically release a dose of toxin lethal for vicinal tumour cells.
7. Lentini R, Yeh Martín N, Mansy SS: **Communicating artificial cells.** *Curr Opin Chem Biol* 2016, **34**:53-61.
 8. Scott SR, Hasty J: **Quorum sensing communication modules for microbial consortia.** *ACS Synth Biol* 2016, **5**:969-977.
 9. Borkowski O, Ceroni F, Stan GB, Ellis T: **Overloaded and stressed: whole-cell considerations for bacterial synthetic biology.** *Curr Opin Microbiol* 2016, **33**:123-130.
- This review provides very clear overview of the issues related to burden created by synthetic circuits.
10. Salehi-Reyhani A, Ces O, Elani Y: **Artificial cell mimics as simplified models for the study of cell biology.** *Exp Biol Med* 2017, **242**:1309-1317.
- This review provides good general description of the field of minimal cell construction.
11. Jia H, Heymann M, Bernhard F, Schwille P, Kai L: **Cell-free protein synthesis in micro compartments: building a minimal cell from biobricks.** *N Biotechnol* 2017, **39**:199-205.
- Another very thorough review about the construction of a minimal cell.
12. Pardee K, Green AA, Ferrante T, Cameron DE, Daleykeyser A, Yin P, Collins JJ: **Paper-based synthetic gene networks.** *Cell* 2014, **159**:940-954.
- This paper describes outstanding applications of TXTL outside the lab for rapid diagnostics.
13. Pardee K, Green AA, Takahashi MK, Braff D, Lambert G, Lee JW, Ferrante T, Ma D, Donghia N, Fan M *et al.*: **Rapid, low-cost detection of Zika virus using programmable biomolecular components.** *Cell* 2016, **165**:1255-1266.
- Follow-up of the Collins lab for detecting Zika virus DNA using TXTL.
14. Hu CY, Takahashi MK, Zhang Y, Lucks JB: **Engineering a functional small RNA negative autoregulation network with model-guided design.** *ACS Synth Biol* 2018, **7**:1507-1518.
 15. Senoussi A, Lee Tin Wah J, Shimizu Y, Robert J, Jaramillo A, Findeiss S, Axmann IM, Estevez-Torres A: **Quantitative characterization of translational riboregulators using an in vitro transcription-translation system.** *ACS Synth Biol* 2018, **7**:1269-1278.
 16. Marshall R, Maxwell CS, Collins SP, Jacobsen T, Luo ML, Begemann MB, Gray BN, January E, Singer A, He Y *et al.*: **Rapid and scalable characterization of CRISPR technologies using an E. coli cell-free transcription-translation system.** *Mol Cell* 2018, **69**:146-157.e3.
 17. Niederholtmeyer H, Stepanova V, Maerkl SJ: **Implementation of cell-free biological networks at steady state.** *Proc Natl Acad Sci U S A* 2013, **110**:15985-15990.
- One of the first paper describing a microfluidic platform for implementing complex gene networks.
18. Karzbrun E, Tayar AM, Noireaux V, Bar-Ziv RH: **Programmable on-chip DNA compartments as artificial cells.** *Science (80-)* 2014, **345**:829-832.
- One of the first paper describing a microfluidic platform for implementing complex gene networks.
19. Niederholtmeyer H, Sun ZZ, Hori Y, Yeung E, Verpoorte A, Murray RM, Maerkl SJ: **Rapid cell-free forward engineering of novel genetic ring oscillators.** *Elife* 2015, **4**:e09771.
- Follow-up paper describing how the microfluidic platform developed by Niederholtmeyer *et al.* is used for the forward engineering of novel networks in bacteria.
20. Del Vecchio D, Dy AJ, Qian Y, Del Vecchio D, Dy AJ, Qian Y: **Control theory meets synthetic biology.** *J R Soc* 2016, **13**:3-43.
 21. Borkowski O, Bricio C, Murgiano M, Rothschild-Mancinelli B, Stan GB, Ellis T: **Cell-free prediction of protein expression costs for growing cells.** *Nat Commun* 2018, **9**.
 22. Ceroni F, Boo A, Furini S, Gorochowski TE, Borkowski O, Ladak YN, Awan AR, Gilbert C, Stan G-B, Ellis TT *et al.*: **Burden-driven feedback control of gene expression.** *Nat Methods* 2018, **15**:387.
 23. Moore SJ, MacDonald JT, Wienecke S, Ishwarbhai A, Tsipa A, Aw R, Kyllis N, Bell DJ, McClymont DW, Jensen K *et al.*: **Rapid acquisition and model-based analysis of cell-free**

- transcription–translation reactions from nonmodel bacteria.** *Proc Natl Acad Sci U S A* 2018, **115**:201715806.
24. Egbert RG, Klavins E: **Fine-tuning gene networks using simple sequence repeats.** *Proc Natl Acad Sci U S A* 2012, **109**:16817–16822.
 25. Shin J, Noireaux V: **Study of messenger RNA inactivation and protein degradation in an Escherichia coli cell-free expression system.** *J Biol Eng* 2010, **4**.
 26. Garamella J, Marshall R, Rustad M, Noireaux V: **The all E. coli TX- toolbox 2.0: a platform for cell-free synthetic biology.** *ACS Synth Biol* 2016, **5**:344–355.
- This article provides essential knowledge for the development of novel circuits in TXTL.
27. Marshall R, Maxwell CS, Collins SP, Beisel CL, Noireaux V: **Short DNA containing χ sites enhances DNA stability and gene expression in E. coli cell-free transcription–translation systems.** *Biotechnol Bioeng* 2017, **114**:2137–2141.
 28. Maddalena LLD, Niederholtmeyer H, Turtola M, Swank ZN, Belogurov GA, Maerkl SJ: **GreA and GreB enhance expression of Escherichia coli RNA polymerase promoters in a reconstituted transcription-translation system.** *ACS Synth Biol* 2016, **5**:929–935.
 29. Shi X, Wu T, Cole CM, Devaraj NK, Joseph S: **Optimization of ClpXP activity and protein synthesis in an E. coli extract-based cell-free expression system.** *Sci Rep* 2018, **8**:3488.
 30. Karim AS, Heggstad JT, Crowe SA, Jewett MC: **Controlling cell-free metabolism through physicochemical perturbations.** *Metab Eng* 2018, **45**:86–94.
 31. Yeung E, Dy AJ, Martin KB, Ng AH, Del Vecchio D, Beck JL, Collins JJ, Murray RM: **Biophysical constraints arising from compositional context in synthetic gene networks.** *Cell Syst* 2017, **5**:11–24.e12.
 32. Failmezger J, Scholz S, Blombach B, Siemann-Herzberg M: **Cell-free protein synthesis from fast-growing Vibrio natriegens.** *Front Microbiol* 2018, **9**:1146.
 33. Des Soye B, Davidson S, Weinstock M, Gibson D, Jewett MC: **Establishing a high-yielding cell-free protein synthesis platform derived from Vibrio natriegens.** *ACS Synth Biol* 2018, **7**:2245–2255.
 34. Kelwick R, Webb AJ, Macdonald JT, Freemont PS: **Development of a Bacillus subtilis cell-free transcription-translation system.** *Metab Eng* 2016, **38**:1.
 35. Li J, Wang H, Jewett MC: **Expanding the palette of Streptomyces-based cell-free protein synthesis systems with enhanced yields.** *Biochem Eng J* 2018, **130**:29–33.
 36. Wu C, Dasgupta A, Shen L, Bell-Pedersen D, Sachs MS: **The cell free protein synthesis system from the model filamentous fungus Neurospora crassa.** *Methods* 2017, **137**:11–19.
 37. Jung WR, Sierecki E, Bastiani M, O'Carroll A, Alexandrov K, Rae J, Johnston W, Hunter DJB, Ferguson C, Gambin Y *et al.*: **Cell-free formation and interactome analysis of caveolae.** *J Cell Biol* 2018, **217**:2141–2165.
 38. Wang S, Majumder S, Emery NJ, Liu AP: **Simultaneous monitoring of transcription and translation in mammalian cell-free expression in bulk and in cell-sized droplets.** *Synth Biol* 2018, **3**:ysy005.
 39. Buxboim A, Bar-Dagan M, Frydman V, Zbaida D, Morpurgo M, Bar-Ziv R: **A single-step photolithographic interface for cell-free gene expression and active biochips.** *Small* 2007, **3**:500–510.
 40. Isalan M, Lemerle C, Serrano L: **Engineering gene networks to emulate Drosophila embryonic pattern formation.** *PLoS Biol* 2005, **3**:e64.
 41. Tayar AM, Karzbrun E, Noireaux V, Bar-Ziv RH: **Propagating gene expression fronts in a one-dimensional coupled system of artificial cells.** *Nat Phys* 2015, **11**:1037–1041.
 42. Tayar AM, Karzbrun E, Noireaux V, Bar-Ziv RH: **Synchrony and pattern formation of coupled genetic oscillators on a chip of artificial cells.** *Proc Natl Acad Sci U S A* 2017, **114**:11609–11614. Important work describing gene pattern formation in flow reactors.
 43. Pardatscher G, Schwarz-Schilling M, Daube SS, Bar-Ziv RH, Simmel FC: **Gene expression on DNA biochips patterned with strand-displacement lithography.** *Angew Chem—Int Ed* 2018, **57**:4783–4786.
 44. Hori Y, Kantak C, Murray RM, Abate AR: **Cell-free extract based optimization of biomolecular circuits with droplet microfluidics.** *Lab Chip* 2017, **17**:3037–3042.
 45. Guan Y, Li Z, Wang S, Barnes PM, Liu X, Xu H, Jin M, Liu AP, Yang Q: **A robust and tunable mitotic oscillator in artificial cells.** *eLife* 2018, **7**:e33549.
 46. Sakamoto R, Noireaux V, Maeda YT: **Anomalous scaling of gene expression in confined cell-free reactions.** *Sci Rep* 2018, **8**:7364.
 47. Sokolova E, Spruijt E, Hansen MMK, Dubuc E, Groen J, Chokkalingam V, Piruska A, Heus HA, Huck WTS: **Enhanced transcription rates in membrane-free protocells formed by coacervation of cell lysate.** *Proc Natl Acad Sci U S A* 2013, **110**:11692–11697.
 48. Hansen MMK, Paffenholz S, Foschepoth D, Heus HA, Thiele J, Huck WTS: **Cell-like nanostructured environments alter diffusion and reaction kinetics in cell-free gene expression.** *ChemBioChem* 2016, **17**:228–232.
 49. Norred SE, Caveney PM, Chauhan G, Collier LK, Collier CP, Abel SM, Simpson ML: **Macromolecular crowding induces spatial correlations that control gene expression bursting patterns.** *ACS Synth Biol* 2018, **7**:1251–1258.
 50. Noireaux V, Libchaber A: **A vesicle bioreactor as a step toward an artificial cell assembly.** *Proc Natl Acad Sci U S A* 2004, **101**:17669–17674.
 51. Majumder S, Garamella J, Wang YL, Denies M, Noireaux V, Liu AP: **Cell-sized mechanosensitive and biosensing compartment programmed with DNA.** *Chem Commun* 2017, **53**:7349–7352.
 52. Krinsky N, Kaduri M, Zinger A, Shainsky-Roitman J, Goldfeder M, Benhar I, Hershkovitz D, Schroeder A: **Synthetic cells synthesize therapeutic proteins inside tumors.** *Adv Healthc Mater* 2018, **7**.
 53. Van Nies P, Westerlaken I, Blanken D, Salas M, Mencia M, Danelon C: **Self-replication of DNA by its encoded proteins in liposome-based synthetic cells.** *Nat Commun* 2018, **9**:1583.
- Authors report the unprecedented self-replication of phage DNA inside liposomes.
54. Banani SF, Lee HO, Hyman AA, Rosen MK: **Biomolecular condensates: organizers of cellular biochemistry.** *Nat Rev Mol Cell Biol* 2017, **18**:285–298.
 55. Alberti S: **Phase separation in biology.** *Curr Biol* 2017, **27**:R1097–R1102.
 56. Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, Schymkowitz J, Shorter J, Wolozin B, Van Den Bosch L *et al.*: **Protein phase separation: a new phase in cell biology.** *Trends Cell Biol* 2018, **28**:420–435.
 57. Torre P, Keating CD, Mansy SS: **Multiphase water-in-oil emulsion droplets for cell-free transcription-translation.** *Langmuir* 2014, **30**:5695–5699.
 58. Dora Tang T-Y, van Swaay D, deMello A, Ross Anderson JL, Mann S: **In vitro gene expression within membrane-free coacervate protocells.** *Chem Commun* 2015, **51**:11429–11432.
 59. Park N, Um SH, Funabashi H, Xu J, Luo D: **A cell-free protein-producing gel.** *Nat Mater* 2009, **8**:432–437.
 60. Thiele J, Ma Y, Foschepoth D, Hansen MMK, Steffen C, Heus HA, Huck WTS: **DNA-functionalized hydrogels for confined membrane-free in vitro transcription/translation.** *Lab Chip* 2014, **14**:2651–2656.
 61. Zhou X, Wu H, Cui M, Lai SN, Zheng B: **Long-lived protein expression in hydrogel particles: towards artificial cells.** *Chem Sci* 2018, **9**:4275–4279.

62. Brown JM, Hazen SL: **Microbial modulation of cardiovascular disease**. *Nat Rev Microbiol* 2018, **16**:171-181.
63. Clemente JC, Ursell LK, Parfrey LW, Knight R: **The impact of the gut microbiota on human health: an integrative view**. *Cell* 2012, **148**:1258-1270.
64. Bullman S, Pedamallu CS, Sicinska E, Clancy TE, Zhang X, Cai D, Neuberger D, Huang K, Guevara F, Nelson T *et al.*: **Analysis of Fusobacterium persistence and antibiotic response in colorectal cancer**. *Science* 2017, **358**:1443-1448.
65. Kim S, Kerns SJ, Ziesack M, Bry L, Gerber GK, Way JC, Silver PA: **Quorum sensing can be repurposed to promote information transfer between bacteria in the mammalian gut**. *ACS Synth Biol* 2018, **7**:2270-2281.
66. Kyllis N, Tuza ZA, Stan GB, Polizzi KM: **Tools for engineering coordinated system behaviour in synthetic microbial consortia**. *Nat Commun* 2018, **9**:2677.
67. Halleran AD, Murray RM: **Cell-free and in vivo characterization of Lux, Las, and Rpa Quorum activation systems in E. coli**. *ACS Synth Biol* 2018, **7**:752-755.
68. Wen KY, Cameron L, Chappell J, Jensen K, Bell DJ, Kelwick R, Kopniczky M, Davies JC, Filloux A, Freemont PS: **A cell-free biosensor for detecting quorum sensing molecules in P. aeruginosa-infected respiratory samples**. *ACS Synth Biol* 2017, **6**:2293-2301.
69. Booth MJ, Schild VR, Graham AD, Olof SN, Bayley H: **Light-activated communication in synthetic tissues**. *Sci Adv* 2016, **2**.
70. Findlay HE, Harris NJ, Booth PJ: **In vitro synthesis of a major facilitator transporter for specific active transport across droplet interface bilayers**. *Sci Rep* 2016, **6**.
71. Tang TYD, Cecchi D, Fracasso G, Accardi D, Coutable-Pennarun A, Mansy SS, Perriman AW, Anderson JLR, Mann S: **Gene-mediated chemical communication in synthetic protocell communities**. *ACS Synth Biol* 2018, **7**:339-346.
72. Adamala KP, Martin-Alarcon DA, Guthrie-Honea KR, Boyden ES: **Engineering genetic circuit interactions within and between synthetic minimal cells**. *Nat Chem* 2017, **9**:431-439.
- This article describes how bacterial and human-based artificial cells can communicate. Authors also report selective fusion of artificial cells upon recognition of membrane markers.
73. Schwarz-Schilling M, Aufinger L, Mückl A, Simmel FC: **Chemical communication between bacteria and cell-free gene expression systems within linear chains of emulsion droplets**. *Integr Biol* 2016, **8**:564-570.
74. Lentini R, Martin NY, Forlin M, Belmonte L, Fontana J, Cornella M, Martini L, Tamburini S, Bentley WE, Jousson O *et al.*: **Two-way chemical communication between artificial and natural cells**. *ACS Cent Sci* 2017, **3**:117-123.
75. Shin J, Noireaux V: **An E. coli cell-free expression toolbox: application to synthetic gene circuits and artificial cells**. *ACS Synth Biol* 2012, **1**:29-41.