

# Saturation Physics in Negative- and Positive-Triangularity Plasmas

**Citation for published version (APA):**

Pueschel, M. J., Duff, J. M., & Ball, J. (2022). Saturation Physics in Negative- and Positive-Triangularity Plasmas. In *48th EPS Conference on Plasma Physics 27 June - 1 July 2022* (Europhysics conference abstracts; Vol. 46A). European Physical Society (EPS). <http://ocs.ciemat.es/EPS2022PAP/pdf/O5.103.pdf>

**Document status and date:**

Published: 01/01/2022

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.

## Saturation Physics in Negative- and Positive-Triangularity Plasmas

M.J. Pueschel<sup>1,2</sup>, J.M. Duff<sup>3</sup>, and J. Ball<sup>4</sup>

<sup>1</sup>*Dutch Institute for Fundamental Energy Research, 5612 AJ Eindhoven, The Netherlands*

<sup>2</sup>*Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands*

<sup>3</sup>*University of Wisconsin-Madison, Madison, Wisconsin 53706, USA*

<sup>4</sup>*Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland*

Strong shaping of tokamak flux surfaces provides an avenue by which turbulence characteristics can be influenced in tokamak plasmas. One particularly pertinent shaping parameter is the triangularity  $\delta$ , which in recent years has received special attention in the context of inward-pointing triangular shapes, or negative  $\delta < 0$ . Here, characteristics of microinstabilities as well as their saturation properties are analyzed in both positive- $\delta$  and negative- $\delta$  configurations.

The potential benefits of finite  $\delta$  are captured in gyrokinetic simulations of ion-temperature-gradient-driven modes. Linearly, substantial negative  $\delta \lesssim 0.5$  leads to a reduction in growth, as local (along the field line) magnetic shear dis-aligns the eigenmode from the region of bad curvature. At large positive  $\delta \gtrsim 0.5$ , the local shear forces the mode to finite radial wavenumber  $k_x$ , reducing transport efficacy. This picture is confirmed by nonlinear simulations, which additionally show that while zonal flows are stronger at positive  $\delta$ , their impact in saturation is comparable regardless of triangularity sign due to higher saturation efficiency at negative  $\delta$ .

The impact of finite  $\beta$  is studied for a pair of TCV discharges at  $\delta < 0$  and  $\delta > 0$ . Notably,  $\delta > 0$  exhibits lower fluxes due to lower normalized equilibrium gradients, and profile flips – where pressure gradients from one discharge are combined with the magnetic geometry of another – are used to ensure comparability. It is found that both the  $\delta > 0$  and the  $\delta < 0$  discharges are near-marginal to decorrelation of field lines: a very moderate increase in  $\beta$  leads to a substantial boost in field-line stochasticity and ensuing zonal-flow erosion [1]. A key consequence is the proximity to a regime of extreme profile stiffness. Notably, even cases where electromagnetic stabilization ensures that the system locally sits below the nonlinear critical gradient, intermittency can still lead to decorrelation. In select cases, transition to a high-flux state is only temporary, and instead a zonal-flow-dominated attractor is eventually reached.

### References

- [1] M.J. Pueschel, P.W. Terry, F. Jenko, D.R. Hatch, W.M. Nevins, T. Görler, and D. Told, *Phys. Rev. Lett.* **110**, 155005 (2013)
- [2] J.M. Duff, B.J. Faber, C.C. Hegna, M.J. Pueschel, and P.W. Terry, *Phys. Plasmas* **29**, 012303 (2022)