

Josef Ladenbauer, Moritz Augustin, Klaus Obermayer

# How adaptation currents and synaptic inhibition change threshold, gain and variability of neuronal spiking

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013 Paris, France. 13-18 July 2013

**Conference paper, Published version**

This version is available at <http://nbn-resolving.de/urn:nbn:de:kobv:83-opus4-69964>.



## **Suggested Citation**

Ladenbauer, Josef ; Augustin, Moritz ; Obermayer, Klaus : How adaptation currents and synaptic inhibition change threshold, gain and variability of neuronal spiking : From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013 Paris, France. 13-18 July 2013. - In: BMC Neuroscience. - ISSN 1471-2202 (online). - 14 (2013), suppl. 1, P299. - doi:10.1186/1471-2202-14-S1-P299.

## **Terms of Use**

This work is licensed under a CC BY 2.0 License (Creative Commons Attribution 2.0 Generic). For more information see <http://creativecommons.org/licenses/by/2.0>.

POSTER PRESENTATION

Open Access

# How adaptation currents and synaptic inhibition change threshold, gain and variability of neuronal spiking


Josef Ladenbauer<sup>1,2\*</sup>, Moritz Augustin<sup>1,2</sup>, Klaus Obermayer<sup>1,2</sup>

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

Many types of neurons show spike rate adaptation, a gradual decrease in spiking activity following a sudden increase in stimulus intensity. This behavior is typically mediated by slow potassium currents through voltage-sensitive low-threshold or calcium-activated high-threshold channels, both of which are susceptible to cholinergic modulation [1]. Such adaptation currents (and changes thereof) contribute to frequency selectivity [2], coding [3] and attention [4]. These effects are likely caused by altering the relationship between synaptic input and spike rate output (I-O curve) as well as the characteristics of inter-spike intervals (ISI). Here we investigate (i) how voltage-dependent subthreshold and spike-dependent adaptation currents change the neuronal I-O curve as well as the ISI distribution for different input statistics and (ii) how these changes compare to those induced by synaptic inhibition.

Based on a population of adaptive exponential integrate-and-fire (aEIF) model neurons receiving noisy external and recurrent synaptic inputs we use the Fokker-Planck equation to compute spike rates and ISI distributions in the limit of a large adaptation timescale.

We show that a subthreshold adaptation current or synaptic inhibition received from independent neurons (external inhibition) change the neuronal I-O curve subtractively. That is, both mechanisms increase the spike threshold. On the other hand, a spike-triggered adaptation current or inhibitory synaptic feedback (recurrent inhibition) change the I-O curve divisively, i.e., they reduce the neuronal gain, see Figure 1A. Both types of adaptation current naturally increase the mean ISI. Surprisingly, they affect spiking variability in opposite ways. Subthreshold adaptation leads to an increase while spike-triggered adaptation causes a decrease of variability, see Figure 1B.



\* Correspondence: jl@ni.tu-berlin.de

<sup>1</sup>Neural Information Processing Group, Berlin Institute of Technology, Berlin, Germany

Full list of author information is available at the end of the article

Both types of synaptic inhibition however increase spiking variability. We simplify the model by neglecting the leak conductance which allows to analytically derive expressions describing these effects. For validation purposes, we show that the effective description of subthreshold and spike-triggered adaptation in the aEIF model corresponds well to the biophysical description of a voltage-sensitive muscarinic and a calcium-activated potassium current in a Hodgkin-Huxley type neuron model, respectively. Our results suggest that neuronal adaptation currents differentially contribute to neuronal threshold and gain control as well as ISI-based coding schemes.

#### Acknowledgements

This work was supported by the DFG Collaborative Research Center SFB910.

#### Author details

<sup>1</sup>Neural Information Processing Group, Berlin Institute of Technology, Berlin, Germany. <sup>2</sup>Bernstein Center for Computational Neuroscience Berlin, Berlin, Germany.

Published: 8 July 2013

#### References

1. McCormick DA: Neurotransmitter actions in the thalamus and cerebral cortex and their role in thalamocortical activity. *Progr Neurobiol* 1992, **39**:337-388.
2. Deemyad T, Kroeger J, Chacron M: Sub and suprathreshold adaptation currents have opposite effects on frequency tuning. *J Physiol* 2012, **590**:4839-4858.
3. Prescott S, Sejnowski T: Spike-rate coding and spike-time coding are affected oppositely by different adaptation mechanisms. *J Neurosci* 2008, **28**(50):13649-13661.
4. Herrero JL, Roberts MJ, Delicato LS, Gieselmann MA, Dayan P, Thiele A: Acetylcholine contributes through muscarinic receptors to attentional modulation in V1. *Nature* 2008, **454**(28):1110-1114.

doi:10.1186/1471-2202-14-S1-P299

Cite this article as: Ladenbauer *et al.*: How adaptation currents and synaptic inhibition change threshold, gain and variability of neuronal spiking. *BMC Neuroscience* 2013 **14**(Suppl 1):P299.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

