

The Abelian sandpile model on an infinite tree

Citation for published version (APA):

Maes, C., Redig, F. H. J., & Saada, E. (2002). The Abelian sandpile model on an infinite tree. *The Annals of Probability*, 30(4), 2081-2107. <https://doi.org/10.1214/aop/1039548382>

DOI:

[10.1214/aop/1039548382](https://doi.org/10.1214/aop/1039548382)

Document status and date:

Published: 01/01/2002

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

THE ABELIAN SANDPILE MODEL ON AN INFINITE TREE

BY CHRISTIAN MAES, FRANK REDIG AND ELLEN SAADA

K. U. Leuven, T. U. Eindhoven and C.N.R.S., Rouen

We consider the standard Abelian sandpile process on the Bethe lattice. We show the existence of the thermodynamic limit for the finite volume stationary measures and the existence of a unique infinite volume Markov process exhibiting features of self-organized criticality.

1. Introduction. Markov processes for spatially extended systems have been around for about 30 years now and interacting particle systems have become a branch of probability theory with an increasing number of connections with the natural and human sciences. While standard techniques and general results have been collected in a number of books such as Liggett (1985), Chen (1992) and Toom (1990) and are capable of treating the infinite volume construction for stochastic systems with locally interacting components, some of the most elementary questions for long range and nonlocal dynamics have remained wide open. We have in mind the class of stochastic interacting systems that during the last decade have invaded the soft condensed matter literature and are sometimes placed under the common denominator of self-organizing systems.

Since the appearance of Bak, Tang and Wiesenfeld (1988), the concept of self-organized criticality (SOC) has excited much interest, and has been applied in a great variety of domains [see, e.g., Turcotte (1999) for an overview]. From the mathematical point of view, the situation is, however, quite unsatisfactory. The models exhibiting SOC are in general very boundary condition dependent [especially the Bak–Tang–Wiesenfeld (BTW) model in dimension 2], which suggests that the definition of an infinite volume dynamics poses a serious problem. Even the existence of a (unique) thermodynamic limit of the finite volume stationary measure is not clear. From the point of view of interacting particle systems no standard theorems are at our disposal. The infinite volume processes we are looking for will be non-Feller and cannot be constructed by monotonicity arguments as in the case of the one-dimensional BTW model [see Maes, Redig, Saada and Van Moffaert (2000)] or the long-range exclusion process [see Liggett (1980)]. On the other hand, to make mathematically exact statements about SOC, it is necessary to have some kind of infinite volume limit, both for statics and for dynamics.

In this paper we continue our study of the BTW model for the case of the Bethe lattice; this is the Abelian sandpile model on an infinite tree. For this system, many

Received December 2000; revised February 2002.

AMS 2000 subject classifications. Primary 82C22; secondary 60K35.

Key words and phrases. Sandpile dynamics, nonlocal interactions, interacting particle systems, thermodynamic limit.

exact results were obtained in [Dhar and Majumdar (1990)]. In contrast to the one-dimensional case this system has a nontrivial stationary measure. We show here that the finite volume stationary measures converge to a unique measure μ which is not Dirac and exhibits all the properties of a SOC state. We then turn to the construction of a *stationary* Markov process starting from this measure μ . The main difficulty to overcome is the strong nonlocality: adding a grain at some lattice site x can influence the configuration far from x . In fact the cluster of sites influenced by adding at some fixed site has to be thought of as a critical percolation cluster which is almost surely finite but not of integrable size. The process we construct is intuitively described as follows: at each site x of the Bethe lattice we have an exponential clock which rings at rate $\varphi(x)$. At the ringing of the clock we add a grain at x . Depending on the addition rate $\varphi(x)$, we show existence of a stationary Markov process which corresponds to this description. We also extend this stationary dynamics to initial configurations which are typical for a measure μ' that is stochastically below μ .

The paper is organized as follows. In Section 2 we introduce standard results on finite volume Abelian sandpile models and summarize some specific results of [Dhar and Majumdar (1990)] for the Bethe lattice which we need for the infinite volume construction. In Section 3 we present the results on the thermodynamic limit of the finite volume stationary measures and on the existence of infinite volume Markovian dynamics. Section 4 is devoted to proofs and contains some additional remarks.

2. Finite volume Abelian sandpiles. In this section we collect some results on Abelian sandpiles on finite graphs which we will need later on. Most of these results are contained in the review paper by Dhar (1999) or in Ivashkevich and Prigogine (1998).

2.1. Toppling matrix. Let V denote a finite set of sites. We will always suppose that V is a nearest neighbor connected subset of \mathbb{Z}^d or of \mathbb{T}_d , the infinite homogeneous tree of degree $d + 1$. Starting from Section 3, we specify to the tree.

A matrix $\Delta = (\Delta_{x,y})_{x,y \in V}$ indexed by the elements of V is called a *toppling matrix* if the following hold:

1. for all $x, y \in V$, $x \neq y$, $\Delta_{x,y} = \Delta_{y,x} \leq 0$;
2. for all $x \in V$, $\Delta_{x,x} \geq 1$;
3. for all $x \in V$, $\sum_{y \in V} \Delta_{x,y} \geq 0$;
4. $\sum_{x,y \in V} \Delta_{x,y} > 0$.

The fourth condition ensures that there are sites (so-called *dissipative sites*) for which the inequality in the third condition is strict. This is fundamental for having a well-defined toppling rule later on.

Starting from Section 3 of this paper we will choose Δ to be the lattice Laplacian with open boundary conditions. More explicitly,

$$(1) \quad \Delta_{x,x} = \begin{cases} 2d, & \text{if } V \subset \mathbb{Z}^d, \\ d+1, & \text{if } V \subset \mathbb{T}_d, \end{cases}$$

$$\Delta_{x,y} = -1 \quad \text{if } x \text{ and } y \text{ are nearest neighbors.}$$

The dissipative sites then correspond to the boundary sites of V . The results on the finite volume Abelian sandpile in this section remain valid for a general toppling matrix Δ .

2.2. Configurations. A height configuration η is a mapping from V to $\mathbb{N} = \{1, 2, \dots\}$ assigning to each site a natural number $\eta(x) \geq 1$ (“the number of sand grains” at site x). A configuration $\eta \in \mathbb{N}^V$ is called *stable* if, for all $x \in V$, $\eta(x) \leq \Delta_{x,x}$. Otherwise η is *unstable*. We denote by Ω_V the set of all stable height configurations. For $\eta \in \mathbb{N}^V$ and $V' \subset V$, $\eta_{V'}$ denotes the restriction of η to V' .

2.3. Toppling rule. The *toppling of a site* x corresponding to the toppling matrix Δ is the mapping

$$T_x : \mathbb{N}^V \times V \rightarrow \mathbb{N}^V$$

defined by

$$(2) \quad T_x(\eta)(y) = \begin{cases} \eta(y) - \Delta_{x,y}, & \text{if } \eta(x) > \Delta_{x,x}, \\ \eta(y), & \text{otherwise.} \end{cases}$$

In words, site x topples if and only if its height is strictly larger than $\Delta_{x,x}$, by transferring $-\Delta_{x,y}$ grains to site $y \neq x$ and losing $\Delta_{x,x}$ grains. Toppling rules commute on unstable configurations. This means, for $x, z \in V$ and η such that $\eta(x) > \Delta_{x,x}$ and $\eta(z) > \Delta_{z,z}$,

$$(3) \quad T_x T_z(\eta) = T_z T_x(\eta).$$

For $\eta \in \mathbb{N}^V$, we say that $\zeta \in \Omega_V$ arises from η by toppling if there exists an n -tuple (x_1, \dots, x_n) of sites in V such that

$$(4) \quad \zeta = \left(\prod_{i=1}^n T_{x_i} \right)(\eta).$$

The *toppling transformation* is the mapping

$$\mathcal{T} : \mathbb{N}^V \rightarrow \Omega_V$$

defined by the requirement that $\mathcal{T}(\eta)$ arises from η by toppling. The fact that stabilization of an unstable configuration is always possible follows from the existence of dissipative sites (only a finite number of sites have to be toppled a

finite number of times). Thus only the fact that \mathcal{T} is *well defined* is less trivial: one has to prove here that for a given unstable configuration every possible stabilization makes *the same sites topple the same number of times*. Moreover, by (3) the order of the T_{x_i} in the product (4) is not important. A complete proof can be found in Meester (2002).

2.4. *Addition operators.* For $\eta \in \mathbb{N}^V$ and $x \in V$, let η^x denote the configuration obtained from η by adding one grain to site x , that is, $\eta^x(y) = \eta(y) + \delta_{x,y}$. The *addition operator* defined by

$$(5) \quad a_x : \Omega_V \rightarrow \Omega_V, \quad \eta \mapsto a_x \eta = \mathcal{T}(\eta^x)$$

represents the effect of adding a grain to the stable configuration η and letting a stable configuration arise by toppling. Because \mathcal{T} is well defined, the composition of addition operators is commutative: for all $\eta \in \Omega_V$, $x, y \in V$,

$$a_x(a_y \eta) = a_y(a_x \eta).$$

2.5. *Finite volume dynamics.* Let p denote a nondegenerate probability measure on V , that is, numbers p_x , $0 < p_x < 1$, with $\sum_{x \in V} p_x = 1$. We define a discrete time Markov chain $\{\eta_n : n \geq 0\}$ on Ω_V by picking a point $x \in V$ according to p at each discrete time step and applying the addition operator a_x to the configuration. This Markov chain has the transition operator

$$(6) \quad Pf(\eta) = \sum_{x \in V} p_x f(a_x \eta).$$

We can equally define a continuous time Markov process $\{\eta_t : t \geq 0\}$ with infinitesimal generator

$$(7) \quad L^\varphi f(\eta) = \sum_{x \in V} \varphi(x) [f(a_x \eta) - f(\eta)],$$

generating a pure jump process on Ω_V , with addition rate $\varphi(x) > 0$ at site x .

2.6. *Recurrent configurations, stationary measure.* We see here that the Markov chain $\{\eta_n, n \geq 0\}$ has only one recurrent class and its stationary measure is the uniform measure on that class.

Let us call \mathcal{R}_V the set of recurrent configurations for $\{\eta_n : n \geq 0\}$, that is, those for which $P_\eta(\eta_n = \eta \text{ infinitely often}) = 1$, where P_η denotes the distribution of $\{\eta_n : n \geq 0\}$ starting from $\eta_0 = \eta \in \Omega_V$. In the following proposition we list some properties of \mathcal{R}_V . For the sake of completeness we include the proof which can also be found in Meester (2002).

PROPOSITION 2.1. (i) \mathcal{R}_V contains only one recurrent class.

(ii) The addition operators a_x generate an Abelian group G of permutations of \mathcal{R}_V .

- (iii) The group G acts transitively on \mathcal{R}_V . In particular $|G| = |\mathcal{R}_V|$.
- (iv) $|\mathcal{R}_V| = \det \Delta$.

PROOF. (i) We write $\eta \leftrightarrow \zeta$ if in the Markov chain ζ can be reached from η with positive probability. Since sand is added with positive probability on all sites ($p_x > 0$), the maximal configuration η_{\max} defined by

$$\eta_{\max}(x) = \Delta_{x,x}$$

can be reached from any other configuration. Hence, if $\eta \in \mathcal{R}_V$, then $\eta \leftrightarrow \eta_{\max}$; therefore $\eta_{\max} \in \mathcal{R}_V$ and $\eta_{\max} \leftrightarrow \eta$ [see, e.g., Chung (1960), page 19].

- (ii) Fix $\eta \in \mathcal{R}_V$; then there exist $n_y \geq 1$ such that

$$\prod_{y \in V} a_y^{n_y} \eta = \eta$$

and

$$g_x = a_x^{n_x - 1} \prod_{y \in V, y \neq x} a_y^{n_y}$$

satisfies $(a_x g_x)(\eta) = (g_x a_x)(\eta) = \eta$. The set

$$\mathcal{R}^x = \{\zeta \in \mathcal{R}_V : (a_x g_x)(\zeta) = \zeta\}$$

is closed under the action of a_x , contains η , hence also η_{\max} : it is a recurrent class. By (i), $\mathcal{R}^x = \mathcal{R}_V$, $a_x g_x$ is the neutral element e and $g_x = a_x^{-1}$ if we restrict a_x to \mathcal{R}_V .

(iii) Fix $\zeta \in \mathcal{R}_V$ and put $\Psi_\zeta : G \rightarrow \mathcal{R}_V; g \mapsto g(\zeta)$. As before $\Psi_\zeta(G)$ is a recurrent class; hence $\Psi_\zeta(G) = \mathcal{R}_V$. If for $g, h \in G$, $\Psi_\zeta(g) = \Psi_\zeta(h)$, then $gh^{-1}(\zeta) = \zeta$, and by commutativity $gh^{-1}(g'\zeta) = g'\zeta$ for any $g' \in G$. Therefore $gh^{-1}(\xi) = \xi$ for all $\xi \in \mathcal{R}_V$; thus $g = h$. This proves that Ψ_ζ is a bijection from G to \mathcal{R}_V .

(iv) Adding $\Delta_{x,x}$ particles at a site $x \in V$ makes the site topple, and $-\Delta_{x,y}$ particles are transferred to y . This gives

$$a_x^{\Delta_{x,x}} = \prod_{y \neq x} a_y^{-\Delta_{x,y}}$$

On \mathcal{R}_V the a_x can be inverted and we obtain the closure relation

$$(8) \quad \prod_{y \in V} a_y^{\Delta_{x,y}} = e.$$

Write

$$\Delta \mathbb{Z}^V = \{\Delta n : n = (n_y)_{y \in V} \in \mathbb{Z}^V\},$$

where

$$(\Delta n)_x = \sum_{y \in V} \Delta_{x,y} n_y.$$

Consider

$$(9) \quad \Psi : \mathbb{Z}^V \rightarrow G : n \mapsto \prod_{x \in V} a_x^{n_x}.$$

The map Ψ is a homomorphism from \mathbb{Z}^V onto G , that is, $\Psi(n + m) = \Psi(n)\Psi(m)$ and $\Psi(\mathbb{Z}^V) = G$. Therefore, G is isomorphic to the quotient $\mathbb{Z}^V / \text{Ker}(\Psi)$. By (8), $\Delta\mathbb{Z}^V$ is contained in $\text{Ker}(\Psi)$. Conversely, let $n \in \text{Ker}(\Psi)$, and put $n = n^+ - n^-$, where $n_x^+ = \max\{n_x, 0\}$, $n_x^- = \max\{-n_x, 0\}$. Since $\Psi(n) = e$, adding to a recurrent $\eta \in \mathcal{R}_V$ according to n^+ has the same effect as adding according to n^- . Therefore, there exist $k^+, k^- \in (\mathbb{Z}^+)^V$, $\zeta \in \mathcal{R}_V$ such that

$$\eta + n^+ - \Delta k^+ = \zeta = \eta + n^- - \Delta k^-$$

and we conclude $n \in \Delta\mathbb{Z}^V$. This shows that G is isomorphic to $\mathbb{Z}^V / \Delta\mathbb{Z}^V$ and the latter group has cardinality $\det \Delta$. \square

As a consequence of the group property of G , the unique stationary measure is uniform on \mathcal{R}_V .

COROLLARY 2.1. (i) *The measure*

$$(10) \quad \mu_V = \sum_{\eta \in \mathcal{R}_V} \frac{1}{|\mathcal{R}_V|} \delta_\eta$$

is invariant under the action of a_x , $x \in V$ (δ_η is the Dirac measure on configuration η).

(ii) *On $L^2(\mu_V)$ the adjoint of a_x is*

$$(11) \quad a_x^* = a_x^{-1}.$$

REMARK. This shows that μ_V is invariant under the Markov processes generated by (6) and (7).

2.7. *Allowed configurations.* Given a configuration $\eta \in \Omega_V$, we say that its restriction η_W to a nonempty subset $W \subset V$ is a *forbidden subconfiguration* if, for all $x \in W$,

$$\eta(x) \leq \sum_{y \in W, y \neq x} (-\Delta_{x,y}).$$

A configuration $\eta \in \Omega_V$ is called *allowed* if it does not contain a forbidden subconfiguration. We denote by \mathcal{A}_V the set of all allowed configurations.

PROPOSITION 2.2.

$$\mathcal{A}_V = \mathcal{R}_V.$$

It is easy to see that toppling or adding cannot create a forbidden subconfiguration, which immediately implies $\mathcal{A}_V \supset \mathcal{R}_V$. For a proof that $\mathcal{A}_V = \mathcal{R}_V$ using spanning trees see Ivashkevich and Priezzhev (1998); a direct proof can be found in Meester (2002). For a generalization to nonsymmetric toppling matrices, see Speer (1993).

The property of having a forbidden subconfiguration in $W \subset V$ only depends on the heights at sites $x \in W$. Therefore $\eta \in \mathcal{R}_V$ implies $\eta_W \in \mathcal{R}_W$. This “consistency” property enables us to define allowed configurations on infinite sets.

2.8. *Expected toppling numbers.* For $x, y \in V$ and $\eta \in \Omega_V$, let $n_V(x, y, \eta)$ denote the number of topplings at site $y \in V$ by adding a grain at $x \in V$, that is, the number of times we have to apply the operator T_y to stabilize η^x . Define

$$(12) \quad G_V(x, y) = \int \mu_V(d\eta) n_V(x, y, \eta).$$

Writing down balance between inflow and outflow at site y , one obtains [cf. Dhar (1990)]

$$\sum_{z \in V} \Delta_{x,z} G_V(z, y) = \delta_{x,y},$$

which yields

$$G_V(x, y) = (\Delta)_{x,y}^{-1}.$$

In the limit $V \uparrow S$ (where S is \mathbb{Z}^d or the infinite tree), G_V converges to the Green’s function of the simple random walk on S .

2.9. *Some specific results for the tree.* When V_n is a binary tree of n generations, many explicit results have been obtained in Dhar and Majumdar (1990). We summarize here the results we need for the construction in infinite volume.

1. When adding a grain on a particular site $0 \in V_n$ of height 3, the set of toppled sites is the connected cluster $C_3(0, \eta)$ of sites including 0 having height 3. This cluster is distributed as a *random animal* (i.e., its distribution only depends on its cardinality, not on its form). Moreover,

$$(13) \quad \lim_{n \uparrow \infty} \mu_{V_n}(|C_3(0, \eta)| = k) \simeq Ck^{-3/2}$$

as k goes to infinity. The notation \simeq means that if we multiply the left-hand side of (13) by $k^{3/2}$, then the limit $k \rightarrow \infty$ is some strictly positive constant C .

2. When adding a grain on site x , the expected number of topplings at site y satisfies

$$(14) \quad \lim_{n \uparrow \infty} \int \mu_{V_n}(d\eta) n_{V_n}(x, y, \eta) = G(x, y),$$

where $G(x, y)$ is the Green’s function of the simple random walk on the infinite tree, that is,

$$(15) \quad G(0, x) = C2^{-|x|},$$

and $|x|$ is the “generation number” of x in the tree.

3. The correlations in the finite volume measures μ_{V_k} can be estimated in terms of the eigenvalues of a product of *transfer matrices*. This formalism is explained in detail in Dhar and Majumdar (1990), Section 5. Let f, g be two local functions whose dependence sets (see below a precise definition) are separated by n generations. To estimate the truncated correlation function

$$(16) \quad \mu_{V_k}(f; g) = \int fg d\mu_{V_k} - \int f d\mu_{V_k} \int g d\mu_{V_k},$$

consider the product of matrices

$$(17) \quad M_n^k = \prod_{i=1}^n \begin{pmatrix} 1 + \gamma_i^{k,n} & 1 + \gamma_i^{k,n} \\ 1 & 2 + \gamma_i^{k,n} \end{pmatrix},$$

where $\gamma_i^{k,n} \in [0, 1]$. The meaning of $\gamma_i^{k,n}$ is explained in Dhar and Majumdar (1990), but we will only use the fact $0 \leq \gamma_i^{k,n} \leq 1$ in Lemma 4.1 below. Let $\lambda_m^{n,k}$ (resp. $\lambda_M^{n,k}$) denote the smallest (resp. largest) eigenvalue of M_n^k . Then

$$(18) \quad \mu_{V_k}(f; g) \leq C(f, g) \frac{\lambda_m^{n,k}}{\lambda_M^{n,k}}.$$

For sites i far from the boundary of V_k , that is, for fixed i and n , in the limit $k \rightarrow \infty$, $\gamma_i^{k,n}$ tends to 1, and the correlations between local functions in the limit $V_k \rightarrow S$ are then governed by the maximal and minimal eigenvalues of $M_n = \begin{pmatrix} 2 & 2 \\ 1 & 3 \end{pmatrix}^n$. We shall need the estimate of a local function with a function living on the boundary of V_k ; therefore we have to use the full expression (17), (18).

3. Main results.

3.1. *Notation, definitions.* From now on, S denotes the infinite rootless binary tree, $V \subset S$ a finite subset of S ; Ω_V is the set of stable configurations in V , that is, $\Omega_V = \{\eta : V \rightarrow \{1, 2, 3\}\}$, and the set of all infinite volume stable configurations is $\Omega = \{1, 2, 3\}^S$. The set Ω is endowed with the product topology, making it into a compact metric space. For $\eta \in \Omega$, η_V is its restriction to V , and for $\eta, \zeta \in \Omega$, $\eta_V \zeta_{V^c}$ denotes the configuration whose restriction to V (resp. V^c) coincides with η_V (resp. ζ_{V^c}). As in the previous section, $\mathcal{R}_V \subset \Omega_V$ is the set of all allowed (or recurrent) configurations in V , and we define

$$(19) \quad \mathcal{R} = \{\eta \in \Omega : \forall V \subset S \text{ finite, } \eta_V \in \mathcal{R}_V\}.$$

A function $f : \Omega \rightarrow \mathbb{R}$ is *local* if there is a finite $V \subset S$ such that $\eta_V = \zeta_V$ implies $f(\eta) = f(\zeta)$. The minimal (in the sense of set ordering) such V is called the *dependence set* of f and is denoted by D_f . A local function can be seen as a function on Ω_V for all $V \supset D_f$, and every function on Ω_V can be seen as a local function on Ω . The set \mathcal{L} of all local functions is uniformly dense in the set $\mathcal{C}(\Omega)$ of all continuous functions on Ω .

Throughtout the paper, we use the following notion of limit by inclusion for a function f on the finite subsets of the tree with values in a metric space (K, d) :

DEFINITION 3.1. Let $\mathcal{S} = \{V \subset S, V \text{ finite}\}$ and $f : \mathcal{S} \rightarrow (K, d)$. Then

$$\lim_{V \uparrow S} f(V) = \kappa$$

if, for all $\varepsilon > 0$, there exists $V_0 \in \mathcal{S}$ such that, for all $V \supset V_0$, $d(f(V), \kappa) < \varepsilon$.

DEFINITION 3.2. A collection of probability measures ν_V on Ω_V is a Cauchy net if, for any local f and for any $\varepsilon > 0$, there exists $V_0 \supset D_f$ such that, for any $V, V' \supset V_0$,

$$\left| \int f(\eta) \nu_V(d\eta) - \int f(\eta) \nu_{V'}(d\eta) \right| \leq \varepsilon.$$

A Cauchy net converges to a probability measure ν in the following sense: the mapping

$$\Psi : \mathcal{L} \rightarrow \mathbb{R}, \quad f \mapsto \Psi(f) = \lim_{V \uparrow S} \int f d\nu_V$$

defines a continuous linear functional on \mathcal{L} [hence on $\mathcal{C}(\Omega)$] which satisfies $\Psi(f) \geq 0$ for $f \geq 0$ and $\Psi(1) = 1$. Thus by the Riesz representation theorem there exists a unique probability measure on Ω such that $\Psi(f) = \int f d\nu$. We write $\nu_V \rightarrow \nu$ and call this ν the *infinite volume limit* of ν_V .

We will also often consider an enumeration of the tree S , $\{x_0, x_1, \dots, x_n, \dots\}$, and put

$$(20) \quad T_n = \{x_0, \dots, x_n\}.$$

3.2. Thermodynamic limit of stationary measures.

THEOREM 3.1. *The set \mathcal{R} defined in (19) is a perfect set; that is, the following hold:*

- (i) \mathcal{R} is compact.
- (ii) The interior of \mathcal{R} is empty.
- (iii) For all $\eta \in \mathcal{R}$ there exists a sequence $\eta_n \neq \eta$, $\eta_n \in \mathcal{R}$, converging to η .

For $\eta \in \Omega$, $x \in S$ we denote by $C_3(x, \eta)$ the nearest neighbor connected cluster of sites containing the site x and having height 3.

THEOREM 3.2. *The finite volume stationary measures μ_V defined in (10) form a Cauchy net. Their infinite volume limit μ satisfies the following:*

- (i) $\mu(\mathcal{R}) = 1$;
- (ii) μ is invariant under tree automorphisms and mixing;
- (iii) $\mu(\eta : |C_3(0, \eta)| < \infty) = 1$;
- (iv) $\int |C_3(0, \eta)| \mu(d\eta) = \infty$.

REMARK. Theorem 3.2(iii) above remains true for the set $C_1(0, \eta)$, the nearest neighbor connected cluster of sites containing the origin and having height 1, and probably also for $C_2(0, \eta)$ but this we have not been able to prove.

3.3. Infinite volume dynamics. The finite volume addition operators $a_{x,V}$ [cf. (5)] can be extended to Ω via

$$(21) \quad a_{x,V} : \Omega \rightarrow \Omega, \quad \eta \mapsto a_{x,V}\eta = (a_{x,V}\eta_V)_V \eta_{V^c}.$$

PROPOSITION 3.1. (i) *There exists a measurable subset Ω' of \mathcal{R} with $\mu(\Omega') = 1$ on which the limit*

$$(22) \quad \lim_{V \uparrow S} a_{x,V}\eta = a_x\eta$$

exists, and $a_x\eta \in \Omega'$. More precisely, Ω' is the subset of configurations for which all the clusters $C_3(x, \eta)$ are finite.

- (ii) *The measure μ of Theorem 3.2 is invariant under the action of a_x .*
- (iii) *For every $\eta \in \Omega'$, $a_x(a_y\eta) = a_y(a_x\eta)$, for all $x, y \in S$.*

We now construct a Markov process on μ -typical infinite volume configurations which can be described intuitively as follows. Let $\varphi : S \rightarrow (0, \infty)$; this function will be the *addition rate function*. To each site $x \in S$ we associate a Poisson process $N_\varphi^{t,x}$ (for different sites these Poisson processes are mutually independent) with rate $\varphi(x)$. At the event times of $N_\varphi^{t,x}$ we “add a grain” at x ; that is, we apply the addition operator a_x to the configuration. Then L_V^φ introduced in (7) generates a pure jump Markov process on Ω . Indeed, this operator is well defined and bounded on any $L^p(\mu)$ space by Proposition 3.1, which implies the following.

PROPOSITION 3.2. L_V^φ is the $L^p(\mu)$ generator of the stationary Markov process defined by

$$\exp(tL_V^\varphi)f = \int \left(\prod_{x \in V} a_x^{N_\varphi^{t,x}} f \right) d\mathbb{P},$$

where \mathbb{P} denotes the joint distribution of the independent Poisson processes $N_\varphi^{t,x}$, and $f \in L^p(\mu)$.

The following condition on the addition rate φ is crucial in our construction. Remember $|x|$ is the generation number of x :

SUMMABILITY CONDITION.

$$(23) \quad \sum_{x \in S} \varphi(x) 2^{-|x|} < \infty.$$

In view of (15) the summability condition ensures that the number of topplings at any site $x \in S$ remains finite almost surely in any finite interval of time when grains are added at intensity φ .

THEOREM 3.3. *If φ satisfies condition (23), then we have the following:*

- (i) *The semigroups $S_V^\varphi(t) = \exp(tL_V^\varphi)$ converge strongly in $L^1(\mu)$ to a semigroup $S_\varphi(t)$.*
- (ii) *$S_\varphi(t)$ is the $L^1(\mu)$ semigroup of a stationary Markov process $\{\eta_t : t \geq 0\}$ on Ω .*
- (iii) *For any $f \in \mathcal{L}$,*

$$\lim_{t \downarrow 0} \frac{S_\varphi(t)f - f}{t} = L^\varphi f = \sum_{x \in S} \varphi(x)[a_x f - f],$$

where the limit is taken in $L^1(\mu)$.

REMARKS. (i) In Proposition 4.1, we show that $S_\varphi(t)$ is a strongly continuous function of φ .

(ii) In Proposition 4.2, we show that condition (23) is in some sense optimal.

THEOREM 3.4. *The process $\{\eta_t : t \geq 0\}$ of Theorem 3.3 admits a cadlag version (right-continuous with left limits).*

The intuitive description of the process $\{\eta_t : t \geq 0\}$ is actually correct under condition (23); that is, the process has a representation in terms of Poisson processes:

THEOREM 3.5. *If φ satisfies condition (23), for $\mu \times \mathbb{P}$ almost every (η, ω) the limit*

$$\lim_{V \uparrow S} \prod_{x \in V} a_x^{N_\varphi^{t,x}(\omega)} \eta = \eta_t$$

exists. The process $\{\eta_t : t \geq 0\}$ is a version of the process of Theorem 3.3; that is, its $L^1(\mu)$ semigroup coincides with $S_\varphi(t)$.

Finally, we can slightly generalize Theorem 3.5 to define the dynamics starting from a measure stochastically below μ . For $\eta, \zeta \in \Omega$, $\eta \leq \zeta$, if for all $x \in S$, $\eta(x) \leq \zeta(x)$. A function $f : \Omega \rightarrow \mathbb{R}$ is *monotone* if $\eta \leq \zeta$ implies $f(\eta) \leq f(\zeta)$. Two probability measures μ and ν satisfy $\mu \leq \nu$ if, for all monotone functions, $\int f d\mu \leq \int f d\nu$.

THEOREM 3.6. *Let $\mu' \leq \mu$. If φ satisfies condition (23), for $\mu' \times \mathbb{P}$ almost every (η, ω) the limit*

$$\lim_{V \uparrow S} \prod_{x \in V} a_x^{N_\varphi^{t,x}(\omega)} \eta = \eta_t$$

exists. The process $\{\eta_t : t \geq 0\}$ is Markovian with η_0 distributed according to μ' .

REMARK. The last theorem implies that $\eta \equiv 1$ can be taken as initial configuration.

4. Proofs. This section is devoted to the proofs of the results described above. Some of them will be put in a slightly more general framework so that they can be applied to other cases (where S is not a binary tree or where we have other addition operators a_x) as soon as the existence of a thermodynamic limit of the finite volume stationary measures is guaranteed. The essential cause of difficulty is the nonlocality of the addition operators. The essential simplification is the Abelian property.

4.1. *Thermodynamic limit of stationary measures.*

PROOF OF THEOREM 3.1. (i), (ii) To see that \mathcal{R} has empty interior, notice that if $\eta \in \mathcal{R}$, there do not exist $x, y \in S$ nearest neighbors such that $\eta(x) = \eta(y) = 1$ (that way, $\eta_{\{x,y\}}$ would be a forbidden subconfiguration). Finally \mathcal{R} is closed as an intersection of closed sets.

(iii) Let η_{\max} be the maximal configuration, $\eta_{\max}(x) = 3$ for all $x \in S$. If $\eta_V \in \mathcal{R}_V$, then $\eta_V(\eta_{\max})_{V^c} \in \mathcal{R}$. Therefore any $\eta \in \mathcal{R}$ containing an infinite number of sites x for which $\eta(x) \neq 3$ has property (iii) of Theorem 3.1. If $\eta \in \mathcal{R}$ contains only a finite number of sites having height 1 or 2, then we choose a sequence $\Sigma = \{x_n : n \in \mathbb{N}\} \subset \{x \in S : \eta(x) = 3 \text{ and } \eta(y) = 3 \text{ for any neighbor of } x\}$ such that two elements of Σ are never nearest neighbors, and $|x_n|$ is strictly increasing in n . We then define $\eta_n(x) = \eta(x)$ for $x \in S \setminus \{x_k \in \Sigma : 0 \leq k \leq n\}$ and $\eta_n(x_k) = 2$ for $x_k \in \Sigma$, $k \geq n + 1$. These η_n belong to \mathcal{R} since they do not contain any forbidden subconfiguration, and $\eta_n \rightarrow \eta$. \square

PROOF OF THEOREM 3.2. We use T_n introduced in (20), but with the x_i such that $n \leq m$ implies that the generation numbers satisfy $|x_n| \leq |x_m|$. Then we have

$$(24) \quad |x_n| \simeq \log_2 n.$$

To prove that the probability measures μ_V form a Cauchy net, it is sufficient to show that for any local function $f : \Omega \rightarrow \mathbb{R}$ we have

$$(25) \quad \sum_n \left| \int f d\mu_{T_n} - \int f d\mu_{T_{n+1}} \right| < \infty.$$

We do it for $f(\eta) = \eta(x_0)$ by giving an upper bound of the difference $\int f d\mu_{T_n} - \int f d\mu_{T_{n+1}}$ by a truncated correlation function [cf. (16)]. Then we estimate the latter by the transfer matrix method (cf. Section 2.9, item 3). We abbreviate in what follows $\mu_n = \mu_{T_n}$, $\mathcal{R}_n = \mathcal{R}_{T_n}$.

LEMMA 4.1.

$$|\mu_{n+1}[\eta(x_0)] - \mu_n[\eta(x_0)]| \leq 3\mu_{n+1}[\eta(x_0); I(\eta(x_{n+1}) = 3)].$$

PROOF. Every $\eta \in \mathcal{R}_n$ can be extended to an element of \mathcal{R}_{n+1} by putting $\eta(x_{n+1}) = 3$: indeed, adding a site with height 3 cannot create a forbidden subconfiguration. Moreover

$$\{\eta_{T_n} : \eta \in \mathcal{R}_{n+1}, \eta(x_{n+1}) = 3\} = \mathcal{R}_n;$$

thus

$$\mu_{n+1}[\eta(x_{n+1}) = 3] = \frac{|\mathcal{R}_n|}{|\mathcal{R}_{n+1}|}.$$

Since, for any n , $|\mathcal{R}_{n+1}| \leq 3|\mathcal{R}_n|$, we have

$$(26) \quad \mu_{n+1}[\eta(x_{n+1}) = 3] \geq \frac{1}{3},$$

$$\begin{aligned} \mu_n(\eta(x_0)) &= \sum_{\eta \in \mathcal{R}_n} \frac{1}{|\mathcal{R}_n|} \eta(x_0) \\ &= \sum_{\eta \in \mathcal{R}_{n+1}} \frac{1}{|\mathcal{R}_{n+1}|} \eta(x_0) I(\eta(x_{n+1}) = 3) \frac{|\mathcal{R}_{n+1}|}{|\mathcal{R}_n|} \\ &= \sum_{\eta \in \mathcal{R}_{n+1}} \frac{1}{|\mathcal{R}_{n+1}|} \eta(x_0) I(\eta(x_{n+1}) = 3) \frac{1}{\mu_{n+1}[\eta(x_{n+1}) = 3]}. \end{aligned}$$

Therefore

$$|\mu_{n+1}(\eta(x_0)) - \mu_n(\eta(x_0))| \leq \frac{\mu_{n+1}[\eta(x_0); I(\eta(x_{n+1}) = 3)]}{\mu_{n+1}[\eta(x_{n+1}) = 3]}.$$

The lemma follows now from (26). \square

Recalling Section 2.9, item 3, we have

$$(27) \quad \mu_n[\eta(x_0); I(\eta(x_n) = 3)] \leq C \frac{\lambda_m^{|x_n|, |x_n|}}{\lambda_M^{|x_n|, |x_n|}}.$$

LEMMA 4.2.

$$\sum_{n=1}^{+\infty} \frac{\lambda_m^{|x_n|, |x_n|}}{\lambda_M^{|x_n|, |x_n|}} < +\infty.$$

PROOF. We abbreviate $\lambda_m^{(n)} = \lambda_m^{|x_n|, |x_n|}$, $\lambda_M^{(n)} = \lambda_M^{|x_n|, |x_n|}$, $M(n) = M_{|x_n|}^{|x_n|}$, $\gamma_i = \gamma_i^{|x_n|, |x_n|}$. In terms of the trace and the determinant of $M(n)$ we have

$$\begin{aligned} \lambda_M^{(n)} &= \frac{1}{2} \left(\text{Tr}(M(n)) + \sqrt{[\text{Tr}(M(n))]^2 - 4 \det(M(n))} \right) \\ \lambda_m^{(n)} &= \frac{1}{2} \left(\text{Tr}(M(n)) - \sqrt{[\text{Tr}(M(n))]^2 - 4 \det(M(n))} \right). \end{aligned}$$

Therefore,

$$\lim_{n \uparrow \infty} \left(\frac{\lambda_m^{(n)}}{\lambda_M^{(n)}} \right) \left(\frac{[\text{Tr}(M(n))]^2}{\det(M(n))} \right) = 1.$$

To prove the lemma we show that [cf. (24)]

$$(28) \quad \left(\frac{\det(M(n))}{[\text{Tr}(M(n))]^2} \right) \leq \left(\frac{4}{9} \right)^{|x_n|}.$$

Use

$$\det(M(n)) = \prod_{i=1}^{|x_n|} (1 + \gamma_i)^2$$

and

$$\begin{aligned} \text{Tr}(M(n)) &\geq \text{Tr} \left(\prod_{i=1}^{|x_n|} \begin{pmatrix} 1 + \gamma_i & 0 \\ 0 & 2 + \gamma_i \end{pmatrix} \right) \\ &= \prod_{i=1}^{|x_n|} (1 + \gamma_i) + \prod_{i=1}^{|x_n|} (2 + \gamma_i), \end{aligned}$$

to estimate [for $1 \leq i \leq |x_n|$, $2(2 + \gamma_i) \geq 3(1 + \gamma_i)$]

$$\begin{aligned} \left(\frac{\det(M(n))}{[\text{Tr}(M(n))]^2} \right) &\leq \left(1 + 2 \prod_{i=1}^{|x_n|} \left[\frac{2 + \gamma_i}{1 + \gamma_i} \right] + \prod_{i=1}^{|x_n|} \left[\frac{2 + \gamma_i}{1 + \gamma_i} \right]^2 \right)^{-1} \\ &\leq \left(1 + 2 \cdot \left(\frac{3}{2} \right)^{|x_n|} + \left(\frac{3}{2} \right)^{2|x_n|} \right)^{-1} \\ &\leq \left(\frac{4}{9} \right)^{|x_n|}. \end{aligned}$$

□

REMARK. Lemmas 4.1 and 4.2 extend to a general local function f with dependence set $D_f \subset T_n$ containing N_0 generations. To obtain Lemma 4.1, replace $\eta(x_0)$ by $f(\eta)$ in its proof. For Lemma 4.2, we have to replace $M_{|x_n|}^{|x_n|}$ by $M_{|x_n|}^{|x_n|-N_0}$, and in the computation $|x_n|$ by $|x_n| - N_0$. Since N_0 is finite, the convergence in (25) is unaffected.

Note that by the same proof we obtain that for any two local functions f, g whose dependence sets D_f and D_g are separated by a distance of n generations we have

$$(29) \quad \mu(f; g) \leq C(f, g) \left(\frac{4}{9}\right)^n.$$

So far we have proved the fact that the uniform measures on \mathcal{R}_n converge to a unique measure μ . Clearly, μ concentrates on \mathcal{R} because for any given $V_0 \subset S$, and $V \supset V_0$, the restriction η_{V_0} is an element of \mathcal{R}_{V_0} with μ_V -probability 1.

To prove point (ii), let $\pi : S \rightarrow S$ be a tree automorphism; that is, π is bijective and x and y are neighbors if and only if $\pi(x)$ and $\pi(y)$ are neighbors. π acts on configurations $\eta \in \Omega$ via $\pi(\eta)(x) = \eta(\pi^{-1}(x))$. First note that $\mathcal{R}_V = \pi^{-1}(\mathcal{R}_{\pi(V)})$. Indeed, if for $W \subset V$, η_W is a forbidden subconfiguration, then $\pi(\eta)_{\pi(V)}$ is a forbidden subconfiguration of $\pi(\eta)$ and vice-versa. This implies that for the finite volume measures the equality $\mu_V \circ \pi = \mu_{\pi(V)}$ holds. Since μ_V is a Cauchy net, the same holds for $\mu_{\pi(V)}$; hence we conclude

$$\mu = \lim_V \mu_V = \lim_V \mu_{\pi(V)} = \lim_V \mu_V \circ \pi = \mu \circ \pi.$$

The mixing property follows from (29).

Points (iii) and (iv) of the theorem follow from the equality

$$\lim_{k \rightarrow \infty} k^{-3/2} \mu(|C_3(0, \eta)| = k) = c > 0,$$

which in turn follows from (13), the convergence of μ_n to μ and the fact that, for any k fixed, the indicator $I(|C_3(0, \eta)| = k)$ is a local function. \square

4.2. Infinite volume toppling operators.

DEFINITION 4.1. Given the finite volume addition operators $a_{x, V}$ [defined in (21)] acting on Ω , we call a configuration $\eta \in \Omega$ *normal* if for every $x \in S$ there exists a minimal finite set $V_x(\eta) \subset S$ such that, for all $V' \supset V \supset V_x(\eta)$,

$$a_{x, V'} \eta = a_{x, V} \eta.$$

In other words, for a normal η , outside $V_x(\eta)$, no sites are affected by adding a grain at x . In our case, when a particle is added at some site $x \in S$, the

cluster of toppled sites coincides with the cluster $C_3(x, \eta)$ of sites having height 3 including x ; thus

$$(30) \quad V_x(\eta) = C_3(x, \eta) \cup \partial_e C_3(x, \eta),$$

where ∂_e denotes the exterior boundary. Notice that for a normal configuration η , by definition,

$$(31) \quad a_x(\eta) = \lim_{V \uparrow S} a_{x,V}(\eta) = a_{x,V_x(\eta)}(\eta).$$

PROOF OF PROPOSITION 3.1. (i) We show that there is a full measure set Ω' of normal configurations. From (13) and Theorem 3.2,

$$\int I(|C_3(x, \eta)| = n) d\mu \simeq Cn^{-3/2}.$$

Therefore μ concentrates on the set Ω' of configurations for which all the clusters $C_3(x, \eta)$ are finite, hence for which η is normal. Since $I(|C_3(x, \eta)| = n)$ is measurable, Ω' is measurable. Moreover this set Ω' is closed under the action of the addition operators a_y , since [cf. (30)]

$$(32) \quad C_3(x, a_y\eta) \subset V_x(\eta) \cup V_y(\eta).$$

(ii) Choose $\varepsilon > 0$; pick a local function f ; fix $V_n \uparrow S$ and n_0 such that $n \geq n_0$ implies

$$(33) \quad \mu\{\eta \in \Omega : V_x(\eta) \not\subset V_n\} \leq \frac{\varepsilon}{4\|f\|_\infty + 1}.$$

This n_0 exists since μ concentrates on normal configurations. We estimate

$$\begin{aligned} & \left| \int f(a_x\eta) d\mu - \int f(\eta) d\mu \right| \\ & \leq \left| \int f(a_{x,V_n}\eta) d\mu - \int f(\eta) d\mu \right| + 2\|f\|_\infty \mu\{\eta \in \Omega : V_x(\eta) \not\subset V_n\} \\ & \leq \lim_m \left| \int f(a_{x,V_n}\eta) d\mu_{V_m} - \int f(\eta) d\mu_{V_m} \right| + \frac{\varepsilon}{2} \\ & \leq \frac{\varepsilon}{2} + 2\|f\|_\infty \lim_m \mu_{V_m}(a_{x,V_n}(\eta) \neq a_{x,V_m}(\eta)) \\ & = \frac{\varepsilon}{2} + 2\|f\|_\infty \left(1 - \lim_m \mu_{V_m}(V_x(\eta) \subset V_n)\right) \\ & = \frac{\varepsilon}{2} + 2\|f\|_\infty (1 - \mu(V_x(\eta) \subset V_n)) \leq \varepsilon. \end{aligned}$$

In the last step we used that the indicator $I(V_x(\eta) \subset V_n)$ is a local function.

(iii) Let $\eta \in \Omega'$, $x, y \in S$ be two different sites and let $V \supset V_x(\eta) \cup V_x(a_x\eta) \cup V_x(a_y\eta)$. Since $a_{x,v}$ and $a_{y,v}$ commute, we have

$$\begin{aligned} a_x(a_y\eta) &= a_x(a_{y,v}\eta) = a_{x,v}(a_{y,v}\eta) \\ &= a_{y,v}(a_{x,v}\eta) = a_{y,v}(a_x\eta) = a_y(a_x\eta). \end{aligned} \quad \square$$

4.3. *Infinite volume semigroup.* We now turn to the proofs of Theorems 3.3 and 3.4.

DEFINITION 4.2. We define the cluster of $\eta \in \Omega$ at $x \in S$ as

$$(34) \quad \mathcal{C}(x, \eta) = \{y \in S : a_y\eta(x) \neq \eta(x)\}$$

and put

$$(35) \quad G_\mu(x, y) = \int I(y \in \mathcal{C}(x, \eta)) d\mu(\eta).$$

Finally, for $\varphi : S \rightarrow [0, \infty)$, write

$$\begin{aligned} \|f\|_\varphi &= \sum_{x \in S} \varphi(x) \int \mu(d\eta) |f(a_x\eta) - f(\eta)|, \\ \mathcal{B}_\varphi &= \{f : \Omega \rightarrow \mathbb{R} : f \text{ bounded, } \|f\|_\varphi < \infty\}. \end{aligned}$$

REMARK. In our case $y \in \mathcal{C}(x, \eta)$ if $y = x$ or if $\eta(y) = 3$ and there exists a nearest neighbor path from y to a neighbor of x along sites having height 3. The estimate for $G_\mu(x, y)$ in Lemma 4.4 below is not dependent on the fact that we are on the tree.

LEMMA 4.3. *If*

$$(36) \quad \sum_{x \in S} \varphi(x) G_\mu(y, x) < \infty \quad \text{for all } y \in S,$$

then all local functions are in \mathcal{B}_φ .

PROOF. Let f be a local function with dependence set D_f . Then $f(a_x\eta) \neq f(\eta)$ if for $y \in D_f$, $a_x\eta(y) \neq \eta(y)$, that is, $x \in \mathcal{C}(y, \eta)$,

$$\begin{aligned} \|f\|_\varphi &= \sum_{x \in S} \varphi(x) \int |a_x f - f| d\mu \\ &= \int \sum_{x \in \bigcup_{y \in D_f} \mathcal{C}(y, \eta)} \varphi(x) |a_x f - f| d\mu \\ &\leq 2\|f\|_\infty \sum_{x \in S} \varphi(x) \int I\left(x \in \bigcup_{y \in D_f} \mathcal{C}(y, \eta)\right) d\mu \\ &\leq 2\|f\|_\infty \sum_{y \in D_f} \sum_{x \in S} G_\mu(y, x) \varphi(x) < \infty. \end{aligned} \quad \square$$

The next lemma provides a link between G_μ and the Green's function for simple random walk on S , that is, between conditions (36) and (23).

LEMMA 4.4.

$$G_\mu(x, y) \leq \sum_{z \sim x} G(y, z) = \delta_{x,y} + 3G(x, y),$$

where $z \sim x$ means that z and x are neighbors.

PROOF. We have to estimate the probability that $a_x \eta(y) \neq \eta(y)$. If by adding a grain at x we influence y , this can only be achieved by the toppling of one of the nearest neighbor sites of y . Since μ concentrates on normal configurations,

$$\begin{aligned} \mu(a_x \eta(y) \neq \eta(y)) &= \lim_{V \uparrow S} \mu(a_x \eta(y) \neq \eta(y), V_x(\eta) \cup V_y(\eta) \subset V) \\ &= \lim_{V \uparrow S} \lim_{W \uparrow S} \mu_W(a_{x,v} \eta(y) \neq \eta(y), V_x(\eta) \cup V_y(\eta) \subset V) \\ &= \lim_{V \uparrow S} \lim_{W \uparrow S} \mu_W(a_{x,w} \eta(y) \neq \eta(y), V_x(\eta) \cup V_y(\eta) \subset V) \\ (37) \quad &\leq \lim_{W \uparrow S} \mu_W(a_{x,w} \eta(y) \neq \eta(y)) \\ &\leq \lim_{W \uparrow S} \mu_W(\exists z \in W, z \sim y, n_W(z, y, \eta) \geq 1) \\ &\leq \lim_{W \uparrow S} \sum_{z \sim y} \int d\mu_W(\eta) n_W(z, y, \eta) \\ &= \sum_{z \sim y} G(z, y), \end{aligned}$$

where we used (12), (14), (30) and (32). \square

The following lemma completes the proof of Theorem 3.3 and shows that \mathcal{B}_φ is a natural core for the domain of the generator of the infinite volume semigroup.

LEMMA 4.5. (i) For $f \in \mathcal{B}_\varphi$ the net

$$(38) \quad S_V^\varphi(t) f = \exp(tL_V^\varphi) f = \exp\left(t \sum_{x \in V} \varphi(x)(a_x - I)\right) f$$

converges in $L^1(\mu)$ (as $V \uparrow S$) to a function $S_\varphi(t) f \in L^1(\mu)$. $f \mapsto S_\varphi(t) f$ defines a semigroup on \mathcal{B}_φ which is a contraction in both $L^1(\mu)$ and \mathcal{B}_φ norms.

(ii) Under condition (36), the semigroup $S_\varphi(t)$ corresponds to a unique Markov process on Ω .

PROOF. We denote by $\|f\|$ the $L^1(\mu)$ -norm of f , and we abbreviate $S_V(t) = S_V^\varphi(t)$, $S(t) = S^\varphi(t)$, $L_V = L_V^\varphi$.

(i) First note that $S_V(t)$ is well defined on $L^1(\mu)$ by Proposition 3.2. By the Abelian property [Proposition 3.1(iii)] we can write, for $V \subset V' \subset S$,

$$\|S_V(t)f - S_{V'}(t)f\| = \|(S_{V' \setminus V}(t) - I)S_V(t)f\|.$$

By Proposition 3.2, $S_V(t)$ is the semigroup of a stationary Markov process and hence a contraction on $L^1(\mu)$. Therefore

$$\begin{aligned} & \|S_V(t)(S_{V' \setminus V}(t) - I)f\| \\ & \leq \|(S_{V' \setminus V}(t) - I)f\| \\ (39) \quad & = \left\| \int_0^t L_{V' \setminus V} S_{V' \setminus V}(s)f \, ds \right\| \\ & \leq \int_0^t \|L_{V' \setminus V} f\| \, ds \\ & \leq t \sum_{x \in V' \setminus V} \varphi(x) \int |a_x - I|f| \, d\mu \rightarrow 0 \quad \text{as } V, V' \uparrow S, \end{aligned}$$

where the last step follows from $f \in \mathcal{B}_\varphi$. Hence $S_V(t)f \rightarrow S(t)f$ in $L^1(\mu)$. We show that $S(t)f \in \mathcal{B}_\varphi$:

$$\begin{aligned} & \sum_{x \in S} \varphi(x) \int |S(t)f(a_x \eta) - S(t)f(\eta)| \mu(d\eta) \\ & \leq \sum_{x \in S} \varphi(x) \int S(t)|a_x f - f| \, d\mu \\ & = \int \sum_{x \in S} \varphi(x) |a_x f - f| \, d\mu = \|f\|_\varphi. \end{aligned}$$

Thus $S(t)$ is also a contraction for the $\|\cdot\|_\varphi$ -norm. We finish with the semigroup property:

$$\begin{aligned} S(t)S(s)f &= \lim_{V \uparrow S} S_V(t)[S(s)f] \\ &= \lim_{V \uparrow S} \lim_{W \uparrow S} S_V(t)S_W(s)f \end{aligned}$$

and

$$S(t+s)f = \lim_{V \uparrow S} S_V(t)S_V(s)f.$$

Then, since $S_V(t)$ is a contraction in $L^1(\mu)$,

$$(40) \quad \|S_V(t)S_W(s)f - S_V(t)S_V(s)f\| \leq \|S_W(s)f - S_V(s)f\|.$$

By (39), the right-hand side of (40) goes to zero as $V, W \uparrow S$.

(ii) If condition (36) is met, then \mathcal{B}_φ contains all local functions by Lemma 4.3. Therefore, by contractivity the semigroup $S(t)$ on \mathcal{B}_φ uniquely extends to a semigroup of contractions on $L^1(\mu)$. Since, by Proposition 3.2, $S_V(t)$ is a Markov semigroup, so is $S(t)$, that is, $S(t)1 = 1$, $S(t)f \geq 0$ if $f \geq 0$. Hence, by Kolmogorov’s theorem there is a unique Markov process with semigroup $S(t)$. \square

REMARK. When $\varphi \equiv 1$, condition (36) is equivalent to

$$\sum_{x \in S} \int \mu(d\eta) I(x \in \mathcal{C}(y, \eta)) = \int |\mathcal{C}(y, \eta)| \mu(d\eta) < +\infty;$$

that is, the clusters must be integrable under μ . For models which exhibit “self-organized criticality,” $\mathcal{C}(y, \eta)$ is usually a “finite but critical percolation cluster,” implying that $\int |\mathcal{C}(y, \eta)| d\mu = \infty$ [cf. Theorem 3.2(iv), because $\mathcal{C}(y, \eta) \supset \partial_e C_3(y, \eta)$]. Therefore this formalism breaks down for addition rate $\varphi \equiv 1$.

The following lemma proves Theorem 3.4.

LEMMA 4.6. *Under condition (36), the process $\{\eta_t : t \geq 0\}$ of Theorem 3.3 is almost surely right-continuous, that is,*

$$(41) \quad \mathbb{P}_\mu \left[\lim_{t \downarrow 0} d(\eta_t, \eta_0) \geq \varepsilon \right] = 0,$$

where \mathbb{P}_μ is its path-space measure, and the distance d is defined below [in (44)].

PROOF. Pick a function $\Psi : S \rightarrow (0, 1)$ such that

$$(42) \quad \sum_{x \in S} \Psi(x) = 1$$

and

$$(43) \quad \sum_{x, y \in S} \varphi(x) G_\mu(x, y) \Psi(y) < \infty.$$

The distance

$$(44) \quad d(\eta, \zeta) = \sum_{x \in S} |\eta(x) - \zeta(x)| \Psi(x)$$

generates the product topology. Denote by \mathbb{E}_μ the expectation w.r.t. \mathbb{P}_μ . For $f_y(\eta) = \eta(y)$,

$$f_y(\eta_t) - f_y(\eta_0) = \int_0^t L^\varphi f_y(\eta_s) ds + M_t^y,$$

where M_t^y is a centered martingale with quadratic variation

$$(45) \quad \mathbb{E}_\mu[(M_t^y)^2] = \mathbb{E}_\mu \left[\int_0^t (L^\varphi f_y^2(\eta_s) - 2f_y(\eta_s)L^\varphi f_y(\eta_s)) ds \right].$$

Using stationarity of η_s and

$$\int d\mu |L^\varphi g| \leq 2\|g\|_\infty \sum_{x \in S} \sum_{y \in D_g} \varphi(x)G_\mu(y, x),$$

for a local bounded function on Ω , we obtain from (45)

$$\mathbb{E}_\mu[(M_t^y)^2] \leq Ct \sum_{x \in S} \varphi(x)G_\mu(y, x).$$

Now we can estimate

$$\begin{aligned} & \mathbb{P}_\mu \left[\exists s \leq t : \sum_{y \in S} |\eta_s(y) - \eta_0(y)| \Psi(y) \geq \varepsilon \right] \\ & \leq \mathbb{P}_\mu \left[\int_0^t ds \sum_{y \in S} |L^\varphi f_y(\eta_s)| \Psi(y) \geq \frac{\varepsilon}{2} \right] \\ & \quad + \mathbb{P}_\mu \left[\sup_{0 \leq s \leq t} \left| \sum_{y \in S} M_s^y \Psi(y) \right| \geq \frac{\varepsilon}{2} \right] \\ & \leq \left(\frac{12t}{\varepsilon} \right) \sum_{x, y \in S} \varphi(x)G_\mu(y, x) \Psi(y) + \left(\frac{2}{\varepsilon} \right)^2 \mathbb{E}_\mu \left| \sum_{y \in S} M_t^y \Psi(y) \right|^2 \\ & \leq \left(\frac{12t}{\varepsilon} \right) \sum_{x, y \in S} \varphi(x)G_\mu(y, x) \Psi(y) + \left(\frac{2}{\varepsilon} \right)^2 \mathbb{E}_\mu \left[\sum_{y \in S} (M_t^y)^2 \Psi(y) \right] \\ & \leq tC_\varepsilon \sum_{x, y \in S} \varphi(x)G_\mu(y, x) \Psi(y). \end{aligned}$$

Here we used Markov's and Doob's inequalities in the second step and the Cauchy–Schwarz inequality combined with (42) in the third step. The result (41) follows. \square

4.4. *Poisson representation.* In this section we prove Theorems 3.5 and 3.6. Intuitively it is clear from the Abelian property that the process for which we showed existence in the previous subsection can be represented as $\prod_{x \in S} a_x^{N_\varphi^{t,x}} \eta$, where $N_\varphi^{t,x}$ are independent Poisson processes of intensity $\varphi(x)$.

We take T_n as in (20). We say that the product $\prod_{x \in S} a_x^{n,x} \eta$ exists if for every $y \in S$ there exists N_y such that, for all $m, n \geq N_y$,

$$\left| \left[\prod_{x \in T_n} a_x^{n,x} \eta \right](y) - \left[\prod_{x \in T_m} a_x^{n,x} \eta \right](y) \right| = 0.$$

This is equivalent to the convergence of the sequence $\prod_{x \in T_n} a_x^{n,x} \eta$ in the product topology.

LEMMA 4.7. *Under condition (36), the product*

$$\prod_{x \in S} a_x^{N_\varphi^{t,x}} \eta = \tilde{\eta}_t$$

exists for μ -almost every realization of $N_\varphi^{t,x}$ and almost every η . The process $\{\tilde{\eta}_t : t \geq 0\}$ is a version of the Markov process of Lemma 4.5.

PROOF. Choose a realization of $N_\varphi^{t,x}$ such that

$$(46) \quad \sum_{x \in S} N_\varphi^{t,x} G_\mu(x, y) < \infty$$

for every y . This happens with probability 1 by condition (36). Define, for $\eta \in \Omega'$,

$$(47) \quad \eta_{T_n}(t) = \prod_{x \in T_n} a_x^{N_\varphi^{t,x}} \eta.$$

Under μ , $\eta_{T_n}(t)$ is stationary in n and t . We have

$$\begin{aligned} & \mu[|(\eta_{T_n}(t))(y) - (\eta_{T_{n+1}}(t))(y)| \geq 1] \\ & \leq \int \left| \left[a_{x_{n+1}}^{N_\varphi^{t,x_{n+1}}} \eta_{T_n}(t) \right](y) - (\eta_{T_n}(t))(y) \right| \mu(d\eta) \\ & = \int \left| \left[a_{x_{n+1}}^{N_\varphi^{t,x_{n+1}}} \eta \right](y) - \eta(y) \right| \mu(d\eta) \\ & \leq \int \sum_{j=1}^{N_\varphi^{t,x_{n+1}}} \left| [a_{x_{n+1}}^j \eta](y) - [a_{x_{n+1}}^{j-1} \eta](y) \right| \mu(d\eta) \\ & \leq 6 N_\varphi^{t,x_{n+1}} G_\mu(x_{n+1}, y). \end{aligned}$$

In the second and last steps we used the invariance of μ under a_x . By the Borel–Cantelli lemma, (46) implies that, for almost every realization of $N_\varphi^{t,x}$,

$$\mu[\exists n_0 : \forall n \geq n_0 (\eta_{T_n}(t))(y) = (\eta_{T_{n_0}}(t))(y)] = 1.$$

This proves μ -a.s. convergence of the product. To see that $\tilde{\eta}_t$ is a version of the Markov process with semigroup $S_\varphi(t)$, combine Proposition 3.2 with Theorem 3.3(i) to get, for any local function f ,

$$\int d\mu \left| \int d\mathbb{P} f(\tilde{\eta}_t) - S_\varphi(t) f \right| = 0.$$

In the preceding argument we used a particular enumeration of the countable set S . But changing it gives again a process with semigroup $S_\varphi(t)$. Therefore the limiting process will not depend (up to sets of measure zero) on the chosen enumeration of S . \square

PROOF OF THEOREM 3.6. For $\eta \in \Omega'$ and $y \in S$ we have the relation [remember (47)]

$$(48) \quad \eta_V(t)(y) = \eta(y) + I(y \in V) \sum_{x \in V} N_\varphi^{t,x} - \Delta n_V^t(y),$$

where $n_V^t(x)$, an integer valued random variable, is the number of topplings at site x in the time interval $[0, t]$, when sand is added in V . For T_n defined in (20) we will first prove that $n_{T_n}^t$ increases $\mu \times \mathbb{P}$ almost surely to an $L^1(\mu \times \mathbb{P})$ random variable n^t , interpreted as the number of topplings in $[0, t]$ when we add grains according to $N_\varphi^{t,x}$. By the Abelian property the sequence $n_{T_k}^t(0)$ is increasing in k . The following estimate is similar to (37):

$$\begin{aligned} & (\mu \times \mathbb{P})(|n_{T_k}^t(0) - n_{T_{k+1}}^t(0)| \geq \varepsilon) \\ &= (\mu \times \mathbb{P})(n_{x_{k+1}}^t(0) \geq \varepsilon) \\ &\leq \frac{1}{\varepsilon} \int n_{x_{k+1}}^t(0) \mu(d\eta) \times \mathbb{P}(d\omega) \\ (49) \quad &\leq \frac{1}{\varepsilon} \lim_{V \uparrow S} \int n_{x_{k+1}}^t(0) I(V_{x_{k+1}}(\eta) \cup V_0(\eta) \subset V) \mu(d\eta) \times \mathbb{P}(d\omega) \\ &\leq \frac{1}{\varepsilon} \lim_{W \uparrow S} \int n_W^t(x_{k+1}, 0, \eta) \mu_W(d\eta) \times \mathbb{P}(d\omega) \\ &\leq \frac{1}{\varepsilon} t\varphi(x_{k+1})G(0, x_{k+1}). \end{aligned}$$

In the fourth line, $n_W^t(x_{k+1}, 0, \eta)$ denotes the number of topplings up to time t at site $0 \in W$ by adding grains at site $x_{k+1} \in W$. By the Borel–Cantelli

lemma, condition (23) implies the a.s. convergence of $n_{T_k}^t(0)$, and analogously of every $n_{T_k}^t(x)$. Pick (η, ω) such that $n_{T_k}^t(\eta, \omega)$ converges, that is, such that $\sup_k n_{T_k}^t(\eta, \omega) = n^t(\eta, \omega)$ is finite [indeed, $n_{T_k}^t(\eta, \omega)$ is an integer]. If $\eta' \leq \eta$, then $n_{T_k}^t(\eta', \omega) \leq n_{T_k}^t(\eta, \omega)$ because we can obtain η from η' by adding sand at sites $x \in S$ for which $\eta'(x) < \eta(x)$ thereby increasing the number of topplings. We thus conclude that the convergence of $n_{T_k}^t(\eta, \omega)$ implies the convergence of $n_{T_k}^t(\eta', \omega)$ for all $\eta' \leq \eta$.

Now let $\mu' \leq \mu$ in the FKG sense. There is a coupling \mathbb{P}_{12} of $\mu' \times \mathbb{P}$ and $\mu \times \mathbb{P}$ such that

$$\mathbb{P}_{12}((\eta_1, \omega_1), (\eta_2, \omega_2)) : \omega_1 = \omega_2, \eta_1 \leq \eta_2 = 1;$$

that is, we use the same Poisson events and couple μ' and μ according to the optimal coupling [see Strassen (1965)]. Then

$$\begin{aligned} & (\mu' \times \mathbb{P})(n_{T_k}^t(\eta, \omega) \rightarrow n^t(\eta, \omega)) \\ &= \mathbb{P}_{12}(n_{T_k}^t(\eta_1, \omega_1) \rightarrow n^t(\eta_1, \omega_1)) \\ &= \mathbb{P}_{12}(n_{T_k}^t(\eta_1, \omega_1) \rightarrow n^t(\eta_1, \omega_1), \\ &\quad n_{T_k}^t(\eta_2, \omega_2) \rightarrow n^t(\eta_2, \omega_2), \omega_1 = \omega_2, \eta_1 \leq \eta_2) \\ &\geq \mathbb{P}_{12}(n_{T_k}^t(\eta_2, \omega_2) \rightarrow n^t(\eta_2, \omega_2)) \\ &= (\mu \times \mathbb{P})(n_{T_k}^t(\eta, \omega) \rightarrow n^t(\eta, \omega)) = 1. \end{aligned}$$

This shows the $\mu' \times \mathbb{P}$ -almost sure convergence of $n_{T_k}^t$; hence by (48) the product $\prod_{x \in S} a_x^{N_{\varphi}^{t,x}(\omega)} \eta'$ converges $\mu' \times \mathbb{P}$ almost surely. \square

As a further result we show that the semigroup $S_{\varphi}(t)$ is continuous as a function of the addition rate φ . We define

$$\ell_1 = \left\{ \varphi : S \rightarrow [0, \infty) : \|\varphi\| = \sum_{x \in S} \varphi(x) G(0, x) < \infty \right\}.$$

It is a complete metric space (as a closed subset of a Banach space) with the following property: If $\varphi_n \in \ell_1$, $\varphi_n \uparrow \varphi$ (pointwise) and $\varphi \in \ell_1$, then $\varphi_n \rightarrow \varphi$ in ℓ_1 .

PROPOSITION 4.1. *The semigroup $S_{\varphi}(t)$ of Theorem 3.3 is a strongly continuous function of φ ; that is, if $\varphi_n \rightarrow \varphi$ in ℓ_1 , then, for any local function f , $S_{\varphi_n}(t)f \rightarrow S_{\varphi}(t)f$.*

PROOF. Let $n_\varphi^t = \lim_{k \rightarrow \infty} n_{T_k}^t$ be the number of topplings in $[0, t]$ from sand addition at rate φ . In the proof of Theorem 3.5 we have shown that this random variable is $\mu \times \mathbb{P}$ almost surely well defined and, after taking limits in (48), satisfies

$$(50) \quad \eta_t = \eta_0 + N_\varphi^t - \Delta n_\varphi^t,$$

where $N_\varphi^t = \lim_{V \uparrow S} \sum_{x \in V} N_\varphi^{t,x}$. Note that if $\psi_1 \leq \psi_2$, the Poisson processes $N_{\psi_1}^t$ and $N_{\psi_2}^t$ can be coupled in such a way that, for all $x \in S$, $N_{\psi_1}^{t,x} \leq N_{\psi_2}^{t,x}$, and hence, by the Abelian property, $n_{\psi_1}^t(x) \leq n_{\psi_2}^t(x)$. Consider a coupling of the four Poisson processes $N_\varphi^t, N_{\varphi \wedge \varphi_n}^t, N_{\varphi \vee \varphi_n}^t$ and $N_{\varphi_n}^t$ under which the inequalities $X_1(t) \geq X_2(t), X_2(t) \leq X_3(t), X_3(t) \geq X_4(t)$ are satisfied with probability 1. Let $\tilde{\mathbb{P}}$ denote the law of the marginal (X_1, X_4) . We have, by a reasoning similar to (49),

$$(51) \quad \begin{aligned} & \int d\mu(\tilde{\mathbb{E}}|n_{\varphi_n}^t(0) - n_\varphi^t(0)|) \\ & \leq \int d\mu(\tilde{\mathbb{E}}(n_{\varphi_n}^t(0) - n_{\varphi_n \wedge \varphi}^t(0)) + \tilde{\mathbb{E}}(n_{\varphi \vee \varphi_n}^t(0) - n_{\varphi \wedge \varphi_n}^t(0)) \\ & \quad + \tilde{\mathbb{E}}(n_{\varphi \vee \varphi_n}^t(0) - n_\varphi^t(0))) \\ & \leq t \sum_{x \in S} |\varphi_n(x) - \varphi(x)| G(0, x), \end{aligned}$$

which tends to zero for $\varphi_n \rightarrow \varphi$ in ℓ_1 . Now take a local function f , and write $\tilde{D}_f = D_f \cup \partial_e D_f$,

$$\begin{aligned} |S_{\varphi_n}(t)(f) - S_\varphi(t)(f)| & \leq \tilde{\mathbb{P}}(n_{\varphi_n}^t(x) \neq n_\varphi^t(x) \text{ for some } x \in \tilde{D}_f) \\ & \leq \sum_{x \in \tilde{D}_f} \tilde{\mathbb{E}}|n_\varphi^t(x) - n_{\varphi_n}^t(x)|. \end{aligned}$$

Combining this with (51) concludes the proof. \square

One might ask whether we can go beyond condition (23), which essentially guarantees that the expected number of topplings stays finite in the addition process. In the following proposition we show that it is impossible to keep integrable toppling numbers and “rate 1” addition. The relation (52) should be regarded as the infinitesimal version of (50), where $\alpha(x)$ replaces the rate $\varphi(x)$. We then show that φ has to depend on x .

PROPOSITION 4.2. *Let $\alpha : S \rightarrow \{0, 1\}$ be a stationary and ergodic random field distributed according to ν . Denote its density by $\int \alpha(0) \nu(d\eta) = \rho$. Suppose there exists a measurable transformation $T : \{0, 1\}^S \times \Omega \rightarrow \Omega$ which satisfies the*

following conditions:

(i) The measure μ of Theorem 3.2 is invariant under $T(\alpha, \cdot)$ for any α .

(ii)

$$(52) \quad T(\alpha, \eta)(x) = \eta(x) + \alpha(x) - \Delta n(\alpha, \eta, x),$$

with $n(\alpha, \eta, \cdot) \in L^1(\mu)$ for ν almost every α .

Then, $\rho = 0$.

PROOF. Taking expectation over μ in (52) gives

$$(53) \quad \Delta \Psi(\alpha, x) = \alpha(x),$$

where $\Psi(\alpha, x) = \int n(\alpha, \eta, x) \mu(d\eta)$. By stationarity of μ and ν , $\Psi(\alpha, x)$ is a stationary random field. Let $(x_t : t \geq 0)$ denote a continuous time simple random walk on S , starting at 0. From (53),

$$\mathbb{E} \Psi(\alpha, x_t) = \Psi(\alpha, 0) + \mathbb{E} \int_0^t \alpha(x_s) ds.$$

Divide this last line by t and let $t \uparrow +\infty$. As ν is ergodic (making the last term equal to ρ) and as the process $\Psi(\alpha, x_t)$ is stationary, we conclude that $\rho = 0$. \square

REFERENCES

- BAK, P., TANG, K. and WIESENFELD, K. (1988). Self-organized criticality. *Phys. Rev. A* **38** 364–374.
- CHEN, M. F. (1992). *From Markov Chains to Non-Equilibrium Particle Systems*. World Scientific, Singapore.
- CHUNG, K. L. (1960). *Markov Chains with Stationary Transition Probabilities*. Springer, Berlin.
- DHAR, D. (1990). Self organised critical state of sandpile automaton models. *Phys. Rev. Lett.* **64** 1613–1616.
- DHAR, D. (1999). The Abelian sandpiles and related models. *Phys. A* **263** 4–25.
- DHAR, D. and MAJUMDAR, S. N. (1990). Abelian sandpile models on the Bethe lattice *J. Phys. A* **23** 4333–4350.
- IVASHKEVICH, E. V. and PRIEZZHEV, V. B. (1998). Introduction to the sandpile model. *Phys. A* **254** 97–116.
- LIGGETT, T. M. (1980). The long range exclusion process. *Ann. Probab.* **8** 861–889.
- LIGGETT, T. M. (1985). *Interacting Particle Systems*. Springer, Berlin.
- MAES, C., REDIG, F., SAADA E. and VAN MOFFAERT, A. (2000). On the thermodynamic limit for a one-dimensional sandpile process. *Markov Process. Related Fields* **6** 1–22.
- MEESTER, R., REDIG, F. and ZNAMENSKI, D. (2002). The Abelian sandpile; a mathematical introduction. *Markov Process. Related Fields* **7** 509–523.
- SPEER, E. (1993). Asymmetric Abelian sandpile models. *J. Statist. Phys.* **71** 61–74.
- STRASSEN, V. (1965). The existence of probability measures with given marginals. *Ann. Math. Statist.* **36** 423–439.

- TOOM, A. L., VASILYEV, N. B., STAVSKAYA, O. N., MITYUSHIN, L. G., KURDYUMOV, G. L. and PIROGOV, S. A. (1990). Discrete local Markov systems. In *Stochastic Cellular Systems: Ergodicity, Memory, Morphogenesis* (R. L. Dobrushin, V. I. Kryukov and A. L. Toom, eds.) 1–182. Manchester Univ. Press.
- TURCOTTE, D. L. (1999). Self-organized criticality. *Rep. Progr. Phys.* **62** 1377–1429.

C. MAES
INSTITUUT VOOR THEORETISCHE FYSICA
K.U. LEUVEN
CELESTIJNENLAAN 200D
B-3001 LEUVEN
BELGIUM
E-MAIL: christian.maes@fys.kuleuven.ac.be

F. REDIG
TECHNISCHE UNIVERSITEIT EINDHOVEN
P.O. BOX 513
5600 MB EINDHOVEN
THE NETHERLANDS
E-MAIL: f.h.j.redig@tue.nl

E. SAADA
CNRS, UMR 6085
UNIVERSITÉ DE ROUEN
76821 MONT-SAINT-AIGNAN CEDEX
FRANCE
E-MAIL: Ellen.Saada@univ-rouen.fr