

Intermittency on catalysts : symmetric exclusion

Citation for published version (APA):

Gärtner, J., Hollander, den, W. T. F., & Maillard, G. (2006). *Intermittency on catalysts : symmetric exclusion*. (Report Eurandom; Vol. 2006015). Eurandom.

Document status and date:

Published: 01/01/2006

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Intermittency on catalysts: symmetric exclusion

J. Gärtner ¹
F. den Hollander ^{2,3}
G. Maillard ⁴

24th May 2006

Abstract

We continue our study of intermittency for the parabolic Anderson equation $\partial u/\partial t = \kappa \Delta u + \xi u$, where $u: \mathbb{Z}^d \times [0, \infty) \rightarrow \mathbb{R}$, κ is the diffusion constant, Δ is the discrete Laplacian, and $\xi: \mathbb{Z}^d \times [0, \infty) \rightarrow \mathbb{R}$ is a space-time random medium. The solution of the equation describes the evolution of a “reactant” u under the influence of a “catalyst” ξ .

In this paper we focus on the case where ξ is exclusion with a symmetric random walk transition kernel, starting from equilibrium with density $\rho \in (0, 1)$. We consider the annealed Lyapunov exponents, i.e., the exponential growth rates of the successive moments of u . We show that these exponents are trivial when the random walk is recurrent, but display an interesting dependence on the diffusion constant κ when the random walk is transient, with qualitatively different behavior in different dimensions. Special attention is given to the asymptotics of the exponents for $\kappa \rightarrow \infty$, which is controlled by moderate deviations of ξ requiring a delicate expansion argument.

In Gärtner and den Hollander [4] the case where ξ is a Poisson field of independent (simple) random walks was studied. The two cases show interesting differences and similarities. Throughout the paper, a comparison of the two cases plays a crucial role.

MSC 2000. Primary 60H25, 82C44; Secondary 60F10, 35B40.

Key words and phrases. Parabolic Anderson model, catalytic random medium, exclusion process, Lyapunov exponents, intermittency, large deviations, graphical representation.

Acknowledgment. GM was supported by a postdoctoral fellowship from the Netherlands Organization for Scientific Research (grant 613.000.307) during his stay at EURANDOM. The research in this paper was partially supported by the ESF Scientific Programme “Random Dynamics in Spatially Extended Systems”. FdH and GM are grateful to the Pacific Institute for the Mathematical Sciences and the Mathematics Department of the University of British Columbia, Vancouver, Canada, for hospitality: FdH January to August 2006, GM mid-January to mid-February 2006 when the work on this paper was completed.

¹Institut für Mathematik, Technische Universität Berlin, Strasse des 17. Juni 136, D-10623 Berlin, Germany, jg@math.tu-berlin.de

²Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands, den-holla@math.leidenuniv.nl

³EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

⁴Institut de Mathématiques, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland, gregory.maillard@epfl.ch

1 Introduction and main results

1.1 Model

The parabolic Anderson equation is the partial differential equation

$$\frac{\partial}{\partial t}u(x, t) = \kappa\Delta u(x, t) + \xi(x, t)u(x, t), \quad x \in \mathbb{Z}^d, t \geq 0. \quad (1.1.1)$$

Here, the u -field is \mathbb{R} -valued, $\kappa \in [0, \infty)$ is the diffusion constant, Δ is the discrete Laplacian, acting on u as

$$\Delta u(x, t) = \sum_{\substack{y \in \mathbb{Z}^d \\ \|y-x\|=1}} [u(y, t) - u(x, t)] \quad (1.1.2)$$

($\|\cdot\|$ is the Euclidian norm), while

$$\xi = \{\xi(x, t): x \in \mathbb{Z}^d, t \geq 0\} \quad (1.1.3)$$

is an \mathbb{R} -valued random field that evolves with time and that drives the equation. As initial condition for (1.1.1) we take

$$u(\cdot, 0) \equiv 1. \quad (1.1.4)$$

In the present paper we focus on the case where ξ is *Symmetric Exclusion* (SE), i.e., ξ takes values in $\{0, 1\}^{\mathbb{Z}^d} \times [0, \infty)$, where $\xi(x, t) = 1$ means that there is a particle at x at time t and $\xi(x, t) = 0$ means that there is none, and particles move around according to a symmetric random walk transition kernel. We choose $\xi(\cdot, 0)$ according to the Bernoulli product measure with density $\rho \in (0, 1)$, i.e., initially each site has a particle with probability ρ and no particle with probability $1 - \rho$, independently for different sites. For this choice, the ξ -field is stationary in time.

One interpretation of (1.1.1) and (1.1.4) comes from population dynamics. Consider a spatially homogeneous system of two types of particles, A (catalyst) and B (reactant), subject to:

- (i) A -particles behave autonomously, according to a prescribed stationary dynamics, with density ρ ;
- (ii) B -particles perform independent random walks with diffusion constant κ and split into two at a rate that is equal to the number of A -particles present at the same location;
- (iii) the initial density of B -particles is 1.

Then

$$u(x, t) = \{ \text{the average number of } B\text{-particles at site } x \text{ at time } t \\ \text{conditioned on the evolution of the } A\text{-particles} \}. \quad (1.1.5)$$

It is possible to add that B -particles die at rate $\delta \in (0, \infty)$. This amounts to the trivial transformation $u(x, t) \rightarrow u(x, t)e^{-\delta t}$.

In Kesten and Sidoravicius [9] and in Gärtner and den Hollander [4], the case was considered where ξ is given by a Poisson field of independent simple random walks. The survival

versus extinction pattern (in [9] for $\delta > 0$) and the annealed Lyapunov exponents (in [4] for $\delta = 0$) were studied, in particular, their dependence on d , κ and the parameters controlling ξ .

Equation (1.1.1) is a discrete heat equation with the ξ -field playing the role of a source. What makes (1.1.1) particularly interesting is that the two terms in the right-hand side *compete with each other*: the diffusion induced by Δ tends to make u flat, while the branching induced by ξ tends to make u irregular. Henceforth we call ξ the “catalyst” and u the “reactant”.

1.2 SE, Lyapunov exponents and comparison with IRW

Throughout the paper, we abbreviate $\Omega = \{0, 1\}^{\mathbb{Z}^d}$ (endowed with the product topology), and we let $p: \mathbb{Z}^d \times \mathbb{Z}^d \rightarrow [0, 1]$ be the transition kernel of an irreducible random walk,

$$\begin{aligned} p(x, y) = p(0, y - x) \geq 0 \quad \forall x, y \in \mathbb{Z}^d, \quad \sum_{y \in \mathbb{Z}^d} p(x, y) = 1 \quad \forall x \in \mathbb{Z}^d, \\ p(x, x) = 0 \quad \forall x \in \mathbb{Z}^d, \quad p(\cdot, \cdot) \text{ generates } \mathbb{Z}^d, \end{aligned} \tag{1.2.1}$$

that is assumed to be *symmetric*,

$$p(x, y) = p(y, x) \quad \forall x, y \in \mathbb{Z}^d. \tag{1.2.2}$$

A special case is *simple* random walk

$$p(x, y) = \begin{cases} \frac{1}{2d} & \text{if } \|x - y\| = 1, \\ 0 & \text{otherwise.} \end{cases} \tag{1.2.3}$$

The exclusion process is the Markov process on Ω whose generator L acts on cylindrical functions f as (see Liggett [12], Chapter VIII)

$$(Lf)(\eta) = \sum_{x, y \in \mathbb{Z}^d} p(x, y) \eta(x) [1 - \eta(y)] [f(\eta^{x,y}) - f(\eta)] = \sum_{\{x, y\} \subset \mathbb{Z}^d} p(x, y) [f(\eta^{x,y}) - f(\eta)], \tag{1.2.4}$$

where the latter sum runs over unoriented bonds $\{x, y\}$ between any pair of sites $x, y \in \mathbb{Z}^d$, and

$$\eta^{x,y}(z) = \begin{cases} \eta(z) & \text{if } z \neq x, y, \\ \eta(y) & \text{if } z = x, \\ \eta(x) & \text{if } z = y. \end{cases} \tag{1.2.5}$$

The first line of (1.2.4) says that a particle at site x jumps to a vacancy at site y at rate $p(x, y)$, the second line says that the states of x and y are interchanged along the bond $\{x, y\}$ at rate $p(x, y)$. For $\rho \in [0, 1]$, let ν_ρ be the Bernoulli product measure on Ω with density ρ . This is an invariant measure for SE. Under (1.2.1–1.2.2), $(\nu_\rho)_{\rho \in [0, 1]}$ are the only extremal equilibria (see Liggett [12], Chapter VIII, Theorem 1.44). We denote by \mathbb{P}_η the law of ξ starting from $\eta \in \Omega$ and write $\mathbb{P}_{\nu_\rho} = \int_\Omega \nu_\rho(d\eta) \mathbb{P}_\eta$.

In the *graphical representation* of SE, space is drawn sideways, time is drawn upwards, and for each pair of sites $x, y \in \mathbb{Z}^d$ links are drawn between x and y at Poisson rate $p(x, y)$. The configuration at time t is obtained from the one at time 0 by transporting the local states along paths that move upwards with time and sideways along links (see Fig. 1).

We will frequently use the following property, which is immediate from the graphical representation:

$$\mathbb{E}_\eta (\xi(y, t)) = \sum_{x \in \mathbb{Z}^d} \eta(x) p_t(x, y), \quad \eta \in \Omega, y \in \mathbb{Z}^d, t \geq 0. \quad (1.2.6)$$

Similar expressions hold for higher order correlations. Here, $p_t(x, y)$ is the probability that the random walk with transition kernel $p(\cdot, \cdot)$ and step rate 1 moves from x to y in time t . The graphical representation shows that the evolution is invariant under time reversal and, in particular, the equilibria $(\nu_\rho)_{\rho \in [0,1]}$ are *reversible*. This fact will turn out to be very important later on.




Fig. 1: Graphical representation. The dashed lines are links. The arrows represent a path from $(x, 0)$ to (y, t) .

By the Feynman-Kac formula, the solution of (1.1.1) and (1.1.4) reads

$$u(x, t) = \mathbb{E}_x \left(\exp \left[\int_0^t ds \xi(X^\kappa(s), t-s) \right] \right), \quad (1.2.7)$$

where X^κ is simple random walk on \mathbb{Z}^d with step rate $2d\kappa$ and \mathbb{E}_x denotes expectation with respect to X^κ given $X^\kappa(0) = x$. We will often write $\xi_t(x)$ and X_t^κ instead of $\xi(x, t)$ and $X^\kappa(t)$, respectively.

For $p \in \mathbb{N}$ and $t > 0$, define

$$\Lambda_p(t) = \frac{1}{pt} \log \mathbb{E}_{\nu_\rho} (u(0, t)^p). \quad (1.2.8)$$

Then

$$\Lambda_p(t) = \frac{1}{pt} \log \mathbb{E}_{\nu_\rho} \left(\mathbb{E}_{0, \dots, 0} \left(\exp \left[\int_0^t ds \sum_{q=1}^p \xi(X_q^\kappa(s), s) \right] \right) \right), \quad (1.2.9)$$

where X_q^κ , $q = 1, \dots, p$, are p independent copies of X^κ , $\mathbb{E}_{0, \dots, 0}$ denotes expectation w.r.t. X_q^κ , $q = 1, \dots, p$, given $X_1^\kappa(0) = \dots = X_p^\kappa(0) = 0$, and the time argument $t-s$ in (1.2.7) is replaced by s in (1.2.9) via the reversibility of ξ starting from ν_ρ . If the last quantity admits a limit as $t \rightarrow \infty$, then we define

$$\lambda_p = \lim_{t \rightarrow \infty} \Lambda_p(t) \quad (1.2.10)$$

to be the p -th *annealed Lyapunov exponent*.

From Hölder's inequality applied to (1.2.8) it follows that $\Lambda_p(t) \geq \Lambda_{p-1}(t)$ for all $t > 0$ and $p \in \mathbb{N} \setminus \{1\}$. Hence $\lambda_p \geq \lambda_{p-1}$ for all $p \in \mathbb{N} \setminus \{1\}$. We say that the system is p -*intermittent*

if $\lambda_p > \lambda_{p-1}$. In the latter case the system is q -intermittent for all $q > p$ as well (cf. Gärtner and Molchanov [6], Section 1.1). We say that the system is *intermittent* if it is p -intermittent for all $p \in \mathbb{N} \setminus \{1\}$. Intermittent means that the u -field develops sparse high peaks dominating the moments in such a way that each moment is dominated by its own collection of peaks (see Gärtner and König [5], Section 1.3, and den Hollander [4], Section 1.2).

Let $(\tilde{\xi}_t)_{t \geq 0}$ be the process of *Independent Random Walks* (IRW) with step rate 1, transition kernel $p(\cdot, \cdot)$ and state space Ω . Let $\mathbb{E}_\eta^{\text{IRW}}$ denote expectation w.r.t. $(\tilde{\xi}_t)_{t \geq 0}$ starting from $\tilde{\xi}_0 = \eta$, and write $\mathbb{E}_{\nu_\rho}^{\text{IRW}} = \int_\Omega \nu_\rho(d\eta) \mathbb{E}_\eta^{\text{IRW}}$. Throughout the paper we will make use of the following inequality comparing SE and IRW. The proof of this inequality is given in Appendix A and uses a lemma due to Landim [11].

Proposition 1.2.1 *For any $K: \mathbb{Z}^d \times [0, \infty) \rightarrow \mathbb{R}$ such that either $K \geq 0$ or $K \leq 0$, any $t \geq 0$ such that $\sum_{z \in \mathbb{Z}^d} \int_0^t ds |K(z, s)| < \infty$ and any $\eta \in \Omega$,*

$$\mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \xi_s(z) \right] \right) \leq \mathbb{E}_\eta^{\text{IRW}} \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \tilde{\xi}_s(z) \right] \right). \quad (1.2.11)$$

This powerful inequality will allow us to obtain bounds that are more easily computable.

1.3 Main theorems

Our first result states that the Lyapunov exponents exist and behave nicely as a function of κ . We write $\lambda_p(\kappa)$ to exhibit the dependence on κ , suppressing d and ρ .

Theorem 1.3.1 *Let $d \geq 1$, $\rho \in (0, 1)$ and $p \in \mathbb{N}$.*

- (i) *For all $\kappa \in [0, \infty)$, the limit in (1.2.10) exists and is finite.*
- (ii) *On $[0, \infty)$, $\kappa \rightarrow \lambda_p(\kappa)$ is continuous, non-increasing and convex.*

Our second result states that the Lyapunov exponents are trivial for recurrent random walk but are non-trivial for transient random walk (see Fig. 2).

Theorem 1.3.2 *Let $d \geq 1$, $\rho \in (0, 1)$ and $p \in \mathbb{N}$.*

- (i) *If $p(\cdot, \cdot)$ is recurrent, then $\lambda_p(\kappa) = 1$ for all $\kappa \in [0, \infty)$.*
- (ii) *If $p(\cdot, \cdot)$ is transient, then $\rho < \lambda_p(\kappa) < 1$ for all $\kappa \in [0, \infty)$. Moreover, $\kappa \mapsto \lambda_p(\kappa)$ is strictly decreasing with $\lim_{\kappa \rightarrow \infty} \lambda_p(\kappa) = \rho$.*




Fig. 2: Qualitative picture of $\kappa \mapsto \lambda_p(\kappa)$ for recurrent, respectively, transient random walk.

Our third result shows that for transient random walk the system is intermittent at $\kappa = 0$.

Theorem 1.3.3 *Let $d \geq 1$ and $\rho \in (0, 1)$. If $p(\cdot, \cdot)$ is transient, then $p \mapsto \lambda_p(0)$ is strictly increasing.*

Our fourth and final result identifies the behavior of the Lyapunov exponents for large κ when $d \geq 4$ and $p(\cdot, \cdot)$ is simple random walk (see Fig. 3).

Theorem 1.3.4 *Assume (1.2.3). Let $d \geq 4$, $\rho \in (0, 1)$ and $p \in \mathbb{N}$. Then*

$$\lim_{\kappa \rightarrow \infty} 2d\kappa[\lambda_p(\kappa) - \rho] = \rho(1 - \rho)G_d \quad (1.3.1)$$

with G_d the Green function at the origin of simple random walk on \mathbb{Z}^d .




Fig. 3: Qualitative picture of $\kappa \mapsto \lambda_p(\kappa)$ for $p = 1, 2, 3$ for simple random walk in $d \geq 4$. The dotted line moving down represents the asymptotics given by the r.h.s. of (1.3.1).

1.4 Discussion

Theorem 1.3.1 gives general properties that need no further comment. We will see that they in fact hold for *any stationary, reversible and bounded ξ* .

The intuition behind Theorem 1.3.2 is the following. If the catalyst is driven by a recurrent random walk, then it suffers from “traffic jams”, i.e., with not too small a probability there is a large region around the origin that the catalyst fully occupies for a long time. Since with not too small a probability the simple random walk (driving the reactant) can stay inside this large region for the same amount of time, the average growth rate of the reactant at the origin is maximal. This phenomenon may be expressed by saying that *for recurrent random walk clumping of the catalyst dominates the growth of the moments*. For transient random walk, on the other hand, clumping of the catalyst is present (the growth rate of the reactant is $> \rho$), but it is not dominant (the growth rate of the reactant is < 1). As the diffusion constant κ of the reactant increases, the effect of the clumping of the catalyst gradually diminishes and the growth rate of the reactant gradually decreases to the density of the catalyst.

Theorem 1.3.3 shows that if the reactant stands still and the catalyst is driven by a transient random walk, then the system is intermittent. Apparently, the successive moments of the reactant, which are equal to the exponential moments of the occupation time of the origin by the catalyst (take (1.2.7) with $\kappa = 0$), are sensitive to successive degrees of clumping. By continuity, *intermittency persists for small κ* .

Theorem 1.3.4 shows that, when the catalyst is driven by simple random walk, *all* Lyapunov exponents decay to ρ as $\kappa \rightarrow \infty$ *in the same manner* when $d \geq 4$. The case $d = 3$ remains open. We conjecture:

Conjecture 1.4.1 *Assume (1.2.3). Let $d = 3$, $\rho \in (0, 1)$ and $p \in \mathbb{N}$. Then*

$$\lim_{\kappa \rightarrow \infty} 2d\kappa[\lambda_p(\kappa) - \rho] = \rho(1 - \rho)G_d + [2d\rho(1 - \rho)]^2\mathcal{P} \quad (1.4.1)$$

with

$$\mathcal{P} = \sup_{\substack{f \in H^1(\mathbb{R}^3) \\ \|f\|_2=1}} \left[\left\| (-\Delta_{\mathbb{R}^3})^{-1/2} f^2 \right\|_2^2 - \|\nabla_{\mathbb{R}^3} f\|_2^2 \right] \in (0, \infty), \quad (1.4.2)$$

where $\nabla_{\mathbb{R}^3}$ and $\Delta_{\mathbb{R}^3}$ are the continuous gradient and Laplacian, $\|\cdot\|_2$ is the $L^2(\mathbb{R}^3)$ -norm, $H^1(\mathbb{R}^3) = \{f: \mathbb{R}^3 \rightarrow \mathbb{R}: f, \nabla_{\mathbb{R}^3} f \in L^2(\mathbb{R}^3)\}$, and

$$\left\| (-\Delta_{\mathbb{R}^3})^{-1/2} f^2 \right\|_2^2 = \int_{\mathbb{R}^3} dx f^2(x) \int_{\mathbb{R}^3} dy f^2(y) \frac{1}{4\pi\|x - y\|}. \quad (1.4.3)$$

In section 1.5 we will explain how this conjecture arises in analogy with the case of IRW studied in Gärtner and den Hollander [4]. If Conjecture 1.4.1 holds true, then in $d = 3$ *intermittency persists for large κ* . It would still remain open whether the same is true for $d \geq 4$. To decide the latter, we need a finer asymptotics for $d \geq 4$. A large diffusion constant of the reactant prevents the solution u to easily localize around the regions where the catalyst clumps, but it is not clear whether this is able to destroy intermittency for $d \geq 4$.

We further conjecture:

Conjecture 1.4.2 *In $d = 3$, the system is intermittent for all $\kappa \in [0, \infty)$.*

Conjecture 1.4.3 *In $d \geq 4$, there exists a strictly increasing sequence $0 < \kappa_2 < \kappa_3 < \dots$ such that for $p = 2, 3, \dots$ the system is p -intermittent if and only if $0 \leq \kappa < \kappa_p$.*

In words, we conjecture that in $d = 3$ the curves in Fig. 3 never merge, whereas for $d \geq 4$ the curves merge successively.

1.5 Heuristics behind Theorem 1.3.4 and Conjecture 1.4.1

The *heuristics* behind Theorem 1.3.4 and Conjecture 1.4.1 is the following. Consider the case $p = 1$. Scaling time by κ in (1.2.9), we have $\lambda_1(\kappa) = \kappa\lambda_1^*(\kappa)$ with

$$\lambda_1^*(\kappa) = \lim_{t \rightarrow \infty} \Lambda_1^*(\kappa; t) \quad \text{and} \quad \Lambda_1^*(\kappa; t) = \frac{1}{t} \log \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \xi \left(X(s), \frac{s}{\kappa} \right) \right] \right), \quad (1.5.1)$$

where $X = X^1$ and we abbreviate

$$\mathbb{E}_{\nu_\rho, 0} = \mathbb{E}_{\nu_\rho} \mathbb{E}_0. \quad (1.5.2)$$

For large κ , the ξ -field in (1.5.1) evolves slowly and therefore does not manage to cooperate with the X -process in determining the growth rate. Also, the prefactor $1/\kappa$ in the exponent is small. As a result, the expectation over the ξ -field can be computed via a *Gaussian*

approximation that becomes sharp in the limit as $\kappa \rightarrow \infty$, i.e.,

$$\begin{aligned} \Lambda_1^*(\kappa; t) - \frac{\rho}{\kappa} &= \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \left[\xi \left(X(s), \frac{s}{\kappa} \right) - \rho \right] \right] \right) \\ &\approx \frac{1}{t} \log \mathbb{E}_0 \left(\exp \left[\frac{1}{2\kappa^2} \int_0^t ds \int_0^t du \mathbb{E}_{\nu_{\rho}} \left(\left[\xi \left(X(s), \frac{s}{\kappa} \right) - \rho \right] \left[\xi \left(X(u), \frac{u}{\kappa} \right) - \rho \right] \right) \right] \right). \end{aligned} \quad (1.5.3)$$

(In essence, what happens here is that the asymptotics for $\kappa \rightarrow \infty$ is driven by moderate deviations of the ξ -field, which fall in the Gaussian regime.) The exponent in the r.h.s. of (1.5.3) equals

$$\frac{1}{\kappa^2} \int_0^t ds \int_s^t du \mathbb{E}_{\nu_{\rho}} \left(\left[\xi \left(X(s), \frac{s}{\kappa} \right) - \rho \right] \left[\xi \left(X(u), \frac{u}{\kappa} \right) - \rho \right] \right). \quad (1.5.4)$$

Now, for $x, y \in \mathbb{Z}^d$ and $b \geq a \geq 0$ we have

$$\begin{aligned} \mathbb{E}_{\nu_{\rho}} \left(\left[\xi(x, a) - \rho \right] \left[\xi(y, b) - \rho \right] \right) &= \mathbb{E}_{\nu_{\rho}} \left(\left[\xi(x, 0) - \rho \right] \left[\xi(y, b-a) - \rho \right] \right) \\ &= \int_{\Omega} \nu_{\rho}(d\eta) \left[\eta(x) - \rho \right] \mathbb{E}_{\eta} \left(\left[\xi(y, b-a) - \rho \right] \right) \\ &= \sum_{z \in \mathbb{Z}^d} p_{b-a}(z, y) \int_{\Omega} \nu_{\rho}(d\eta) \left[\eta(x) - \rho \right] \left[\eta(z) - \rho \right] \\ &= \rho(1 - \rho) p_{b-a}(x, y), \end{aligned} \quad (1.5.5)$$

where the first equality uses the stationarity of ξ , the third equality uses (1.2.6) from the graphical representation, and the fourth equality uses that ν_{ρ} is Bernoulli. Substituting (1.5.5) into (1.5.4), we get that the r.h.s. of (1.5.3) equals

$$\frac{1}{t} \log \mathbb{E}_0 \left(\exp \left[\frac{\rho(1 - \rho)}{\kappa^2} \int_0^t ds \int_s^t du p_{\frac{u-s}{\kappa}}(X(s), X(u)) \right] \right). \quad (1.5.6)$$

This is precisely the integral that was investigated in Gärtner and den Hollander [4] (see Sections 5–8 and equations (1.5.4–1.5.11) of that paper). Therefore the limit

$$\lim_{\kappa \rightarrow \infty} \kappa [\lambda_1(\kappa) - \rho] = \lim_{\kappa \rightarrow \infty} \kappa^2 \lim_{t \rightarrow \infty} \left[\Lambda_1^*(\kappa; t) - \frac{\rho}{\kappa} \right] = \lim_{\kappa \rightarrow \infty} \kappa^2 \lim_{t \rightarrow \infty} (1.5.6) \quad (1.5.7)$$

can be read off from [4] and yields (1.3.1) for $d \geq 4$ and (1.4.1) for $d = 3$. A similar heuristics applies for $p > 1$.

The r.h.s. of (1.3.1), which is valid for $d \geq 4$, is obtained from the above computations by moving the expectation in (1.5.6) into the exponent. Indeed,

$$\mathbb{E}_0 \left(p_{\frac{u-s}{\kappa}}(X(s), X(u)) \right) = \sum_{x, y \in \mathbb{Z}^d} p_{2ds}(0, x) p_{2d(u-s)}(x, y) p_{\frac{u-s}{\kappa}}(x, y) = p_{2d(u-s)(1 + \frac{1}{2d\kappa})}(0, 0) \quad (1.5.8)$$

and hence

$$\int_0^t ds \int_s^t du \mathbb{E}_0 \left(p_{\frac{u-s}{\kappa}}(X(s), X(u)) \right) = \int_0^t ds \int_0^{t-s} dv p_{2dv(1 + \frac{1}{2d\kappa})}(0, 0) \sim t \frac{1}{2d(1 + \frac{1}{2d\kappa})} G_d. \quad (1.5.9)$$

Thus we see that the result in Theorem 1.3.4 comes from a second order asymptotics on ξ and a first order asymptotics on X . Despite this simple fact, it turns out to be hard to make the above heuristics rigorous. For $d = 3$, on the other hand, we expect the first order asymptotics on X to fail, leading to the more complicated behavior in (1.4.1) .

Remark 1: In (1.1.1), the ξ -field may be multiplied by a coupling constant $\gamma \in (0, \infty)$. This produces no change in Theorems 1.3.1, 1.3.2(i) and 1.3.3. In Theorem 1.3.2(ii), $(\rho, 1)$ becomes $(\gamma\rho, \gamma)$, while in the r.h.s. of Theorem 1.3.4 and Conjecture 1.4.1, $\rho(1 - \rho)$ gets multiplied by γ^2 . Similarly, if the simple random walk in Theorem 1.3.4 is replaced by a random walk with transition kernel $p(\cdot, \cdot)$ satisfying (1.2.1–1.2.2), then we expect that in (1.3.1) and (1.4.1) G_d becomes the Green function at the origin of this random walk and a factor $1/\sigma^4$ appears in front of the last term in the r.h.s. of (1.4.1) with σ^2 the variance of $p(\cdot, \cdot)$.

Remark 2: In Gärtner and den Hollander [4] the catalyst was γ times a Poisson field with density ρ of independent simple random walks stepping at rate $2d\theta$, where $\gamma, \rho, \theta \in (0, \infty)$ are parameters. It was found that the Lyapunov exponents are infinite in $d = 1, 2$ for all p and in $d \geq 3$ for $p \geq 2d\theta/\gamma G_d$, irrespective of κ and ρ . In $d \geq 3$ for $p < 2d\theta/\gamma G_d$, on the other hand, the Lyapunov exponents are finite for all κ , and exhibit a dichotomy similar to the one expressed by Theorem 1.3.4 and Conjecture 1.4.1. Apparently, in this regime the two types of catalyst are qualitatively similar. Remarkably, the *same* asymptotic behavior for large κ was found (with $\rho\gamma^2$ replacing $\rho(1 - \rho)$ in (1.3.1)), and the *same* variational formula as in (1.4.2) was seen to play a central role in $d = 3$. [Note: In [4] the symbols ν, ρ, G_d were used instead of $\rho, \theta, G_d/2d$.]

1.6 Outline

In Section 2 we derive a variational formula for λ_p from which Theorem 1.3.1 follows immediately. The arguments that will be used to derive this variational formula apply to an arbitrary bounded, stationary and *reversible* catalyst. Thus, the properties in Theorem 1.3.1 are quite general. In Section 3 we do a range of estimates, either directly on (1.2.9) or on the variational formula for λ_p derived in Section 2, to prove Theorems 1.3.2 and 1.3.3. Here, the special properties of SE, in particular, its space-time correlation structure expressed through the graphical representation (see Fig. 1), are crucial. These results hold for an arbitrary random walk subject to (1.2.1–1.2.2). Finally, in Section 4 we prove Theorem 1.3.4, which is restricted to simple random walk. The analysis consists of a long series of estimates, taking up half of the paper and, in essence, showing that the problem reduces to understanding the asymptotic behavior of (1.5.6). This reduction is important, because it explains why there is some degree of *universality* in the behavior for $\kappa \rightarrow \infty$ under different types of catalysts: apparently, the Gaussian approximation and the two-point correlation function in space and time determine the asymptotics (recall the heuristic argument in Section 1.5).

2 Lyapunov exponents: general properties

In this section we prove Theorem 1.3.1. In Section 2.1 we formulate a large deviation principle for the occupation time of the origin in SE due to Landim [11], which will be needed in Section 3.2. In Section 2.2 we extend the line of thought in [11] and derive a variational formula for λ_p from which Theorem 1.3.1 will follow immediately.

2.1 Large deviations for the occupation time of the origin

Kipnis [10], building on techniques developed by Arratia [1], proved that the *occupation time of the origin up to time t* ,

$$T_t = \int_0^t \xi(0, s) ds, \quad (2.1.1)$$

satisfies a strong law of large numbers and a central limit theorem. Landim [11] subsequently proved that T_t satisfies a *large deviation principle*, i.e.,

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_{\nu_\rho}(T_t/t \in F) &\leq - \inf_{\alpha \in F} \Psi_d(\alpha), & F \subseteq [0, 1] \text{ closed,} \\ \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_{\nu_\rho}(T_t/t \in G) &\geq - \inf_{\alpha \in G} \Psi_d(\alpha), & G \subseteq [0, 1] \text{ open,} \end{aligned} \quad (2.1.2)$$

with $\Psi_d: [0, 1] \rightarrow [0, \infty)$ a rate function that is given by the following formulas. Define the Dirichlet form associated with the generator L of SE given in (1.2.4),

$$\mathcal{E}(f) = (-Lf, f)_{L^2(\nu_\rho)} = \int_{\Omega} \nu_\rho(d\eta) \frac{1}{2} \sum_{\{x,y\} \subset \mathbb{Z}^d} p(x, y) [f(\eta^{x,y}) - f(\eta)]^2, \quad f \in L^2(\nu_\rho). \quad (2.1.3)$$

Next, define $I_d: \mathcal{M}_1(\Omega) \rightarrow [0, \infty]$ by

$$I_d(\mu) = \lim_{\varepsilon \downarrow 0} \inf_{\substack{\nu \in B(\mu, \varepsilon) \\ \nu \ll \nu_\rho}} \mathcal{E} \left(\sqrt{\frac{d\nu}{d\nu_\rho}} \right), \quad (2.1.4)$$

where $\mathcal{M}_1(\Omega)$ is the set of probability measures on Ω (endowed with the Prokhorov metric), and $B(\mu, \varepsilon)$ is the open ball of radius ε centered at μ . Then

$$\Psi_d(\alpha) = \inf_{\substack{\mu \in \mathcal{M}_1(\Omega) \\ \int_{\Omega} \eta(0) \mu(d\eta) = \alpha}} I_d(\mu). \quad (2.1.5)$$

The function I_d is convex and lower semi-continuous. Since Ψ_d is a minimum of I_d under a linear constraint, it inherits these properties (see e.g. den Hollander [7], Theorem III.32).

Remark: The function $\tilde{I}_d(\mu)$ defined by $\mathcal{E}(\sqrt{d\mu/d\nu_\rho})$ if $\mu \ll \nu_\rho$ and ∞ otherwise is not lower semi-continuous, and therefore does not qualify as a large deviation rate function. This is why the regularization in (2.1.4) is necessary (see also Deuschel and Stroock [3], Section 5.3).

Landim [11] showed that Ψ_d has a unique zero at ρ and satisfies the quadratic lower bound

$$\Psi_d(\alpha) \geq \frac{1}{2G_d} (\sqrt{\alpha} - \sqrt{\rho})^2 \quad (2.1.6)$$

with G_d the Green function at the origin of the random walk with transition kernel $p(\cdot, \cdot)$. This bound was obtained with the help of Proposition 1.2.1 with $K(z, s) = \delta_0(z)$, which implies that the occupation time for SE is stochastically smaller than the occupation time for IRW with the same density (see [11], Proposition 4.1). For the latter the rate function can be computed and equals the lower bound in (2.1.6) for $\alpha \geq \rho$. The same lower bound holds for $\alpha \leq \rho$, which can be seen by interchanging the role of the states 0 and 1.

Thus, as seen from (2.1.6), for transient random walk the rate function Ψ_d is non-trivial. For recurrent random walk Ψ_d turns out to be zero, so a different scaling is needed in (2.1.2) to get a non-trivial large deviation principle. This is carried out in Landim [11] for $d = 1$ and in Chang, Landim and Lee [2] for $d = 2$ (when $p(\cdot, \cdot)$ has positive and finite variance). We shall, however, not need this refinement.

2.2 Variational formula for $\lambda_p(\kappa)$: proof of Theorem 1.3.1

Return to (1.2.9). In this section we show that, by considering ξ and $X_1^\kappa, \dots, X_p^\kappa$ as a joint random process and exploiting the reversibility of ξ , we can use the spectral theorem to express the Lyapunov exponents in terms of a variational formula. From the latter it will follow that $\kappa \mapsto \lambda_p(\kappa)$ is continuous, non-increasing and convex on $[0, \infty)$.

Define

$$Y(t) = (\xi(t), X_1^\kappa(t), \dots, X_p^\kappa(t)), \quad t \geq 0, \quad (2.2.1)$$

and

$$V(\eta, x_1, \dots, x_p) = \sum_{i=1}^p \eta(x_i), \quad \eta \in \Omega, x_1, \dots, x_p \in \mathbb{Z}^d. \quad (2.2.2)$$

Then we may write (1.2.9) as

$$\Lambda_p(t) = \frac{1}{pt} \log \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \right). \quad (2.2.3)$$

The random process $Y = (Y(t))_{t \geq 0}$ takes values in $\Omega \times (\mathbb{Z}^d)^p$ and has generator

$$G^\kappa = L + \kappa \sum_{i=1}^p \Delta_i \quad (2.2.4)$$

in $L^2(\nu_\rho \otimes m^p)$ (endowed with the inner product (\cdot, \cdot)), with L given by (1.2.4), Δ_i the discrete Laplacian acting on the i -th spatial coordinate, and m the counting measure on \mathbb{Z}^d . Let

$$G_V^\kappa = G^\kappa + V. \quad (2.2.5)$$

By (1.2.2), this is a *self-adjoint* operator. Our claim is that λ_p equals $\frac{1}{p}$ times the upper boundary of the spectrum of G_V^κ .

Proposition 2.2.1 $\lambda_p = \frac{1}{p} \mu_p$ with $\mu_p = \sup \text{Sp}(G_V^\kappa)$.

Proof. The proof is standard. Let $(\mathcal{P}_t)_{t \geq 0}$ denote the semigroup generated by G_V^κ .

Upper bound: Let $Q_{t \log t} = [-t \log t, t \log t]^d \cap \mathbb{Z}^d$. By a standard large deviation estimate for simple random walk, we have

$$\begin{aligned} & \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \right) \\ &= \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \mathbf{1} \{X_i^\kappa(t) \in Q_{t \log t} \text{ for } i = 1, \dots, p\} \right) + R_t \end{aligned} \quad (2.2.6)$$

with $\lim_{t \rightarrow \infty} \frac{1}{t} \log R_t = -\infty$. Thus it suffices to focus on the term with the indicator.

Estimate, with the help of the spectral theorem (Kato [8], Section VI.5),

$$\begin{aligned} & \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \mathbf{1} \{X_i^\kappa(t) \in Q_{t \log t} \text{ for } i = 1, \dots, p\} \right) \\ & \leq \sum_{x_1, \dots, x_p \in Q_{t \log t}} \mathbb{E}_{\nu_\rho, x_1, \dots, x_p} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \mathbf{1} \{X_i^\kappa(t) \in Q_{t \log t} \text{ for } i = 1, \dots, p\} \right) \\ & = \left(\mathbf{1}_{(Q_{t \log t})^p}, \mathcal{P}_t \mathbf{1}_{(Q_{t \log t})^p} \right) = \int_{(-\infty, \mu_p]} e^{\mu t} d \|E_\mu \mathbf{1}_{(Q_{t \log t})^p}\|_{L^2(\nu_\rho \otimes m^p)}^2 \\ & \leq e^{\mu_p t} \| \mathbf{1}_{(Q_{t \log t})^p} \|_{L^2(\nu_\rho \otimes m^p)}^2, \end{aligned} \quad (2.2.7)$$

where $\mathbb{1}_{(Q_{t \log t})^p}$ is the indicator function of $(Q_{t \log t})^p \subset (\mathbb{Z}^d)^p$ and $(E_\mu)_{\mu \in \mathbb{R}}$ denotes the spectral family of orthogonal projection operators associated with G_V^κ . Since $\|\mathbb{1}_{(Q_{t \log t})^p}\|_{L^2(\nu_\rho \otimes m^p)}^2 = |Q_{t \log t}|^p$ does not increase exponentially fast, it follows from (1.2.10), (2.2.3) and (2.2.6–2.2.7) that $\lambda_p \leq \frac{1}{p}\mu_p$.

Lower bound: For every $\delta > 0$ there exists an $f_\delta \in L^2(\nu_\rho \otimes m^p)$ such that

$$(E_{\mu_p} - E_{\mu_p - \delta})f_\delta \neq 0 \quad (2.2.8)$$

(see Kato [8], Section VI.2; the spectrum of G_V^κ coincides with the set of μ 's for which $E_{\mu+\delta} - E_{\mu-\delta} \neq 0$ for all $\delta > 0$). Approximating f_δ by bounded functions, we may without loss of generality assume that $0 \leq f_\delta \leq 1$. Similarly, approximating f_δ by bounded functions with finite support in the spatial variables, we may assume without loss of generality that there exists a finite $K_\delta \subset \mathbb{Z}^d$ such that

$$0 \leq f_\delta \leq \mathbb{1}_{(K_\delta)^p}. \quad (2.2.9)$$

First estimate

$$\begin{aligned} & \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_0^t V(Y(s)) ds \right] \right) \\ & \geq \sum_{x_1, \dots, x_p \in K_\delta} \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\exp \left[\int_1^t V(Y(s)) ds \right] \mathbb{1}\{X_1^\kappa(1) = x_1, \dots, X_p^\kappa(1) = x_p\} \right) \\ & \geq \sum_{x_1, \dots, x_p \in K_\delta} \mathbb{E}_{\nu_\rho, 0, \dots, 0} \left(\mathbb{1}\{X_1^\kappa(1) = x_1, \dots, X_p^\kappa(1) = x_p\} \right. \\ & \quad \left. \times \mathbb{E}_{\xi(1), x_1, \dots, x_p} \left(\exp \left[\int_0^{t-1} V(Y(s)) ds \right] \right) \right) \\ & = \sum_{x_1, \dots, x_p \in K_\delta} p_1^\kappa(0, x_1) \dots p_1^\kappa(0, x_p) \mathbb{E}_{\nu_\rho, x_1, \dots, x_p} \left(\exp \left[\int_0^{t-1} V(Y(s)) ds \right] \right) \\ & \geq C_\delta^p \sum_{x_1, \dots, x_p \in K_\delta} \mathbb{E}_{\nu_\rho, x_1, \dots, x_p} \left(\exp \left[\int_0^{t-1} V(Y(s)) ds \right] \right), \end{aligned} \quad (2.2.10)$$

where $p_t^\kappa(x, y) = \mathbb{P}_x(X^\kappa(t) = y)$ and $C_\delta = \min_{x \in K_\delta} p_1^\kappa(0, x) > 0$. The equality in (2.2.10) uses that ν_ρ is invariant for the exclusion dynamics. Next estimate

$$\begin{aligned} \text{r.h.s. (2.2.10)} & \geq C_\delta^p \int_\Omega \nu_\rho(d\eta) \sum_{x_1, \dots, x_p \in \mathbb{Z}^d} f_\delta(\eta, x_1, \dots, x_p) \\ & \quad \times \mathbb{E}_{\eta, x_1, \dots, x_p} \left(\exp \left[\int_0^{t-1} V(Y(s)) ds \right] \right) f_\delta(Y(t-1)) \\ & = C_\delta^p (f_\delta, \mathcal{P}_{t-1} f_\delta) \geq \frac{C_\delta^p}{|K_\delta|^p} \int_{(\mu_p - \delta, \mu_p]} e^{\mu(t-1)} d\|E_\mu f_\delta\|_{L^2(\nu_\rho \otimes m^p)}^2 \\ & \geq C_\delta^p e^{(\mu_p - \delta)(t-1)} \|(E_{\mu_p} - E_{\mu_p - \delta})f_\delta\|_{L^2(\nu_\rho \otimes m^p)}^2, \end{aligned} \quad (2.2.11)$$

where the first inequality uses (2.2.9). Combine (2.2.10–2.2.11) with (2.2.8), and recall (2.2.3), to get $\lambda_p \geq \frac{1}{p}(\mu_p - \delta)$. Let $\delta \downarrow 0$, to obtain $\lambda_p \geq \frac{1}{p}\mu_p$. \blacksquare

The Rayleigh-Ritz formula for μ_p applied to Proposition 2.2.1 gives (recall (1.2.4), (2.2.2) and (2.2.4–2.2.5)):

Proposition 2.2.2 For all $p \in \mathbb{N}$,

$$\lambda_p = \frac{1}{p} \mu_p = \frac{1}{p} \sup_{\|f\|_{L^2(\nu_\rho \otimes m^p)}=1} (G_V^\kappa f, f) \quad (2.2.12)$$

with

$$(G_V^\kappa f, f) = A_1(f) - A_2(f) - \kappa A_3(f), \quad (2.2.13)$$

where

$$\begin{aligned} A_1(f) &= \int_{\Omega} \nu_\rho(d\eta) \sum_{z_1, \dots, z_p \in \mathbb{Z}^d} \left[\sum_{i=1}^p \eta(z_i) \right] f(\eta, z_1, \dots, z_p)^2, \\ A_2(f) &= \int_{\Omega} \nu_\rho(d\eta) \sum_{z_1, \dots, z_p \in \mathbb{Z}^d} \frac{1}{2} \sum_{\{x, y\} \subset \mathbb{Z}^d} p(x, y) [f(\eta^{x, y}, z_1, \dots, z_p) - f(\eta, z_1, \dots, z_p)]^2, \\ A_3(f) &= \int_{\Omega} \nu_\rho(d\eta) \sum_{z_1, \dots, z_p \in \mathbb{Z}^d} \frac{1}{2} \sum_{i=1}^p \sum_{\substack{y_i \in \mathbb{Z}^d \\ \|y_i - z_i\|=1}} [f(\eta, z_1, \dots, z_p)|_{z_i \rightarrow y_i} - f(\eta, z_1, \dots, z_p)]^2, \end{aligned} \quad (2.2.14)$$

and $z_i \rightarrow y_i$ means that the argument z_i is replaced by y_i .

We are now ready to give the proof of Theorem 1.3.1.

Proof. The existence of λ_p was established in Proposition 2.2.1. By (2.2.13–2.2.14), the r.h.s. of (2.2.12) is a supremum over functions that are linear and non-increasing in κ . Consequently, $\kappa \mapsto \lambda_p(\kappa)$ is lower semi-continuous, convex and non-increasing on $[0, \infty)$ (and, hence, also continuous). ■

The variational formula in Proposition 2.2.2 is useful to deduce qualitative properties of λ_p , as demonstrated above. Unfortunately, it is not clear how to deduce from it more detailed information about the Lyapunov exponents. To achieve the latter, we resort in Sections 3 and 4 to different techniques, only occasionally making use of Proposition 2.2.2.

3 Lyapunov exponents: recurrent vs. transient random walk

In this section we prove Theorems 1.3.2 and 1.3.3. In Section 3.1 we consider recurrent random walk, in Section 3.2 transient random walk.

3.1 Recurrent random walk: proof of Theorem 1.3.2(i)

The key to the proof of Theorem 1.3.2(i) is the following.

Lemma 3.1.1 *If $p(\cdot, \cdot)$ is recurrent, then for any finite box $Q \subset \mathbb{Z}^d$,*

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_{\nu_\rho} \left(\xi(x, s) = 1 \ \forall s \in [0, t] \ \forall x \in Q \right) = 0. \quad (3.1.1)$$

Proof. In the spirit of Arratia [1], Section 3, we argue as follows. Let

$$H_t^Q = \left\{ x \in \mathbb{Z}^d : \text{there is a path from } (x, 0) \text{ to } Q \times [0, t] \text{ in the graphical representation} \right\}. \quad (3.1.2)$$




Fig. 4: A path from $(x, 0)$ to $Q \times [0, t]$ (recall Fig. 1).

Note that $H_0^Q = Q$ and that $t \mapsto H_t^Q$ is non-decreasing. Denote by \mathcal{P} and \mathcal{E} , respectively, probability and expectation associated with the graphical representation. Then

$$\mathbb{P}_{\nu_\rho} \left(\xi(x, s) = 1 \ \forall s \in [0, t] \ \forall x \in Q \right) = (\mathcal{P} \otimes \nu_\rho) \left(H_t^Q \subseteq \xi(0) \right), \quad (3.1.3)$$

where $\xi(0) = \{x \in \mathbb{Z}^d : \xi(x, 0) = 1\}$ is the set of initial locations of the particles. Indeed, (3.1.3) holds because if $\xi(x, 0) = 0$ for some $x \in H_t^Q$, then this 0 will propagate into Q prior to time t (see Fig. 4).

By Jensen's inequality,

$$(\mathcal{P} \otimes \nu_\rho) \left(H_t^Q \subseteq \xi(0) \right) = \mathcal{E} \left(\rho^{|H_t^Q|} \right) \geq \rho^{\mathcal{E}|H_t^Q|}. \quad (3.1.4)$$

Moreover, $H_t^Q \subseteq \cup_{y \in Q} H_t^{\{y\}}$, and hence

$$\mathcal{E}|H_t^Q| \leq |Q| \mathcal{E}|H_t^{\{0\}}|. \quad (3.1.5)$$

Furthermore, we have

$$\mathcal{E}|H_t^{\{0\}}| = \mathbb{E}_0^{p(\cdot, \cdot)} R_t, \quad (3.1.6)$$

where R_t is the range after time t of the random walk with transition kernel $p(\cdot, \cdot)$ driving ξ and $\mathbb{E}_0^{p(\cdot, \cdot)}$ denotes expectation w.r.t. this random walk starting from 0. Indeed, by time reversal, the probability that there is a path from $(x, 0)$ to $\{0\} \times [0, t]$ in the graphical representation is equal to the probability that the random walk starting from 0 hits x prior to time t . It follows from (3.1.3–3.1.6) that

$$\frac{1}{t} \log \mathbb{P}_{\nu_\rho} \left(\xi(x, s) = 1 \ \forall s \in [0, t] \ \forall x \in Q \right) \geq -|Q| \log \left(\frac{1}{\rho} \right) \left\{ \frac{1}{t} \mathbb{E}_0^{p(\cdot, \cdot)} R_t \right\}. \quad (3.1.7)$$

Finally, since $\lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}_0^{p(\cdot, \cdot)} R_t = 0$ when $p(\cdot, \cdot)$ is recurrent (see Spitzer [13], Chapter 1, Section 4), we get (3.1.1). \blacksquare

We are now ready to give the proof of Theorem 1.3.2(i).

Proof. Since $p \mapsto \lambda_p$ is non-decreasing and $\lambda_p \leq 1$ for all $p \in \mathbb{N}$, it suffices to give the proof for $p = 1$. For $p = 1$, (1.2.9) gives

$$\Lambda_1(t) = \frac{1}{t} \log \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\int_0^t \xi(X^\kappa(s), s) ds \right] \right). \quad (3.1.8)$$

By restricting X^κ to stay inside a finite box $Q \subset \mathbb{Z}^d$ up to time t and requiring ξ to be 1 throughout this box up to time t , we obtain

$$\begin{aligned} & \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\int_0^t \xi(X^\kappa(s), s) ds \right] \right) \\ & \geq e^t \mathbb{P}_{\nu_\rho} \left(\xi(x, s) = 1 \forall s \in [0, t] \forall x \in Q \right) \mathbb{P}_0 \left(X^\kappa(s) \in Q \forall s \in [0, t] \right). \end{aligned} \quad (3.1.9)$$

For the first factor, we apply (3.1.1). For the second factor, we have

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_0 \left(X^\kappa(s) \in Q \forall s \in [0, t] \right) = -\lambda^\kappa(Q) \quad (3.1.10)$$

with $\lambda^\kappa(Q) > 0$ the principal Dirichlet eigenvalue on Q of $-\kappa\Delta$, the generator of the simple random walk X^κ . Combining (3.1.1) and (3.1.8–3.1.10), we arrive at

$$\lambda_1 = \lim_{t \rightarrow \infty} \Lambda_1(t) \geq 1 - \lambda^\kappa(Q). \quad (3.1.11)$$

Finally, let $Q \rightarrow \mathbb{Z}^d$ and use that $\lim_{Q \rightarrow \mathbb{Z}^d} \lambda^\kappa(Q) = 0$ for any κ , to arrive at $\lambda_1 \geq 1$. Since, trivially, $\lambda_1 \leq 1$, we get $\lambda_1 = 1$. \blacksquare

3.2 Transient random walk: proof of Theorems 1.3.2(ii) and 1.3.3

Theorem 1.3.2(ii) is proved in Sections 3.2.1 and 3.2.3–3.2.5, Theorem 1.3.3 in Section 3.2.2. In Section 3.2.6 we make a link between Section 2.1 and Proposition 2.2.2 for $\kappa = 0$ that provides further background for Theorem 1.3.3. Throughout the present section we assume that the random walk kernel $p(\cdot, \cdot)$ is transient.

3.2.1 Proof of the lower bound in Theorem 1.3.2(ii)

Proposition 3.2.1 $\lambda_p(\kappa) > \rho$ for all $\kappa \in [0, \infty)$ and $p \in \mathbb{N}$.

Proof. Since $p \mapsto \lambda_p(\kappa)$ is non-decreasing for all κ , it suffices to give the proof for $p = 1$. For every $\epsilon > 0$ there exists a function $\phi_\epsilon: \mathbb{Z}^d \rightarrow \mathbb{R}$ such that

$$\sum_{x \in \mathbb{Z}^d} \phi_\epsilon(x)^2 = 1 \quad \text{and} \quad \sum_{\substack{x, y \in \mathbb{Z}^d \\ \|x-y\|=1}} [\phi_\epsilon(x) - \phi_\epsilon(y)]^2 \leq \epsilon^2. \quad (3.2.1)$$

Let

$$f_\epsilon(\eta, x) = \frac{1 + \epsilon\eta(x)}{[1 + (2\epsilon + \epsilon^2)\rho]^{1/2}} \phi_\epsilon(x), \quad \eta \in \Omega, x \in \mathbb{Z}^d. \quad (3.2.2)$$

Then

$$\|f_\epsilon\|_{L^2(\nu_\rho \otimes m)}^2 = \int_\Omega \nu_\rho(d\eta) \sum_{x \in \mathbb{Z}^d} \frac{[1 + \epsilon\eta(x)]^2}{1 + (2\epsilon + \epsilon^2)\rho} \phi_\epsilon(x)^2 = \sum_{x \in \mathbb{Z}^d} \phi_\epsilon(x)^2 = 1. \quad (3.2.3)$$

Therefore we may use f_ϵ as a test function in (2.2.12) in Proposition 2.2.2. This gives

$$\lambda_1 = \mu_1 \geq \frac{1}{1 + (2\epsilon + \epsilon^2)\rho} (I - II - \kappa III) \quad (3.2.4)$$

with

$$I = \int_{\Omega} \nu_\rho(d\eta) \sum_{z \in \mathbb{Z}^d} \eta(z) [1 + \epsilon\eta(z)]^2 \phi_\epsilon(z)^2 = (1 + 2\epsilon + \epsilon^2)\rho \sum_{z \in \mathbb{Z}^d} \phi_\epsilon(z)^2 = (1 + 2\epsilon + \epsilon^2)\rho \quad (3.2.5)$$

and

$$\begin{aligned} II &= \int_{\Omega} \nu_\rho(d\eta) \sum_{z \in \mathbb{Z}^d} \frac{1}{4} \sum_{x, y \in \mathbb{Z}^d} p(x, y) \epsilon^2 [\eta^{x, y}(z) - \eta(z)]^2 \phi_\epsilon(z)^2 \\ &= \frac{1}{2} \int_{\Omega} \nu_\rho(d\eta) \sum_{x, y \in \mathbb{Z}^d} p(x, y) \epsilon^2 [\eta(x) - \eta(y)]^2 \phi_\epsilon(x)^2 \\ &= \epsilon^2 \rho (1 - \rho) \sum_{\substack{x, y \in \mathbb{Z}^d \\ x \neq y}} p(x, y) \phi_\epsilon(x)^2 \leq \epsilon^2 \rho (1 - \rho) \end{aligned} \quad (3.2.6)$$

and

$$\begin{aligned} III &= \frac{1}{2} \int_{\Omega} \nu_\rho(d\eta) \sum_{\substack{x, y \in \mathbb{Z}^d \\ \|x-y\|=1}} \{ [1 + \epsilon\eta(x)]\phi_\epsilon(x) - [1 + \epsilon\eta(y)]\phi_\epsilon(y) \}^2 \\ &= \frac{1}{2} \sum_{\substack{x, y \in \mathbb{Z}^d \\ \|x-y\|=1}} \{ [1 + (2\epsilon + \epsilon^2)\rho][\phi_\epsilon(x)^2 + \phi_\epsilon(y)^2] - 2(1 + \epsilon\rho)^2 \phi_\epsilon(x)\phi_\epsilon(y) \} \\ &= \frac{1}{2} [1 + (2\epsilon + \epsilon^2)\rho] \sum_{\substack{x, y \in \mathbb{Z}^d \\ \|x-y\|=1}} [\phi_\epsilon(x) - \phi_\epsilon(y)]^2 + \epsilon^2 \rho (1 - \rho) \sum_{\substack{x, y \in \mathbb{Z}^d \\ \|x-y\|=1}} \phi_\epsilon(x)\phi_\epsilon(y) \\ &\leq \frac{1}{2} [1 + (2\epsilon + \epsilon^2)\rho] \epsilon^2 + 2d\epsilon^2 \rho (1 - \rho). \end{aligned} \quad (3.2.7)$$

In the last line we use that $\phi_\epsilon(x)\phi_\epsilon(y) \leq \frac{1}{2}\phi_\epsilon(x)^2 + \frac{1}{2}\phi_\epsilon(y)^2$. Combining (3.2.4-3.2.7), we find

$$\lambda_1 = \mu_1 \geq \rho \frac{1 + 2\epsilon + O(\epsilon^2)}{1 + 2\epsilon\rho + O(\epsilon^2)}. \quad (3.2.8)$$

Because $\rho \in (0, 1)$, it follows that for ϵ small enough the r.h.s. is strictly larger than ρ . ■

3.2.2 Proof of Theorem 1.3.3

Proof. For $\kappa = 0$, (1.2.9) reduces to

$$\Lambda_p(t) = \frac{1}{pt} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[p \int_0^t \xi(0, s) ds \right] \right) = \frac{1}{pt} \log \mathbb{E}_{\nu_\rho} (\exp [pT_t]) \quad (3.2.9)$$

(recall (2.1.1)). In order to compute $\lambda_p(0) = \lim_{t \rightarrow \infty} \Lambda_p(t)$, we may use the large deviation principle for $(T_t)_{t \geq 0}$ cited in Section 2.1 due to Landim [11]. Indeed, by applying Varadhan's Lemma (see e.g. den Hollander [7], Theorem III.13) to (3.2.9), we get

$$\lambda_p(0) = \frac{1}{p} \max_{\alpha \in [0, 1]} [p\alpha - \Psi_d(\alpha)] \quad (3.2.10)$$

with Ψ_d the rate function given in (2.1.5). Since Ψ_d is lower semi-continuous, (3.2.10) has at least one maximizer α_p :

$$\lambda_p(0) = \alpha_p - \frac{1}{p}\Psi_d(\alpha_p). \quad (3.2.11)$$

By Proposition 3.2.1 for $\kappa = 0$, we have $\lambda_p(0) > \rho$. Hence $\alpha_p > \rho$ (because $\Psi_d(\rho) = 0$). Since $p(\cdot, \cdot)$ is transient, we may use the quadratic lower bound in (2.1.6) to see that $\Psi_d(\alpha_p) > 0$. Therefore we get from (3.2.10–3.2.11) that

$$\lambda_{p+1}(0) \geq \frac{1}{p+1} [\alpha_p(p+1) - \Psi_d(\alpha_p)] = \alpha_p - \frac{1}{p+1}\Psi_d(\alpha_p) > \alpha_p - \frac{1}{p}\Psi_d(\alpha_p) = \lambda_p(0). \quad (3.2.12)$$

Since p is arbitrary, this completes the proof of Theorem 1.3.3. \blacksquare

3.2.3 Proof of the upper bound in Theorem 1.3.2(ii)

Proposition 3.2.2 $\lambda_p(\kappa) < 1$ for all $\kappa \in [0, \infty)$ and $p \in \mathbb{N}$.

Proof. By Theorem 1.3.3, which was proved in Section 3.2.2, we know that $p \mapsto \lambda_p(0)$ is strictly increasing. Since $\lambda_p(0) \leq 1$ for all $p \in \mathbb{N}$, it therefore follows that $\lambda_p(0) < 1$ for all $p \in \mathbb{N}$. Moreover, by Theorem 1.3.1(ii), which was proved in Section 2.2, we know that $\kappa \mapsto \lambda_p(\kappa)$ is non-increasing. It therefore follows that $\lambda_p(\kappa) < 1$ for all $\kappa \in [0, \infty)$ and $p \in \mathbb{N}$. \blacksquare

3.2.4 Proof of the asymptotics in Theorem 1.3.2(ii)

The proof of the next proposition is somewhat delicate.

Proposition 3.2.3 $\lim_{\kappa \rightarrow \infty} \lambda_p(\kappa) = \rho$ for all $p \in \mathbb{N}$.

Proof. We give the proof for $p = 1$. The generalization to arbitrary p is straightforward and will be explained at the end. We need a cube $Q = [-R, R]^d \cap \mathbb{Z}^d$ of length $2R$, centered at the origin and $\delta \in (0, 1)$. Limits are taken in the order

$$t \rightarrow \infty, \kappa \rightarrow \infty, \delta \downarrow 0, Q \uparrow \mathbb{Z}^d. \quad (3.2.13)$$

The proof proceeds in 4 steps, each containing a lemma.

Step 1: Let $X^{\kappa, Q}$ be simple random walk on Q obtained from X^κ by suppressing jumps outside of Q . Then $(\xi_t, X_t^{\kappa, Q})_{t \geq 0}$ is a Markov process on $\Omega \times Q$ with self-adjoint generator in $L^2(\nu_\rho \otimes m_Q)$, where m_Q is the counting measure on Q .

Lemma 3.2.4 For all Q finite (centered and cubic) and $\kappa \in [0, \infty)$,

$$\mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\int_0^t ds \xi(X_s^\kappa, s) \right] \right) \leq e^{o(t)} \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\int_0^t ds \xi(X_s^{\kappa, Q}, s) \right] \right), \quad t \rightarrow \infty. \quad (3.2.14)$$

Proof. We consider the partition of \mathbb{Z}^d into cubes $Q_z = 2Rz + Q$, $z \in \mathbb{Z}^d$. The Lyapunov exponent $\lambda_1(\kappa)$ associated with X^κ is given by the variational formula (2.2.12–2.2.14) for $p = 1$. It can be estimated from above by splitting the sums over \mathbb{Z}^d in (2.2.14) into separate sums over the individual cubes Q_z and suppressing in $A_3(f)$ the summands on pairs of lattice

sites belonging to different cubes. The resulting expression is easily seen to coincide with the original variational expression (2.2.12), except that the supremum is restricted in addition to functions f with spatial support contained in Q . But this is precisely the Lyapunov exponent $\lambda_1^Q(\kappa)$ associated with $X^{\kappa,Q}$. Hence, $\lambda_1(\kappa) \leq \lambda_1^Q(\kappa)$, and this implies (3.2.14). \blacksquare

Step 2: For large κ the random walk $X^{\kappa,Q}$ moves fast through the finite box Q and therefore samples it in a way that is close to the uniform distribution.

Lemma 3.2.5 *For all Q finite and $\delta \in (0, 1)$, there exist $\varepsilon = \varepsilon(\kappa, \delta, Q)$ and $N_0 = N_0(\delta, \varepsilon)$, satisfying $\lim_{\kappa \rightarrow \infty} \varepsilon(\kappa, \delta, Q) = 0$ and $\lim_{\delta, \varepsilon \downarrow 0} N_0(\delta, \varepsilon) = N_0 > 1$, such that*

$$\begin{aligned} \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\int_0^t ds \xi(X_s^{\kappa,Q}, s) \right] \right) &\leq o(1) + \exp \left[\left(\left(1 + \frac{1+\varepsilon}{1-\delta} \right) \delta N_0 |Q| + \frac{\delta + \varepsilon}{1-\delta} \right) (t + \delta) \right] \\ &\quad \times \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\int_0^{t+\delta} ds \frac{1}{|Q|} \sum_{y \in Q} \xi(y, s) \right] \right), \quad t \rightarrow \infty. \end{aligned} \quad (3.2.15)$$

Proof. We split time into intervals of length $\delta > 0$. Let I_k be the indicator of the event that ξ has a jump time in Q during the k -th time interval. If $I_k = 0$, then $\xi_s = \xi_{(k-1)\delta}$ for all $s \in [(k-1)\delta, k\delta)$. Hence,

$$\int_{(k-1)\delta}^{k\delta} ds \xi_s(X_s^{\kappa,Q}) \leq \int_{(k-1)\delta}^{k\delta} ds \xi_{(k-1)\delta}(X_s^{\kappa,Q}) + \delta I_k \quad (3.2.16)$$

and, consequently, we have for all $x \in \mathbb{Z}^d$ and $k = 1, \dots, \lceil t/\delta \rceil$,

$$\mathbb{E}_x \left(\exp \left[\int_0^\delta ds \xi_{(k-1)\delta+s}(X_s^{\kappa,Q}) \right] \right) \leq e^{\delta I_k} \mathbb{E}_x \left(\exp \left[\int_0^\delta ds \eta(X_s^{\kappa,Q}) \right] \right), \quad (3.2.17)$$

where we abbreviate $\xi_{(k-1)\delta} = \eta$. Next, we do a Taylor expansion and use the Markov property of $X^{\kappa,Q}$, to obtain ($s_0 = 0$)

$$\begin{aligned} \mathbb{E}_x \left(\exp \left[\int_0^\delta ds \eta(X_s^{\kappa,Q}) \right] \right) &= \sum_{n=0}^{\infty} \left(\prod_{l=1}^n \int_{s_{l-1}}^\delta ds_l \right) \mathbb{E}_x \left(\prod_{m=1}^n \eta(X_{s_m}^{\kappa,Q}) \right) \\ &\leq \sum_{n=0}^{\infty} \left(\prod_{l=1}^n \int_{s_{l-1}}^\delta ds_l \right) \left(\prod_{m=1}^n \max_{x \in Q} \mathbb{E}_x \left(\eta(X_{s_m - s_{m-1}}^{\kappa,Q}) \right) \right) \\ &\leq \sum_{n=0}^{\infty} \left\{ \int_0^\delta ds \max_{x \in Q} \mathbb{E}_x \left(\eta(X_s^{\kappa,Q}) \right) \right\}^n \leq \exp \left[\frac{1}{1-\delta} \int_0^\delta ds \max_{x \in Q} \mathbb{E}_x \left(\eta(X_s^{\kappa,Q}) \right) \right] \\ &\leq \exp \left[\frac{1}{1-\delta} \sum_{y \in Q} \eta(y) \int_0^\delta ds \max_{x \in Q} \mathbb{E}_x \left(\delta_y(X_s^{\kappa,Q}) \right) \right], \end{aligned} \quad (3.2.18)$$

where we use that $\max_{x \in Q} \mathbb{E}_x \left(\int_0^\delta ds \eta(X_s^{\kappa,Q}) \right) \leq \delta$. Now, let $p_s^{\kappa,Q}(\cdot, \cdot)$ denote the transition kernel of $X^{\kappa,Q}$. Note that

$$\lim_{\kappa \rightarrow \infty} p_s^{\kappa,Q}(x, y) = \frac{1}{|Q|} \quad \text{for all } s > 0, \quad Q \text{ finite and } x, y \in Q. \quad (3.2.19)$$

Hence

$$\lim_{\kappa \rightarrow \infty} \mathbb{E}_x \left(\delta_y (X_s^{\kappa, Q}) \right) = \frac{1}{|Q|} \quad \text{for all } s > 0, Q \text{ finite and } x, y \in Q. \quad (3.2.20)$$

Therefore, by the Lebesgue dominated convergence theorem, we have

$$\lim_{\kappa \rightarrow \infty} \int_0^\delta ds \max_{x \in Q} \mathbb{E}_x \left(\delta_y (X_s^{\kappa, Q}) \right) = \delta \frac{1}{|Q|} \quad \text{for all } \delta > 0, Q \text{ finite and } y \in Q. \quad (3.2.21)$$

This implies that the expression in the exponent in the r.h.s. of (3.2.18) converges to

$$\frac{\delta}{1 - \delta} \frac{1}{|Q|} \sum_{y \in Q} \eta(y), \quad (3.2.22)$$

uniformly in $\eta \in \Omega$. Combining the latter with (3.2.18), we see that there exists some $\varepsilon = \varepsilon(\kappa, \delta, Q)$, satisfying $\lim_{\kappa \rightarrow \infty} \varepsilon(\kappa, \delta, Q) = 0$, such that for all $x \in Q$,

$$\mathbb{E}_x \left(\exp \left[\int_0^\delta ds \eta (X_s^{\kappa, Q}) \right] \right) \leq \exp \left[\frac{1 + \varepsilon}{1 - \delta} \delta \frac{1}{|Q|} \sum_{y \in Q} \eta(y) \right] \quad \text{for all } \delta \in (0, 1) \text{ and } Q \text{ finite.} \quad (3.2.23)$$

Next, similarly as in (3.2.16), we have

$$\delta \frac{1}{|Q|} \sum_{y \in Q} \xi_{(k-1)\delta}(y) \leq \int_{(k-1)\delta}^{k\delta} ds \frac{1}{|Q|} \sum_{y \in Q} \xi_s(y) + \delta I_k. \quad (3.2.24)$$

Applying the Markov property to $X^{\kappa, Q}$, and using (3.2.16) and (3.2.23-3.2.24), we find that

$$\begin{aligned} \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\int_0^t ds \xi (X_s^{\kappa, Q}, s) \right] \right) &\leq \mathbb{E}_{\nu_\rho} \left(\exp \left[\left(1 + \frac{1 + \varepsilon}{1 - \delta} \right) \delta N_{t+\delta} + \frac{\delta + \varepsilon}{1 - \delta} (t + \delta) \right] \right. \\ &\quad \left. \times \exp \left[\int_0^{t+\delta} ds \frac{1}{|Q|} \sum_{y \in Q} \xi_s(y) \right] \right), \end{aligned} \quad (3.2.25)$$

where $N_{t+\delta}$ is the total number of jumps that ξ makes inside Q up to time $t + \delta$. The second term in the r.h.s. of (3.2.25) equals the second term in the r.h.s. of (3.2.15). The first term will be negligible on an exponential scale for $\delta \downarrow 0$, because, as can be seen from the graphical representation, $N_{t+\delta}$ is stochastically smaller than the total number of jumps up to time $t + \delta$ of a Poisson process with rate $|Q \cup \partial Q|$. Indeed, abbreviating

$$a = \left(1 + \frac{1 + \varepsilon}{1 - \delta} \right) \delta, \quad b = \frac{\delta + \varepsilon}{1 - \delta}, \quad M_{t+\delta} = \int_0^{t+\delta} ds \frac{1}{|Q|} \sum_{y \in Q} \xi_s(y), \quad (3.2.26)$$

we estimate, for each N ,

$$\begin{aligned} \text{r.h.s. (3.2.25)} &= \mathbb{E}_{\nu_\rho} \left(e^{aN_{t+\delta} + b(t+\delta) + M_{t+\delta}} \right) \\ &\leq e^{(b+1)(t+\delta)} \mathbb{E}_{\nu_\rho} \left(e^{aN_{t+\delta}} 1_{\{N_{t+\delta} \geq N|Q|(t+\delta)\}} \right) + e^{(aN|Q| + b)(t+\delta)} \mathbb{E}_{\nu_\rho} \left(e^{M_{t+\delta}} \right). \end{aligned} \quad (3.2.27)$$

For $N \geq N_0 = N_0(a, b)$, the first term tends to zero as $t \rightarrow \infty$ and can be discarded. Hence

$$\text{r.h.s. (3.2.25)} \leq e^{(aN_0|Q| + b)(t+\delta)} \mathbb{E}_{\nu_\rho} \left(e^{bM_{t+\delta}} \right), \quad (3.2.28)$$

which is the desired bound in (3.2.15). Note that $a \downarrow 0$, $b \downarrow 1$ as $\delta, \varepsilon \downarrow 0$ and hence $N_0(a, b) \downarrow N_0 > 1$. \blacksquare

Step 3: By combining Lemmas 3.2.4–3.2.5, we now know that for any Q finite,

$$\lim_{\kappa \rightarrow \infty} \lambda_1(\kappa) \leq \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\int_0^t ds \frac{1}{|Q|} \sum_{y \in Q} \xi_s(y) \right] \right), \quad (3.2.29)$$

where we have taken the limits $\kappa \rightarrow \infty$ and $\delta \downarrow 0$. According to Proposition 1.2.1 (with $K(z, s) = (1/|Q|)1_Q(z)$),

$$\mathbb{E}_{\nu_\rho} \left(\exp \left[\int_0^t ds \frac{1}{|Q|} \sum_{y \in Q} \xi_s(y) \right] \right) \leq \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[\int_0^t ds \frac{1}{|Q|} \sum_{y \in Q} \tilde{\xi}_s(y) \right] \right), \quad (3.2.30)$$

where $(\tilde{\xi}_t)_{t \geq 0}$ is the process of Independent Random Walks on \mathbb{Z}^d with step rate 1 and transition kernel $p(\cdot, \cdot)$, and $\mathbb{E}_{\nu_\rho}^{\text{IRW}} = \int_\Omega \nu_\rho(d\eta) \mathbb{E}_\eta^{\text{IRW}}$. The r.h.s. can be computed and estimated as follows. Write

$$(\Delta^{(p)}f)(x) = \sum_{y \in \mathbb{Z}^d} p(x, y)[f(y) - f(x)], \quad x \in \mathbb{Z}^d, \quad (3.2.31)$$

to denote the generator of the random walk with step rate 1 and transition kernel $p(\cdot, \cdot)$.

Lemma 3.2.6 *For all Q finite,*

$$\text{r.h.s. (3.2.30)} \leq e^{\rho t} \exp \left[\int_0^t ds \frac{1}{|Q|} \sum_{x \in Q} w^Q(x, s) \right], \quad (3.2.32)$$

where $w^Q: \mathbb{Z}^d \times [0, \infty) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial w^Q}{\partial t}(x, t) = \Delta^{(p)}w^Q(x, t) + \left\{ \frac{1}{|Q|}1_Q(x) \right\} [w^Q(x, t) + 1], \quad w^Q(\cdot, 0) \equiv 0, \quad (3.2.33)$$

which has the representation

$$w^Q(x, t) = \mathbb{E}_x^{\text{RW}} \left(\exp \left[\int_0^t ds \frac{1}{|Q|}1_Q(Y_s) \right] \right) - 1 \geq 0, \quad (3.2.34)$$

where $Y = (Y_t)_{t \geq 0}$ is the single random walk with step rate 1 and transition kernel $p(\cdot, \cdot)$, and \mathbb{E}_x^{RW} denotes the expectation w.r.t. to Y starting from $Y_0 = x$.

Proof. Let

$$A_\eta = \{x \in \mathbb{Z}^d: \eta(x) = 1\}, \quad \eta \in \Omega. \quad (3.2.35)$$

Then

$$\begin{aligned} \text{r.h.s. (3.2.30)} &= \int_\Omega \nu_\rho(d\eta) \mathbb{E}_\eta^{\text{IRW}} \left(\exp \left[\int_0^t ds \frac{1}{|Q|} \sum_{x \in A_\eta} \sum_{y \in Q} 1_y(\tilde{\xi}_{s,x}) \right] \right) \\ &= \int_\Omega \nu_\rho(d\eta) \prod_{x \in A_\eta} \mathbb{E}_x^{\text{RW}} \left(\exp \left[\int_0^t ds \frac{1}{|Q|}1_Q(Y_s) \right] \right), \end{aligned} \quad (3.2.36)$$

where $\tilde{\xi}_{s,x}$ is the position at time s of the random walk starting from $\tilde{\xi}_{0,x} = x$ (in the process of Independent Random Walks $\tilde{\xi} = (\tilde{\xi}_t)_{t \geq 0}$). Let

$$v^Q(x, t) = \mathbb{E}_x^{\text{RW}} \left(\exp \left[\int_0^t ds \frac{1}{|Q|} 1_Q(Y_s) \right] \right). \quad (3.2.37)$$

By the Feynman-Kac formula, $v^Q(x, t)$ is the solution of the Cauchy problem

$$\frac{\partial v^Q}{\partial t}(x, t) = \Delta^{(p)} v^Q(x, t) + \left\{ \frac{1}{|Q|} 1_Q(x) \right\} v^Q(x, t), \quad v^Q(\cdot, 0) \equiv 1. \quad (3.2.38)$$

Now put

$$w^Q(x, t) = v^Q(x, t) - 1. \quad (3.2.39)$$

Then (3.2.38) can be rewritten as (3.2.33). Combining (3.2.36–3.2.37) and (3.2.39), we get

$$\begin{aligned} \text{r.h.s. (3.2.30)} &= \int_{\Omega} \nu_{\rho}(d\eta) \prod_{x \in A_{\eta}} (1 + w^Q(x, t)) = \int_{\Omega} \nu_{\rho}(d\eta) \prod_{x \in \mathbb{Z}^d} (1 + \eta(x) w^Q(x, t)) \\ &= \prod_{x \in \mathbb{Z}^d} (1 + \rho w^Q(x, t)) \leq \exp \left[\rho \sum_{x \in \mathbb{Z}^d} w^Q(x, t) \right], \end{aligned} \quad (3.2.40)$$

where we use that ν_{ρ} is the Bernoulli product measure with density ρ . Summing (3.2.33) over \mathbb{Z}^d , we have

$$\frac{\partial}{\partial t} \left(\sum_{x \in \mathbb{Z}^d} w^Q(x, t) \right) = \sum_{x \in Q} \frac{1}{|Q|} w^Q(x, t) + 1. \quad (3.2.41)$$

Integrating (3.2.41) w.r.t. time, we get

$$\sum_{x \in \mathbb{Z}^d} w^Q(x, t) = \int_0^t ds \sum_{x \in Q} \frac{1}{|Q|} w^Q(x, s) + t. \quad (3.2.42)$$

Combining (3.2.40) and (3.2.42), we get the claim. \blacksquare

Step 4: The proof is completed by showing the following:

Lemma 3.2.7

$$\lim_{Q \uparrow \mathbb{Z}^d} \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t ds \frac{1}{|Q|} \sum_{x \in Q} w^Q(x, s) = 0. \quad (3.2.43)$$

Proof. Let \mathcal{G} denote the Green operator acting on functions $V: \mathbb{Z}^d \rightarrow [0, \infty)$ as

$$(\mathcal{G}V)(x) = \sum_{y \in \mathbb{Z}^d} G(x, y) V(y), \quad x \in \mathbb{Z}^d, \quad (3.2.44)$$

where $G(x, y) = \int_0^{\infty} dt p_t(x, y)$ denotes the Green kernel on \mathbb{Z}^d . We have

$$\left\| \mathcal{G} \left(\frac{1}{|Q|} 1_{|Q|} \right) \right\|_{\infty} = \sup_{x \in \mathbb{Z}^d} \sum_{y \in Q} G(x, y) \frac{1}{|Q|}. \quad (3.2.45)$$

The r.h.s. tends to zero as $Q \uparrow \mathbb{Z}^d$, because $G(x, y)$ tends to zero as $\|x - y\| \rightarrow \infty$. Hence Lemma 8.2.1 in Gärtner and den Hollander [4] can be applied to (3.2.34) for Q large enough, to yield

$$\sup_{\substack{x \in \mathbb{Z}^d \\ s \geq 0}} w^Q(x, s) \leq \varepsilon(Q) \downarrow 0 \quad \text{as } Q \uparrow \mathbb{Z}^d, \quad (3.2.46)$$

which proves (3.2.43). ■

Combine (3.2.29–3.2.30), (3.2.32) and (3.2.43) to get the claim in Proposition 3.2.3.

This completes the proof of Proposition 3.2.3 for $p = 1$. The generalization to arbitrary p is straightforward and runs as follows. Return to (1.2.9). Separate the p terms under the sum with the help of Hölder's inequality with weights $1/p$. Next, use (3.2.14) for each of the p factors, leading to $\frac{1}{p} \log$ of the r.h.s. of (3.2.14) with an extra factor p in the exponent. Then proceed as before, which leads to Lemma 3.2.6 but with w^Q the solution of (3.2.33) with $\frac{p}{|Q|} 1_Q(x)$ between braces. Then again proceed as before, which leads to (3.2.40) but with an extra factor p in the r.h.s. of (3.2.42). The latter gives a factor $e^{p\rho t}$ replacing $e^{\rho t}$ in (3.2.32). Now use Lemma 3.2.7 to get the claim. ■

3.2.5 Proof of the strict monotonicity in Theorem 1.3.2(ii)

By Theorem 1.3.1(ii), $\kappa \mapsto \lambda_p(\kappa)$ is convex. Because of Proposition 3.2.1 and Proposition 3.2.3, it must be strictly decreasing. This completes the proof of Theorem 1.3.2(ii).

3.2.6 Relation between Proposition 2.2.2 and (3.2.10)

Let $\kappa = 0$ and $p = 1$. The generalization to arbitrary p is straightforward.

For $\kappa = 0$ and $p = 1$, (2.2.12–2.2.14) in Proposition 2.2.2 read

$$\lambda_1(0) = \sup_{\|f\|_{L^2(\nu_{\rho \otimes m})} = 1} \sum_{z \in \mathbb{Z}^d} \left[\int_{\Omega} \nu_{\rho}(d\eta) \eta(z) f(\eta, z)^2 - \mathcal{E}(f(\cdot, z)) \right], \quad (3.2.47)$$

where we recall (2.1.3). Split the supremum into two parts,

$$\lambda_1(0) = \sup_{\|g\|_{L^2(m)} = 1} \sup_{\|f_z\|_{L^2(\nu_{\rho})} = 1 \forall z \in \mathbb{Z}^d} \sum_{z \in \mathbb{Z}^d} g^2(z) \left[\int_{\Omega} \nu_{\rho}(d\eta) \eta(z) f_z(\eta)^2 - \mathcal{E}(f_z) \right], \quad (3.2.48)$$

where $f_z(\eta) = f(\eta, z)/g(z)$ with $g(z)^2 = \int_{\Omega} \nu_{\rho}(d\eta) f(\eta, z)^2$. The second supremum in (3.2.48), which runs over a family of functions indexed by z , can be brought under the sum,

$$\lambda_1(0) = \sup_{\|g\|_{L^2(m)} = 1} \sum_{z \in \mathbb{Z}^d} g^2(z) \sup_{\|f_z\|_{L^2(\nu_{\rho})} = 1} \left[\int_{\Omega} \nu_{\rho}(d\eta) \eta(z) f_z(\eta)^2 - \mathcal{E}(f_z) \right]. \quad (3.2.49)$$

By the shift-invariance of ν_{ρ} , we may replace $\eta(z)$ by $\eta(0)$ under the second supremum in (3.2.49), in which case the latter no longer depends on z , and we get

$$\begin{aligned} \lambda_1(0) &= \sup_{\|f\|_{L^2(\nu_{\rho})} = 1} \left[\int_{\Omega} \nu_{\rho}(d\eta) \eta(0) f(\eta)^2 - \mathcal{E}(f) \right] \\ &= \sup_{\|f\|_{L^2(\nu_{\rho})} = 1} \left[\int_{\Omega} \nu_{\rho}(d\eta) \eta(0) f(\eta)^2 - I_d(f^2 \nu_{\rho}) \right]. \end{aligned} \quad (3.2.50)$$

Here note that the smoothing in (2.1.4) can be removed under the supremum in (3.2.50) (recall the remark made below (2.1.5)). But the r.h.s. of (3.2.50) is precisely the r.h.s. of (3.2.10) for $p = 1$, where we recall (2.1.4–2.1.5) and put $f(\eta)^2 = (d\mu/d\nu_\rho)(\eta)$.

4 Lyapunov exponents: transient simple random walk

This section is devoted to the proof of Theorem 1.3.4, where $d \geq 4$ and $p(\cdot, \cdot)$ is simple random walk given by (1.2.3), i.e., ξ is simple symmetric exclusion (SSE). *The proof is long and technical*, taking up half of the present paper. In Sections 4.2–4.7, we give the proof for $p = 1$. In Section 4.8, we indicate how to extend the proof to arbitrary p .

4.1 Outline

In Section 4.2, we do an appropriate scaling in κ . In Section 4.3, we introduce an SSE+RW generator and an auxiliary exponential martingale. In Section 4.4, we compute upper and lower bounds for the l.h.s. of (1.3.1) in terms of certain key quantities, and we complete the proof of Theorem 1.3.4 (for $p = 1$) subject to two propositions, whose proof is given in Sections 4.6–4.7. In Section 4.5, we list some preparatory facts that are needed as we go along.

As before, we write $X_s^\kappa, \xi_s(x)$ instead of $X^\kappa(s), \xi(x, s)$. We abbreviate

$$1[\kappa] = 1 + \frac{1}{2d\kappa}, \quad (4.1.1)$$

and write $\{a, b\}$ to denote the unoriented bond between nearest-neighbor sites $a, b \in \mathbb{Z}^d$ (recall (1.2.3)–(1.2.4)). Three parameters will be important: t, κ and T . We will take limits in the following order:

$$t \rightarrow \infty, \quad \kappa \rightarrow \infty, \quad T \rightarrow \infty. \quad (4.1.2)$$

4.2 Scaling

We have $X_t^\kappa = X_{\kappa t}$, $t \geq 0$, where $X = (X_t)_{t \geq 0}$ is simple random walk with step rate $2d$, being independent of $(\xi_t)_{t \geq 0}$. We therefore have

$$\mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\int_0^t ds \xi_s(X_s^\kappa) \right] \right) = \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{\kappa} \int_0^{\kappa t} ds \xi_{\frac{s}{\kappa}}(X_s) \right] \right). \quad (4.2.1)$$

Define the scaled Lyapunov exponent (recall (1.2.9–1.2.10))

$$\lambda_1^*(\kappa) = \lim_{t \rightarrow \infty} \Lambda_1^*(\kappa; t) \quad \text{with} \quad \Lambda_1^*(\kappa; t) = \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \xi_{\frac{s}{\kappa}}(X_s) \right] \right). \quad (4.2.2)$$

Then $\lambda_1(\kappa) = \kappa \lambda_1^*(\kappa)$. Therefore, in what follows we will focus on the quantity

$$\lambda_1^*(\kappa) - \frac{\rho}{\kappa} = \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(X_s) - \rho \right) \right] \right) \quad (4.2.3)$$

and compute its asymptotic behavior for large κ . We must show that

$$\lim_{\kappa \rightarrow \infty} 2d\kappa^2 \left[\lambda_1^*(\kappa) - \frac{\rho}{\kappa} \right] = \rho(1 - \rho)G_d. \quad (4.2.4)$$

4.3 SSE+RW generator and an auxiliary exponential martingale

For $t \geq 0$, let

$$Z_t = (\xi_{\frac{t}{\kappa}}, X_t) \quad (4.3.1)$$

and denote by $\mathbb{P}_{\eta,x}$ the law of Z starting from $Z_0 = (\eta, x)$. Then $Z = (Z_t)_{t \geq 0}$ is a Markov process on $\Omega \times \mathbb{Z}^d$ with generator

$$\mathcal{A} = \frac{1}{\kappa}L + \Delta \quad (4.3.2)$$

(acting on the Banach space of bounded continuous functions on $\Omega \times \mathbb{Z}^d$, equipped with the supremum norm). Let $(\mathcal{P}_t)_{t \geq 0}$ be the semigroup generated by \mathcal{A} . The following lemma will be crucial to rewrite the expectation in the r.h.s. of (4.2.3) in a more manageable form.

Lemma 4.3.1 *Fix $\kappa > 0$ and $r > 0$. For all $t \geq 0$ and all bounded continuous functions $\psi: \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$ such that ψ and $\exp[(r/\kappa)\psi]$ belong to the domain of \mathcal{A} , define*

$$\begin{aligned} M_t^r &= \frac{r}{\kappa} \left[\psi(Z_t) - \psi(Z_0) - \int_0^t ds \mathcal{A}\psi(Z_s) \right], \\ N_t^r &= \exp \left[M_t^r - \int_0^t ds \left[\left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) - \mathcal{A} \left(\frac{r}{\kappa}\psi \right) \right] (Z_s) \right]. \end{aligned} \quad (4.3.3)$$

Then:

- (i) $M^r = (M_t^r)_{t \geq 0}$ is a $\mathbb{P}_{\eta,x}$ -martingale for all (η, x) .
- (ii) For $t \geq 0$, let $\mathcal{P}_t^{\text{new}}$ be the operator defined by

$$(\mathcal{P}_t^{\text{new}} f)(\eta, x) = e^{-\frac{r}{\kappa}\psi(\eta,x)} \mathbb{E}_{\eta,x} \left(\exp \left[- \int_0^t ds \left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) (Z_s) \right] \left(e^{\frac{r}{\kappa}\psi} f \right) (Z_t) \right) \quad (4.3.4)$$

for bounded continuous $f: \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$. Then $(\mathcal{P}_t^{\text{new}})_{t \geq 0}$ is a strongly continuous semigroup with generator

$$(\mathcal{A}^{\text{new}} f)(\eta, x) = \left[e^{-\frac{r}{\kappa}\psi} \mathcal{A} \left(e^{\frac{r}{\kappa}\psi} f \right) - \left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) f \right] (\eta, x). \quad (4.3.5)$$

- (iii) $N^r = (N_t^r)_{t \geq 0}$ is a $\mathbb{P}_{\eta,x}$ -martingale for all (η, x) .
- (iv) Define a new path measure $\mathbb{P}_{\eta,x}^{\text{new}}$ by putting

$$\frac{d\mathbb{P}_{\eta,x}^{\text{new}}}{d\mathbb{P}_{\eta,x}}((Z_s)_{0 \leq s \leq t}) = N_t^r, \quad t \geq 0. \quad (4.3.6)$$

Then, under $\mathbb{P}_{\eta,x}^{\text{new}}$, $(Z_t)_{t \geq 0}$ is a Markov process with semigroup $(\mathcal{P}_t^{\text{new}})_{t \geq 0}$.

Proof. The proof is standard.

- (i) This follows from the fact that \mathcal{A} is a Markov generator and ψ belongs to its domain (see Liggett [12], Chapter I, Section 5).
- (ii) Let $\eta \in \Omega$, $x \in \mathbb{Z}^d$ and $f: \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$ bounded measurable. Rewrite (4.3.4) as

$$\begin{aligned} (\mathcal{P}_t^{\text{new}} f)(\eta, x) &= \mathbb{E}_{\eta,x} \left(\exp \left[\frac{r}{\kappa}\psi(Z_t) - \frac{r}{\kappa}\psi(Z_0) - \int_0^t ds \left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) (Z_s) \right] f(Z_t) \right) \\ &= \mathbb{E}_{\eta,x} (N_t^r f(Z_t)). \end{aligned} \quad (4.3.7)$$

This gives

$$(\mathcal{P}_0^{\text{new}} f)(\eta, x) = f(\eta, x) \quad (4.3.8)$$

and

$$\begin{aligned} (\mathcal{P}_{t_1+t_2}^{\text{new}} f)(\eta, x) &= \mathbb{E}_{\eta, x} (N_{t_1+t_2}^r f(Z_{t_1+t_2})) = \mathbb{E}_{\eta, x} \left(N_{t_1}^r \frac{N_{t_1+t_2}^r}{N_{t_1}^r} f(Z_{t_1+t_2}) \right) \\ &= \mathbb{E}_{\eta, x} \left(N_{t_1}^r \mathbb{E}_{Z_{t_1}} (N_{t_2}^r f(Z_{t_2})) \right) = (\mathcal{P}_{t_1}^{\text{new}} (\mathcal{P}_{t_2}^{\text{new}} f))(\eta, x), \end{aligned} \quad (4.3.9)$$

where we use the Markov property of Z at time t_1 (under $\mathbb{P}_{\eta, x}$) together with the fact that $N_{t_1+t_2}^r/N_{t_1}^r$ only depends on Z_t for $t \in [t_1, t_1+t_2]$. Equations (4.3.8–4.3.9) show that $(\mathcal{P}_t^{\text{new}})_{t \geq 0}$ is a semigroup which is easily seen to be strongly continuous.

Taking the derivative of (4.3.4) in the norm w.r.t. t at $t = 0$, we get (4.3.5). Next, if $f \equiv 1$, then (4.3.5) gives $\mathcal{A}^{\text{new}} 1 = 0$. This last equality implies that

$$\frac{1}{\lambda} (\lambda Id - \mathcal{A}^{\text{new}}) 1 = 1 \quad \forall \lambda > 0. \quad (4.3.10)$$

Since $\lambda Id - \mathcal{A}^{\text{new}}$ is invertible, we get

$$(\lambda Id - \mathcal{A}^{\text{new}})^{-1} 1 = \frac{1}{\lambda} \quad \forall \lambda > 0, \quad (4.3.11)$$

i.e.,

$$\int_0^\infty dt e^{-\lambda t} \mathcal{P}_t^{\text{new}} 1 = \frac{1}{\lambda} \quad \forall \lambda > 0. \quad (4.3.12)$$

Inverting this Laplace transform, we see that

$$\mathcal{P}_t^{\text{new}} 1 = 1 \quad \forall t \geq 0. \quad (4.3.13)$$

(iii) Fix $t \geq 0$ and $h > 0$. Since N_t^r is \mathcal{F}_t -measurable, with \mathcal{F}_t the σ -algebra generated by $(Z_s)_{0 \leq s \leq t}$, we have

$$\begin{aligned} &\mathbb{E}_{\eta, x} (N_{t+h}^r | \mathcal{F}_t) \\ &= N_t^r \mathbb{E}_{\eta, x} \left(\exp \left[M_{t+h}^r - M_t^r - \int_t^{t+h} ds \left[\left(e^{-\frac{r}{\kappa} \psi} \mathcal{A} e^{\frac{r}{\kappa} \psi} \right) - \mathcal{A} \left(\frac{r}{\kappa} \psi \right) \right] (Z_s) \right] \middle| \mathcal{F}_t \right). \end{aligned} \quad (4.3.14)$$

Applying the Markov property of Z at time t , we get

$$\begin{aligned} \mathbb{E}_{\eta, x} (N_{t+h}^r | \mathcal{F}_t) &= N_t^r \mathbb{E}_{Z_t} \left(\exp \left[\frac{r}{\kappa} \psi(Z_h) - \frac{r}{\kappa} \psi(Z_0) - \int_0^h ds \left(e^{-\frac{r}{\kappa} \psi} \mathcal{A} e^{\frac{r}{\kappa} \psi} \right) (Z_s) \right] \right) \\ &= N_t^r (\mathcal{P}_h^{\text{new}} 1)(Z_t) = N_t^r, \end{aligned} \quad (4.3.15)$$

where the third equality uses (4.3.13).

(iv) This follows from (iii) via a calculation similar to (4.3.9). ■

4.4 Proof of Theorem 1.3.4

In this section we compute upper and lower bounds for the r.h.s. of (4.2.3) in terms of certain key quantities (Proposition 4.4.1 below). We then state two propositions for these quantities (Propositions 4.4.2–4.4.3 below), from which Theorem 1.3.4 will follow. The proof of these two propositions is given in Sections 4.6–4.7.

For $T > 0$, let $\psi: \Omega \times \mathbb{Z}^d$ be defined by

$$\psi(\eta, x) = \int_0^T ds (\mathcal{P}_s \phi)(\eta, x) \quad \text{with} \quad \phi(\eta, x) = \eta(x) - \rho, \quad (4.4.1)$$

where $(\mathcal{P}_t)_{t \geq 0}$ is the semigroup generated by \mathcal{A} (recall (4.3.2)). We have

$$\psi(\eta, x) = \int_0^T ds \mathbb{E}_{\eta, x} (\phi(Z_s)) = \int_0^T ds \mathbb{E}_\eta \sum_{y \in \mathbb{Z}^d} p_{2ds}(y, x) \left(\xi_{\frac{s}{\kappa}}(y) - \rho \right), \quad (4.4.2)$$

where $p_t(x, y)$ is the probability that simple random walk with step rate 1 moves from x to y in time t (recall that we assume (1.2.3)). Using (1.2.6), we obtain the representation

$$\psi(\eta, x) = \int_0^T ds \sum_{z \in \mathbb{Z}^d} p_{2ds1[\kappa]}(z, x) [\eta(z) - \rho], \quad (4.4.3)$$

where $1[\kappa]$ is given by (4.1.1). Note that ψ depends on κ and T . We suppress this dependence. Similarly,

$$-\mathcal{A}\psi = \int_0^T ds (-\mathcal{A}\mathcal{P}_s \phi) = \phi - \mathcal{P}_T \phi, \quad (4.4.4)$$

with

$$(\mathcal{P}_T \phi)(\eta, x) = \mathbb{E}_{\eta, x} (\phi(Z_T)) = \mathbb{E}_{\eta, x} \left(\xi_{\frac{T}{\kappa}}(X_T) - \rho \right) = \sum_{z \in \mathbb{Z}^d} p_{2dT1[\kappa]}(z, x) [\eta(z) - \rho]. \quad (4.4.5)$$

The auxiliary function ψ will play a key role throughout the remaining sections. The integral in (4.4.1) is a regularization that is useful when dealing with central limit type behavior of Markov processes (see e.g. Kipnis [10]). Heuristically, $T = \infty$ corresponds to $-\mathcal{A}\psi = \phi$. Later we will let $T \rightarrow \infty$.

The following proposition serves as the starting point of our asymptotic analysis.

Proposition 4.4.1 *For any $\kappa, T > 0$,*

$$\lambda_1^*(\kappa) - \frac{\rho}{\kappa} \begin{matrix} \leq \\ \geq \end{matrix} I_1^{r,q}(\kappa, T) + I_2^{r,q}(\kappa, T), \quad (4.4.6)$$

where

$$\begin{aligned} I_1^{r,q}(\kappa, T) &= \frac{1}{2q} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{2q}{r} \int_0^t ds \left[\left(e^{-\frac{r}{\kappa} \psi} \mathcal{A} e^{\frac{r}{\kappa} \psi} \right) - \mathcal{A} \left(\frac{r}{\kappa} \psi \right) \right] (Z_s) \right] \right), \\ I_2^{r,q}(\kappa, T) &= \frac{1}{2q} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{2q}{\kappa} \int_0^t ds (\mathcal{P}_T \phi)(Z_s) \right] \right), \end{aligned} \quad (4.4.7)$$

and $1/r + 1/q = 1$ for any $r, q > 1$ in the first inequality and any $q < 0 < r < 1$ in the second inequality.

Proof. Recall (4.2.3). From the first line of (4.3.3) and (4.4.4) it follows that

$$\frac{1}{r}M_t^r + \frac{1}{\kappa}\psi(Z_0) - \frac{1}{\kappa}\psi(Z_t) = \frac{1}{\kappa} \int_0^t ds [(-\mathcal{A})\psi](Z_s) = \frac{1}{\kappa} \int_0^t ds \phi(Z_s) - \frac{1}{\kappa} \int_0^t ds (\mathcal{P}_T\phi)(Z_s). \quad (4.4.8)$$

Hence

$$\begin{aligned} \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \phi(Z_s) \right] \right) \\ = \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{1}{r}M_t^r + \frac{1}{\kappa}\psi(Z_0) - \frac{1}{\kappa}\psi(Z_t) + \frac{1}{\kappa} \int_0^t ds (\mathcal{P}_T\phi)(Z_s) \right] \right) \\ = \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[U_t^r + \frac{1}{r}V_t^r \right] \right) \end{aligned} \quad (4.4.9)$$

with

$$U_t^r = \frac{1}{r} \int_0^t ds \left[\left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) - \mathcal{A} \left(\frac{r}{\kappa}\psi \right) \right] (Z_s) + \frac{1}{\kappa} \left(\psi(Z_0) - \psi(Z_t) \right) + \frac{1}{\kappa} \int_0^t ds (\mathcal{P}_T\phi)(Z_s) \quad (4.4.10)$$

and

$$V_t^r = M_t^r - \int_0^t ds \left[\left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) - \mathcal{A} \left(\frac{r}{\kappa}\psi \right) \right] (Z_s). \quad (4.4.11)$$

By Hölder's inequality, with $r, q > 1$ such that $1/r + 1/q = 1$, it follows from (4.4.9) that

$$\begin{aligned} \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \phi(Z_s) \right] \right) &\leq \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [V_t^r]) \right)^{1/r} \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [qU_t^r]) \right)^{1/q} \\ &= \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [qU_t^r]) \right)^{1/q}, \end{aligned} \quad (4.4.12)$$

where the second line of (4.4.12) comes from the fact that $N_t^r = \exp[V_t^r]$ is a martingale, by Lemma 4.3.1(iii). Similarly, by the reverse of Hölder's inequality, with $q < 0 < r < 1$ such that $1/r + 1/q = 1$, it follows from (4.4.9) that

$$\begin{aligned} \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \phi(Z_s) \right] \right) &\geq \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [V_t^r]) \right)^{1/r} \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [qU_t^r]) \right)^{1/q} \\ &= \left(\mathbb{E}_{\nu_{\rho,0}} (\exp [qU_t^r]) \right)^{1/q}. \end{aligned} \quad (4.4.13)$$

The middle term in the r.h.s. of (4.4.10) can be discarded, because (4.4.3) shows that $-\rho T \leq \psi \leq (1-\rho)T$. Apply the Cauchy-Schwarz inequality to the r.h.s. of (4.4.12–4.4.13) to separate the other two terms in the r.h.s. of (4.4.10). \blacksquare

Note that in the r.h.s. of (4.4.7) the prefactors of the logarithms and the prefactors in the exponents are *both positive for the upper bound and both negative for the lower bound*. This will be important later on.

The following two propositions will be proved in Sections 4.6–4.7, respectively. Abbreviate

$$\limsup_{t,\kappa,T \rightarrow \infty} = \limsup_{T \rightarrow \infty} \limsup_{\kappa \rightarrow \infty} \limsup_{t \rightarrow \infty}. \quad (4.4.14)$$

Proposition 4.4.2 *If $d \geq 3$, then for any $\alpha \in \mathbb{R}$ and $r > 0$,*

$$\limsup_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_{\rho, 0}} \left(\exp \left[\frac{\alpha}{r} \int_0^t ds \left[e^{-\frac{r}{\kappa} \psi} \mathcal{A} e^{\frac{r}{\kappa} \psi} - \mathcal{A} \left(\frac{r}{\kappa} \psi \right) \right] (Z_s) \right] \right) \leq \alpha r \rho (1 - \rho) \frac{1}{2d} G_d. \quad (4.4.15)$$

Proposition 4.4.3 *If $d \geq 4$, then for any $\alpha \in \mathbb{R}$,*

$$\limsup_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_{\rho, 0}} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds (\mathcal{P}_T \phi) (Z_s) \right] \right) \leq 0. \quad (4.4.16)$$

Picking $\alpha = 2q$ in Proposition 4.4.2, we see that the first term in the r.h.s. of (4.4.6) satisfies the bounds

$$\begin{aligned} \limsup_{T \rightarrow \infty} \limsup_{\kappa \rightarrow \infty} \kappa^2 I_1^{r, q}(\kappa, T) &\leq r \rho (1 - \rho) \frac{1}{2d} G_d \quad \text{if } r > 1, \\ \liminf_{T \rightarrow \infty} \liminf_{\kappa \rightarrow \infty} \kappa^2 I_1^{r, q}(\kappa, T) &\geq r \rho (1 - \rho) \frac{1}{2d} G_d \quad \text{if } r < 1. \end{aligned} \quad (4.4.17)$$

Letting r tend to 1, we obtain

$$\lim_{T \rightarrow \infty} \lim_{\kappa \rightarrow \infty} \kappa^2 I_1^{r, q}(\kappa, T) = \rho (1 - \rho) \frac{1}{2d} G_d. \quad (4.4.18)$$

Picking $\alpha = 2q$ in Proposition 4.4.3, we see that the second term in the r.h.s. of (4.4.6) satisfies

$$\limsup_{T \rightarrow \infty} \limsup_{\kappa \rightarrow \infty} \kappa^2 I_2^{r, q}(\kappa, T) = 0 \quad \text{if } d \geq 4. \quad (4.4.19)$$

Combining (4.4.18–4.4.19), we see that we have completed the proof of Theorem 1.3.4 for $d \geq 4$.

In order to prove Conjecture 1.4.1, we would have to extend Proposition 4.4.3 to $d = 3$ and show that it contributes the second term in the r.h.s. of (4.4.16) rather than being negligible.

4.5 Preparatory facts and notation

In order to estimate $I_1^{r, q}(\kappa, T)$ and $I_2^{r, q}(\kappa, T)$, we need a number of preparatory facts. These are listed in Lemmas 4.5.1–4.5.4 below.

It follows from (4.4.3) that

$$\psi(\eta, b) - \psi(\eta, a) = \int_0^T ds \sum_{z \in \mathbb{Z}^d} (p_{2ds1[\kappa]}(z, b) - p_{2ds1[\kappa]}(z, a)) [\eta(z) - \rho] \quad (4.5.1)$$

and

$$\begin{aligned} \psi(\eta^{a, b}, x) - \psi(\eta, x) &= \int_0^T ds \sum_{z \in \mathbb{Z}^d} p_{2ds1[\kappa]}(z, x) [\eta^{a, b}(z) - \eta(z)] \\ &= \int_0^T ds (p_{2ds1[\kappa]}(b, x) - p_{2ds1[\kappa]}(a, x)) [\eta(a) - \eta(b)], \end{aligned} \quad (4.5.2)$$

where we recall the definitions of $1[\kappa]$ and $\eta^{a, b}$ in (4.1.1) and (1.2.5), respectively. We need bounds on both these differences.

Lemma 4.5.1 For any $\eta \in \Omega$, $a, b, x \in \mathbb{Z}^d$ and $\kappa, T > 0$,

$$|\psi(\eta, b) - \psi(\eta, a)| \leq 2T, \quad (4.5.3)$$

$$\left| \psi(\eta^{a,b}, x) - \psi(\eta, x) \right| \leq 2G_d < \infty, \quad (4.5.4)$$

and

$$\sum_{\{a,b\}} \left(\psi(\eta^{a,b}, x) - \psi(\eta, x) \right)^2 \leq \frac{1}{2d} G_d < \infty, \quad (4.5.5)$$

where G_d is the Green function at the origin of simple random walk.

Proof. The bound in (4.5.3) is immediate from (4.5.1). By (4.5.2), we have

$$\left| \psi(\eta^{a,b}, x) - \psi(\eta, x) \right| \leq \int_0^T ds |p_{2ds1[\kappa]}(b, x) - p_{2ds1[\kappa]}(a, x)|. \quad (4.5.6)$$

Using the bound $p_t(x, y) \leq p_t(0, 0)$ (which is immediate from the Fourier representation of the transition kernel), we get

$$\left| \psi(\eta^{a,b}, x) - \psi(\eta, x) \right| \leq 2 \int_0^\infty ds p_{2ds1[\kappa]}(0, 0) \leq 2G_d. \quad (4.5.7)$$

Again by (4.5.2), we have

$$\begin{aligned} \sum_{\{a,b\}} \left(\psi(\eta^{a,b}, x) - \psi(\eta, x) \right)^2 &= \sum_{\{a,b\}} [\eta(a) - \eta(b)]^2 \left(\int_0^T ds \left(p_{2ds1[\kappa]}(b, x) - p_{2ds1[\kappa]}(a, x) \right) \right)^2 \\ &\leq 2 \int_0^T du \int_u^T dv \sum_{\{a,b\}} \left(p_{2du1[\kappa]}(b, x) - p_{2du1[\kappa]}(a, x) \right) \left(p_{2dv1[\kappa]}(b, x) - p_{2dv1[\kappa]}(a, x) \right) \\ &= -2 \int_0^T du \int_u^T dv \sum_{a \in \mathbb{Z}^d} p_{2du1[\kappa]}(a, x) \left[\Delta_1 p_{2dv1[\kappa]}(a, x) \right] \\ &= -\frac{2}{1[\kappa]} \int_0^T du \int_u^T dv \sum_{a \in \mathbb{Z}^d} p_{2du1[\kappa]}(a, x) \left[\frac{\partial}{\partial v} p_{2dv1[\kappa]}(a, x) \right] \\ &= -\frac{2}{1[\kappa]} \int_0^T du \sum_{a \in \mathbb{Z}^d} p_{2du1[\kappa]}(a, x) (p_{2dT1[\kappa]}(a, x) - p_{2du1[\kappa]}(a, x)) \\ &\leq \frac{2}{1[\kappa]} \int_0^T du \sum_{a \in \mathbb{Z}^d} p_{2du1[\kappa]}^2(a, x) \\ &\leq \frac{2}{1[\kappa]} \int_0^\infty du p_{4du1[\kappa]}(0, 0) = \frac{1}{2d(1[\kappa])^2} G_d(0) \leq \frac{1}{2d} G_d, \end{aligned} \quad (4.5.8)$$

where Δ_1 denotes the discrete Laplacian acting on the first coordinate, and in the fifth line we use that $(\partial/\partial t)p_t = (1/2d)\Delta_1 p_t$. \blacksquare

For $x \in \mathbb{Z}^d$, let $\tau_x: \Omega \rightarrow \Omega$ be the x -shift on Ω defined by

$$(\tau_x \eta)(z) = \eta(z + x), \quad \eta \in \Omega, z \in \mathbb{Z}^d. \quad (4.5.9)$$

Lemma 4.5.2 For any bounded measurable $W: \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$,

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\int_0^t ds W(\xi_{\frac{s}{\kappa}}, X_s) \right] \right) \\ & \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\int_0^t ds W(\xi_{\frac{s}{\kappa}}, 0) \right] \right), \end{aligned} \quad (4.5.10)$$

provided

$$W(\eta, x) = W(\tau_x \eta, 0) \quad \forall \eta \in \Omega, x \in \mathbb{Z}^d. \quad (4.5.11)$$

Proof. The proof uses arguments similar to those in Sections 2.2 and 3.2.6. Recall (4.3.1). Proposition 2.2.2 with $p = 1$, applied to the self-adjoint operator $G_W^{\kappa} = \frac{1}{\kappa}L + \Delta + W$ (instead of G_V^{κ} in (2.2.4–2.2.5)), gives

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\int_0^t ds W(Z_s) \right] \right) = \sup_{\|f\|_{L^2(\nu_{\rho} \otimes m)}=1} \left(B_1(f) - \frac{1}{\kappa} B_2(f) - B_3(f) \right) \quad (4.5.12)$$

with

$$\begin{aligned} B_1(f) &= \int_{\Omega} \nu_{\rho}(d\eta) \sum_{z \in \mathbb{Z}^d} W(\eta, z) f(\eta, z)^2, \\ B_2(f) &= \int_{\Omega} \nu_{\rho}(d\eta) \sum_{z \in \mathbb{Z}^d} \frac{1}{2} \sum_{\{x,y\} \subset \mathbb{Z}^d} p(x,y) [f(\eta^{x,y}, z) - f(\eta, z)]^2, \\ B_3(f) &= \int_{\Omega} \nu_{\rho}(d\eta) \sum_{z \in \mathbb{Z}^d} \frac{1}{2} \sum_{\substack{y \in \mathbb{Z}^d \\ \|y-z\|=1}} [f(\eta, y) - f(\eta, z)]^2. \end{aligned} \quad (4.5.13)$$

An upper bound is obtained by dropping $B_3(f)$, i.e., the part associated with the simple random walk X . After that, split the supremum into two parts,

$$\begin{aligned} & \sup_{\|f\|_{L^2(\nu_{\rho} \otimes m)}=1} \left(B_1(f) - B_2(f) \right) \\ &= \sup_{\|g\|_{L^2(m)}=1} \sup_{\|f_z\|_{L^2(\nu_{\rho})}=1 \forall z \in \mathbb{Z}^d} \sum_{z \in \mathbb{Z}^d} g(z)^2 \int_{\Omega} \nu_{\rho}(d\eta) \\ & \quad \times \left(W(\eta, z) f_z(\eta)^2 - \frac{1}{2} \sum_{\{x,y\} \subset \mathbb{Z}^d} p(x,y) [f_z(\eta^{x,y}) - f_z(\eta)]^2 \right), \end{aligned} \quad (4.5.14)$$

where $f_z(\eta) = f(\eta, z)/g(z)$ with $g(z)^2 = \int_{\Omega} \nu_{\rho}(d\eta) f(\eta, z)^2$. The second supremum in (4.5.14), which runs over a family of functions indexed by z , can be brought under the sum. This gives

$$\begin{aligned} \text{r.h.s. (4.5.14)} &= \sup_{\|g\|_{L^2(m)}=1} \sum_{z \in \mathbb{Z}^d} g(z)^2 \sup_{\|f_z\|_{L^2(\nu_{\rho})}=1} \int_{\Omega} \nu_{\rho}(d\eta) \\ & \quad \times \left(W(\eta, z) f_z(\eta)^2 - \frac{1}{2} \sum_{\{x,y\} \subset \mathbb{Z}^d} p(x,y) [f_z(\eta^{x,y}) - f_z(\eta)]^2 \right). \end{aligned} \quad (4.5.15)$$

By (4.5.11) and the shift-invariance of ν_ρ , we may replace z by 0 under the second supremum in (4.5.15), in which case the latter no longer depends on z , and we get

$$\begin{aligned} \text{r.h.s. (4.5.15)} &= \sup_{\|f\|_{L^2(\nu_\rho)}=1} \int_{\Omega} \nu_\rho(d\eta) \left[W(\eta, 0) f(\eta)^2 - \frac{1}{2} \sum_{\{x,y\} \subset \mathbb{Z}^d} p(x,y) [f(\eta^{x,y}) - f(\eta)]^2 \right] \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\int_0^t ds W\left(\xi_{\frac{s}{\kappa}}, 0\right) \right] \right), \end{aligned} \quad (4.5.16)$$

where the second equality comes from the analogue of Proposition 2.2.2 with self-adjoint operator $\frac{1}{\kappa}L + W(\cdot, 0)$ (instead of G_V^κ). \blacksquare

Lemma 4.5.3 *For any $\rho \in (0, 1)$,*

$$\max_{\beta \in [0,1]} \left[\gamma\beta - \frac{1}{2G_d} (\sqrt{\beta} - \sqrt{\rho})^2 \right] = \frac{\rho\gamma}{1 - 2G_d\gamma} \sim \rho\gamma \quad \text{as } \gamma \downarrow 0. \quad (4.5.17)$$

Proof. A straightforward computation shows that the maximum in (4.5.17) is attained at

$$\beta = \frac{\rho}{(1 - 2G_d\gamma)^2} \in [0, 1] \quad (4.5.18)$$

for small enough γ . Substitution yields the claim. \blacksquare

Lemma 4.5.4 *There exists $C > 0$ such that, for all $t \geq 0$ and $x, y \in \mathbb{Z}^d$,*

$$p_t(x, y) \leq \frac{C}{(1+t)^{\frac{d}{2}}}. \quad (4.5.19)$$

Proof. This is a standard fact. Indeed, we can decompose the transition kernel of simple random walk with step rate 1 as

$$p_{dt}(x, y) = \prod_{j=1}^d p_t^{(1)}(x^j, y^j), \quad x = (x^1, \dots, x^d), \quad y = (y^1, \dots, y^d), \quad (4.5.20)$$

where $p_t^{(1)}(x, y)$ is the transition kernel of 1-dimensional simple random walk with step rate 1. In Fourier representation,

$$p_t^{(1)}(x, y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} dk e^{ik \cdot (y-x)} e^{-t\hat{\varphi}(k)}, \quad \hat{\varphi}(k) = 1 - \cos k. \quad (4.5.21)$$

The bound in (4.5.19) follows from (4.5.20) and

$$p_t^{(1)}(x, y) \leq p_t^{(1)}(0, 0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} dk e^{-t\hat{\varphi}(k)} \leq \frac{C}{(1+t)^{\frac{1}{2}}}, \quad t \geq 0, \quad x, y \in \mathbb{Z}^d. \quad (4.5.22)$$

\blacksquare

4.6 Proof of Proposition 4.4.2

The proof of Proposition 4.4.2 is given in Section 4.6.1 subject to four lemmas. The latter will be proved in Sections 4.6.2–4.6.5, respectively. All results are valid for $d \geq 3$.

4.6.1 Proof of Proposition 4.4.2

Lemma 4.6.1 *Uniformly in $\eta \in \Omega$ and $x \in \mathbb{Z}^d$, as $\kappa \rightarrow \infty$,*

$$\left[\left(e^{-\frac{r}{\kappa}\psi} \mathcal{A} e^{\frac{r}{\kappa}\psi} \right) - \mathcal{A} \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) = \frac{r^2}{2\kappa^2} \sum_{e: \|e\|=1} \left(\psi(\eta, x+e) - \psi(\eta, x) \right)^2 + O\left(\left(\frac{1}{\kappa} \right)^3 \right). \quad (4.6.1)$$

Lemma 4.6.2 *For any $\kappa, T > 0$, $\alpha \in \mathbb{R}$ and $r > 0$,*

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{\alpha r}{2\kappa^2} \int_0^t ds \sum_{e: \|e\|=1} \left(\psi \left(\xi_{\frac{s}{\kappa}}, X_s + e \right) - \psi \left(\xi_{\frac{s}{\kappa}}, X_s \right) \right)^2 \right] \right) \\ & \leq \limsup_{t \rightarrow \infty} \frac{1}{2t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{z \in \mathbb{Z}^d} K_{\text{diag}}^{\kappa, T}(z) \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right)^2 \right] \right) \\ & \quad + \limsup_{t \rightarrow \infty} \frac{1}{2t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right), \end{aligned} \quad (4.6.2)$$

where

$$\begin{aligned} K_{\text{diag}}^{\kappa, T}(z) &= \sum_{e: \|e\|=1} \left(\chi(z+e) - \chi(z) \right)^2, \\ K_{\text{off}}^{\kappa, T}(z_1, z_2) &= \sum_{e: \|e\|=1} \left(\chi(z_1+e) - \chi(z_1) \right) \left(\chi(z_2+e) - \chi(z_2) \right), \quad z_1 \neq z_2, \end{aligned} \quad (4.6.3)$$

with

$$\chi(z) = \int_0^T du p_{2du1[\kappa]}(0, z). \quad (4.6.4)$$

Lemma 4.6.3 *For any $\alpha \in \mathbb{R}$ and $r > 0$,*

$$\limsup_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{z \in \mathbb{Z}^d} K_{\text{diag}}^{\kappa, T}(z) \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right)^2 \right] \right) \leq \alpha r \rho (1 - \rho) \frac{1}{d} G_d. \quad (4.6.5)$$

Lemma 4.6.4 *For any $\alpha \in \mathbb{R}$ and $r > 0$,*

$$\limsup_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \leq 0. \quad (4.6.6)$$

Combining Lemmas 4.6.1–4.6.4, we obtain the claim in Proposition 4.4.2.

4.6.2 Proof of Lemma 4.6.1

Lemma 4.6.1 is immediate from (4.3.2) and the following two lemmas.

Lemma 4.6.5 *Uniformly in $\eta \in \Omega$ and $x \in \mathbb{Z}^d$, as $\kappa \rightarrow \infty$,*

$$\frac{1}{\kappa} \left[\left(e^{-\frac{r}{\kappa}\psi} L e^{\frac{r}{\kappa}\psi} \right) - L \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) = O \left(\frac{1}{\kappa^3} \right). \quad (4.6.7)$$

Lemma 4.6.6 *Uniformly in $\eta \in \Omega$ and $x \in \mathbb{Z}^d$, as $\kappa \rightarrow \infty$,*

$$\left[\left(e^{-\frac{r}{\kappa}\psi} \Delta e^{\frac{r}{\kappa}\psi} \right) - \Delta \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) = \frac{r^2}{2\kappa^2} \sum_{e: \|e\|=1} \left(\psi(\eta, x+e) - \psi(\eta, x) \right)^2 + O \left(\frac{1}{\kappa^3} \right). \quad (4.6.8)$$

Proof of Lemma 4.6.5. By (1.2.3–1.2.4), we have

$$\begin{aligned} & \left[\left(e^{-\frac{r}{\kappa}\psi} L e^{\frac{r}{\kappa}\psi} \right) - L \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) \\ &= \frac{1}{2d} \sum_{\{a,b\}} \left(e^{\frac{r}{\kappa}[\psi(\eta^{a,b},x) - \psi(\eta,x)]} - 1 - \frac{r}{\kappa}[\psi(\eta^{a,b},x) - \psi(\eta,x)] \right). \end{aligned} \quad (4.6.9)$$

Taylor expansion of the r.h.s. of (4.6.9) gives that uniformly in $\eta \in \Omega$ and $x \in \mathbb{Z}^d$,

$$\frac{1}{\kappa} \left[\left(e^{-\frac{r}{\kappa}\psi} L e^{\frac{r}{\kappa}\psi} \right) - L \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) = \frac{r^2}{4d\kappa^3} \sum_{\{a,b\}} \left(\psi(\eta^{a,b},x) - \psi(\eta,x) \right)^2 e^{o(1)} = O \left(\frac{1}{\kappa^3} \right), \quad (4.6.10)$$

where we use (4.5.4–4.5.5). ■

Proof of Lemma 4.6.6. By (1.1.2), we have

$$\begin{aligned} & \left[\left(e^{-\frac{r}{\kappa}\psi} \Delta e^{\frac{r}{\kappa}\psi} \right) - \Delta \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) \\ &= \sum_{e: \|e\|=1} \left(e^{\frac{r}{\kappa}[\psi(\eta,x+e) - \psi(\eta,x)]} - 1 - \frac{r}{\kappa}[\psi(\eta,x+e) - \psi(\eta,x)] \right). \end{aligned} \quad (4.6.11)$$

Taylor expansion of the r.h.s. of (4.6.11) gives that uniformly in $\eta \in \Omega$ and $x \in \mathbb{Z}^d$,

$$\left[\left(e^{-\frac{r}{\kappa}\psi} \Delta e^{\frac{r}{\kappa}\psi} \right) - \Delta \left(\frac{r}{\kappa}\psi \right) \right] (\eta, x) = \frac{r^2}{2\kappa^2} \sum_{e: \|e\|=1} \left(\psi(\eta, x+e) - \psi(\eta, x) \right)^2 + R_{\kappa,T}(\eta, x) \quad (4.6.12)$$

with

$$|R_{\kappa,T}(\eta, x)| \leq \frac{r^3}{6\kappa^3} \sum_{e: \|e\|=1} \left| \psi(\eta, x+e) - \psi(\eta, x) \right|^3 e^{o(1)} \leq \frac{8dr^3}{3\kappa^3} T^3 e^{o(1)}, \quad (4.6.13)$$

where we use (4.5.3). Combining (4.6.12–4.6.13), we arrive at (4.6.8). ■

4.6.3 Proof of Lemmas 4.6.2

Proof. By (4.4.3), we have for all $\eta \in \Omega$ and $x \in \mathbb{Z}^d$,

$$\begin{aligned}
& \sum_{e: \|e\|=1} \left(\psi(\eta, x + e) - \psi(\eta, x) \right)^2 \\
&= \sum_{e: \|e\|=1} \sum_{z_1, z_2 \in \mathbb{Z}^d} \int_0^T du \int_0^T dv \left(p_{2du1[\kappa]}(z_1, x + e) - p_{2du1[\kappa]}(z_1, x) \right) \\
&\quad \times \left(p_{2dv1[\kappa]}(z_2, x + e) - p_{2dv1[\kappa]}(z_2, x) \right) [\eta(z_1) - \rho] [\eta(z_2) - \rho] \\
&= \sum_{z_1, z_2 \in \mathbb{Z}^d} K^{\kappa, T}(z_1, z_2) [\eta(z_1 + x) - \rho] [\eta(z_2 + x) - \rho],
\end{aligned} \tag{4.6.14}$$

where $K^{\kappa, T}: \mathbb{Z}^d \times \mathbb{Z}^d \mapsto \mathbb{R}$ is given by

$$K^{\kappa, T}(z_1, z_2) = \sum_{e: \|e\|=1} \left(\chi(z_1 + e) - \chi(z_1) \right) \left(\chi(z_2 + e) - \chi(z_2) \right). \tag{4.6.15}$$

Therefore, for all $\kappa, T > 0$,

$$\begin{aligned}
& \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho, 0}} \left(\exp \left[\frac{\alpha r}{2\kappa^2} \int_0^t ds \sum_{e: \|e\|=1} \left(\psi\left(\xi_{\frac{s}{\kappa}}, X_s + e\right) - \psi\left(\xi_{\frac{s}{\kappa}}, X_s\right) \right)^2 \right] \right) \\
&= \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho, 0}} \left(\exp \left[\frac{\alpha r}{2\kappa^2} \int_0^t ds \sum_{z_1, z_2 \in \mathbb{Z}^d} K^{\kappa, T}(z_1, z_2) \right. \right. \\
&\quad \left. \left. \times \left(\xi_{\frac{s}{\kappa}}(z_1 + X_s) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2 + X_s) - \rho \right) \right] \right) \\
&\leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho}} \left(\exp \left[\frac{\alpha r}{2\kappa^2} \int_0^t ds \sum_{z_1, z_2 \in \mathbb{Z}^d} K^{\kappa, T}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right),
\end{aligned} \tag{4.6.16}$$

where in the last line we use Lemma 4.5.2 with

$$W(\eta, x) = \frac{\alpha r}{2\kappa^2} \sum_{z_1, z_2 \in \mathbb{Z}^d} K^{\kappa, T}(z_1, z_2) [\eta(z_1 + x) - \rho] [\eta(z_2 + x) - \rho], \tag{4.6.17}$$

which satisfies $W(\eta, x) = W(\tau_x \eta, 0)$ as required in (4.5.11). Splitting the sum in the r.h.s. of (4.6.16) into its diagonal and off-diagonal part and using the Cauchy-Schwarz inequality, we arrive at (4.6.2). \blacksquare

4.6.4 Proof of Lemma 4.6.3

The proof of Lemma 4.6.3 is based on the following two lemmas. Recall (2.1.1).

Lemma 4.6.7 *For any $T > 0$ there exists $C_T > 0$, satisfying $\lim_{T \rightarrow \infty} C_T = 0$, such that*

$$\lim_{\kappa \rightarrow \infty} \|K_{\text{diag}}^{\kappa, T}\|_1 = \frac{1}{d} G_d + C_T. \tag{4.6.18}$$

Lemma 4.6.8 For any $T > 0$, $\alpha \in \mathbb{R}$ and $r > 0$,

$$\limsup_{t, \kappa \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 T_{t/\kappa} \right] \right) \leq \alpha r \rho (1 - 2\rho) \lim_{\kappa \rightarrow \infty} \|K_{\text{diag}}^{\kappa, T}\|_1. \quad (4.6.19)$$

Before giving the proofs of Lemmas 4.6.7–4.6.8, we first prove Lemma 4.6.3.

Proof of Lemma 4.6.3. By Jensen's inequality, we have

$$\begin{aligned} & \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{z \in \mathbb{Z}^d} K_{\text{diag}}^{\kappa, T}(z) \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right)^2 \right] \right) \\ & \leq \sum_{z \in \mathbb{Z}^d} \frac{K_{\text{diag}}^{\kappa, T}(z)}{\|K_{\text{diag}}^{\kappa, T}\|_1} \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa^2} \|K_{\text{diag}}^{\kappa, T}\|_1 \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right)^2 \right] \right) \\ & = \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa^2} \|K_{\text{diag}}^{\kappa, T}\|_1 \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(0) - \rho \right)^2 \right] \right) \\ & = \exp \left[\frac{\alpha r}{\kappa^2} \rho^2 \|K_{\text{diag}}^{\kappa, T}\|_1 t \right] \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 \int_0^{t/\kappa} ds \xi_s(0) \right] \right), \end{aligned} \quad (4.6.20)$$

where the first equality uses the shift-invariance of ν_ρ . Therefore

$$\begin{aligned} & \lim_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{z \in \mathbb{Z}^d} K_{\text{diag}}^{\kappa, T}(z) \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right)^2 \right] \right) \\ & \leq \lim_{\kappa, T \rightarrow \infty} \alpha r \rho^2 \|K_{\text{diag}}^{\kappa, T}\|_1 + \lim_{t, \kappa, T \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 T_{\frac{t}{\kappa}} \right] \right). \end{aligned} \quad (4.6.21)$$

Now use Lemmas 4.6.7–4.6.8 to obtain (4.6.5). ■

Proof of Lemma 4.6.7. By (4.6.3), we have

$$\begin{aligned} \|K_{\text{diag}}^{\kappa, T}\|_1 & = 2 \sum_{\{x, y\}} \int_0^T du \int_u^T dv \left(p_{2du1[\kappa]}(0, y) - p_{2du1[\kappa]}(0, x) \right) \left(p_{2dv1[\kappa]}(0, y) - p_{2dv1[\kappa]}(0, x) \right) \\ & = -4 \int_0^T du \int_u^T dv \sum_{x \in \mathbb{Z}^d} p_{2du1[\kappa]}(0, x) \left[\Delta_1 p_{2dv1[\kappa]}(0, x) \right] \\ & = -\frac{4}{1[\kappa]} \int_0^T du \int_u^T dv \sum_{x \in \mathbb{Z}^d} p_{2du1[\kappa]}(0, x) \left[\frac{\partial}{\partial v} p_{2dv1[\kappa]}(0, x) \right], \end{aligned} \quad (4.6.22)$$

where we recall the remark below (4.5.8). After performing the integration w.r.t. the variable v , we get

$$\begin{aligned} \|K_{\text{diag}}^{\kappa, T}\|_1 & = \frac{4}{1[\kappa]} \left(\int_0^T du \sum_{x \in \mathbb{Z}^d} p_{2du1[\kappa]}^2(0, x) - \int_0^T du \sum_{x \in \mathbb{Z}^d} p_{2du1[\kappa]}(0, x) p_{2dT1[\kappa]}(0, x) \right) \\ & = \frac{4}{1[\kappa]} \left(\int_0^T du p_{4du1[\kappa]}(0, 0) - \int_0^T du p_{2d(u+T)1[\kappa]}(0, 0) \right). \end{aligned} \quad (4.6.23)$$

Hence

$$\lim_{\kappa \rightarrow \infty} \|K_{\text{diag}}^{\kappa, T}\|_1 = \frac{1}{d} \left(\int_0^{2dT} du p_u(0, 0) - \int_{2dT}^{4dT} du p_u(0, 0) \right), \quad (4.6.24)$$

which gives (4.6.18). \blacksquare

Proof of Lemma 4.6.8. To derive (4.6.19), we use the large deviation principle for $(T_t)_{t \geq 0}$ stated in Section 2.1. By Varadhan's Lemma we have, for all $\kappa, T > 0$,

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 T_{t/\kappa} \right] \right) \\ = \frac{1}{\kappa} \max_{\beta \in [0, 1]} \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 \beta - \Psi_d(\beta) \right] \\ \leq \frac{1}{\kappa} \max_{\beta \in [0, 1]} \left[\frac{\alpha r}{\kappa} (1 - 2\rho) \|K_{\text{diag}}^{\kappa, T}\|_1 \beta - \frac{1}{2G_d} (\sqrt{\beta} - \sqrt{\rho})^2 \right], \end{aligned} \quad (4.6.25)$$

where we use the quadratic lower bound in (2.1.6). By Lemma 4.6.7, $(1/\kappa) \|K_{\text{diag}}^{\kappa, T}\|_1 \downarrow 0$ as $\kappa \rightarrow \infty$ for any $T > 0$. Hence Lemma 4.5.3 can be applied to get (4.6.19). \blacksquare

4.6.5 Proof of Lemma 4.6.4

The proof of Lemma 4.6.4 is based on the following two lemmas. Recall (4.6.3). For $z_1, z_2 \in \mathbb{Z}^d$ with $z_1 \neq z_2$ and $\gamma \in \mathbb{R}$, let

$$\begin{aligned} h_{\gamma, \kappa}(z_1, z_2) &= \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\gamma}{\kappa^2} \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \\ &= \limsup_{t \rightarrow \infty} \frac{1}{\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\gamma}{\kappa} \int_0^t ds \left(\xi_s(z_1) - \rho \right) \left(\xi_s(z_2) - \rho \right) \right] \right). \end{aligned} \quad (4.6.26)$$

Lemma 4.6.9 For all $\kappa, T > 0$,

$$\|K_{\text{off}}^{\kappa, T}\|_1 \leq 8dT^2. \quad (4.6.27)$$

Lemma 4.6.10 For any $z_1, z_2 \in \mathbb{Z}^d$ with $z_1 \neq z_2$ and any $\gamma \in \mathbb{R}$,

$$\limsup_{\kappa \rightarrow \infty} \kappa^2 h_{\gamma, \kappa}(z_1, z_2) \leq 0. \quad (4.6.28)$$

Before giving the proof of Lemmas 4.6.9–4.6.10, we first prove Lemma 4.6.4.

Proof of Lemma 4.6.4. Let $K_{\text{off}}^{\kappa, T; +}$ and $K_{\text{off}}^{\kappa, T; -}$ denote, respectively, the positive and negative part of $K_{\text{off}}^{\kappa, T}$. By the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \\ \leq \frac{1}{2} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{2\alpha r}{\kappa^2} \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \\ + \frac{1}{2} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[-\frac{2\alpha r}{\kappa^2} \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; -}(z_1, z_2) \int_0^t ds \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right). \end{aligned} \quad (4.6.29)$$

We estimate the first term in the r.h.s. of (4.6.29). For $R > 0$, let

$$B_R = \{(z_1, z_2) \in \mathbb{Z}^d \times \mathbb{Z}^d : \|z_1\| + \|z_2\| \leq R\}. \quad (4.6.30)$$

Then

$$\begin{aligned} & \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{2\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \\ & \leq \exp \left[\frac{2|\alpha|r t}{\kappa^2} \sum_{\substack{(z_1, z_2) \in B_R^c \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) \right] \\ & \quad \times \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{2\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{(z_1, z_2) \in B_R \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right). \end{aligned} \quad (4.6.31)$$

Applying Jensen's inequality, we get

$$\begin{aligned} & \frac{\kappa^2}{t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{2\alpha r}{\kappa^2} \int_0^t ds \sum_{\substack{z_1, z_2 \in \mathbb{Z}^d \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) \left(\xi_{\frac{s}{\kappa}}(z_1) - \rho \right) \left(\xi_{\frac{s}{\kappa}}(z_2) - \rho \right) \right] \right) \\ & \leq 2|\alpha|r \sum_{\substack{(z_1, z_2) \in B_R^c \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2) + \frac{\kappa}{t/\kappa} \log \sum_{\substack{(z_1, z_2) \in B_R \\ z_1 \neq z_2}} \frac{K_{\text{off}}^{\kappa, T; +}(z_1, z_2)}{\|K_{\text{off}; R}^{\kappa, T; +}\|_1} \\ & \quad \times \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{2\alpha r}{\kappa} \|K_{\text{off}; R}^{\kappa, T; +}\|_1 \int_0^{t/\kappa} ds \left(\xi_s(z_1) - \rho \right) \left(\xi_s(z_2) - \rho \right) \right] \right), \end{aligned} \quad (4.6.32)$$

where

$$\|K_{\text{off}; R}^{\kappa, T; +}\|_1 = \sum_{\substack{(z_1, z_2) \in B_R \\ z_1 \neq z_2}} K_{\text{off}}^{\kappa, T; +}(z_1, z_2). \quad (4.6.33)$$

By Lemma 4.6.10 (with $\gamma = 2\alpha r \|K_{\text{off}; R}^{\kappa, T; +}\|_1$), the second term in the r.h.s. of (4.6.32) is asymptotically bounded by above by zero (as $t \rightarrow \infty$) for any $\kappa, T > 0$, $\alpha \in \mathbb{R}$ and $r > 0$, and any R finite. The first term in the r.h.s. of (4.6.32) does not depend on t and, by Lemma 4.6.9, tends to zero as $R \rightarrow \infty$. This shows that the first term in the r.h.s of (4.6.29) yields a zero contribution. The same is true for the second term by the same argument. This completes the proof of (4.6.6). \blacksquare

Proof of Lemma 4.6.9. The claim follows from (4.6.3–4.6.4). \blacksquare

Proof of Lemma 4.6.10. The proof of Lemma 4.6.10 is long, since it is based on three further lemmas. Let $z_1, z_2 \in \mathbb{Z}^d$ with $z_1 \neq z_2$. Without loss of generality, we may assume that

$$z_1 \in H^- \quad \text{and} \quad z_2 \in H^+ \quad (4.6.34)$$

with

$$H^- = \{z \in \mathbb{Z}^d : z^1 \leq 0\} \quad \text{and} \quad H^+ = \{z \in \mathbb{Z}^d : z^1 > 0\}. \quad (4.6.35)$$

Let

$$\begin{aligned}
h_{\gamma,\kappa}^-(z_1) &= \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \tilde{\xi}_s^-(z_1) \right] \right), \\
h_{\gamma,\kappa}^+(z_2) &= \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \tilde{\xi}_s^+(z_2) \right] \right), \\
h_{\gamma,\kappa}^\pm(z_1, z_2) &= \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds \tilde{\xi}_s^-(z_1) \tilde{\xi}_s^+(z_2) \right] \right),
\end{aligned} \tag{4.6.36}$$

where $(\tilde{\xi}_t^-)_{t \geq 0}$ and $(\tilde{\xi}_t^+)_{t \geq 0}$ are independent IRW's on H^- and H^+ , respectively, with transition kernels $p^-(\cdot, \cdot)$ and $p^+(\cdot, \cdot)$ corresponding to simple random walks stepping at rate 1 such that steps outside H^- and H^+ , respectively, are suppressed.

Lemma 4.6.11 For all $\kappa > 0$, $z_1 \in H^-$, $z_2 \in H^+$ and $\gamma \in \mathbb{R}$,

$$h_{\gamma,\kappa}(z_1, z_2) \leq \frac{\gamma}{\kappa^2} \rho^2 + h_{\gamma,\kappa}^-(z_1) + h_{\gamma,\kappa}^+(z_2) + h_{\gamma,\kappa}^\pm(z_1, z_2). \tag{4.6.37}$$

Lemma 4.6.12 For all $\gamma \in \mathbb{R}$,

$$\limsup_{\kappa \rightarrow \infty} \kappa^2 \sup_{z_1 \in H^-} h_{\gamma,\kappa}^-(z_1) \leq -\gamma \rho^2 \quad \text{and} \quad \limsup_{\kappa \rightarrow \infty} \kappa^2 \sup_{z_2 \in H^+} h_{\gamma,\kappa}^+(z_2) \leq -\gamma \rho^2. \tag{4.6.38}$$

Lemma 4.6.13 For all $\gamma \in \mathbb{R}$,

$$\limsup_{\kappa \rightarrow \infty} \kappa^2 \sup_{\substack{z_1 \in H^- \\ z_2 \in H^+}} h_{\gamma,\kappa}^\pm(z_1, z_2) \leq \gamma \rho^2. \tag{4.6.39}$$

Combining (4.6.37–4.6.39), we get (4.6.28). ■

Proof of Lemma 4.6.11. Similarly as in the proof of Lemma 4.5.2, by cutting the bonds connecting H^- and H^+ in the analogue of the variational formula of Proposition 2.2.2, we get

$$\begin{aligned}
h_{\gamma,\kappa}(z_1, z_2) &\leq \limsup_{t \rightarrow \infty} \frac{1}{\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\gamma}{\kappa} \int_0^t ds (\xi_s^-(z_1) - \rho)(\xi_s^+(z_2) - \rho) \right] \right) \\
&= \limsup_{t \rightarrow \infty} \frac{1}{\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{\gamma}{\kappa} \int_0^t ds \left(\rho^2 - \rho \xi_s^-(z_1) + \rho \xi_s^+(z_2) + \xi_s^-(z_1) \xi_s^+(z_2) \right) \right] \right),
\end{aligned} \tag{4.6.40}$$

where $(\xi_t^-)_{t \geq 0}$ and $(\xi_t^+)_{t \geq 0}$ are independent exclusion processes in H^- and H^+ , respectively, obtained from $(\xi_t)_{t \geq 0}$ by suppressing jumps between H^- and H^+ . Applying Hölder's inequality in the r.h.s. of (4.6.40) to separate terms, we obtain

$$\begin{aligned}
h_{\gamma,\kappa}(z_1, z_2) &\leq \frac{\gamma}{\kappa^2} \rho^2 + \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \xi_s^-(z_1) \right] \right) \\
&\quad + \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \xi_s^+(z_2) \right] \right) \\
&\quad + \limsup_{t \rightarrow \infty} \frac{1}{3\kappa t} \log \mathbb{E}_{\nu_\rho} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds \xi_s^-(z_1) \xi_s^+(z_2) \right] \right).
\end{aligned} \tag{4.6.41}$$

In order to get (4.6.37), we apply Proposition 1.2.1 to the last three terms in the r.h.s. of (4.6.41). For the first two terms, pick, respectively $K(z, s) = -(3\gamma/\kappa)\rho 1_{z_1}$ and $K(z, s) = -(3\gamma/\kappa)\rho 1_{z_2}(z)$. For the last term, we have to apply Proposition 1.2.1 twice, once for the exclusion process $(\xi_t^+)_{t \geq 0}$ on H^+ with $K(z, s) = -(3\gamma/\kappa)\xi_s^-(z_1) 1_{z_2}(z)$ and once for the exclusion process $(\xi_t^-)_{t \geq 0}$ on H^- with $K(z, s) = -(3\gamma/\kappa)\xi_s^-(z_2) 1_{z_1}(z)$. Here, we in fact apply a *modification* of Proposition 1.2.1 by considering $(\xi_t^-)_{t \geq 0}$ and $(\xi_t^+)_{t \geq 0}$ on \mathbb{Z}^d with particles not moving on H^+ and H^- , respectively. See the proof of Proposition 1.2.1 in Appendix A to verify that this modification holds true. \blacksquare

Proof of Lemma 4.6.12. We prove the second line of (4.6.38). The first line follows by symmetry.

Let

$$H_\eta^+ = \{x \in H^+ : \eta(x) = 1\}, \quad \eta \in \Omega. \quad (4.6.42)$$

Fix $z \in H^+$. Then

$$\mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \tilde{\xi}_s^+(z) \right] \right) = \int_\Omega \nu_\rho(d\eta) \prod_{x \in H_\eta^+} \mathbb{E}_x^{\text{RW},+} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds 1_z(Y_s^+) \right] \right), \quad (4.6.43)$$

where $\mathbb{E}_x^{\text{RW},+}$ is expectation w.r.t. simple random walk $Y^+ = (Y_t^+)_{t \geq 0}$ on H^+ with transition kernel $p^+(\cdot, \cdot)$ and step rate 1 starting from $Y_0^+ = x \in H^+$. Using that ν_ρ is the Bernoulli product measure with density ρ , we get

$$\begin{aligned} & \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds \tilde{\xi}_s^+(z) \right] \right) \\ &= \int_\Omega \nu_\rho(d\eta) \prod_{x \in H^+} \mathbb{E}_x^{\text{RW},+} \left(\exp \left[-\eta(x) \frac{3\gamma}{\kappa} \rho \int_0^t ds 1_z(Y_s^+) \right] \right) \\ &= \prod_{x \in H^+} \left(1 - \rho + \rho v(x, t) \right) \leq \exp \left[\rho \sum_{x \in H^+} (v(x, t) - 1) \right] \end{aligned} \quad (4.6.44)$$

with

$$v(x, t) = \mathbb{E}_x^{\text{RW},+} \left(\exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds 1_z(Y_s^+) \right] \right). \quad (4.6.45)$$

By the Feynman-Kac formula, $v: H^+ \times [0, \infty) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial}{\partial t} v(x, t) = \frac{1}{2d} \Delta^+ v(x, t) - \left\{ \frac{3\gamma}{\kappa} \rho 1_z(x) \right\} v(x, t), \quad v(\cdot, 0) \equiv 1, \quad (4.6.46)$$

where

$$\Delta^+ v(x, t) = \sum_{\substack{y \in H^+ \\ \|y-x\|=1}} [v(y, t) - v(x, t)], \quad x \in H^+. \quad (4.6.47)$$

Put

$$w(x, t) = v(x, t) - 1. \quad (4.6.48)$$

Then $w: H^+ \times [0, \infty) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial w}{\partial t}(x, t) = \frac{1}{2d} \Delta^+ w(x, t) - \left\{ \frac{3\gamma}{\kappa} \rho 1_z(x) \right\} [w(x, t) + 1], \quad w(\cdot, 0) \equiv 0. \quad (4.6.49)$$

Since $\sum_{x \in H^+} \Delta^+ f(x) = 0$ for all $f: H^+ \rightarrow \mathbb{R}$, (4.6.49) gives

$$\frac{\partial}{\partial t} \sum_{x \in H^+} w(x, t) = -\frac{3\gamma}{\kappa} \rho [w(z, t) + 1]. \quad (4.6.50)$$

After integrating (4.6.50) w.r.t. t , we obtain

$$\sum_{x \in H^+} w(x, t) = -\frac{3\gamma}{\kappa} \rho t - \frac{3\gamma}{\kappa} \rho \int_0^t ds w(z, s). \quad (4.6.51)$$

Combining (4.6.36), (4.6.44), (4.6.48) and (4.6.51), we arrive at

$$h_{\gamma, \kappa}^+(z) \leq -\frac{\gamma}{\kappa^2} \rho^2 \left(1 + \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t ds w(z, s) \right). \quad (4.6.52)$$

The limit in the r.h.s. exists since, by (4.6.45) and (4.6.48), $w(z, t)$ is monotone in t .

We will complete the proof by showing that the second term in the r.h.s. of (4.6.52) tends to zero as $\kappa \rightarrow \infty$. This will rely on the following lemma, the proof of which is deferred to the end of this section.

Lemma 4.6.14 *Let $G^+(x, y)$ be the Green kernel on H^+ associated with $p_t^+(x, y)$. Then $\|G^+\|_\infty \leq 2G_d < \infty$.*

Return to (4.6.45). If $\gamma > 0$, then by Jensen's inequality we have

$$1 \geq v(x, t) \geq \exp \left[-\frac{3\gamma}{\kappa} \rho \int_0^t ds p_s^+(x, z) \right] \geq \exp \left[-\frac{3\gamma}{\kappa} \rho \|G^+\|_\infty \right], \quad (4.6.53)$$

where $\|G^+\|_\infty < \infty$ by Lemma 4.6.14. To deal with the case $\gamma \leq 0$, let \mathcal{G}^+ denote the Green operator acting on functions $V: H^+ \rightarrow [0, \infty)$ as

$$(\mathcal{G}^+ V)(x) = \sum_{y \in H^+} G^+(x, y) V(y), \quad x \in H^+. \quad (4.6.54)$$

We have

$$\left\| \mathcal{G}^+ \left(\frac{3\gamma}{\kappa} \rho 1_z \right) \right\|_\infty \leq \frac{3|\gamma|}{\kappa} \rho \|G^+\|_\infty. \quad (4.6.55)$$

The r.h.s. tends to zero as $\kappa \rightarrow \infty$. Hence Lemma 8.2.1 in Gärtner and den Hollander [4] can be applied to (4.6.45) for κ large enough, to yield

$$1 \leq v(x, t) \leq \frac{1}{1 - \frac{3|\gamma|}{\kappa} \rho \|G^+\|_\infty} \downarrow 1 \quad \text{as } \kappa \rightarrow \infty. \quad (4.6.56)$$

Therefore, combining (4.6.53) and (4.6.56), we see that for all $\gamma \in \mathbb{R}$ and $\delta \in (0, 1)$ there exists $\kappa_0 = \kappa_0(\gamma, \delta)$ such that

$$\|v - 1\|_\infty \leq \delta \quad \forall \kappa > \kappa_0. \quad (4.6.57)$$

By (4.6.48–4.6.49), we have

$$w(z, t) = -\frac{3\gamma}{\kappa} \rho \int_0^t ds \mathbb{E}_z^{\text{RW}, +} \left(1_z(Y_s^+) v(Y_s^+, t - s) \right). \quad (4.6.58)$$

Via (4.6.57) it therefore follows that

$$-\frac{3\gamma}{\kappa}\rho(1 \pm \delta)G^+(z, z) \leq \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t ds w(z, s) \leq -\frac{3\gamma}{\kappa}\rho(1 \mp \delta)G^+(z, z) \quad \forall \kappa > \kappa_0, \quad (4.6.59)$$

where the choice of + or - in front of δ depends on the sign of γ . The latter shows that the second term in the r.h.s. of (4.6.52) is $O(1/\kappa)$. This proves (4.6.38). \blacksquare

Proof of Lemma 4.6.13. The proof is similar to that of Lemma 4.6.12. Let

$$\begin{aligned} H_\eta^+ &= \{x \in H^+ : \eta(x) = 1\}, \quad \eta \in \Omega, \\ H_\eta^- &= \{x \in H^- : \eta(x) = 1\}, \quad \eta \in \Omega. \end{aligned} \quad (4.6.60)$$

Fix $z_1 \in H^-$ and $z_2 \in H^+$. Then

$$\begin{aligned} &\mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds \tilde{\xi}_s^-(z_1) \tilde{\xi}_s^+(z_2) \right] \right) \\ &= \int_\Omega \nu_\rho(d\eta) \prod_{x \in H_\eta^-} \prod_{y \in H_\eta^+} \mathbb{E}_x^{\text{RW}, -} \mathbb{E}_y^{\text{RW}, +} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds 1_{(z_1, z_2)}(Y_s^-, Y_s^+) \right] \right), \end{aligned} \quad (4.6.61)$$

where Y^- on H^- and Y^+ on H^+ are simple random walks with step rate 1 and transition kernel $p^-(\cdot, \cdot)$ and $p^+(\cdot, \cdot)$ starting from $Y_0^- = x \in H^-$ and $Y_0^+ = y \in H^+$, respectively. Using that ν_ρ is the Bernoulli product measure with density ρ , we get

$$\begin{aligned} &\mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds \tilde{\xi}_s^-(z_1) \tilde{\xi}_s^+(z_2) \right] \right) \\ &= \int_\Omega \nu_\rho(d\eta) \prod_{x \in H^-} \prod_{y \in H^+} \mathbb{E}_x^{\text{RW}, -} \mathbb{E}_y^{\text{RW}, +} \left(\exp \left[\eta(x)\eta(y) \frac{3\gamma}{\kappa} \int_0^t ds 1_{(z_1, z_2)}(Y_s^-, Y_s^+) \right] \right) \\ &= \prod_{x \in H^-} \prod_{y \in H^+} \left(1 - \rho^2 + \rho^2 v(z_1, z_2; t) \right) \leq \exp \left[\rho^2 \sum_{x \in H^-} \sum_{y \in H^+} (v(z_1, z_2; t) - 1) \right] \end{aligned} \quad (4.6.62)$$

with

$$v(z_1, z_2; t) = \mathbb{E}_x^{\text{RW}, -} \mathbb{E}_y^{\text{RW}, +} \left(\exp \left[\frac{3\gamma}{\kappa} \int_0^t ds \int_0^t ds 1_{(z_1, z_2)}(Y_s^-, Y_s^+) \right] \right). \quad (4.6.63)$$

By the Feynman-Kac formula, $v: (H^- \times H^+) \times [0, \infty) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial}{\partial t} v(x, y; t) = \frac{1}{2d} (\Delta^- + \Delta^+) v(x, y; t) + \left\{ \frac{3\gamma}{\kappa} 1_{z_1, z_2}(x, y) \right\} v(x, y; t), \quad v(\cdot, \cdot; 0) \equiv 1, \quad (4.6.64)$$

where

$$\begin{aligned} \Delta^- v(x; t) &= \sum_{\substack{y \in H^- \\ \|y-x\|=1}} [v(y, t) - v(x, t)], \quad x \in H^-, \\ \Delta^+ v(x; t) &= \sum_{\substack{y \in H^+ \\ \|y-x\|=1}} [v(y, t) - v(x, t)], \quad x \in H^+. \end{aligned} \quad (4.6.65)$$

Put

$$w(x, y; t) = v(x, y; t) - 1. \quad (4.6.66)$$

Then, $w: (H^- \times H^+) \times [0, \infty) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial w}{\partial t}(x, y; t) = \frac{1}{2d}(\Delta^- + \Delta^+)w(x, y; t) + \left\{ \frac{3\gamma}{\kappa} 1_{(z_1, z_2)}(x, y) \right\} [w(x, y; t) + 1], \quad w(\cdot, \cdot; 0) \equiv 0. \quad (4.6.67)$$

By (4.6.65) and (4.6.67),

$$\frac{\partial}{\partial t} \sum_{x \in H^-} \sum_{y \in H^+} w(x, y; t) = \frac{3\gamma}{\kappa} [w(z_1, z_2; t) + 1]. \quad (4.6.68)$$

After integrating (4.6.68) w.r.t. t , we obtain

$$\sum_{x \in H^-} \sum_{y \in H^+} w(x, y; t) = \frac{3\gamma}{\kappa} t + \frac{3\gamma}{\kappa} \int_0^t ds w(z_1, z_2; s). \quad (4.6.69)$$

Combining (4.6.36), (4.6.62), (4.6.66) and (4.6.69), we arrive at

$$h_{\gamma, \kappa}^{\pm}(z_1, z_2) \leq \frac{\gamma}{\kappa^2} \rho^2 \left(1 + \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t ds w(z_1, z_2; s) \right). \quad (4.6.70)$$

The limit in the r.h.s. exists, since $w(z_1, z_2; s)$ is monotone in s .

We will complete the proof by showing that the second term in the r.h.s. of (4.6.70) tends to zero as $\kappa \rightarrow \infty$. Return to (4.6.63). If $\gamma \leq 0$, then by Jensen's inequality we have

$$1 \geq v(x, y; t) \geq \exp \left[-\frac{3|\gamma|}{\kappa} \int_0^\infty ds p_s^-(x, z_1) p_s^+(y, z_2) \right] \geq \exp \left[-\frac{3|\gamma|}{\kappa} \left(\|G^-\|_\infty \wedge \|G^+\|_\infty \right) \right], \quad (4.6.71)$$

where $\|G^-\|_\infty, \|G^+\|_\infty < \infty$ by Lemma 4.6.14. To deal with the case $\gamma > 0$, let \mathcal{G}^\pm denote the Green operator acting on functions $V: H^- \times H^+ \rightarrow [0, \infty)$ as

$$(\mathcal{G}^\pm V)(x, y) = \sum_{\substack{a \in H^- \\ b \in H^+}} G^\pm(x, y; a, b) V(a, b), \quad x \in H^-, y \in H^+, \quad (4.6.72)$$

where

$$G^\pm(x, y; a, b) = \int_0^\infty ds p_s^-(x, a) p_s^+(y, b). \quad (4.6.73)$$

We have

$$\left\| \mathcal{G}^\pm \left(\frac{3\gamma}{\kappa} 1_{(z_1, z_2)} \right) \right\|_\infty \leq \frac{3\gamma}{\kappa} \|G^\pm\|_\infty \leq \frac{3\gamma}{\kappa} \left(\|G^-\|_\infty \wedge \|G^+\|_\infty \right). \quad (4.6.74)$$

The r.h.s. tends to zero as $\kappa \rightarrow \infty$. Hence Lemma 8.2.1 in Gärtner and den Hollander [4] can be applied to (4.6.63) for κ large enough, to yield

$$1 \leq v(x, t) \leq \frac{1}{1 - \frac{3\gamma}{\kappa} \left(\|G^-\|_\infty \wedge \|G^+\|_\infty \right)} \downarrow 1 \quad \text{as } \kappa \rightarrow \infty. \quad (4.6.75)$$

Therefore, combining (4.6.71) and (4.6.75), we see that for all $\gamma \in \mathbb{R}$ and $\delta > 0$ there exists $\kappa_0 = \kappa_0(\gamma, \delta)$ such that

$$\|v - 1\|_\infty \leq \delta \quad \forall \kappa > \kappa_0. \quad (4.6.76)$$

By (4.6.66–4.6.67), we have

$$w(z_1, z_2; t) = \frac{3\gamma}{\kappa} \int_0^t ds \mathbb{E}_{z_1}^{\text{RW}, -} \mathbb{E}_{z_2}^{\text{RW}, +} \left(1_{(z_1, z_2)}(Y_s^-, Y_s^+) v(Y_s^-, Y_s^+; t-s) \right). \quad (4.6.77)$$

Via (4.6.76) it therefore follows that for all $\kappa > \kappa_0$,

$$\frac{3\gamma}{\kappa} (1 \pm \delta) G^\pm(z_1, z_1; z_2, z_2) \leq \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t ds w(z_1, z_2; s) \leq \frac{3\gamma}{\kappa} (1 \mp \delta) G^\pm(z_1, z_1; z_2, z_2). \quad (4.6.78)$$

Combining (4.6.70) and (4.6.78), we arrive at (4.6.39). \blacksquare

Proof of Lemma 4.6.14. We have $G^+(x, y) = \sum_{n=0}^{\infty} p_n^+(x, y)$, $x, y \in H^+$, with $p_n^+(x, y)$ the n -step transition probability of simple random walk on H^+ whose steps outside H^+ are suppressed (i.e., the walk pauses when it attempts to leave H^+). Let $p_n(x, y)$ be the n -step transition probability of simple random walk on \mathbb{Z}^d . Then

$$p_n^+(x, y) \leq 2p_n(x, y), \quad x, y \in H^+, n \in \mathbb{N}_0. \quad (4.6.79)$$

Indeed, if we reflect simple random walk in the $(d-1)$ -dimensional hyperplane between H^+ and its complement, then we obtain precisely the random walk that pauses when it attempts to leave H^+ . Hence, we have $p_n^+(x, y) = p_n(x, y) + p_n(x, y^*)$, $x, y \in H^+$, $n \in \mathbb{N}_0$, with y^* the reflection image of y . Since $p_n(x, y^*) \leq p_n(x, y)$, $x, y \in H^+$, the claim in (4.6.79) follows. Sum on n , to get $G^+(x, y) \leq 2G(x, y)$, $x, y \in H^+$. Now use that $G(x, y) \leq G(0, 0) = G_d$, $x, y \in \mathbb{Z}^d$. \blacksquare

4.7 Proof of Proposition 4.4.3

The proof of Proposition 4.4.3 is given in Section 4.7.1 subject to three lemmas. The latter are proved in Sections 4.7.2–4.7.4, respectively. The first two lemmas are valid for $d \geq 3$, the third for $d \geq 4$.

4.7.1 Proof of Proposition 4.4.3

Lemma 4.7.1 *For all $t \geq 0$, $\kappa, T > 0$ and $\alpha \in \mathbb{R}$,*

$$\mathbb{E}_{\nu_{\rho, 0}} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds (\mathcal{P}_T \phi)(Z_s) \right] \right) \leq \mathbb{E}_0 \left(\exp \left[\frac{\alpha}{\kappa} \rho \int_0^t ds \sum_{x \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(X_{t-s}, x) w^{(t)}(x, s) \right] \right), \quad (4.7.1)$$

where $w^{(t)}: \mathbb{Z}^d \times [0, t) \rightarrow \mathbb{R}$ is the solution of the Cauchy problem

$$\frac{\partial w^{(t)}}{\partial s}(x, s) = \frac{1}{2d\kappa} \Delta w^{(t)}(x, s) + \frac{\alpha}{\kappa} p_{2dT_1[\kappa]}(X_{t-s}, x) [w^{(t)}(x, s) + 1], \quad w^{(t)}(\cdot, 0) \equiv 0. \quad (4.7.2)$$

Lemma 4.7.2 *For all $t \geq 0$, $\kappa > 0$, T large enough and $\alpha \in \mathbb{R}$,*

$$\begin{aligned} & \mathbb{E}_0 \left(\exp \left[\frac{\alpha}{\kappa} \rho \int_0^t ds \sum_{x \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(X_{t-s}, x) w^{(t)}(x, s) \right] \right) \\ & \leq \mathbb{E}_0 \left(\exp \left[\frac{2\alpha^2}{\kappa^2} \rho \int_0^t ds \int_s^t du p_{\frac{u-s}{\kappa} + 4dT_1[\kappa]}(X_u, X_s) \right] \right). \end{aligned} \quad (4.7.3)$$

Lemma 4.7.3 *If $d \geq 4$, then for any $\alpha \in \mathbb{R}$,*

$$\lim_{T, \kappa, t \rightarrow \infty} \frac{\kappa^2}{t} \log \mathbb{E}_0 \left(\exp \left[\frac{2\alpha^2}{\kappa^2} \rho \int_0^t ds \int_s^t du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \right] \right) = 0. \quad (4.7.4)$$

Lemmas 4.7.1–4.7.3 clearly imply (4.4.16).

4.7.2 Proof of Lemma 4.7.1

For all $t \geq 0$, $\kappa, T > 0$ and $\alpha \in \mathbb{R}$, let $v^{(t)}: \mathbb{Z}^d \times [0, t] \rightarrow \mathbb{R}$ be such that

$$v^{(t)}(x, s) = w^{(t)}(x, s) + 1, \quad (4.7.5)$$

where $w^{(t)}$ is defined by (4.7.2). Then $v^{(t)}$ is the solution of the Cauchy problem

$$\frac{\partial v^{(t)}}{\partial s}(x, s) = \frac{1}{2d\kappa} \Delta v^{(t)}(x, s) + \frac{\alpha}{\kappa} p_{2dT1[\kappa]}(X_{t-s}, x) v^{(t)}(x, s), \quad v^{(t)}(\cdot, 0) \equiv 1, \quad (4.7.6)$$

and has the representation

$$v^{(t)}(x, s) = \mathbb{E}_x^{\text{RW}} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^s du p_{2dT1[\kappa]} \left(X_{t-s+u}, Y_{\frac{u}{\kappa}} \right) \right] \right). \quad (4.7.7)$$

Proof. By (4.3.1) and (4.4.5), we have

$$\mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds (\mathcal{P}_T \phi)(Z_s) \right] \right) = \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\frac{\alpha}{\kappa} \sum_{z \in \mathbb{Z}^d} \int_0^t ds p_{2dT1[\kappa]}(X_s, z) \left(\xi_{\frac{s}{\kappa}}(z) - \rho \right) \right] \right). \quad (4.7.8)$$

Therefore, by Proposition 1.2.1 (with $K(z, s) = \alpha p_{2dT1[\kappa]}(X_{\kappa s}, z)$), we get

$$\begin{aligned} & \mathbb{E}_{\nu_\rho, 0} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds (\mathcal{P}_T \phi)(Z_s) \right] \right) \\ & \leq \mathbb{E}_0 \mathbb{E}_{\nu_\rho}^{\text{IRW}} \left(\exp \left[\frac{\alpha}{\kappa} \sum_{z \in \mathbb{Z}^d} p_{2dT1[\kappa]}(X_s, z) \int_0^t ds \left(\tilde{\xi}_{\frac{s}{\kappa}}(z) - \rho \right) \right] \right) \\ & \leq \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \mathbb{E}_0 \int_\Omega \nu_\rho(d\eta) \prod_{x \in A_\eta} \mathbb{E}_x^{\text{RW}} \left(\exp \left[\frac{\alpha}{\kappa} \sum_{z \in \mathbb{Z}^d} p_{2dT1[\kappa]}(X_s, z) \int_0^t ds \delta_z \left(Y_{\frac{s}{\kappa}} \right) \right] \right) \\ & = \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \mathbb{E}_0 \int_\Omega \nu_\rho(d\eta) \prod_{x \in A_\eta} \mathbb{E}_x^{\text{RW}} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds p_{2dT1[\kappa]} \left(X_s, Y_{\frac{s}{\kappa}} \right) \right] \right), \end{aligned} \quad (4.7.9)$$

where $A_\eta = \{x \in \mathbb{Z}^d : \eta(x) = 1\}$ and \mathbb{E}_x^{RW} is expectation w.r.t. to simple random walk $Y = (Y_t)_{t \geq 0}$ on \mathbb{Z}^d with step rate 1 starting from $Y_0 = x$. Using that ν_ρ is the Bernoulli

product measure with density ρ , we get

$$\begin{aligned}
& \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{\alpha}{\kappa} \int_0^t ds (\mathcal{P}_T \phi)(Z_s) \right] \right) \\
& \leq \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \int_{\Omega} \nu_{\rho}(d\eta) \mathbb{E}_0 \left(\prod_{x \in \mathbb{Z}^d} \mathbb{E}_x^{\text{RW}} \left(\exp \left[\eta(x) \frac{\alpha}{\kappa} \int_0^t ds p_{2dT_1[\kappa]}(X_s, Y_{\frac{s}{\kappa}}) \right] \right) \right) \\
& = \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \int_{\Omega} \nu_{\rho}(d\eta) \mathbb{E}_0 \left(\prod_{x \in \mathbb{Z}^d} \left[1 + \eta(x) w^{(t)}(x, t) \right] \right) \\
& = \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \mathbb{E}_0 \left(\prod_{x \in \mathbb{Z}^d} \left[1 + \rho w^{(t)}(x, t) \right] \right) \leq \exp \left[-\frac{\alpha}{\kappa} \rho t \right] \mathbb{E}_0 \left(\exp \left[\rho \sum_{x \in \mathbb{Z}^d} w^{(t)}(x, t) \right] \right),
\end{aligned} \tag{4.7.10}$$

where $w^{(t)}: \mathbb{Z}^d \times [0, t] \rightarrow \mathbb{R}$ solves (4.7.2). From (4.7.2) we deduce that

$$\frac{\partial}{\partial s} \sum_{x \in \mathbb{Z}^d} w^{(t)}(x, s) = \frac{\alpha}{\kappa} \sum_{x \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(X_{t-s}, x) [1 + w^{(t)}(x, s)]. \tag{4.7.11}$$

Integrating (4.7.11) w.r.t. s and inserting the result into (4.7.10), we get (4.7.1). ■

4.7.3 Proof of Lemma 4.7.2

Next, we consider $v^{(t)}$ and $w^{(t)}$ as defined in (4.7.5–4.7.7), but with $|\alpha|$ instead of α .

Proof. We begin by showing that, for T large enough and all x, s, t and $X_{(\cdot)}$, we have $v^{(t)}(x, s) \leq 2$.

Do a Taylor expansion, to obtain ($s_0 = 0$)

$$v^{(t)}(x, s) = \sum_{n=0}^{\infty} \left(\frac{|\alpha|}{\kappa} \right)^n \left(\prod_{l=1}^n \int_{s_{l-1}}^s ds_l \right) \mathbb{E}_x^{\text{RW}} \left(\prod_{m=1}^n p_{2dT_1[\kappa]}(X_{t-s+s_m}, Y_{\frac{s_m}{\kappa}}) \right). \tag{4.7.12}$$

In Fourier representation the transition kernel of simple random walk with step rate 1 reads

$$p_s(x, y) = \oint dk e^{-ik \cdot (y-x)} e^{-s\widehat{\varphi}(k)}, \tag{4.7.13}$$

where $\oint dk = (2\pi)^{-d} \int_{[-\pi, \pi]^d} dk$ and

$$\widehat{\varphi}(k) = \frac{1}{2d} \sum_{\substack{x \in \mathbb{Z}^d \\ \|x\|=1}} (1 - e^{ik \cdot x}), \quad k \in [-\pi, \pi]^d. \tag{4.7.14}$$

Combining (4.7.12–4.7.13), we get

$$\begin{aligned}
v^{(t)}(x, s) &= \sum_{n=0}^{\infty} \left(\frac{|\alpha|}{\kappa} \right)^n \left(\prod_{l=1}^n \int_{s_{l-1}}^s ds_l \right) \left(\prod_{m=1}^n \int \phi dk_m \right) \\
&\quad \times \mathbb{E}_x^{\text{RW}} \left(\exp \left[i \sum_{p=1}^n \left(Y_{\frac{s_p}{\kappa}} - X_{t-s+s_p} \right) \cdot k_p \right] \exp \left[- \left(2dT1[\kappa] \right) \sum_{q=1}^n \widehat{\varphi}(k_q) \right] \right) \\
&= \sum_{n=0}^{\infty} \left(\frac{|\alpha|}{\kappa} \right)^n \left(\prod_{l=1}^n \int_{s_{l-1}}^s ds_l \right) \left(\prod_{m=1}^n \int \phi dk_m \right) \exp \left[- i \sum_{p=1}^n \left(X_{t-s+s_p} - x \right) \cdot k_p \right] \\
&\quad \times \exp \left[- \left(2dT1[\kappa] \right) \sum_{q=1}^n \widehat{\varphi}(k_q) \right] \mathbb{E}_0^{\text{RW}} \left(\exp \left[i \sum_{r=1}^n Y_{\frac{s_r}{\kappa}} \cdot k_r \right] \right), \tag{4.7.15}
\end{aligned}$$

where in the last line we did a spatial shift of Y by x . Because Y has independent increments, we have

$$\begin{aligned}
\mathbb{E}_0^{\text{RW}} \left(\exp \left[i \sum_{r=1}^n Y_{\frac{s_r}{\kappa}} \cdot k_r \right] \right) &= \mathbb{E}_0^{\text{RW}} \left(\exp \left[i \sum_{r=1}^n (k_r + \dots + k_n) \cdot \left(Y_{\frac{s_r}{\kappa}} - Y_{\frac{s_{r-1}}{\kappa}} \right) \right] \right) \\
&= \prod_{r=1}^n \mathbb{E}_0^{\text{RW}} \left(\exp \left[i (k_r + \dots + k_n) \cdot Y_{\frac{s_r - s_{r-1}}{\kappa}} \right] \right) \\
&= \prod_{r=1}^n \sum_{z \in \mathbb{Z}^d} p_{\frac{s_r - s_{r-1}}{\kappa}}(0, z) \exp \left[i (k_r + \dots + k_n) \cdot z \right] \\
&= \prod_{r=1}^n \exp \left[- \frac{s_r - s_{r-1}}{\kappa} \widehat{\varphi}(k_r + \dots + k_n) \right], \tag{4.7.16}
\end{aligned}$$

where the last line uses (4.7.13). Since the r.h.s. is non-negative, taking the modulus of the r.h.s. of (4.7.15), we obtain

$$\begin{aligned}
v^{(t)}(x, s) &\leq \sum_{n=0}^{\infty} \left(\frac{|\alpha|}{\kappa} \right)^n \left(\prod_{l=1}^n \int_{s_{l-1}}^s ds_l \right) \left(\prod_{m=1}^n \int \phi dk_m \right) \\
&\quad \times \exp \left[- \left(2dT1[\kappa] \right) \sum_{q=1}^n \widehat{\varphi}(k_q) \right] \mathbb{E}_0^{\text{RW}} \left(\exp \left[i \sum_{r=1}^n Y_{\frac{s_r}{\kappa}} \cdot k_r \right] \right) \\
&= \sum_{n=0}^{\infty} \left(\frac{|\alpha|}{\kappa} \right)^n \left(\prod_{l=1}^n \int_{s_{l-1}}^s ds_l \right) \mathbb{E}_0^{\text{RW}} \left(\prod_{m=1}^n p_{2dT1[\kappa]} \left(0, Y_{\frac{s_m}{\kappa}} \right) \right), \tag{4.7.17}
\end{aligned}$$

where the last line uses (4.7.13). Thus

$$v^{(t)}(x, s) \leq \mathbb{E}_0^{\text{RW}} \left(\exp \left[\frac{|\alpha|}{\kappa} \int_0^s du p_{2dT1[\kappa]} \left(0, Y_{\frac{u}{\kappa}} \right) \right] \right) \leq \mathbb{E}_0^{\text{RW}} \left(\exp \left[|\alpha| \int_0^\infty du p_{2dT1[\kappa]} \left(0, Y_u \right) \right] \right). \tag{4.7.18}$$

Next, let \mathcal{G} denote the Green operator acting on functions $V: \mathbb{Z}^d \rightarrow [0, \infty)$ as

$$(\mathcal{G}V)(x) = \sum_{y \in \mathbb{Z}^d} G(x, y)V(y), \quad x \in \mathbb{Z}^d. \tag{4.7.19}$$

With p_t denoting the function $p_t(0, \cdot)$, we have

$$\left\| \mathcal{G}(|\alpha| p_{2dT_1[\kappa]}) \right\|_\infty = |\alpha| \sup_{x \in \mathbb{Z}^d} \int_0^\infty ds \sum_{y \in \mathbb{Z}^d} p_s(x, y) p_{2dT_1[\kappa]}(0, y) \leq |\alpha| G_{2dT_1[\kappa]} \quad (4.7.20)$$

with

$$G_t = \int_t^\infty ds p_s(0, 0) \quad (4.7.21)$$

the truncated Green function at the origin. The r.h.s. of (4.7.20) tends to zero as $T \rightarrow \infty$. Hence Lemma 8.2.1 in Gärtner and den Hollander [4] can be applied to the r.h.s. of (4.7.18) for T large enough, to yield

$$v^{(t)}(x, s) \leq \frac{1}{1 - \left\| \mathcal{G}(|\alpha| p_{2dT_1[\kappa]}) \right\|_\infty} \downarrow 1 \quad \text{as } T \rightarrow \infty, \quad \text{uniformly in } \kappa > 0. \quad (4.7.22)$$

Thus, for T large enough and all x, s, t, κ and $X_{(\cdot)}$, we have $v^{(t)}(x, s) \leq 2$, as claimed earlier. For such T , recalling (4.7.5), we conclude from (4.7.2) that $w^{(t)} \leq \bar{w}^{(t)}$, where $\bar{w}^{(t)}$ solves

$$\frac{\partial \bar{w}^{(t)}}{\partial s}(x, s) = \frac{1}{2d\kappa} \Delta \bar{w}^{(t)}(x, s) + \frac{2|\alpha|}{\kappa} p_{2dT_1[\kappa]}(X_{t-s}, x), \quad \bar{w}^{(t)}(\cdot, 0) \equiv 0, \quad (4.7.23)$$

The latter has the representation

$$\bar{w}^{(t)}(x, s) = \frac{2|\alpha|}{\kappa} \int_0^s du \sum_{z \in \mathbb{Z}^d} p_{\frac{s-u}{\kappa}}(x, z) p_{2dT_1[\kappa]}(X_{t-u}, z) = \frac{2|\alpha|}{\kappa} \int_0^s du p_{\frac{s-u}{\kappa} + 2dT_1[\kappa]}(x, X_{t-u}). \quad (4.7.24)$$

Hence,

$$\begin{aligned} & \mathbb{E}_0 \left(\exp \left[\frac{\alpha}{\kappa} \rho \int_0^t ds \sum_{x \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(X_{t-s}, x) w^{(t)}(x, s) \right] \right) \\ & \leq \mathbb{E}_0 \left(\exp \left[\frac{|\alpha|}{\kappa} \rho \int_0^t ds \sum_{x \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(X_{t-s}, x) \bar{w}^{(t)}(x, s) \right] \right) \\ & = \mathbb{E}_0 \left(\exp \left[\frac{2\alpha^2}{\kappa^2} \rho \int_0^t ds \int_0^s du p_{\frac{u-s}{\kappa} + 4dT_1[\kappa]}(X_{t-s}, X_{t-u}) \right] \right), \end{aligned} \quad (4.7.25)$$

which proves the claim in (4.7.3). ■

4.7.4 Proof of Lemma 4.7.3

The proof of Lemma 4.7.3 is based on the following lemma. For $t \geq 0$, $\alpha \in \mathbb{R}$ and $a, \kappa, T > 0$, let

$$\Lambda_\alpha(t; a, \kappa, T) = \frac{1}{2t} \log \mathbb{E}_0 \left(\exp \left[\frac{4\alpha^2}{\kappa^2} \rho \int_0^t ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT_1[\kappa]}(X_u, X_s) \right] \right) \quad (4.7.26)$$

and

$$\lambda_\alpha(a, \kappa, T) = \limsup_{t \rightarrow \infty} \Lambda_\alpha(t; a, \kappa, T). \quad (4.7.27)$$

Lemma 4.7.4 *If $d \geq 4$, then for any $\alpha \in \mathbb{R}$ and $a, T > 0$,*

$$\limsup_{\kappa \rightarrow \infty} \kappa^2 \lambda_\alpha(a, \kappa, T) \leq 2\alpha^2 \rho G_{4dT}, \quad (4.7.28)$$

where G_t is the truncated Green function at the origin defined by (4.7.21). Before giving the proof of Lemma 4.7.4, we first prove Lemma 4.7.3.

Proof of Lemma 4.7.3. Return to (4.7.4). By the Cauchy-Schwarz inequality, we have

$$\begin{aligned} & \frac{\kappa^2}{t} \log \mathbb{E}_0 \left(\exp \left[\frac{2\alpha^2}{\kappa^2} \rho \int_0^t ds \int_s^t du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \right] \right) \\ & \leq \frac{\kappa^2}{2t} \log \mathbb{E}_0 \left(\exp \left[\frac{4\alpha^2}{\kappa^2} \rho \int_0^t ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \right] \right) \\ & \quad + \frac{\kappa^2}{2t} \log \mathbb{E}_0 \left(\exp \left[\frac{4\alpha^2}{\kappa^2} \rho \int_0^t ds \int_{s+a\kappa^3}^\infty du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \right] \right). \end{aligned} \quad (4.7.29)$$

Moreover, by Lemma 4.5.4 and the fact that $d \geq 3$, we have

$$\begin{aligned} \frac{1}{\kappa^2} \int_0^t ds \int_{s+a\kappa^3}^\infty du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) & \leq \frac{1}{\kappa^2} \int_0^t ds \int_{s+a\kappa^3}^\infty du p_{\frac{u-s}{\kappa}}(0, 0) \\ & \leq \frac{C}{\kappa} t \int_{a\kappa^2}^\infty du \frac{1}{(1+u)^{\frac{d}{2}}} \leq \frac{\tilde{C}}{a^{\frac{1}{2}} \kappa^2} t \end{aligned} \quad (4.7.30)$$

with $C, \tilde{C} > 0$. Combining (4.7.29–4.7.30) and Lemma 4.7.4, and letting $a \rightarrow \infty$, we get (4.7.4). \blacksquare

The proof of Lemma 4.7.4 is based on one further lemma. For $\gamma \geq 0$ and $a, \kappa, T > 0$, let

$$\Lambda_\gamma(a, T) = \limsup_{\kappa \rightarrow \infty} \frac{1}{a\kappa} \log \mathbb{E}_0 \left(\frac{\gamma}{\kappa^2} \int_0^{a\kappa^3} ds \int_s^\infty du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_s, X_u) \right). \quad (4.7.31)$$

Lemma 4.7.5 *If $d \geq 4$, then for any $\gamma \geq 0$ and $a, T > 0$,*

$$\Lambda_\gamma(a, T) \leq \gamma G_{4dT}. \quad (4.7.32)$$

Before giving the proof of Lemma 4.7.5, we first prove Lemma 4.7.4.

Proof of Lemma 4.7.4. Split the integral in the exponent in the r.h.s. of (4.7.26) as follows:

$$\begin{aligned} & \int_0^t ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \\ & \leq \left(\sum_{\substack{k=1 \\ \text{even}}}^{\lceil t/a\kappa^3 \rceil} + \sum_{\substack{k=1 \\ \text{odd}}}^{\lceil t/a\kappa^3 \rceil} \right) \int_{(k-1)a\kappa^3}^{ka\kappa^3} ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s). \end{aligned} \quad (4.7.33)$$

Note that in each of the two sums, the summands are i.i.d. Hence, substituting (4.7.33) into (4.7.26) and applying the Cauchy-Schwarz inequality, we get

$$\Lambda_\alpha(t; a, \kappa, T) \leq \frac{\lceil t/a\kappa^3 \rceil}{4t} \log \mathbb{E}_0 \left(\exp \left[\frac{8\alpha^2}{\kappa^2} \rho \int_0^{a\kappa^3} ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT1[\kappa]}(X_u, X_s) \right] \right). \quad (4.7.34)$$

Letting $t \rightarrow \infty$ and recalling (4.7.27), we arrive at

$$\lambda_\alpha(a, \kappa, T) \leq \frac{1}{4a\kappa^3} \log E_0 \left(\exp \left[\frac{8\alpha^2}{\kappa^2} \rho \int_0^{a\kappa^3} ds \int_s^{s+a\kappa^3} du p_{\frac{u-s}{\kappa} + 4dT_1[\kappa]}(X_u, X_s) \right] \right). \quad (4.7.35)$$

Combining this with Lemma 4.7.5 (with $\gamma = 8\alpha^2\rho$), we obtain (4.7.28). \blacksquare

The proof of Lemma 4.7.5 is based on two further lemmas.

Lemma 4.7.6 *For any $\beta > 0$ and $M \in \mathbb{N}$,*

$$\begin{aligned} & E_0 \left(\exp \left[\beta \sum_{k=1}^M \int_0^\infty ds p_{\frac{s}{\kappa} + 4dT_1[\kappa]}(U_{k-1}(0), U_{k-1}(s)) \right] \right) \\ & \leq \prod_{k=1}^M \max_{y_1, \dots, y_{k-1} \in \mathbb{Z}^d} E_0 \left(\exp \left[\beta \sum_{l=0}^{k-1} \int_0^\infty ds p_{\frac{a\kappa^2}{M}l + \frac{s}{\kappa} + 4dT_1[\kappa]}(0, X_s + y_l) \right] \right), \end{aligned} \quad (4.7.36)$$

where $U_k(t) = X(\frac{k}{M}a\kappa^3 + s)$, $k \in \mathbb{N}_0$ and $y_0 = 0$.

Lemma 4.7.7 *For any $\beta > 0$, $M \in \mathbb{N}$, $k \in \mathbb{N}_0$, and $y_0, \dots, y_k \in \mathbb{Z}^d$,*

$$E_0 \left(\exp \left[\beta \sum_{l=0}^{k-1} \int_0^\infty ds p_{\frac{a\kappa^2}{M}l + \frac{s}{\kappa} + 4dT_1[\kappa]}(0, X_s + y_l) \right] \right) \leq \exp \left[\frac{\beta \sum_{l=0}^k G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]}}{1 - \beta \sum_{l=0}^k G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]}} \right], \quad (4.7.37)$$

(recall (4.7.21)), provided that

$$\beta \sum_{l=0}^k G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]} < 1. \quad (4.7.38)$$

The proofs of Lemmas 4.7.6–4.7.7 are similar to those of Lemmas 6.3.1–6.3.2 in Gärtner and den Hollander [4]. We refrain from spelling out the details. We conclude by proving Lemma 4.7.5.

Proof of Lemma 4.7.5. As in the proof of Lemma 6.2.1 in Gärtner and den Hollander [4], using Lemmas 4.7.6–4.7.7 we obtain

$$\begin{aligned} & \frac{1}{a\kappa} \log E_0 \left(\exp \left[\frac{\gamma}{\kappa^2} \int_0^{a\kappa^3} ds \int_s^\infty du p_{\frac{u-s}{\kappa} + 4dT_1[\kappa]}(X_s, X_u) \right] \right) \\ & \leq \frac{\gamma \sum_{l=0}^{M-1} G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]}}{1 - \gamma \frac{a\kappa}{M} \sum_{l=0}^{M-1} G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]}}, \end{aligned} \quad (4.7.39)$$

provided that

$$\gamma \frac{a\kappa}{M} \sum_{l=0}^{M-1} G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]} < 1. \quad (4.7.40)$$

But (recall (4.7.21))

$$\sum_{l=0}^{M-1} G_{\frac{a\kappa^2}{M}l + 4dT_1[\kappa]} \leq G_{4dT_1[\kappa]} + \sum_{l=1}^{M-1} G_{\frac{a\kappa^2}{M}l}. \quad (4.7.41)$$

From Lemma 4.5.4 we get $G_t \leq C/t^{\frac{d}{2}-1}$. Therefore

$$\frac{\kappa}{M} \sum_{l=1}^{M-1} G_{\frac{a\kappa^2}{M}l} \leq \begin{cases} \frac{C_3}{a^{\frac{1}{2}}} & \text{if } d = 3, \\ \frac{C_4}{a} \frac{1}{\kappa} \log M & \text{if } d = 4, \\ \frac{C_d}{a^{\frac{d}{2}-1}} \frac{M^{\frac{d}{2}-2}}{\kappa^{d-3}} & \text{if } d \geq 5, \end{cases} \quad (4.7.42)$$

for some $C_d > 0$, $d \geq 3$. Hence, picking $1 \ll M \leq C\kappa^2$, (4.7.40) holds for κ large enough when $d \geq 4$, and so the claim (4.7.32) follows from (4.7.39) and (4.7.41–4.7.42). \blacksquare

4.8 Extension to arbitrary p

In Sections 4.2–4.7 we proved Theorem 1.3.4 for $p = 1$. We briefly indicate how the proof can be extended to arbitrary p .

As in (4.2.2), after time rescaling we have, for any $p \in \mathbb{N}$,

$$\lambda_p^*(\kappa) = \lim_{t \rightarrow \infty} \Lambda_p^*(\kappa; t) \quad \text{with} \quad \Lambda_p^*(\kappa; t) = \frac{1}{t} \log \mathbb{E}_{\nu_{\rho, 0, \dots, 0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \sum_{k=1}^p \xi_{\frac{s}{\kappa}}(X_k(s)) \right] \right). \quad (4.8.1)$$

We are interested in the quantity

$$\lambda_p^*(\kappa) - \frac{\rho}{\kappa} = \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho, 0, \dots, 0}} \left(\exp \left[\frac{1}{\kappa} \int_0^t ds \sum_{k=1}^p \left(\xi_{\frac{s}{\kappa}}(X_k(s)) - \rho \right) \right] \right). \quad (4.8.2)$$

As in (4.4.1), for $T > 0$ let $\psi_p: \Omega \times (\mathbb{Z}^d)^p$ be defined by

$$\psi(\eta, x_1, \dots, x_p) = \int_0^T ds \left(\mathcal{P}_s^{(p)} \phi_p \right) (\eta, x_1, \dots, x_p) \quad \text{with} \quad \phi_p(\eta, x_1, \dots, x_p) = \sum_{k=1}^p [\eta(x_k) - \rho], \quad (4.8.3)$$

where $(\mathcal{P}_s^{(p)})_{s \geq 0}$ is the semigroup with generator (compare with (4.3.2))

$$\mathcal{A}^{(p)} = \frac{1}{\kappa} L + \sum_{k=1}^p \Delta_k. \quad (4.8.4)$$

Using (1.2.6), we obtain the representation (compare with (4.4.3))

$$\psi_p(\eta, x_1, \dots, x_p) = \int_0^T ds \sum_{z \in \mathbb{Z}^d} \sum_{k=1}^p p_{2ds1[\kappa]}(z, x_k) [\eta(z) - \rho] = \sum_{k=1}^p \psi(\eta, x_k). \quad (4.8.5)$$

Let (compare with (4.3.1))

$$Z_s^{(p)} = \left(\xi_{\frac{s}{\kappa}}, X_1(s), \dots, X_p(s) \right). \quad (4.8.6)$$

First, we have the analogue of Proposition 4.4.1:

Proposition 4.8.1 For any $p \in \mathbb{N}$, $\kappa, T > 0$,

$$\begin{aligned} \lambda_p^*(\kappa) - \frac{\rho}{\kappa} &\leq \frac{1}{2q} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{2q}{r} \int_0^t ds \left[\left(e^{-\frac{r}{\kappa} \psi_p} \mathcal{A} e^{\frac{r}{\kappa} \psi_p} \right) - \mathcal{A} \left(\frac{r}{\kappa} \psi_p \right) \right] (Z_s^{(p)}) \right] \right) \\ &\geq \frac{1}{4q} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_{\nu_{\rho,0}} \left(\exp \left[\frac{4q}{\kappa} \int_0^t ds \left(\mathcal{P}_T^{(p)} \phi_p \right) (Z_s^{(p)}) \right] \right), \end{aligned} \quad (4.8.7)$$

where $1/r + 1/q = 1$, for any $r, q > 1$ in the first inequality and any $q < 0 < r < 1$ in the second inequality.

Next, using (4.8.5), the bound

$$\left(\psi_p(\eta^{a,b}, x_1, \dots, x_p) - \psi_p(\eta, x_1, \dots, x_p) \right)^2 \leq p \sum_{k=1}^p \left(\psi(\eta^{a,b}, x_k) - \psi(\eta, x_k) \right)^2, \quad (4.8.8)$$

and the estimate in (4.5.4), we also have the analogue of Lemma 4.6.1:

Lemma 4.8.2 Uniformly in $\eta \in \Omega$ and $x_1, \dots, x_p \in \mathbb{Z}^d$,

$$\begin{aligned} &\left[\left(e^{-\frac{r}{\kappa} \psi_p} \mathcal{A} e^{\frac{r}{\kappa} \psi_p} \right) - \mathcal{A} \left(\frac{r}{\kappa} \psi_p \right) \right] (\eta, x_1, \dots, x_p) \\ &= \frac{r^2}{2\kappa^2} \sum_{k=1}^p \sum_{e: \|e\|=1} \left(\psi(\eta, x_k + e) - \psi(\eta, x_k) \right)^2 + O\left(\left(\frac{1}{\kappa} \right)^3 \right). \end{aligned} \quad (4.8.9)$$

Using Hölder's inequality to separate terms, we may therefore reduce to the case $p = 1$ and deal with the first term in the r.h.s. of (4.8.7) to get the analogue of Proposition 4.4.2.

For the second term in (4.8.7), using (1.2.6) we have

$$\left(\mathcal{P}_T^{(p)} \phi_p \right) (\eta, x_1, \dots, x_p) = \sum_{k=1}^p \sum_{z \in \mathbb{Z}^d} p_{2dT_1[\kappa]}(z, x_k) [\eta(z) - \rho] = \sum_{k=1}^p \left(\mathcal{P}_T \phi \right) (\eta, x_k). \quad (4.8.10)$$

Using Hölder's inequality to separate terms, we may therefore again reduce to the case $p = 1$ and deal with the second term in the r.h.s. of (4.8.7) to get the analogue of Proposition 4.4.3.

A Appendix

In this appendix we give the proof of Proposition 1.2.1.

Proof. Fix $t \geq 0$, $\eta \in \Omega$ and $K: \mathbb{Z}^d \times [0, \infty) \rightarrow \mathbb{R}$ such that $S = \sum_{z \in \mathbb{Z}^d} \int_0^t ds |K(z, s)| < \infty$. First consider the case $K \geq 0$. Since the ξ -process and the $\tilde{\xi}$ -process are both monotone in their initial configuration (as is evident from the graphical representation described in Section 1.2), it suffices to show that

$$\mathbb{E}_{\eta} \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \xi_s(z) \right] \right) \leq \mathbb{E}_{\eta}^{\text{IRW}} \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \tilde{\xi}_s(z) \right] \right), \quad (\text{A.0.11})$$

for all $\eta \in \Omega$ such that $|\{x \in \mathbb{Z}^d : \eta(x) = 1\}| < \infty$. This goes as follows.

Since $\xi_s(z) \in \{0, 1\}$, we may write for any $r \in \mathbb{R} \setminus \{0\}$,

$$\mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \xi_s(z) \right] \right) = \mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \frac{e^{r \xi_s(z)} - 1}{e^r - 1} \right] \right). \quad (\text{A.0.12})$$

By Taylor expansion, we get

$$\begin{aligned} & \mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \frac{e^{r \xi_s(z)} - 1}{e^r - 1} \right] \right) \\ &= \exp \left[\frac{-t}{e^r - 1} S \right] \mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \frac{e^{r \xi_s(z)}}{e^r - 1} \right] \right) \\ &= \exp \left[\frac{-t}{e^r - 1} S \right] \\ & \quad \times \sum_{n=0}^{\infty} \left(\frac{1}{e^r - 1} \right)^n \frac{1}{n!} \left(\prod_{j=1}^n \int_0^t ds_j \sum_{z_j \in \mathbb{Z}^d} \right) \left(\prod_{j=1}^n K(z_j, s_j) \right) \mathbb{E}_\eta \left(\exp \left[r \sum_{j=1}^n \xi_{s_j}(z_j) \right] \right). \end{aligned} \quad (\text{A.0.13})$$

According to Lemma 4.1 in Landim [11], we have for any $r \in \mathbb{R}$,

$$\mathbb{E}_\eta \left(\exp \left[r \sum_{j=1}^n \xi_{s_j}(z_j) \right] \right) \leq \mathbb{E}_\eta^{\text{IRW}} \left(\exp \left[r \sum_{j=1}^n \tilde{\xi}_{s_j}(z_j) \right] \right). \quad (\text{A.0.14})$$

Picking $r \geq 0$, combining (A.0.12–A.0.14), and using the analogue of (A.0.13) for $(\tilde{\xi}_t)_{t \geq 0}$, we obtain

$$\mathbb{E}_\eta \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \xi_s(z) \right] \right) \leq \mathbb{E}_\eta^{\text{IRW}} \left(\exp \left[\sum_{z \in \mathbb{Z}^d} \int_0^t ds K(z, s) \frac{e^{r \tilde{\xi}_s(z)} - 1}{e^r - 1} \right] \right). \quad (\text{A.0.15})$$

Now let $r \downarrow 0$ and use the dominated convergence theorem to arrive at (A.0.11).

For the case $K \leq 0$ we can use the same argument with

$$-\xi_s = \frac{e^{-r \xi_s} - 1}{1 - e^{-r}}. \quad (\text{A.0.16})$$

■

References

- [1] R. Arratia, Symmetric exclusion processes: a comparison inequality and a large deviation result, *Ann. Probab.* 13 (1985) 53–61.
- [2] C.-C. Chang, C. Landim and T.-Y. Lee, Occupation time large deviations of two-dimensional symmetric simple exclusion process, *Ann. Probab.* 32 (2004) 661–691.
- [3] J. D. Deuschel and D. W. Stroock, *Large Deviations*, Academic Press, London, 1989.

- [4] J. Gärtner and F. den Hollander, Intermittency in a catalytic random medium, EURANDOM Report 2004-019. To appear in *Ann. Prob.*
- [5] J. Gärtner and W. König, The parabolic Anderson model, in: *Interacting Stochastic Systems* (J.-D. Deuschel and A. Greven, eds.), Springer, Berlin, 2005, pp. 153–179.
- [6] J. Gärtner and S.A. Molchanov, Parabolic problems for the Anderson Hamiltonian. I. Intermittency and related topics. *Commun. Math. Phys.* 132 (1990) 613–655.
- [7] F. den Hollander, *Large Deviations*, Fields Institute Monographs 14, American Mathematical Society, Providence, RI, 2000.
- [8] T. Kato, *Perturbation Theory for Linear Operators* (2nd. ed.), Springer, New York, 1976.
- [9] H. Kesten and V. Sidoravicius, Branching random walk with catalysts, *Electr. J. Probab.* 8 (2003) Paper no. 5, pp. 1–51.
- [10] C. Kipnis, Fluctuations des temps d’occupation d’un site dans l’exclusion simple symétrique, *Ann. Inst. H. Poincaré Probab. Statist.* 23 (1987) 21–35.
- [11] C. Landim, Occupation time large deviations for the symmetric simple exclusion process, *Ann. Probab.* 20 (1992) 206–231.
- [12] T.M. Liggett, *Interacting Particle Systems*, Grundlehren der Mathematischen Wissenschaften 276, Springer, New York, 1985.
- [13] F. Spitzer, *Principles of Random Walk* (2nd. ed.), Springer, Berlin, 1976.