

## Direct epoxidation of propene with vapor hydrogen peroxide

**Citation for published version (APA):**

Perez Ferrandez, D. M., Croon, de, M. H. J. M., Schouten, J. C., & Nijhuis, T. A. (2013). Direct epoxidation of propene with vapor hydrogen peroxide. In *Proceedings of the 9th European Conference of Chemical Engineering (ECCE9)*, 21-25 April 2013, The Hague, The Netherlands

**Document status and date:**

Published: 01/01/2013

**Document Version:**

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.tue.nl/taverne](http://www.tue.nl/taverne)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[openaccess@tue.nl](mailto:openaccess@tue.nl)

providing details and we will investigate your claim.



## View abstract data

**Abstract title** MAXWELL-STEFAN MODELLING OF ION TRANSPORT IN A NAFION MEMBRANE

**Author** Moshtarikhah , Shohreh, Technical University of Eindhoven, Eindhoven, Nederland (Presenting author)

**Co-author(s)** Keurentjes, Jos, Technical University of Eindhoven, Eindhoven, Nederland  
Schouten, Jaap, Technical University of Eindhoven, Eindhoven, Nederland  
Schaaf, John van der, Technical University of Eindhoven, Eindhoven, Nederland

### Background

Cation-exchange membranes are used in electrolysis processes to separate anodic and cathodic products due to their high selectivity of positive ions. The principle of all cation-exchange membranes is the combination of mass and electrical charge transport by cations through the perm-selective membrane applying electrical potential gradient. A nafion membrane with sulfonic groups separating sodium chloride and sodium hydroxide of higher concentration solutions is studied in this paper.

### Aims

The aim is to investigate the membrane resistance to transport of ions in a nafion membrane by mathematical modeling of ion transport through the membrane under high current and concentration conditions.

### Methods

For prediction of ion transport inside the membrane appropriate mass transport and equilibrium models are required. The mass transfer can be modeled using either the Nernst-Planck or the Maxwell-Stefan equations [1]. The Nernst-planck equation neglects the interactions of ions and according to literature it is valid for system of dilute concentrations [2]. The Maxwell-Stefan equation takes the interaction of different species into account. However, it requires reliable data for diffusion coefficients [3]. In this paper, a mathematical model of ion transport in the membrane is developed based on the Maxwell-Stefan equation. In addition, the boundary conditions are determined with the Donnan equilibrium [4,5]. Solving the Maxwell-Stefan equations requires the selection of a component that is not calculated from the molar flux equations, but from the Gibbs-Duhem rule that the sum of mole fractions is unity. For charged components an additional condition is the electroneutrality.

### Results

We show that the Maxwell-Stefan equation is sensitive for the choice of component that is eliminated (Figure 1-2). When no current is applied  $\text{Na}^+$  ions are expected to diffuse from cathode to anode and the current density should force them toward the cathode. Also, increasing the current density causes higher driving force for  $\text{OH}^-$  ions to diffuse back and less  $\text{Cl}^-$  ions to diffuse into the membrane.

### Summary/Conclusion

The model in case of  $\text{Na}^+$  elimination predicts that increasing the current density results in more transport of  $\text{Na}^+$  ions from cathode to anode compartment. This can be due to the fact that the potential gradient which forces  $\text{Na}^+$  ions to transport toward cathode is not predicted by the electroneutrality condition. Also, the back migration of  $\text{OH}^-$  is decreasing when current is applied to the system. In case of  $\text{OH}^-$  elimination the transport of ions is as expected but increasing the current density can limit the transport of sodium ions.

[1] R. Krishna, Multicomponent mass transfer, John Wiley and Sons , 1993.

[2] S.T.P. Psaltis, et al, J. Electrochem. Soc. 158 (2010) A33-A42.

[3] J.H.G. Van der Stegen, The state of the art of modern chlor alkali electrolysis with membranecells, PhD thesis, Twente University, 2000.

[4] H. Strathmann, Ion-exchange membrane separation processes, first ed., Elsevier, 2004.

[5] M. Higa, et al, Simulation of the transport of ions against their concentration gradient across charged membrane, J. Mem. Sci. 37 (1987) 251-266.







Figure 1. Mole fraction profile of  $Na^+$ ,  $Cl^-$  and  $OH^-$  over anolyte ( $NaCl$ ), membrane and catholyte ( $NaOH$ ) ( $Na^+$  elimination)




Figure 2. Mole fraction profile of  $Na^+$ ,  $Cl^-$  and  $OH^-$  over anolyte (NaCl), membrane and catholyte (NaOH) ( $OH^-$  elimination)