

Development of a mobile phone application for stimulation of personal mobility for COPD patients

Citation for published version (APA): Vorrink, S. N. W., Kort, H. S. M., Troosters, T., & Lammers, J. J. (2014). Development of a mobile phone application for stimulation of personal mobility for COPD patients. Gerontechnology, 13(2), 87-88. https://doi.org/10.4017/gt.2014.13.02.208.00

DOI: 10.4017/gt.2014.13.02.208.00

Document status and date:

Published: 01/01/2014

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

• A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.

• The final author version and the galley proof are versions of the publication after peer review.

• The final published version features the final layout of the paper including the volume, issue and page numbers.

Link to publication

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- · Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

the recordings. **Results & Discussion** Peak sound levels are mostly caused by the slamming of doors (e.g. closets) and activities of residents and professional caregivers. Averaged over the five sleeping rooms the results show in the night period a mean A-weighted background noise level of 32.1 dB. The maximum A-weighted peak levels go up to 97.8 dB. During the day in the common

rooms a mean Aweighted background noise level of 55.3 dB was measured with a maximum A-weighted peak levels up to 115.0 dB.

References

 World Health Organization, International Classification of Functioning, Disabilities and Health. Geneva: WHO; 2001 Table 1. Measurement results of five common rooms and five sleeping rooms; Aeq=A-weighted background noise level; Peak.max=Maximum peak sound level; Peak. 5min= 5min Mean sound level

	Common rooms		Sleeping rooms	
Parameter (dB)	Mean	Range	Mean	Range
L _{Aeg day}	55,3	53,4–57,0	42,7	41,0–45,6
L _{Aeq night}	32,2	29,6–33,7	32,1	28,5–36,5
L _{A,peak,max day}	108,6	101,9–115,0	107,5	99,2–117,0
L _{A,peak,max night}	92,1	83,9–105,4	89,0	83,7–97,8
LA,peak,5min,>80,avg day	90,0	88,6–91,6	88,9	86,9–92,6
LA,peak,5min,>80,avg night	86,1	83,5–89,2	85,2	83,2–87,7

 Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB ,Blazer DG. Sleep complaints among elderly persons: An epidemiologic study of three communities. Sleep 1995;18(6):425-432
 Keywords: housing & daily activities, health, sound levels, sound sources, older adults

Address: Acoustics Laboratory, Eindhoven University of Technology, Eindhoven, Netherlands *E*: n.h.a.m.v.hout@tue.nl

S.N.W. VORRINK, H.S.M. KORT, T. TROOSTERS, J-W.J. LAMMERS. **Development of a mobile** *phone application for stimulation of personal mobility for COPD patients.* Gerontechnology 2014;13(2):87-88; doi:10.4017/gt.2014.13.02.208.00 **Purpose** Chronic Obstructive Pulmonary Disease (COPD) is a disabling airway disease with variable extrapulmonary effects that may contribute to disease severity in individual patients¹. Patients with COPD show reduced levels of spontaneous daily physical activity (DPA) compared with healthy controls². This results in a higher risk of hospital admission and shorter survival³. Pulmonary rehabilitation can help to



Figure 1: Three versions of the app, adjusted from left to right; the newest version is on the right

improve the DPA level, but the benefits obtained decline after 1-2 years⁴. **Purpose** In order to maintain DPA in COPD patients after rehabilitation, we developed a mobile phone application⁵. It measures DPA as steps per day, measured by the accelerometer of the smartphone, and shows the information to the patient via the display of the mobile phone. A physiotherapist can monitor the patient via a secure website, on which DPA measurements are visible for all patients. Here DPA goals can be adjusted and text messages sent. This presentation focuses on the development of the mobile phone application (app). Method A list of requirements for the app was created by the research group, keeping the main research purpose in mind. An algorithm was written that measures steps by looking at sinus movements. Preliminary testing was performed by carrying the smartphone with the app for 3 days in combination with a validated accelerometer (Figure 1, T=1). Subsequently adjustments were made and testing was repeated. Results have been discussed in an interactive workshop. Development of the app was iterative (SCRUM methodology⁶). Subsequently, three pilot studies were performed to test the mobile phone application for usability, user friendliness, and reliability, and were assessed with questionnaires. The application was tested with students and COPD patients, who wore the phone in their pocket. Pilot studies 1, 2, and 3 (Figure 1, T=1, T=2, and T=3, respectively) lasted for a week (n=10 students), 3 days (n=3), and 3 weeks (n=7), respectively. Feedback was collected in an interactive workshop. Subjects also wore a validated accelerometer (Sensewear) during the period of the study in order to compare these data with the DPA measurements of the mobile phone application. Mix methods have been employed for quantitative and qualitative data analysis in SPSS. Results & Discussion The findings from the interactive workshop indicated that the design for description of the data needed to be explored. It also provided insight in the feedback to be given in text to persuade users to be physically active and which type of widgets motivate users to increase their physical activity. The application was found to be useful and easy to learn.

References

- Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodriguez-Roisin R, van Weel C, Zielinski J. Global Initiative for Chronic Obstructive Lung Disease. American Journal of Respiratory and Critical Care Medicine 2007;176(6):532-555
- Pitta F, Troosters T, Spruit MA, Probst VS, Decramer M, Gosselink R. American Journal of Respiratory and Critical Care Medicine 2005;171(9):972-7; doi:10.1164/rccm.200407-855OC
- 3. Pitta F, Troosters T, Probst VS, Spruit MA, Decramer M, Gosselink R. European Respiratory Journal 2006;27(5):1040-1055; doi:10.1183/09031936.06.00064105
- Foglio K, Bianchi L, Bruletti G, Porta R, Vitacca M, Balbi B, Ambrosino N. Respiratory Medicine 2007;101(9):1961-1970; doi:10.1016/j.rmed.2007.04.007
- Vorrink S, Kort H, Troosters T, Lammers J-W. Gerontechnology 2012;11(2):423; doi:10.4017/gt.2012.11.02.114.00
- Schwaber K. SCRUM Development Process. Business Object Design and Implementation. London: Springer 1997; pp 117-134

Keywords: housing & daily activities, health and self-esteem, COPD, physical activity *Address*: Bolognalaan 101, 3584 CJ Utrecht, Netherlands; *E*: sigrid.vorrink@hu.nl

A. NEVEN, T. BELLEMANS, D. JANSSENS, G. WETS. Application to support elderly while taking public transport services. Gerontechnology 2014; 13(2):88-89; doi:10.4017/gt.2014.13.02.150.00 **Purpose** On-demand responsive transport (DRT) services are frequently offered in the context of door-to-door transportation of the elderly with their social participation as the ultimate goal. In Flanders, however, the supply of accessible transportation is fragmented. There are many different types of transportation provided, but many of these are especially expensive. Therefore, we aimed to support the elderly in using regular public transport (PT) services through coaching sessions and personalized monitoring during their trips. We also aimed to meet the specific transportation requests of elderly that cannot be met by using the regular PT network, in order to provide a wide inclusive mobility system in Flanders. **Method** In order to monitor elderly during their PT trips, the smartphone application 'Viamigo' was developed (in cooperation with Thomas More). This app allows the registration of newly learned PT routes by means of their GPS coordinates. The routes can be monitored from a distance by a coach (e.g. family member or care giver), who can receive a signal when the user gets too far from