

Thermophysical properties of novel ethyl-2-alkylpyridinium based ionic liquids

Citation for published version (APA):

Alves da Rocha, M. A., Vilas, M., Rodrigues, A. S. M. C., Tojo, E., & Santos, L. M. N. B. F. (2014). Thermophysical properties of novel ethyl-2-alkylpyridinium based ionic liquids. In M. Kroon, C. Peters, & T. Vlugt (Eds.), *27th European Symposium on Applied Thermodynamics(ESAT2014)*, 6-9 July 2014, Eindhoven, The Netherlands

Document status and date:

Published: 01/01/2014

Document Version:

Accepted manuscript including changes made at the peer-review stage

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Thermophysical Properties of Novel Ethyl-2-alkylpyridinium based Ionic Liquids

Marisa A. A. Rocha^{1,2}, Miguel Vilas³, Ana S. M. C. Rodrigues¹

Emilia Tojo³, Luís M. N. B. F. Santos¹

1 CIQ, Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade do Porto, R. Campo Alegre 687, P-4169-007, Porto, Portugal

2 Department of Chemical Engineering and Chemistry, Eindhoven University of Technology, Den Dolech 2, 5600 MB Eindhoven, The Netherlands

3 Department of Organic Chemistry, Faculty of Chemistry of the University of Vigo, Marcosende, 36310, Vigo, Spain

Keywords: Pyridinium based ionic liquids; thermophysical properties.
Corresponding author: Marisa A. A. Rocha, M.A.A.Rocha@tue.nl

Abstract

Densities and viscosities and their dependence on temperature were measured for the 1-ethyl-2-alkylpyridinium bis(trifluoromethanesulfonyl)imide, $[\text{C}_{N-2}^1\text{C}_2\text{Py}][\text{NTf}_2]$ ($N=4, 5, 6, 7, 8, 9, 10$) ionic liquid series. Molar volume, thermal expansion coefficients, VTF parameters, as well as, the energy barrier related to the fluid shear stress at $T = 298.15$ K, were derived. High precision heat capacities, at $T = 298.15$ K, were measured using a heat capacity drop calorimeter.

The effect and trend of the thermophysical properties along the alkyl chain length in the position 2 of the core 1-ethylpyridinium cation, are used to explore the nanostructuring effect of ILs. A trend shift on the thermophysical properties at the $[\text{C}_5^1\text{C}_2\text{Py}][\text{NTf}_2]$ was found, which presents an indication and support for the initiation of the nanostructuring in the pyridinium based ionic liquids. The results and insights are in agreement with the previously studied alkyimidazolium ionic liquids series.[1-6] The observed trend shift in the viscosity and in the heat capacities of the pyridinium ionic liquids is more pronounced in the pyridinium than in the imidazolium ionic liquids series. This work confirms and is an additional support, for the molecular understanding of the trend shift observed in the thermophysical properties of ionic liquids, as well as, a contribution for the understanding of the relation between the structural organization of the ionic liquids, their physico-chemical properties, tuning and design.

References

- [1] M. A. A. Rocha, C.F. R. A. C. Lima, L. R. Gomes, B. Schröder, J. A. P. Coutinho, I. M. Marrucho, J. M. S. S. Esperança, L. P. N. Rebelo, K. Shimizu, J. N. Canongia Lopes, L. M. N. B. F. Santos, *J. Phys. Chem. B*, 2011, 115, 10919–10926.
- [2] K. Shimizu, C. E. S. Bernardes, J. N. Canongia Lopes, *J. Phys. Chem. B*, 2014, 118, 567–576.
- [3] M. A. A. Rocha, C. M. S. S. Neves, M. G. Freire, O. Russina, A. Triolo, J. A. P. Coutinho, L. M. N. B. F. Santos, *J. Phys. Chem. B*, 2013, 117, 10889–10897.
- [4] M. A. A. Rocha, J. A. P. Coutinho, L. M. N. B. F. Santos, *J. Chem. Phys.*, 2013, 139, 104502.
- [5] O. Russina, A. Triolo, L. Gontrani, R. Caminiti, D. Xiao, L. G. Hines Jr, R. A. Bartsch, E. L. Quitevis, N. Plechkova, K. R. Seddon, *J. Phys.: Condens. Matter*, 2009, 21, 424121.
- [6] K. Shimizu, M. F. Costa Gomes, A. A.H. Pádua, L. P.N. Rebelo, J. N. Canongia Lopes, *J. Mol. Struct.: THEOCHEM*, 2010, 946, 70–76