

The influence of an electrostatic field on cyclotron resonance behaviour of a plasma

Citation for published version (APA):

Schram, D. C., Manintveld, P., & Oord, E. (1970). The influence of an electrostatic field on cyclotron resonance behaviour of a plasma. In *Contributions to the 4th European Conference on Controlled Fusion and Plasma Physics* (pp. 107). National Nuclear Energy Committee.

Document status and date:

Published: 01/01/1970

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

H.F. PLASMAS AND HEATING

THE INFLUENCE OF AN ELECTROSTATIC FIELD ON CYCLOTRON RESONANCE BEHAVIOUR OF A PLASMA

by
D.C. Schram, P. Manintveld and E. Oord
Association Euratom-FOM
FOM-Instituut voor Plasma-Fysica
Rijnhuizen, Jutphaas, The Netherlands

Abstract

The theoretically predicted influence of an electrostatic field on the energy gain of electrons at e.c.r. is confirmed experimentally by measuring the loss tangent of the plasma as a function of an applied D.C. voltage. The applicability of this effect as a heating scheme is discussed in general terms.

In a recent publication¹⁾ the influence of an electrostatic field on the energy gain of charged particles at cyclotron resonance has been studied. Here a circularly polarized TEM wave (ω, k) was assumed to propagate along a uniform static magnetic field. In the following $g=eE^v/m\omega c$, $a=eE_0/m\omega v_p$, $\beta=aB_0/m\omega-1$, are parameters defining the e.m. wave, the e.s. field, and the static magnetic field resp.; $\alpha=n_e^2/mc_0\omega^2=\omega_p^2/\omega^2$, is a parameter for the electron number density, n_e . Close to the resonance even a weak external effect, here an e.s. field, has a considerable influence on the motion of single electrons, resulting in a change of the characteristics of plasma.

In the absence of an e.s. field, the so-called "undisturbed case", the energy of the particle is a periodic function of time, Fig. 1 curve "0", and an oscillation time t_{OS} can be defined. This remains valid if the relativistic mass variation and axial velocity variation due to the Lorentz force are taken into account³⁾

In the presence of an e.s. field two regimes can be distinguished, depending on the time scale on which the e.s. field causes a certain change of the resonance function, $r(t)=(\omega-kv_z-\Omega(t))\omega^2$:
- "strong external effect", the particle is pulled out of resonance within the first oscillation period. A stepwise energy increase is found (curves "1.03", "2" of Fig. 1).
- "weak external effect", the particle remains close to resonance: a continuous increase in energy is added to the undisturbed oscillatory behaviour (Fig. 1, curves ".1", ".5").
At the transition, $a=a_{synchr.}$, defined as the synchronous case, a continuous increase in energy occurs (Fig. 1, curve "1").




Fig. 1.

Numerically calculated gain of kinetic energy as a function of time; $c^2/v^2=0.75$, $g=2.10^{-3}$, $v(t=0)=0$, $\beta=0$, $a_{synchr.}=1.2 \times 10^{-4}$. The quantity $a/a_{synchr.}$ is curve parameter.

The reactive and resistive behaviour of the plasma has been studied for the "undisturbed case" in Ref. 3. Using the same calibration for the electric field strength of the e.m. wave as in Ref. 3, we find for $\beta=0$ (at cyclotron resonance), $a=0$ (no e.s. field): $\frac{1-\epsilon}{\alpha}/\beta=0, a=0 \approx 900 P_{in}^{-1/3}$ where P_{in} is the incident power of the e.m. wave in Watts. If an e.s. field is applied a part of the electrons is accelerated and a part is decelerated. This leads to resonance functions of opposite sign and, consequently, to dielectric currents of opposite phase. For high values of the e.s. field parameter a , the permittivity, ϵ , tends to the vacuum value 1.

The loss tangent of the plasma can be calculated if the time of escape of the particles is known. Experimentally³⁾

$$t_{es} = 1/2 t_{OS} / \beta=0, a=0 \text{ yielding } \frac{tg\delta}{\alpha} / \beta=0, a=0 \approx 620 P_{in}^{-1/3}$$

Evidently, the loss tangent will show a weak maximum for $a=a_{synchr.}$. For values of $a > a_{synchr.}$ the step in energy decreases with increasing a . Consequently, if the escape time is assumed to be independent of a , the loss tangent decreases.

The experimental facility shown in Figs. 9, 10 of Ref. 3 has been used for the measurements. Two electrodes were mounted on the axis and a negative D.C. voltage was applied. The magnetic field strength of the static magnetic field was chosen such that the cyclotron frequency was equal to the applied frequency ($\beta=0$). The loss tangent, the permittivity, and the density of the plasma were measured as functions of the applied D.C. voltage for several values of P_{in} . The result for $P_{in}=20W$ are plotted in Fig. 2 (normalized to the density); the loss tangent shows a maximum at $V=2.2$ Volt, taken to be $V_{synchr.}$. Values of $V_{synchr.}$ for various values of P_{in} are plotted in Fig. 3.




Fig. 2.

Measured loss tangent and dielectric constant as functions of the applied D.C. voltage. $V_{synchr.}$ is the value of V where $tg\delta/\alpha$ is max.




Fig. 3.

$V_{synchr.}$ as a function of P_{in} . The dashed line is the theoretically predicted curve for a Debye length $\lambda_D \approx 1$ cm.

As predicted in Ref. 1 we find $V_{synchr.} = P_{in}^{2/3}$. Taking the field calibration of Ref. 3, we find theoretically $a_{synchr.} = .92 \times 10^{-6} \times P_{in}^{2/3}$. The e.s. field is assumed to be constant over a shielding length λ_D of 1 cm, which corresponds to the measured $n_e = 3 \times 10^{17}/cc$ and the estimated parallel electron energy of 50 eV. The experimental result, $V_{synchr.} = .27 \times P_{in}^{2/3}$, is in agreement with theoretical predictions.

Discussion

The mechanisms discussed here may, in more general terms, be applied to heat plasmas. A discussion of both cases, weak and strong external effects, may be of interest.

If the external effect is weak, the energy grows without any limitation but time. However, even if the density is small the plasma can become extremely reactive, while for higher densities cyclotron absorption will damp the wave before large energies are reached. As a heating scheme it is only attractive in very tenuous plasmas of large dimensions, e.g. extra terrestrial plasmas.

In a strong e.s. field a quasi-collisional energy gain occurs. The reactive components of the current are relatively low and can be chosen zero by choosing a value of β slightly different from zero. The permittivity of the plasma is close to the vacuum value. This mechanism could be applied as a heating mechanism in which the heating rate can be chosen. It should be remarked that the external field need not be static: e.g. an e.s. field varying slowly with respect to t_{OS} leads to repetitive heating. Such electrostatic waves can be introduced into the plasma. Frequency modulation is an attractive alternative as an external mechanism.

Acknowledgement

Discussions with Dr. L.Th.M. Ornstein and Ir. T.J. Schep have been of great help to the authors.

This work was performed under the association agreement of Euratom and FOM with financial support from ZWO and Euratom.

References

- 1) D.C. Schram and G.P. Beukema, Physica 42 (1969), 277-290.
- 2) D.C. Schram, W. Strijland and L.Th.M. Ornstein, Contributions III Eur. Conf. Contr. Fusion & Plasma Phys., p.71, Utrecht, 1969.
- 3) D.C. Schram, Physica 46 (1970), 97-118.