

Approximating the Moments of Generalized Gaussian Distributions via Bell's Polynomials †

Citation for published version (APA):

Caratelli, D., Sabbadini, R., & Ricci, P. E. (2023). Approximating the Moments of Generalized Gaussian Distributions via Bell's Polynomials †. *Axioms*, 12(2), Article 206. <https://doi.org/10.3390/axioms12020206>

Document license:

CC BY

DOI:

[10.3390/axioms12020206](https://doi.org/10.3390/axioms12020206)

Document status and date:

Published: 01/02/2023

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy



If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Article

Approximating the Moments of Generalized Gaussian Distributions via Bell's Polynomials [†]

Diego Caratelli ^{1,2} , Ruben Sabbadini ³ and Paolo Emilio Ricci ^{4,*} ¹ The Antenna Company, High Tech Campus 41, 5656 AE Eindhoven, The Netherlands² Department of Electrical Engineering, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands³ Liceo Scientifico Statale Farnesina, Via dei Giuochi Istmici, 64, 00135 Roma, Italy⁴ Department of Mathematics, International Telematic University UniNettuno, Corso Vittorio Emanuele II, 39, 00186 Roma, Italy

* Correspondence: paoloemilio.ricci@uninettunouniversity.net

[†] Dedicated to Prof. Hari M. Srivastava on the Occasion of His 82th Birthday.

Abstract: Bell's polynomials are used in many different fields of mathematics, ranging from number theory to operator theory. This paper shows a relevant application in probability theory aimed at computing the moments of generalized Gaussian distributions. To this end, a table containing the first values of the complete Bell's polynomials is provided. Furthermore, a dedicated code for approximating the moments of the general distributions in terms of complete Bell's polynomials is detailed. Several test cases concerning different nested functions are discussed.

Keywords: moments of a continuous distribution; Bell's polynomials; generalized Gaussian distributions

MSC: 05A10; 11B83; 60E05; 62E17



Citation: Caratelli, D.; Sabbadini, R.; Ricci, P.E. Approximating the Moments of Generalized Gaussian Distributions via Bell's Polynomials. *Axioms* **2023**, *12*, 206. <https://doi.org/10.3390/axioms12020206>

Academic Editors: Juan J. Nieto and Sevtap Sümer Eker

Received: 3 January 2023

Revised: 8 February 2023

Accepted: 13 February 2023

Published: 15 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Bell's polynomials were initially introduced to represent successive derivatives of composite functions [1]. However, as it has been pointed out in [2], such polynomials, being related to partitions of integers [3], find immediate applications in combinatorial mathematics [4]. They also intervene in formulas for representing symmetric functions of the zeros of analytic functions, thus generalizing the classical Newton–Girard formulas. Using these results, the reduction formulas of the orthogonal invariants of a strictly positive compact operator were expressed. In this way, Robert's formulas [5] and other expressions related to those can be derived.

Generalized classes of Bell's polynomials have been proposed in the literature, e.g., by Fujiwara [6], Kim [7], and Rai, Singh [8]. Further generalizations, including the multi-dimensional case, can be found in papers by Bernardini, Natalini, Ricci and Natalini, and Ricci (see [9], and the references therein).

There are many areas in which the adoption of Bell's polynomials has allowed useful approximations to be obtained in applied mathematics, such as those concerning the Laplace transform of composite analytic functions (see [10], and the references therein).

In this paper, we discuss another application to the probability theory [11] by deriving some formulas useful to calculate the moments of generalized Gaussian distributions. To this end, judicious use is made of the extensions of Bell's polynomials presented in previous papers.

This article is organized as follows: In Section 2, Bell's polynomials and their main properties are briefly recalled. In Section 3, Bell's polynomials are used to approximate the primitives of composite exponential functions. Finally, after recalling, in Section 4, the generalization of Bell's polynomials to the case of multi-nested functions, the moments

of generalized Gaussian distributions are approximated in Section 5. Some examples illustrated using version 13.2 of the computer algebra software Mathematic[®] are shown in Section 6.

2. Recalling the Bell’s Polynomials

Consider the composite function $\Phi(t) := f(g(t))$, where $x = g(t)$ and $y = f(x)$ are differentiable functions (up to a sufficiently large order), defined in suitable intervals of the real axis so that $\Phi(t)$ can be differentiated n times with respect to t by using the chain rule. Here, and in what follows, we use the following notation:

$$\Phi_m := D_t^m \Phi(t), \quad f_h := D_x^h f(x)|_{x=g(t)}, \quad g_k := D_t^k g(t).$$

Then, the n -th derivative of $\Phi(t)$ is represented by

$$\Phi_n = Y_n(f_1, g_1; f_2, g_2; \dots; f_n, g_n) = \sum_{k=1}^n B_{n,k}(g_1, g_2, \dots, g_n) f_k, \tag{1}$$

where Y_n denotes the general Bell’s polynomial, and the coefficients $B_{n,k}$ for all $k = 1, \dots, n$, are homogeneous polynomials of the variables g_1, g_2, \dots, g_n , of degree k and *isobaric* of weight n (i.e., they are a linear combination of monomials $g_1^{k_1} g_2^{k_2} \dots g_n^{k_n}$ having the same degree $k_1 + 2k_2 + \dots + nk_n = n$).

Bell’s polynomials satisfy the recurrence relation

$$\begin{cases} Y_0 := f_1; \\ Y_{n+1}(f_1, g_1; \dots; f_n, g_n; f_{n+1}, g_{n+1}) = \\ = \sum_{k=0}^n \binom{n}{k} Y_{n-k}(f_2, g_1; f_3, g_2; \dots; f_{n-k+1}, g_{n-k}) g_{k+1}. \end{cases} \tag{2}$$

and they can be expressed explicitly using the Faà di Bruno’s formula [12,13]

$$Y_n(f_1, g_1; f_2, g_2; \dots; f_n, g_n) = \sum_{\pi(n)} \frac{n!}{r_1! r_2! \dots r_n!} f_r \left[\frac{g_1}{1!} \right]^{r_1} \left[\frac{g_2}{2!} \right]^{r_2} \dots \left[\frac{g_n}{n!} \right]^{r_n}, \tag{3}$$

where the sum runs over all the partitions $\pi(n)$ of the integer n , r_i denotes the number of parts of size i , and $r = r_1 + r_2 + \dots + r_n$ denotes the number of parts of the considered partition [2].

The coefficients $B_{n,k}$ in Equation (1) depend only on the values $g_1, g_2, \dots, g_{n-k+1}$, and satisfy the recursion

$$B_{n,k}(g_1, g_2, \dots, g_{n-k+1}) = \sum_{h=0}^{n-k} \binom{n-1}{h} B_{n-h-1, k-1}(g_1, g_2, \dots, g_{n-k-h+1}) g_{h+1}. \tag{4}$$

In the particular case where $f(x) = e^x$, that is equivalent to considering the composite function $e^{g(t)}$, and upon assuming that $g(0) = 0$, we obtain the following simplified expression for the right-hand side of (1):

$$\sum_{k=1}^n B_{n,k}(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_{n-k+1}) \overset{\circ}{f}_k = \sum_{k=1}^n B_{n,k}(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_{n-k+1}) = B_n(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_n), \tag{5}$$

where the B_n are the *complete Bell’s polynomials*. It results in $B_0(g_0) := f(g(0))$, and the first few values of B_n , for $n = 1, 2, \dots, 5$ are given by

$$B_1(g_1) = g_1, \quad B_2(g_1, g_2) = g_1^2 + g_2, \quad B_3(g_1, g_2, g_3) = g_1^3 + 3g_1g_2 + g_3. \tag{6}$$

Further values are reported in Appendix A.

The values of the complete Bell’s polynomials for particular choices of the parameter can be found in [14].

The complete Bell’s polynomials satisfy the identity (see e.g., [15])

$$B_{n+1}(g_1, \dots, g_{n+1}) = \sum_{k=0}^n \binom{n}{k} B_{n-k}(g_1, \dots, g_{n-k}) g_{k+1}. \tag{7}$$

3. Primitives of Exponential Functions

Let $f(g(t))$ be a composite function that is analytic in a neighborhood of the origin. The relevant Taylor’s expansion is given by

$$f(g(t)) = \sum_{n=0}^{\infty} a_n \frac{t^n}{n!}, \quad a_n = D_t^n [f(g(t))]_{t=0}, \tag{8}$$

where

$$a_0 = f(\overset{\circ}{g}_0),$$

$$a_n = D_t^n [f(g(t))]_{t=0} = \sum_{k=1}^n B_{n,k}(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_{n-k+1}) \overset{\circ}{f}_k, \quad (n \geq 1), \tag{9}$$

and

$$\overset{\circ}{f}_k := D_x^k f(x)|_{x=g(0)}, \quad \overset{\circ}{g}_h := D_t^h g(t)|_{t=0}. \tag{10}$$

Then, the primitive of the composite function $f(g(t))$ writes

$$\int_0^x f(g(t))dt = xf(g(0)) + \sum_{n=1}^{\infty} \left(\sum_{k=1}^n B_{n,k}(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_{n-k+1}) \overset{\circ}{f}_k \right) \frac{x^{n+1}}{(n+1)!}, \tag{11}$$

Remark 1. Note that here and in what follows we avoid adding a constant to indefinite integrals, based on the assumption that the relevant primitives vanish at the origin.

In the case of exponential functions, Equation (11) becomes

$$\int_0^x \exp(g(t))dt = \sum_{n=0}^{\infty} B_n(\overset{\circ}{g}_1, \overset{\circ}{g}_2, \dots, \overset{\circ}{g}_n) \frac{x^{n+1}}{(n+1)!}, \tag{12}$$

where we have assumed, for convenience, $B_0 := 1$.

Then, in particular, it results

$$\int_0^x \exp(-t^2)dt = \frac{\sqrt{\pi}}{2} \operatorname{erf}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)n!} = \sum_{n=0}^{\infty} B_n(0, -2, 0, \dots, 0) \frac{x^{n+1}}{(n+1)!}. \tag{13}$$

As a consequence, we find

$$B_{2k}(0, -2, 0, \dots, 0) = (-1)^k, \quad B_{2k+1}(0, -2, 0, \dots, 0) = 0, \quad (k \geq 0). \tag{14}$$

4. An Extension of Bell’s Polynomials

We consider the second-order Bell’s polynomials, $Y_n^{[2]}(f_1, g_1, h_1; f_1, g_1, h_1; \dots; f_n, g_n, h_n)$, defined by the n -th derivative of the composite function $\Phi(t) := f[g(h(t))]$.

Consider the functions $x = h(t)$, $z = g(x)$, $y = f(z)$, and suppose that $h(t)$, $g(x)$, and $f(z)$ are n times differentiable with respect to their variables, so that the composite function $\Phi(t) := f[g(h(t))]$ can be differentiated n times with respect to t , by using the chain rule.

Similarly to the previous section, we use the following notation:

$$\Phi_j := D_t^j \Phi(t), \quad f_h := D_y^h f(y)|_{y=g(x)}, \quad g_k := D_x^k g(x)|_{x=h(t)}, \quad h_r := D_t^r h(t).$$

Then, the n -th derivative of $\Phi(t)$ can be represented by

$$\Phi_n = Y_n^{[2]}(f_1, g_1, h_1; f_2, g_2, h_2; \dots; f_n, g_n, h_n) = Y_n^{[2]}([f, g, h]_n), \tag{15}$$

where $Y_n^{[2]}$ denotes the general second-order Bell’s polynomial.

The first few polynomials are given by

$$\begin{aligned} Y_1^{[2]}([f, g, h]_1) &= f_1 g_1 h_1; \\ Y_2^{[2]}([f, g, h]_2) &= f_1 g_1 h_2 + f_1 g_2 h_1^2 + f_2 g_1^2 h_1^2; \\ Y_3^{[2]}([f, g, h]_3) &= f_1 g_1 h_3 + f_1 g_3 h_1^3 + 3f_1 g_2 h_1 h_2 + 3f_2 g_1 g_2 h_1^3 + f_3 g_1^3 h_1^3. \end{aligned} \tag{16}$$

A more extended table can be found in [10].

The connection between $Y_n^{[2]}$ and the ordinary Bell’s polynomials is discussed below. In [16], the following theorem has been proven.

Theorem 1. For every integer n , the polynomials $Y_n^{[2]}$ are represented in terms of the ordinary Bell’s polynomials by the following equation:

$$\begin{aligned} Y_n^{[2]}(f_1, g_1, h_1; \dots; f_n, g_n, h_n) &= \\ &= Y_n(f_1, Y_1(g_1, h_1); f_2, Y_2(g_1, h_1; g_2, h_2); \dots; f_n, Y_n(g_1, h_1; g_2, h_2; \dots; g_n, h_n)) \end{aligned} \tag{17}$$

From here, it has been shown that the second-order Bell’s polynomials can be computed recursively, avoiding the use of computationally cumbersome expressions such as the Faà di Bruno formula, on the basis of the following theorem.

Theorem 2. The second-order Bell’s polynomials verify the recursion

$$\begin{aligned} Y_0^{[2]} &= f_1; \\ Y_{n+1}^{[2]}(f_1, g_1, h_1; \dots; f_{n+1}, g_{n+1}, h_{n+1}) &= \sum_{k=0}^n \binom{n}{k} Y_{n-k}^{[2]}(f_2, g_1, h_1; f_3, g_2, h_2; \dots \\ &\dots; f_{n-k+1}, g_{n-k}, h_{n-k}) Y_{k+1}(g_1, h_1; \dots; g_{k+1}, h_{k+1}). \end{aligned} \tag{18}$$

The proof of this result is provided in [16].

Primitives of Nested Functions

Considering the second-order nested functions of the type $f[g(h(t))]$, the preceding Equation (11) writes

$$\int_0^x f[g(h(t))] dt = x f[g(h(0))] + \sum_{n=1}^{\infty} Y_n^{[2]}(f_1, g_1, h_1; f_2, g_2, h_2; \dots; f_n, g_n, h_n) \frac{x^{n+1}}{(n+1)!}, \tag{19}$$

and in the case of a nested exponential function, with $g(h(0)) = 0$,

$$\int_0^x \exp[g(h(t))] dt = x + \sum_{n=1}^{\infty} Y_n^{[2]}(1, g_1, h_1; 1, g_2, h_2; \dots; 1, g_n, h_n) \frac{x^{n+1}}{(n+1)!}. \tag{20}$$

5. Moments of Generalized Gaussian Distributions

Consider the analytic monotonically increasing function $h(t)$ and consider the nested exponential function $\exp[-(h(t))^2]$.

The normalization constant of the generalized Gaussian distribution writes

$$A = \int_{-\infty}^{+\infty} \exp[-(h(t))^2] dt, \tag{21}$$

the mean value is

$$\mu = \mu_1 = \frac{1}{A} \int_{-\infty}^{+\infty} t \exp[-(h(t))^2] dt, \tag{22}$$

and the second moment

$$\mu_2 = \frac{1}{A} \int_{-\infty}^{+\infty} t^2 \exp[-(h(t))^2] dt, \tag{23}$$

so that we find the variance

$$\sigma^2 = \mu_2 - \mu^2. \tag{24}$$

Note that, thanks to the rapid decay of the exponential function, the integrals in (21)–(23) can be estimated with a pretty negligible numerical error using a suitable Gaussian quadrature rule on a bounded interval of the real axis.

Then, assuming $y = f(z) = \exp(z)$; $z = g(x) = -x^2$; $x = h(t)$, the n th moment is given by

$$\mu_n = \frac{1}{A} \int_{-\infty}^{+\infty} t^n \exp[-(h(t))^2] dt \simeq \frac{1}{A} \int_{\mu-3\sigma}^{\mu+3\sigma} t^n \exp[-(h(t))^2] dt, \tag{25}$$

and using the second order Bell’s polynomials (17), and assuming $Y_0^{[2]} := 1$, we find

$$\begin{aligned} \mu_n \simeq \frac{1}{A} \sum_{k=0}^{\infty} Y_k^{[2]}((1, 0, \overset{\circ}{h}_1; 1, -2, \overset{\circ}{h}_2; 1, 0, \overset{\circ}{h}_3; \dots; \dots; 1, 0, \overset{\circ}{h}_k)) \cdot \\ \cdot \frac{(\mu + 3\sigma)^{n+k+1} - (\mu - 3\sigma)^{n+k+1}}{(n + k + 1) k!}. \end{aligned} \tag{26}$$

6. Particular Examples

6.1. Example 1

Consider the generalized Gaussian distribution defined by the function

$$F(t) = \exp[-(\sinh(t))^2]. \tag{27}$$

We find $\mu = \mu_1 = 0$ and $\sigma^2 = 0.55296$.

Its graph is shown in Figure 1.

The percentage deviation between $F(t)$ and the relevant expansion $\tilde{F}(t)$ in terms of Bell’s polynomials is shown in Figure 2. As can be seen, $F(t)$ is nicely approximated by $\tilde{F}(t)$ in the domain where $F(t)$ does not assume negligibly small values.

The table containing the first few moments of the considered function $F(t)$, as computed using the Gauss–Kronrod quadrature rule (G-KQR), μ_k , and the proposed Bell’s method, $\tilde{\mu}_k$, is reported in Figure 3.




Figure 1. Graph of the function $F(t) = \exp[-(\sinh(t))^2]$.




Figure 2. Relative percentage error of the Bell's approximation of $F(t)$ in (27).

$\mu[1] = 0$	$\tilde{\mu}[1] = 0.$
$\mu[2] = 0.305765$	$\tilde{\mu}[2] = 0.30574$
$\mu[3] = 0$	$\tilde{\mu}[3] = 0.$
$\mu[4] = 0.23272$	$\tilde{\mu}[4] = 0.232633$
$\mu[5] = 0$	$\tilde{\mu}[5] = 0.$
$\mu[6] = 0.256353$	$\tilde{\mu}[6] = 0.256048$
$\mu[7] = 0$	$\tilde{\mu}[7] = 0.$
$\mu[8] = 0.352805$	$\tilde{\mu}[8] = 0.351738$
$\mu[9] = 0$	$\tilde{\mu}[9] = 0.$
$\mu[10] = 0.567349$	$\tilde{\mu}[10] = 0.56361$

Figure 3. Moments of $F(t)$ in (27) as computed using the G-KQR (left) and the proposed Bell's polynomial approximation (right) methods.

6.2. Example 2

Consider the generalized Gaussian distribution defined by the function

$$F(t) = \exp[-(t + t^3/4\sqrt{3} + t^5/240)^2]. \tag{28}$$

We find $\mu = \mu_1 = 0$ and $\sigma^2 = 0.566431$.

Its graph is shown in Figure 4.




Figure 4. Graph of the function $F(t) = \exp[-(t + t^3/4\sqrt{3} + t^5/240)^2]$.

The percentage deviation between $F(t)$ and the relevant expansion $\tilde{F}(t)$ in terms of Bell’s polynomials is shown in Figure 5. As can be seen, $F(t)$ is nicely approximated by $\tilde{F}(t)$ in the domain where $F(t)$ does not assume negligibly small values.




Figure 5. Relative percentage error of the Bell’s approximation of $F(t)$ in (28).

The table containing the first few moments of the considered function $F(t)$, as computed using the Gauss–Kronrod quadrature rule (G-KQR), μ_k , and the proposed Bell’s method, $\tilde{\mu}_k$, is reported in Figure 6.

$\mu[1] = 0$	$\tilde{\mu}[1] = 0.$
$\mu[2] = 0.320844$	$\tilde{\mu}[2] = 0.320833$
$\mu[3] = 0$	$\tilde{\mu}[3] = 0.$
$\mu[4] = 0.259794$	$\tilde{\mu}[4] = 0.25975$
$\mu[5] = 0$	$\tilde{\mu}[5] = 0.$
$\mu[6] = 0.307347$	$\tilde{\mu}[6] = 0.307162$
$\mu[7] = 0$	$\tilde{\mu}[7] = 0.$
$\mu[8] = 0.457552$	$\tilde{\mu}[8] = 0.456789$
$\mu[9] = 0$	$\tilde{\mu}[9] = 0.$
$\mu[10] = 0.800564$	$\tilde{\mu}[10] = 0.797407$

Figure 6. Moments of $F(t)$ in (28) as computed using the G-KQR (left) and the proposed Bell’s polynomial approximation (right) methods.

6.3. Example 3

Consider the generalized Gaussian distribution defined by the function

$$\psi(t) := \exp \left[- \left(\frac{1}{2} \left(t - \frac{\cos(\pi/8)}{8} \right)^2 \sin(\pi/8) + \frac{\cos(\pi/8)}{6!} \left(t - \frac{\sin(\pi/8)}{8} \right)^6 + \frac{t^8}{4! 8! \sqrt{2}} \left(t - \frac{1}{8\sqrt{2}} \right)^4 \right)^2 \right]. \tag{29}$$

We find $\mu = \mu_1 = 0.096855$ and $\sigma^2 = 1.02151$. Its graph is shown in Figure 7.




Figure 7. Graph of the function $\psi(t)$ in (29).

The percentage deviation between $\psi(t)$ and the relevant expansion $\tilde{\psi}(t)$ in terms of Bell's polynomials is shown in Figure 8. As can be seen, $\psi(t)$ is nicely approximated by $\tilde{\psi}(t)$ in the domain where $\psi(t)$ does not assume negligibly small values.




Figure 8. Relative percentage error of the Bell's approximation of $\psi(t)$ in (29).

The table containing the first few moments of the considered function $\psi(t)$, as computed using the Gauss–Kronrod quadrature rule (G-KQR), μ_k , and the proposed Bell's method, $\tilde{\mu}_k$, is reported in Figure 9.

$\mu[1] = 0.096855$	$\tilde{\mu}[1] = 0.0965639$
$\mu[2] = 1.05286$	$\tilde{\mu}[2] = 1.05375$
$\mu[3] = 0.298461$	$\tilde{\mu}[3] = 0.2967$
$\mu[4] = 2.25481$	$\tilde{\mu}[4] = 2.25053$
$\mu[5] = 1.03878$	$\tilde{\mu}[5] = 1.02607$
$\mu[6] = 6.27311$	$\tilde{\mu}[6] = 6.22373$
$\mu[7] = 3.94173$	$\tilde{\mu}[7] = 3.85297$
$\mu[8] = 20.3358$	$\tilde{\mu}[8] = 19.9709$
$\mu[9] = 15.9893$	$\tilde{\mu}[9] = 15.3913$
$\mu[10] = 73.2833$	$\tilde{\mu}[10] = 70.8689$

Figure 9. Moments of $\psi(t)$ in (29) as computed using the G-KQR (left) and the proposed Bell’s polynomial approximation (right) methods.

6.4. Example 4

Consider the generalized Gaussian distribution defined by the function

$$F(t) = \exp[-(\sinh(2t))^2]. \tag{30}$$

We find $\mu = \mu_1 = 0$ and $\sigma^2 = 0.27648$.
Its graph is shown in Figure 10.




Figure 10. Graph of the function $F(t) = \exp[-(\sinh(2t))^2]$.

The percentage deviation between $F(t)$ and the relevant expansion $\tilde{F}(t)$ in terms of Bell’s polynomials is shown in Figure 11. As can be seen, $F(t)$ is nicely approximated by $\tilde{F}(t)$ in the domain where $F(t)$ does not assume negligibly small values.




Figure 11. Relative percentage error of the Bell’s approximation of $F(t)$ in (30).

The table containing the first few moments of the considered function $F(t)$, as computed using the Gauss–Kronrod quadrature rule (G-KQR), μ_k , and the proposed Bell’s method, $\tilde{\mu}_k$, is reported in Figure 12.

$\mu[1] = 0$	$\tilde{\mu}[1] = 0.$
$\mu[2] = 0.0764412$	$\tilde{\mu}[2] = 0.0764349$
$\mu[3] = 0$	$\tilde{\mu}[3] = 0.$
$\mu[4] = 0.014545$	$\tilde{\mu}[4] = 0.0145395$
$\mu[5] = 0$	$\tilde{\mu}[5] = 0.$
$\mu[6] = 0.00400552$	$\tilde{\mu}[6] = 0.00400075$
$\mu[7] = 0$	$\tilde{\mu}[7] = 0.$
$\mu[8] = 0.00137814$	$\tilde{\mu}[8] = 0.00137398$
$\mu[9] = 0$	$\tilde{\mu}[9] = 0.$
$\mu[10] = 0.000554051$	$\tilde{\mu}[10] = 0.0005504$

Figure 12. Moments of $F(t)$ in (30) as computed using the G-KQR (left) and the proposed Bell’s polynomial approximation (right) methods.

Different generalized Gaussian distributions can be evaluated using the approach illustrated in Section 5. The same applies to the multimodal distributions considered hereafter.

- Assuming $\exp[-(5x + (x - 1/2)^2 + x^6)^2]$, we find the graph in Figure 13.




Figure 13. Graph of the function $\exp[-(5x + (x - 1/2)^2 + x^6)^2]$.

- Assuming $\exp[-\sinh^2(x^3 + 3x^2 - 2)]$, we find the graph in Figure 14.




Figure 14. Graph of the function $\exp[-\sinh^2(x^3 + 3x^2 - 2)]$.

In the next example, we evaluate the moments of a bi-modal distribution. In this and other similar cases, one can verify that achieving the desired precision requires a drastic increase in the number of expansion terms, thus affecting the overall computational times.

6.5. Example 5

Consider the generalized Gaussian distribution defined by the function

$$F(t) = \exp[-(\sinh(t^4 - 1/2))^2]. \tag{31}$$

We find $\mu = \mu_1 = 0$ and $\sigma^2 = 0.64156$.
 Its graph is shown in Figure 15.




Figure 15. Graph of the function $F(t) = \exp[-(\sinh(t^4 - 1/2))^2]$.

The percentage deviation between $F(t)$ and the relevant expansion $\tilde{F}(t)$ in terms of Bell’s polynomials is shown in Figure 16. As can be seen, $F(t)$ is nicely approximated by $\tilde{F}(t)$ in the domain where $F(t)$ does not assume negligibly small values.




Figure 16. Relative percentage error of the Bell’s approximation of $F(t)$ in (31).

The table containing the first few moments of the considered function $F(t)$, as computed using the Gauss–Kronrod quadrature rule (G-KQR), μ_k , and the proposed Bell’s method, $\tilde{\mu}_k$, is reported in Figure 17.

$\mu[1] = 0$	$\tilde{\mu}[1] = 0$
$\mu[2] = 0.411599$	$\tilde{\mu}[2] = 0.411555$
$\mu[3] = 0$	$\tilde{\mu}[3] = 2.3822 \times 10^{-10}$
$\mu[4] = 0.291238$	$\tilde{\mu}[4] = 0.291172$
$\mu[5] = 0$	$\tilde{\mu}[5] = 4.61546 \times 10^{-10}$
$\mu[6] = 0.244599$	$\tilde{\mu}[6] = 0.244502$
$\mu[7] = 0$	$\tilde{\mu}[7] = 7.23248 \times 10^{-10}$
$\mu[8] = 0.225444$	$\tilde{\mu}[8] = 0.225301$
$\mu[9] = 0$	$\tilde{\mu}[9] = 1.07339 \times 10^{-9}$
$\mu[10] = 0.22094$	$\tilde{\mu}[10] = 0.220728$

Figure 17. Moments of $F(t)$ in (31) as computed using the G-KQR (left) and the proposed Bell's polynomial approximation (right) methods

7. Conclusions

We have shown how Bell's polynomials can be adopted to evaluate the moments of generalized Gaussian distributions originated by the composition of the classical Gauss exponential. The possibility of computing integrals of composite and higher-order nested analytic functions by means of Bell's polynomials allows for tackling the considered application in a very effective way.

By using the generalization of the Gaussian distribution, as proposed in this research study and illustrated with several examples, one can define a wide variety of curves starting from arbitrary coefficients of series expansions. Different bell-shaped functions can thus be obtained that can be adapted to represent the most diverse probabilistic distributions. Numerical calculations using the computer algebra program Mathematica have shown that our approach for approximating moments is accurate and efficient.

Author Contributions: Conceptualization, P.E.R., D.C. and R.S.; Data curation, R.S.; Investigation, P.E.R. and D.C.; Methodology, P.E.R.; Software, D.C.; Supervision, R.S.; Validation, D.C.; Writing—original draft, P.E.R. and D.C.. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

$$B_1 = g_1,$$

$$B_2 = g_1^2 + g_2,$$

$$B_3 = g_1^3 + 3g_1g_2 + g_3,$$

$$B_4 = g_1^4 + 6g_1^2g_2 + 4g_1g_3 + 3g_2^2 + g_4,$$

$$B_5 = g_1^5 + 10g_1^3g_2 + 15g_1g_2^2 + 10g_1^2g_3 + 10g_2g_3 + 5g_1g_4 + g_5,$$

$$B_6 = g_1^6 + 15g_1^4g_2 + 45g_1^2g_2^2 + 15g_2^3 + 20g_1^3g_3 + 60g_1g_2g_3 + 10g_3^2 + 15g_1^2g_4 + 15g_2g_4 + 6g_1g_5 + g_6,$$

$$B_7 = g_1^7 + 21g_1^5g_2 + 105g_1^3g_2^2 + 105g_1g_2^3 + 35g_1^4g_3 + 210g_1^2g_2g_3 + 105g_2^2g_3 + 70g_1g_3^2 + 35g_1^3g_4 + 105g_1g_2g_4 + 35g_3g_4 + 21g_1^2g_5 + 21g_2g_5 + 7g_1g_6 + g_7,$$

$$B_8 = g_1^8 + 28g_1^6g_2 + 210g_1^4g_2^2 + 420g_1^2g_2^3 + 105g_2^4 + 56g_1^5g_3 + 560g_1^3g_2g_3 + 840g_1g_2^2g_3 + 280g_1^2g_3^2 + 280g_2g_3^2 + 70g_1^4g_4 + 420g_1^2g_2g_4 + 210g_2^2g_4 + 280g_1g_3g_4 + 35g_4^2 + 56g_1^3g_5 + 168g_1g_2g_5 + 56g_3g_5 + 28g_1^2g_6 + 28g_2g_6 + 8g_1g_7 + g_8,$$

$$B_9 = g_1^9 + 36g_1^7g_2 + 378g_1^5g_2^2 + 1260g_1^3g_2^3 + 945g_1g_2^4 + 84g_1^6g_3 + 1260g_1^4g_2g_3 + 3780g_1^2g_2^2g_3 + 1260g_2^3g_3 + 840g_1^3g_3^2 + 2520g_1g_2g_3^2 + 280g_3^3 + 126g_1^5g_4 + 1260g_1^3g_2g_4 + 1890g_1g_2^2g_4 + 1260g_1^2g_3g_4 + 1260g_2g_3g_4 + 315g_1g_4^2 + 126g_1^4g_5 + 756g_1^2g_2g_5 + 378g_2^2g_5 + 504g_1g_3g_5 + 126g_4g_5 + 84g_1^3g_6 + 252g_1g_2g_6 + 84g_3g_6 + 36g_1^2g_7 + 36g_2g_7 + 9g_1g_8 + g_9,$$

$$B_{10} = g_1^{10} + 45g_1^8g_2 + 630g_1^6g_2^2 + 3150g_1^4g_2^3 + 4725g_1^2g_2^4 + 945g_2^5 + 120g_1^7g_3 + 2520g_1^5g_2g_3 + 12600g_1^3g_2^2g_3 + 12600g_1g_2^3g_3 + 2100g_1^4g_3^2 + 12600g_1^2g_2g_3^2 + 6300g_2^2g_3^2 + 2800g_1g_3^3 + 210g_1^6g_4 + 3150g_1^4g_2g_4 + 9450g_1^2g_2^2g_4 + 3150g_2^3g_4 + 4200g_1^3g_3g_4 + 12600g_1g_2g_3g_4 + 2100g_3^2g_4 + 1575g_1^2g_4^2 + 1575g_2g_4^2 + 252g_1^5g_5 + 2520g_1^3g_2g_5 + 3780g_1g_2^2g_5 + 2520g_1^2g_3g_5 + 2520g_2g_3g_5 + 1260g_1g_4g_5 + 126g_5^2 + 210g_1^4g_6 + 1260g_1^2g_2g_6 + 630g_2^2g_6 + 840g_1g_3g_6 + 210g_4g_6 + 120g_1^3g_7 + 360g_1g_2g_7 + 120g_3g_7 + 45g_1^2g_8 + 45g_2g_8 + 10g_1g_9 + g_{10}.$$

References

- Bell, E.T. Exponential polynomials. *Ann. Math.* **1934**, *35*, 258–277. [[CrossRef](#)]
- Riordan, J. *An Introduction to Combinatorial Analysis*; John Wiley & Sons: Hoboken, NJ, USA, 1958.
- Faà di Bruno, F. *Théorie des Formes Binaires*; Brero: Turin, Italy, 1876.
- Comtet, L. *Advanced Combinatorics: The Art of Finite and Infinite Expansions*; Springer: Berlin, Germany, 1974.
- Robert, D. Invariants orthogonaux pour certaines classes d'opérateurs. *Annales Mathém. Pures Appl.* **1973**, *52*, 81–114.
- Fujiwara, D. Generalized Bell's polynomials. *Sugaku* **1990**, *42*, 89–90.
- Kim, H.K. Fully degenerate Bell polynomials associated with degenerate Poisson random variables. *Open Math.* **2021**, *19*, 284–296. [[CrossRef](#)]
- Rai, P.N.; Singh, S.N. Generalization of Bell polynomials and related operatorial formula (in Hindi). *Vijnana Parishad Anusandhan Patrika* **1982**, *25*, 251–258.
- Natalini, P.; Ricci, P.E. Higher order Bell polynomials and the relevant integer sequences. *Appl. Anal. Discrete Math.* **2017**, *11*, 327–339. [[CrossRef](#)]
- Caratelli, D.; Srivastava, R.; Ricci, P.E. The Laplace Transform of composite functions and Bivariate Bell's polynomials. *Axioms* **2022**, *11*, 591. [[CrossRef](#)]
- Duran, U.; Acikgoz, M. A new approach to the Poisson distribution. *J. Ineq. Special Func.* **2020**, *11*, 1–11.
- Roman, S.M.; Rota, G.C. The umbral calculus. *Adv. Math.* **1978**, *27*, 95–188. [[CrossRef](#)]
- Roman, S.M. The Faà di Bruno Formula. *Amer. Math. Monthly* **1980**, *87*, 805–809. [[CrossRef](#)]
- Qi, F.; Niu, D.-W.; Lim, D.; Yao, Y.-H. Special values of the Bell's polynomials of the second kind for some sequences and functions. *J. Math. Anal. Appl.* **2020**, *491*, 124382. [[CrossRef](#)]
- Orozco López, R. Solution of the Differential Equation $y^{(k)} = e^{ay}$, Special Values of Bell's polynomials, and (k, a) -Autonomous Coefficients. *J. Integer Seq.* **2021**, *24*, 21.8.6
- Natalini, P.; Ricci, P.E. An Extension of the Bell's polynomials. *Comput. Math. Appl.* **2004**, *47*, 719–725. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.