

2023

## The Relationship Between Diet and Lifestyle Behaviours in a Sample of Higher Education Students; A Cross-Sectional Study

Stephen Doak

John Kearney

Jacqueline M. McCormack

*See next page for additional authors*

Follow this and additional works at: <https://arrow.tudublin.ie/scschbioart>



Part of the [Public Health Commons](#)



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 4.0 International License](#).  
Funder: Stephen Doak was a recipient of the IT Sligo Presidents Bursary award; no external funding was sought.

---

**Authors**

Stephen Doak, John Kearney, Jacqueline M. McCormack, and Laura Keaver



## Original article

## The relationship between diet and lifestyle behaviours in a sample of higher education students; a cross-sectional study

Stephen Doak<sup>a</sup>, John M. Kearney<sup>b</sup>, Jacqueline M. McCormack<sup>c</sup>, Laura Keaver<sup>a,\*</sup><sup>a</sup> Department of Health and Nutritional Science, Atlantic Technological University, Ash Lane, Sligo F91 YW50, Ireland<sup>b</sup> School of Biological and Health Sciences, Technological University Dublin, Dublin City Campus, Grangegorman, Dublin 7, Ireland<sup>c</sup> Vice President for Equality, Diversity & Inclusion and Online Development, Atlantic Technological University, Ash Lane, Sligo F91 YW50, Ireland

## ARTICLE INFO

## Article history:

Received 29 August 2022

Accepted 28 January 2023

## Keywords:

Diet  
Fruit  
Vegetables  
Life style  
Sedentary behaviour  
Body mass index  
Alcohol drinking in college  
Stress  
Sleep quality  
Smoking  
Associations  
Students

## SUMMARY

**Background & aims:** Transitioning into higher education (HE) impacts health behaviours. Poor dietary and lifestyle behaviours may correlate and increase risk of co-morbidities. The introduction of the Okanagan Charter detailed the important role of health promotion within a HE setting. The aim of this study was to assess the relationship between dietary quality and lifestyle behaviours of students attending HE.

**Methods:** Full-time students, aged 18+, were eligible to participate in this online cross-sectional study. Self-reported questions were asked in relation to demographics, body mass index (BMI), smoking, and COVID-19. A food frequency questionnaire measured dietary quality along with tools assessing alcohol use, sleep quality, perceived stress, and physical activity. Statistical analyses were performed using chi-square, one-way ANOVA, independent sample t-tests, Pearson's correlation, and multivariate linear regression.

**Results:** Evidence of a correlation between poor diet quality and having a higher BMI ( $p = 0.040$ ), higher alcohol consumption ( $p = <0.001$ ), poorer sleep quality ( $p = 0.003$ ), higher stress levels ( $p = 0.006$ ) and smoking ( $p = 0.001$ ) was found. Low fruit and vegetable consumption were associated with higher BMI ( $p = 0.013$ ), higher alcohol consumption ( $p = <0.001$ ), lower physical activity levels ( $p = 0.006$ ), higher stress levels ( $p = <0.001$ ), smoking ( $p = <0.001$ ) and being male ( $p = 0.002$ ).

**Conclusions:** This study provides data on the association between dietary quality and lifestyle behaviours among HE students and will inform healthy campus initiatives.

© 2023 The Authors. Published by Elsevier Ltd on behalf of European Society for Clinical Nutrition and Metabolism. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

An unhealthy diet is a contributor to higher BMI, increased NCDs risk, decreased productivity, and premature morbidity and mortality [1–6], leading to obesity, diabetes, CVD and hypertension [6–8] through inflammation [6], and not consuming substantial amounts of micronutrients [7] and food compounds e.g., phytochemicals [8]. In 2019, globally NCDs accounted for 7.9 million deaths and 187.7 million Disability-adjusted life years (DALYs) attributable to an unhealthy diet [9]. A healthy diet is defined as consuming food in a pattern that is beneficial to health, or at least not harmful [10,11], common features include a higher proportion of plant-based foods, F&V, whole grains, legumes, seeds, nuts, polyunsaturated fatty acids; an unhealthy diet often contains high amounts of sodium, trans fats, and red meat [8–14]. Predictors of dietary change include health-related lifestyle behaviours [1] that can be defined as daily behavioural choices that affect one's overall

health [15]. PA, diet, alcohol consumption and smoking remain the most studied modifiable lifestyle behaviours and are associated with all-cause mortality [16–19].

Many students do not achieve the RDA for whole grains [20] and over 82% do not adhere to F&V guidelines [21–23] increasing the likelihood of overweight and obesity (OWO) and other NCDs [24]. Among Irish students the median number of daily F&V servings was three, 38% were OWO, 9% reach PA guidelines, and 25–33% smoked tobacco [25,26]. Hazardous alcohol consumption is the main substance used by higher education (HE) students [27] with 54–66% of Irish students deemed hazardous alcohol consumers [25,27] and 25–34% are physically inactive [25,28]. Students eating behaviours are associated with poor sleep quality correlating with unhealthy lifestyle behaviours [29–31]. These behaviours affect the health of students, as does smoking, OWO and stress [32,33].

Happier students have been shown to be more active [34], and those stressed have a high BMI [35]. Students experience elevated levels of stress [36,37], which is associated with an unhealthy diet and poor academic performance [38,39]; increasing during HE [40]. Academic impairment increases college dropout rates, therefore, a

\* Corresponding author.

E-mail address: [laura.keaver@atu.ie](mailto:laura.keaver@atu.ie) (L. Keaver).

### Abbreviations

BMI	Body Mass Index
CVD	Cardiovascular Disease
F&V	Fruit and Vegetables
HE	Higher Education
HEI	Higher Education Institute
GDP	Gross Domestic Product
NCD	Non-Communicable Diseases
OWO	Overweight and Obesity
PA	Physical Activity
RDA	Recommended Daily Amounts

prominent issue for HE Institutes (HEIs) to consider [41]. An international study found that poorer dietary behaviours were associated with poorer sleep quality and less PA, but not alcohol misuse [34,42]. Globally among students a correlation between higher diet quality and both higher PA levels and lower alcohol consumption was found [43]. Smoking status has a more inconclusive relationship with diet; sleep appears to be trending towards a correlation [43].

Unhealthy lifestyle behaviours decrease the number of disease-free years a person has, and increases the risk of NCDs, mortality, and weight gain [1,2,6,19,44]. The life course approach to health has been supported in recent years [1,3,45]. Although healthy diet and lifestyle behaviour interventions of college students are understudied [43], their development is imperative for the improvement of student's health; their potential impact on subsequent stages of life can reduce the onset of NCDs improve health later in life [1,3,45]. There are 231,710 students enrolled in Irish HEIs which has been identified as a key setting to target and implement health-promoting programmes and research [46–48]. During the transition into HE lifelong health-related behaviours are established and are likely to continue throughout life, affecting future health status [49–51].

The association between diet and lifestyle behaviours of HE students is an area with little evidence in Ireland or globally and their relationship is unclear within this cohort [43,52]. This study aimed to identify the relationship between dietary quality and lifestyle behaviours of students and has the potential to shape the development and content of healthy campus policy in HEIs.

## 2. Methods

Data were collected as part of a cross-sectional web-based survey, hosted by Qualtrics<sup>®</sup> during the 2020–21 academic year. A hyperlink was circulated through class representative systems, students' unions, social media, wellness officers and email. Students registered for full-time undergraduate or graduate programmes, aged 18 and over, were eligible to participate from three HEIs in the North-West of Ireland. Students provided informed consent and participation was voluntary and anonymous with the opportunity to exit the survey any time prior to submission. The median time for completion was 14 min, 52.5 s. A total of 956 students responded to the survey, with 682 (71.3% completion rate) used in the final analyses.

### 2.1. Measures and materials

The questionnaire consisted of forty-four questions and available in [supplementary material 1](#). Demographic characteristics assessed included age, gender, year of study, area of study, living situation,

current employment situation, dietary preferences (e.g., vegan, vegetarian, gluten-free, etc.) and dietary supplementation use (e.g., creatine, protein powder, multivitamins, etc.). A question was adapted from a previous Irish study of students [28] asking "How often do you prepare food from fresh ingredients rather than pre-prepared food?" with answers categorised on a Likert scale, as 'never', '1–2 times a week', '3–4 times a week', 'every day' or 'multiple times a day'. Self-reported general health and diet were measured using single item questions adapted from previous research of Irish students [25]. The questions were "In general, would you say your health/diet is:" with answers on a 5-point Likert Scale: 'very good', 'good', 'neither good nor poor', 'poor' and 'very poor'. BMI was estimated from self-reported height and weight with categories defined according to the World Health Organization criteria. Underweight, normal weight, overweight and obesity defined as a BMI of  $\leq 18.4$ , 18.5–24.99, 25–29.99 and  $\geq 30$  kg/m<sup>2</sup>, respectively [53]. Due to the unfeasibility of measuring anthropometric data during COVID-19, self-reported measures were used, a validated method for students [54]. Smoking status was assessed with one question "Which of the following best describes your smoking (cigarettes/vape) habits?" with four potential answers; 'non-smoker', 'ex-smoker', 'smoker' and 'social smoker (e.g., not every day)'.

A Short-Form Food Frequency Questionnaire (SFFQ) was used in this study which is reliable and valid in comparison to a longer FFQ [55] and has been previously used in a student population [56]. The SFFQ measures five components: fruit, vegetable, oily fish, fat, and non-milk extrinsic sugar providing a total diet quality score (DQS) ranging from 5 to 15, with fifteen deemed optimum diet quality [55]. Daily F&V consumption was determined by calculating the total fruit, fruit juice, salad, and vegetables consumed, with fruit juice limited to 1.0 portion, as per recommendations [57]. Four validated and reliable tools were used to assess behaviours: Alcohol Use Disorders Identification Test (AUDIT-C) questionnaire [27,58], Pittsburgh Sleep Quality Index (PSQI) [59–61], Perceived Stress Scale-10 (PSS-10) [43,62], and International Physical Activity Questionnaire-Short Form (IPAQ-SF) [63–65]. Habitual sleep duration was extracted from the PSQI; participants were classified as meeting sleep duration guidelines if they slept a total of seven or more hours a night [61]. To coincide with previous Irish research, those that report 'high' activity levels (1500 MET vigorous or 3000 MET moderate minutes of activity a week) are deemed to be meeting Irish national PA guidelines [65,66]. Seven questions were adapted from previous research about the impact of the COVID-19 Pandemic [43].

### 2.2. Statistical analyses

Data was exported directly to Microsoft Excel (Microsoft Inc., Redmond, WA, USA) and then to IBM Statistical Package for the Social Sciences [SPSS] Version 26.0 (IBM Inc., Armonk, NY, USA). Descriptive statistics were performed, and data were presented as count (%) or mean  $\pm$  standard deviations (SD). Chi-square tests of association (for categorical variables) and independent t-tests were used to check for associations between two categorical variables. All cells had an expected count of greater than or equal to five. Z-tests for differences in the two proportions determined if a statistically significant difference existed between the proportions of a column for each category. Pearson's product-moment correlations were run to assess the relationship between variables. A one-way ANOVA was performed to assess the relationship between smoking and both diet quality and F&V consumption. Multivariate linear regression was used to identify significant predictors of diet quality and F&V consumption. The significance levels for all tests were set at an alpha level of  $P < 0.05$ .

### 3. Results

The mean age of participants was 24.97 ± 8.90 and the majority were female (n = 455, 66.8%). The mean BMI was 24.67 ± 4.49 with (n = 355, 54.9%) classified as a healthy weight, 40.2% (n = 260) were OWO and 4.9% (n = 32) underweight. Only 39.6% (n = 269) of students reported being somewhat or very satisfied with their current weight, 45.2% (n = 307) indicated they are not. Further demographic information can be found in [Supplementary Material 2](#).

#### 3.1. Dietary intake and lifestyle behaviours

The sample in this study had an overall diet quality score of 9.97 ± 1.89. A small number (n = 147, 21.6%) of students consume the RDA of at least five servings of F&V; the average daily F&V consumption was 3.17 ± 3.07 (253.6g). A total of 26.7% (n = 181) outline that they have a dietary requirement/restriction and 52.5% (n = 355) of participants use dietary supplements. Hazardous alcohol consumption is seen in 39.0% (n = 266) of students surveyed. PSQI scores indicate that 45.2% (n = 304) have poor sleep quality and 32.9% (n = 219) do not meet sleep recommendations. Mean PSS-10 scores were 16.78 ± 5.37 with over two-thirds (n = 500, 73.3%) deemed moderate-highly stressed. Mean IPAQ scores were 2780.05 ± 2562.40 with 38.3% (n = 255) meeting Irish PA guidelines. Over a quarter (n = 182, 26.7%) reported being smokers. Further diet and lifestyle behaviour information can be found in [Supplementary Material 3](#).

#### 3.2. The Impact of COVID-19 on Diet and Lifestyle Behaviours

When asked about the impact of COVID-19 on dietary behaviours 29.5% (n = 195) reported that their diet is less healthy and 26.4% (n = 174) reported eating more healthily. Almost half (n = 332, 48.8%) reported drinking less, and sleep was negatively impacted in 39.0% (n = 266) of students. Most students (n = 411, 62.1%) feel more stressed than usual, and 54.8% (n = 372) reported less PA, as seen in supplementary material 4.

#### 3.3. Diet quality and lifestyle behaviours

A Pearson's product-moment correlation, as seen in [Table 1](#), was run and found a statistically significant, small (weak) negative correlations between diet quality and BMI, alcohol consumption, sleep quality, and stress and none between diet quality and PA (p = 0.101). A one-way ANOVA found a statistically significant difference among the four smoking groups (F(3, 677) = 5.592, p = 0.001). Tukey post hoc test analysis revealed a statistically significant increase in the diet quality of ex-smokers compared to non-smokers (0.813, 95% CI (0.13 to 1.50), p = 0.012), as well as from ex-smokers compared to smokers (1.34, 95% CI (0.49 to 2.19), p < 0.001), but no other group differences were statistically significant (not in table).

**Table 1**  
Diet quality and its association with lifestyle behaviours-correlation co-efficient.

	Mean	SD	Correlations	p-Value
Diet Quality	9.97	1.89		
BMI	24.67	4.49	-0.081*	0.040
AUDIT-C Score	3.85	2.70	-0.163***	<0.001
PSQI Score	4.70	2.46	-0.115**	0.003
PSS-10 Score	16.78	5.37	-0.104**	0.006
IPAQ Score	2780.05	2562.40	0.064	0.101

\*Correlation Significant at 0.05 level (two tailed); \*\*Correlation Significant at 0.01 level (two tailed); \*\*\*Correlation Significant at 0.001 level (two tailed).

**Table 2**  
Multiple regression results for diet quality.

Diet Quality	B	SE B	β	R <sup>2</sup>	ΔR <sup>2</sup>
Model				0.036	0.033***
Constant	10.739***	10.388	0.179		
AUDIT-C	-0.108***	-0.162	0.027	-0.153***	
PSQI	-0.075*	-0.133	0.030	-0.097**	

\*Correlation Significant at 0.05 level (two tailed); \*\*Correlation Significant at 0.01 level (two tailed); \*\*\*Correlation Significant at 0.001 level (two tailed); B = unstandardized regression coefficient; CI = confidence interval; L = Limit; SE B = standard error of the coefficient; β = standardized coefficient; R<sup>2</sup> = coefficient of determination ΔR<sup>2</sup> = adjusted R<sup>2</sup>.

**Table 3**  
F&V consumptions association with lifestyle behaviours - correlation co-efficient.

Variable	Mean	SD	Correlation	p-Value
F&V Consumption	3.17	3.07		
BMI	24.67	4.49	-0.098*	0.013
AUDIT-C Score	3.85	2.70	-0.149***	<0.001
PSQI Score	4.70	2.46	-0.052	0.176
PSS-10 Score	16.78	5.37	-0.158***	<0.001
IPAQ Score	2780.05	2562.40	0.107**	0.006

\*Correlation Significant at 0.05 level (two tailed); \*\*Correlation Significant at 0.01 level (two tailed); \*\*\*Correlation Significant at 0.001 level (two tailed).

A multiple regression was run to predict diet quality from BMI, AUDIT-C, PSQI and PSS-10 scores. BMI (p = 0.119) and PSS-10 scores (p = 0.083) were removed from the model and all assumptions were met. The model statistically significantly predicted diet quality (F(2, 669) = 12.580, p < 0.001), adj. R<sup>2</sup> = -0.033), seen in [Table 2](#). A higher diet quality is associated with having better sleep quality and lower alcohol use, accounting for 3.6% variation. For every one-point increase in AUDIT-C and PSQI scores diet quality scores reduce by 0.108 and 0.075, respectively.

#### 3.4. F&V consumption and lifestyle behaviours

A Pearson's product-moment correlation, as seen in [Table 3](#), was run and found a statistically significant, small (weak) negative correlation between F&V consumption and BMI, alcohol consumption, stress, and PA. There was no statistically significant correlation between F&V consumption and sleep quality (p = 0.176). A one-way ANOVA found a statistically significant difference among the four smoking groups (F(3, 671) = 7.839, p < 0.001). Games-Howell post hoc test analysis revealed a statistically significant increase in F&V consumption of non-smokers compared to smokers (1.23, 95% CI (0.50 to 1.96), p < 0.001), as well as from ex-smokers compared to smokers (1.86, 95% CI (0.57 to 3.14), p = 0.002), but no other group differences were statistically significant (not in table).

A multiple regression was run to predict F&V consumption from gender, AUDIT-C scores, PSQI scores and PSS-10 scores. BMI (p = 0.060) and IPAQ scores (p = 0.068) were removed from the

**Table 4**  
Multiple regression results for F&V consumption.

F&V Consumption	B	SE B	β	R <sup>2</sup>	ΔR <sup>2</sup>
Model				0.055	0.050**
Constant	2.889**	2.081	0.411		
Gender	0.766**	0.404	0.184	0.162**	
AUDIT-C	-0.080*	-0.142	0.032	-0.097**	
PSS-10	-0.067**	-0.099	0.016	-0.157*	

\*Correlation Significant at 0.05 level (two tailed); \*\*Correlation Significant at 0.01 level (two tailed); B = unstandardized regression coefficient; CI = confidence interval; L = Limit; SE B = standard error of the coefficient; β = standardized coefficient; R<sup>2</sup> = coefficient of determination R<sup>2</sup> = adjusted R<sup>2</sup>.



model and all assumptions were met. The model statistically significantly predicted F&V consumption ( $F(3, 642) = 12.390, p < 0.001$ ),  $adj. R^2 = .05$ ), as seen in Table 4. Consuming more F&V is associated with being female, having a lower AUDIT-C and PSS-10 score accounting for 5.5% variation. For every one-point increase in AUDIT-C and PSS-10 score, F&V consumption decreases by 0.080 and 0.067 portions per day, respectively. Females consume 0.766 F&V portions more per day than males.

#### 4. Discussion

This study found a higher prevalence of students consuming the RDA of F&V (22%) than others [21–23]. Although results are not directly comparable, other studies found that the mean results of students' diet quality were close to the midpoint of potential scores [29,42]. The mean BMI ( $24.67 \pm 4.49$ ) and the amount classified as OWO (40%) within this study is similar to the global range [67–69], but lower than the 60% of Irish adults deemed OWO [70]. This study found that BMI is associated with both diet quality and F&V consumption and a lower percentage of obese students met F&V guidelines, a trend well documented in studies of adults including students [43,71,72]. Improving dietary behaviours could be a positive approach to reducing students BMI [47].

Irish studies previously found that 26–64% of students were classified as meeting PA guidelines [73–77], however, PA levels in this study sample (38.3%) were considerably lower. This could be due to the reduction of PA seen in students due to COVID-19 with the closure of facilities, sports being cancelled and lockdown measures [78,79]. In this study PA is associated with F&V consumption but not diet quality. Three studies that assessed overall diet quality in a recent SR did not find an association with PA, however, all eleven papers that assessed diet quality via F&V consumption found some form of a correlation [43]. This suggests F&V consumption is a good predictor of diet quality in students. Academic pressure and stress are predictors of PA [32]; the high level of stress among this cohort could be why PA levels are low [21,33,74].

The mean PSS-10 score of  $16.78 \pm 5.37$  found is similar or less than recent studies [42,80–82]. However, 73% of participants are experiencing moderate to high levels of stress [83], much higher than the 17% in a study of 3440 Irish students [84]. Stress levels of students are higher than that of the general population impacting their academic performance and health [37,85]. This study found that stress is associated with both diet quality and F&V consumption, the strength of which can be seen in the regression model. Students have been found to both increase and reduce food consumption because of stress [38], causing weight change [37]. The association between stress levels of students and dietary habits has been well documented [38–40,86]. Healthier diet and lifestyle behaviours are associated with lower levels of stress in students [33], including sleep [42].

Almost half of students within this study had poor sleep quality; similar to other studies deeming 30–64% as poor sleepers [42,82,87]. Poor sleep quality is a risk factor for adverse health outcomes [88]; adherence to a healthier diet is associated with higher sleep quality [89,90]. Six of ten papers identified in a recent SR found a significant association between diet and sleep [43]. This study found an association between sleep quality and diet quality, however, not between sleep and F&V consumption, despite a "healthy" diet, generally categorised as including high F&V consumption [89]. Di Benedetto et al., noted that theirs and other studies found no association between sleep quality and F&V consumption, however, they did find an association between overall diet and sleep quality [91]. This indicates that more thorough dietary assessment may yield more conclusive results, with the

strength of association between diet and sleep quality seen in the regression model. Poor sleep is identified as a common health problem among students, impacted by academic demands [92] and lifestyle behaviours e.g., smoking [93].

Over one-quarter of students self-reported as smokers, a rate similar to the 17–33% in other studies [25–28,73,76] yet much higher than the 17% of Irish adults deemed smokers [70]. Smoking patterns among young Irish people differ from national rates with a high prevalence of social smokers, also seen in this study [94]. In a recent SR 39% of papers found a significant relationship between poor diet quality and being a smoker [43]. This study found a higher diet quality among ex-smokers, whereas high F&V consumption was found among both ex- and non-smokers. Many studies assessing overall diet quality find no association with smoking and potentially why F&V is more conclusive within this study [43]. Studies of the general population have found that non- or ex-smokers engage in healthier diet behaviours [95]. Although smokers tend to have a lower quality diet this relationship may be dependent on the amount of tobacco consumed [96]. A low number of students identified as smokers and ex-smokers and frequency was not assessed, potentially why associations were not found. Higher rates of alcohol consumption within a student cohort may be a reason for higher smoking prevalence [94].

In this study, 39% were deemed hazardous alcohol consumers, lower than the 61–84% previously found among Irish students [27,49], yet, similar to recent European research that found 37% of 2191 students are hazardous consumers [97]. Alcohol consumption among Irish adults varied since COVID-19 with some finding a decrease of 66% [98] and others finding an increase [99]. Studies of HE students have found a reduction in alcohol consumption during the COVID-19 pandemic [36]. Recent publications, found that poorer dietary behaviours were not correlated with alcohol misuse [35,42]. Studies finding no correlation between alcohol consumption and diet may be due to alcohol being a societal norm to the extent that it may be independent of other factors [100]. The association between alcohol and both diet quality and F&V in this study is strengthened by its inclusion in both multiple regression models. This association could be strengthened by the fact that although alcohol was reportedly consumed less than usual due to COVID-19 by almost half of students, there is still a high number of hazardous consumers, which has been found in other studies of students during COVID-19 measures [101].

##### 4.1. Strengths and limitations

The high number of responses and the number of variables explored heighten the robustness of results, strengthened using validated measures. This study included students from multiple HEIs and both undergraduate and graduate students, increasing generalizability of results. Limitations must be considered, firstly, this study had a cross-sectional design, therefore, causal relationships could not be identified. Results cannot be generalized to all students as a convenience sample was used, however, results are important for informing local policies and initiatives. Variables were all measured using subjective, self-report tools, limiting the validity and reliability of the data e.g., PA questions were incorrectly completed by many participants, making analysing difficult. Although the SFFQ is validated, the 'gold standard' of dietary surveillance is a 7-day weighed food diary [102], however, it has been deemed a burdensome and impractical method [103].

##### 4.2. Future use of findings

HE students are attending a setting that often has high-quality facilities, technology, and highly educated staff, therefore, ideal

for health promotion interventions and campaigns [104]. Recent SRs have been conducted to assess the effectiveness of health interventions in a HE setting with the majority focusing on a singular health outcome [52]. Several interventions are effective including in-person interventions, media approaches and nutrition labelling to improve dietary habits, face-to-face programs, internet-based approaches, personalised interventions and cognitive behavioural therapy to improve lifestyle behaviours [52]. The results from this study can be used to develop interventions that combine diet and lifestyle behaviours.

Future studies with a prospective design are needed to better understand these relationships and allow for the assessment of temporal sequence, eliminate recall bias and comparing multiple behaviours [105]. Future research could benefit from including objective measures, to reduce the likelihood of over- or under-estimation and misinterpreting behaviours [106]. Those who obtain a higher education level are more likely to have a healthier diet [107], therefore, maintaining their status as a student and targeting students is important for population health and can potentially improve a countries GDP [107]. HEIs should be proactive in their approach to the health of students and create an interdisciplinary health promotion team to create educational programmes that support a healthy lifestyle [35,42].

## 5. Conclusion

There is evidence of a correlation between diet quality and BMI, alcohol consumption, sleep quality, smoking and stress. F&V consumption is associated with BMI, smoking, alcohol consumption, PA, and stress. This study supports the need for healthy campus committees to plan interventions and educational programmes on diet, health, and lifestyle behaviours. These could help create healthier institutes and graduates, while improving supports and services for students. The results indicate that improving the diet quality of students may also improve their lifestyle behaviours and vice versa, with academic achievement potentially benefiting as a result.

## Ethical approval

The study was conducted in accordance with the Declaration of Helsinki. Ethical approval was received by [removed for blind peer review] prior to data collection.

## Authors' contributions

SD, LK, JMM and JMK conceptualized and designed the study. SD and LK drafted the introduction section. SD conducted the survey, collected, extracted and analysed the data. SD and LK drafted the methods, results, discussion and conclusion section. All authors contributed to the writing and editing of the manuscript. All authors read and approved the final manuscript.

## Funding

Stephen Doak was a recipient of the IT Sligo Presidents Bursary award; no external funding was sought.

## Declaration of competing interest

Authors state no conflicts of interest.

## Acknowledgments

The authors wish to thank Jason Quinn, Keelan Kennoy and Ryan O'Reilly from ITS Students' Union, Yvonne Roache from ITS, Colin

Kearney and Victor O'Loughlin from GMIT Students' Union and Mary Hernandez from LYIT Students' Union for helping with survey promotion. Thank you to Padraig McGrouarty for his help with statistical analysis. In addition, the authors thank the students who made the study possible by volunteering their time.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.clnesp.2023.01.036>.

## References

- [1] Thorpe MG, Milte CM, Crawford D, McNaughton SA. Education and lifestyle predict change in dietary patterns and diet quality of adults 55 years and over. *Nutr J* 2019;18:67. <https://doi.org/10.1186/s12937-019-0495-6>.
- [2] Gakidou E, Afshin A, Abajobir AA, Abate KH, Abbafati C, Abbas KM, et al. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;390:1345–422. [https://doi.org/10.1016/S0140-6736\(17\)32366-8](https://doi.org/10.1016/S0140-6736(17)32366-8).
- [3] Jacob CM, Baird J, Barker M, Cooper C, Hanson M. The importance of a life-course approach to health: chronic disease risk from preconception through adolescence and adulthood: white 211. World Health Organization; 2017. p. 41. paper. <https://www.who.int/lifecourse/publications/life-course-approach-to-health.pdf>.
- [4] Khandelwal S, Kurpad A, Narayan KMV. Global non-communicable diseases—the nutrition conundrum. *Front Public Health* 2018;6:9. <https://doi.org/10.3389/fpubh.2018.00009>.
- [5] Stephen AM, Champ MM, Cloran SJ, Fleith M, van Lieshout L, Mejbourn H, et al. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev* 2017;30(2):149–90. <https://doi.org/10.1017/S095442241700004X>.
- [6] Hariharan R, Odjidja EN, Scott D, Shivappa N, Hébert JR, Hodge A, et al. The dietary inflammatory index, obesity, type 2 diabetes, and cardiovascular risk factors and diseases. *Obes Rev* 2022;23(1):e13349. <https://doi.org/10.1111/obr.13349>.
- [7] Mansouri M, Pahlavani N, Sharifi F, Varmaghani M, Shokri A, Yaghubi H, et al. Dairy consumption in relation to hypertension among a large population of university students: the MEPHASOUS study. *Diabetes Metab Syndr Obes* 2020;13:1633–42. <https://doi.org/10.2147/DMSO.S248592>.
- [8] Nattagh-Eshstivani E, Barghchi H, Pahlavani N, Barati M, Amiri Y, Fadel A, et al. Biological and pharmacological effects and nutritional impact of phytosterols: a comprehensive review. *Phytother Res* 2022;36(1):299–322. <https://doi.org/10.1002/ptr.7312>.
- [9] Qiao J, Lin X, Wu Y, Huang X, Pan X, Xu J, et al. Global burden of non-communicable diseases attributable to dietary risks in 1990–2019. *J Hum Nutr Diet* 2022;35:202–13. <https://doi.org/10.1111/jhn.12904>.
- [10] de Ridder D, Kroese F, Evers C, Adriaanse M, Gillebaart M. Healthy diet: health impact, prevalence, correlates, and interventions. *Psychol Health* 2017;32:907–41. <https://doi.org/10.1080/08870446.2017.1316849>.
- [11] Cena H, Calder PC. Defining a healthy diet: evidence for the role of contemporary dietary patterns in health and disease. *Nutrients* 2020;12:334. <https://doi.org/10.3390/nu12020334>.
- [12] Pistollato F, Iglesias RC, Ruiz R, Aparicio S, Crespo J, Lopez LD, et al. Nutritional patterns associated with the maintenance of neurocognitive functions and the risk of dementia and Alzheimer's disease: a focus on human studies. *Pharmacol Res* 2018;131:32–43. <https://doi.org/10.1016/j.phrs.2018.03.012>.
- [13] Turner-McGrievy G, Wirth MD, Hill KL, Dear ER, Hébert JR. Examining commonalities and differences in food groups, nutrients, and diet quality among popular diets. *Clin Nutr ESPEN* 2021;41:377–85. <https://doi.org/10.1016/j.clnesp.2020.10.017>.
- [14] Tapsell LC. Dietary behaviour changes to improve nutritional quality and health outcomes. *Chronic Dis Transl Med* 2017;3(3):154–8. <https://doi.org/10.1016/j.cdtm.2017.06.005>.
- [15] Almutairi KM, Alonazi WB, Vinluan JM, Almigbal TH, Batais MA, Alodhayani AA, et al. Health promoting lifestyle of university students in Saudi Arabia: a cross-sectional assessment. *BMC Publ Health* 2018;18:1093. <https://doi.org/10.1186/s12889-018-5999-z>.
- [16] Mulder M, Ranchor AV, Sanderman R, Bouma J, van den Heuvel WJA. The stability of lifestyle behaviour. *Int J Epidemiol* 1998;27:199–207. <https://doi.org/10.1093/ije/27.2.199>.
- [17] Chudasama YV, Khunti K, Gillies CL, Dhalwani NN, Davies MJ, Yates T, et al. Healthy lifestyle and life expectancy in people with multimorbidity in the UK Biobank: a longitudinal cohort study. *PLoS Med* 2020;17:e1003332. <https://doi.org/10.1371/journal.pmed.1003332>.
- [18] Larsson SC, Kaluza J, Wolk A. Combined impact of healthy lifestyle factors on lifespan: two prospective cohorts. *J Intern Med* 2017;282:209–19. <https://doi.org/10.1111/joim.12637>.

- [19] Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: a systematic review and meta-analysis. *Prev Med* 2012;55:163–70. <https://doi.org/10.1016/j.ypmed.2012.06.017>.
- [20] Bernardo GL, Jomori MM, Fernandes AC, Proenca RP. Food intake of university students. *Rev Nutr* 2017;30:847–65. <https://doi.org/10.1590/1678-98652017000600016>.
- [21] Dinger MK, Brittain DR, Hutchinson SR. Associations between physical activity and health-related factors in a national sample of college students. *J Am Coll Health* 2014;62:67–74. <https://doi.org/10.1080/07448481.2013.849710>.
- [22] Peltzer K, Pengpid S. Correlates of healthy fruit and vegetable diet in students in low, middle and high income countries. *Int J Publ Health* 2015;60:79–90. <https://doi.org/10.1007/s00038-014-0631-1>.
- [23] Deliens T, Verhoeven H, De Bourdeaudhuij I, Huybrechts I, Mullie P, Clarys P, et al. Factors associated with fruit and vegetable and total fat intake in university students: a cross-sectional explanatory study. *Nutr Diet* 2018;75:151–8. <https://doi.org/10.1111/1747-0080.12399>.
- [24] Stephen AM, Champ MM, Cloran SJ, Fleith M, van Lieshout L, Meijborn H, et al. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutr Res Rev* 2017;30:149–90. <https://doi.org/10.1017/S095442241700004X>.
- [25] Bickerdike A, Dinneen J, O'Neill C. A healthy CIT: an investigation into student health metrics, lifestyle behaviours and the predictors of positive mental health in an Irish higher education setting. *Int J Environ Res Publ Health* 2019;16:4318. <https://doi.org/10.3390/ijerph16224318>.
- [26] Mullaney MI, Corish CA, Loxley A. Exploring the nutrition and lifestyle knowledge, attitudes, and behaviour of student home economics teachers: baseline findings from a 4-year longitudinal study. *Int J Consum Stud* 2008;32:314–22. <https://doi.org/10.1111/j.1470-6431.2007.00650.x>.
- [27] Davoren MP, Shiely F, Byrne M, Perry IJ. Hazardous alcohol consumption among university students in Ireland: a cross-sectional study. *BMJ Open* 2015;5:e006045. <https://doi.org/10.1136/bmjopen-2014-006045>.
- [28] Murphy JJ, MacDonncha C, Murphy MH, Murphy N, Timperio A, Leech RM, et al. Identification of health-related behavioural clusters and their association with demographic characteristics in Irish university students. *BMC Publ Health* 2019;19:121. <https://doi.org/10.1186/s12889-019-6453-6>.
- [29] Aslan Çin NN, Yardımcı H. Association of total energy intake, diet quality and sleep disorders in university-term female students. *Sleep Biol Rhythm* 2021;19:313–23. <https://doi.org/10.1007/s41105-021-00320-1>.
- [30] Wang F, Biró É. Determinants of sleep quality in college students: a literature review. *Explore* 2021;17:170–7. <https://doi.org/10.1016/j.explore.2020.11.003>.
- [31] Fatima Y, Doi SAR, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias adjusted meta-analysis. *Obes Rev* 2015;16:137–49. <https://doi.org/10.1111/obr.12245>.
- [32] Aceijas C, Waldhäusl S, Lambert N, Cassar S, Bello-Corassa R. Determinants of health-related lifestyles among university students. *Perspect Public Health* 2017;137:227–36. <https://doi.org/10.1177/1757913916666875>.
- [33] Hanawi SA, Saat NZM, Zulkafly M, Hazlenah H, Taibukahn NH, Yoganathan D, et al. Impact of a healthy lifestyle on the psychological well-being of university students. *Int J Pharm Res Allied Sci* 2020;9:1–7. ISSN: 2277-3657.
- [34] Murphy MH, Carlin A, Woods C, Neville A, MacDonncha C, Ferguson K, et al. Active students are healthier and happier than their inactive peers: the results of a large representative cross-sectional study of university students in Ireland. *J Phys Activ Health* 2018;15:737–46. <https://doi.org/10.1123/jpah.2017-0432>.
- [35] Cena H, Porri D, De Giuseppe R, Kalmpourtzidou A, Salvatore FP, El Ghoch M, et al. How healthy are health-related behaviors in university students: the HOLISTIC study. *Nutrients* 2021;13:675. <https://doi.org/10.3390/nu13020675>.
- [36] Du C, Zan MCH, Cho MJ, Fenton JJ, Hsiao PY, Hsiao R, et al. Health behaviors of higher education students from 7 countries: poorer sleep quality during the COVID-19 pandemic predicts higher dietary risk. *Clocks Sleep* 2021;3:12–30. <https://doi.org/10.3390/clocksleepp3010002>.
- [37] Haidar SA, de Vries NK, Karavetian M, El-Rassi R. Stress, anxiety, and weight gain among university and college students: a systematic review. *J Acad Nutr Diet* 2018;18:261–74. <https://doi.org/10.1016/j.jand.2017.10.015>.
- [38] Elshurbjy AJ, Ellulu MS. Association between stress and dietary behaviors among university students: mini-review. *Med Clin Arch* 2017;1. <https://doi.org/10.15761/mca.1000108>.
- [39] Sprake EF, Russell JM, Cecil JE, Cooper RJ, Grabowski P, Pourshahidi LK, et al. Dietary patterns of university students in the UK: a cross-sectional study. *Nutr J* 2018;17:90. <https://doi.org/10.1186/s12937-018-0398-y>.
- [40] Kessler R, Wang P. The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annu Rev Publ Health* 2008;29:115–26. <https://doi.org/10.1146/annurev.publhealth.29.020907.090847>.
- [41] Turner K, McCarthy VL. Stress and anxiety among nursing students: a review of intervention strategies in literature between 2009 and 2015. *Nurse Educ Pract* 2017;22:21–9. <https://doi.org/10.1016/j.nepr.2016.11.002>.
- [42] Du C, Zan MCH, Cho MJ, Fenton JJ, Hsiao PY, Hsiao R, et al. The effects of sleep quality and resilience on perceived stress, dietary behaviors, and alcohol misuse: a mediation-moderation analysis of higher education students from Asia, Europe, and North America during the COVID-19 pandemic. *Nutrients* 2021;13:442. <https://doi.org/10.3390/nu13020442>.
- [43] Doak S, O'Callaghan N, Kearney J, McCormack JM, Keaver L. Lifestyle behaviors associated with dietary quality in higher education students: a systematic review. *Intern J Health Nutr Exer Sci* 2022;1:15–82. [www.gaics.org/userdata/upload/file/JNES2022\\_1\\_002.pdf](http://www.gaics.org/userdata/upload/file/JNES2022_1_002.pdf).
- [44] Nyberg ST, Singh-Manoux A, Pentti J, Madsen IEH, Sabia S, Alfredsson L, et al. Association of healthy lifestyle with years lived without major chronic diseases. *JAMA Intern Med* 2020;180:760–8. <https://doi.org/10.1001/jamainternmed.2020.0618>.
- [45] Mikkelsen B, Williams J, Rakovac I, Wickramasinghe K, Hennis A, Shin H-R, et al. Life course approach to prevention and control of non-communicable diseases. *BMJ* 2019;364:l257. <https://doi.org/10.1136/bmj.l257>.
- [46] Department of Health. Healthy Ireland - get Ireland active! National physical activity plan for Ireland. 2016. <https://assets.gov.ie/7563/23f51643fd1d4ad7abf529e58c8d8041.pdf> (Accessed November 12th, 2021).
- [47] Health Service Executive. Healthy eating and active living programme: national implementation plan 2017–2020. 2016. <https://www.hse.ie/eng/about/who/healthwellbeing/our-priority-programmes/health/health-docs/health-programme-national-implementation-plan-2017-2020.pdf> (Accessed November 12th, 2021).
- [48] Higher Education Authority. Higher education: key facts and figures. 2018. <https://hea.ie/assets/uploads/2019/01/Higher-Education-Authority-Key-Facts-Figures-2017-18.pdf> (Accessed November 12th, 2021).
- [49] Davoren MP, Demant J, Shiely F, Perry IJ. Alcohol consumption among university students in Ireland and the United Kingdom from 2002 to 2014: a systematic review. *BMC Publ Health* 2016;16:173. <https://doi.org/10.1186/s12889-016-2843-1>.
- [50] Fazzino TL, Serwatka C, Schneider H, Sullivan D. A systematic review of the methodology used to study weight change among young adults attending college. *Eat Behav* 2019;35:101333. <https://doi.org/10.1016/j.eatbeh.2019.101333>.
- [51] Gherasim A, Arhire LI, Niță O, Popa AD, Graur M, Mihalache L. The relationship between lifestyle components and dietary patterns. *Proc Nutr Soc* 2020;79:311–23. <https://doi.org/10.1017/S0029665120006898>.
- [52] Dietz P, Reichel JL, Edelmann D, Werner AM, Tibubos AN, Schäfer M, et al. A systematic umbrella review on the epidemiology of modifiable health influencing factors and on health promoting interventions among university students. *Front Public Health* 2020;8:137. <https://doi.org/10.3389/fpubh.2020.00137>.
- [53] World Health Organisation. Body mass index - BMI. 2021. <https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi> (Accessed on December 2nd 2021).
- [54] Leone RJ, Morgan AL, Ludy MJ. Validation of self-reported anthropometrics in female college Freshmen. *Int J Exerc Sci* 2016;9:47–55. <https://doi.org/10.1249/01.MSS.0000495302.06939.B7>.
- [55] Cleghorn CL, Harrison RA, Ransley JK, Wilkinson S, Thomas J, Cade JE. Can a dietary quality score derived from a short-form FFQ assess dietary quality in UK adult population surveys? *Publ Health Nutr* 2016;19:2915–23. <https://doi.org/10.1017/S1368980016001099>.
- [56] Hawkins LK, Farrow C, Thomas JM. Do perceived norms of social media users' eating habits and preferences predict our own food consumption and BMI? *Appetite* 2020;149:104611. <https://doi.org/10.1016/j.appet.2020.104611>.
- [57] Healthy Ireland. Healthy food for life: food pyramid questions and answers. 2016. <https://www.hse.ie/eng/about/who/healthwellbeing/our-priority-programmes/health/food-pyramid-images/foodforlifefoodpyramidqas2016.pdf> (Accessed December 2nd, 2021).
- [58] Bush K, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. *Arch Intern Med* 1998;158:1789–95. <https://doi.org/10.1001/archinte.158.16.1789>.
- [59] Dietz JR, Taylor DJ, Sethi K, Kelly K, Bramoweth AD, Roane BM. Psychometric evaluation of the PSQI in U.S. College students. *J Clin Sleep Med* 2016;12:1121–9. <https://doi.org/10.5664/jcsm.6050>.
- [60] Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatr Res* 1989;28:193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4).
- [61] Hirshkowitz M, Whitton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health* 2015;1:40–3. <https://doi.org/10.1016/j.sleh.2014.12.010>.
- [62] Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385–96. <https://doi.org/10.2307/2136404>.
- [63] Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35:1381–95. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>.
- [64] Murphy JJ, Murphy MH, MacDonncha C, Murphy N, Neville AM, Woods CB. Validity and reliability of three self-report instruments for assessing Attainment of physical activity guidelines in university students. *Meas Phys Educ Exerc Sci* 2017;21:134–41. <https://doi.org/10.1080/1091367X.2017.1297711>.
- [65] International Physical Activity Questionnaire (IPAQ). Guidelines for data processing and analysis of the IPAQ. 2005. [www.ipaq.ki.se](http://www.ipaq.ki.se) (Accessed on December 2nd 2021).



- [66] Department of Health. Be Well - PA guidelines. 2021. <https://www.gov.ie/en/publication/06de8b-be-well/#physical-activity-guidelines> (Accessed December 2nd, 2021).
- [67] Munt AE, Partridge SR, Allman-Farinelli M. The barriers and enablers of healthy eating among young adults: a missing piece of the obesity puzzle: a scoping review. *Obes Rev* 2017;18:1–17. <https://doi.org/10.1111/obr.12472>.
- [68] Odlaug BL, Lust K, Wimmelmann CL, Chamberlain SR, Mortensen EL, Derbyshire K, et al. Prevalence and correlates of being overweight or obese in college. *Psychiatr Res* 2015;227:58–64. <https://doi.org/10.1016/j.psychres.2015.01.029>.
- [69] Peltzer K, Pengpid S, Samuels T, Özcan NK, Mantilla C, Rahamefy OH, et al. Prevalence of overweight/obesity and its associated factors among university students from 22 countries. *Int J Environ Res Publ Health* 2014;11:7425–41. <https://doi.org/10.3390/ijerph110707425>.
- [70] Department of Health. Healthy Ireland: summary report 2019. 2019. <https://assets.gov.ie/41141/e5d6fea3a59a4720b081893e11fe299e.pdf> (Accessed November 12th, 2021).
- [71] Tam CF, Xi E, Chan V, Gouzoubachian A. An inverse correlation between fruit and vegetable consumption and BMI among college female and male students. *Coll Stud J* 2017;51:407–23. <https://link.gale.com/apps/doc/A507012102/AONE?u>.
- [72] Asghari G, Mirmiran P, Yuzbashian E, Azizi F. A systematic review of diet quality indices in relation to obesity. *Br J Nutr* 2017;117:1055–65. <https://doi.org/10.1017/S0007114517000915>.
- [73] Davoren MP, Fitzgerald E, Shiely F, Pery JJ. Positive mental health and well-being among a third level student population. *PLoS One* 2013;8:e74921. <https://doi.org/10.1371/journal.pone.0074921>.
- [74] Macilwrait P, Bennett D. Burnout and physical activity in medical students. *Ir Med J* 2018;111:707. <http://imj.ie/burnout-and-physical-activity-in-medical-students/>.
- [75] Murphy MH, Murphy N, MacDonncha C, Woods C, Byrne N, Ferguson K, et al. Student activity and sports study Ireland. 2015. <http://www.studentsport.ie/wp-content/uploads/2016/02/SASSI-Full-Report-Without-Appendices.pdf> (Accessed December 1st, 2021).
- [76] Patterson E, McGeough D, Cannon E, Hagströmer M, Bergman P, Kearney J, et al. Self-efficacy, stages of change and physical activity in Irish college students. *J Public Health* 2006;14:81–6. <https://doi.org/10.1007/s10389-006-0028-6>.
- [77] Walsh LM, Callaghan HP, Keaver LM. Physical activity knowledge, attitudes and behaviours among Irish nursing students. *Int J Health Promot Educ* 2020;59:145–55. <https://doi.org/10.1080/14635240.2020.1729221>.
- [78] Luciano F, Cenacchi V, Vegro V, Pavei G. COVID-19 lockdown: physical activity, sedentary behaviour and sleep in Italian medicine students. *Eur J Sport Sci* 2021;21:1459–68. <https://doi.org/10.1080/17461391.2020.1842910>.
- [79] Rivera PA, Nys BL, Fiestas F. Impact of COVID-19 induced lockdown on physical activity and sedentary behavior among university students: a systematic review. *Medwave* 2021;21:e8456. <https://doi.org/10.5867/medwave.2021.08.8456>.
- [80] Anwer S, Manzar MD, Alghadir AH, Salahuddin M, Abdul Hameed U. Psychometric analysis of the perceived stress scale among healthy university students. *Neuropsychiatric Dis Treat* 2020;16:2389–96. <https://doi.org/10.2147/NDT.S268582>.
- [81] Zhan H, Zheng C, Zhang X, Yang M, Zhang L, Jia X. Chinese college students' stress and anxiety levels under COVID-19. *Front Psychiatr* 2021;12:615390. <https://doi.org/10.3389/fpsy.2021.615390>.
- [82] Benham G. Stress and sleep in college students prior to and during the COVID-19 pandemic. *Stress Health* 2020;37:504–15. <https://doi.org/10.1002/smi.3016>.
- [83] Othman N, Ahmad F, El Morr C, Ritvo P. Perceived impact of contextual determinants on depression, anxiety and stress: a survey with university students. *Int J Ment Health Syst* 2019;13:17. <https://doi.org/10.1186/s13033-019-0275-x>.
- [84] Price A, Smith HA. USI national report on student mental health in third level education. 2019. <https://usi.ie/wp-content/uploads/2019/08/WEB-USI-MH-report-1.pdf> (Accessed December 2nd, 2021).
- [85] Pascoe MC, Hettrick SE, Parker AG. The impact of stress on students in secondary school and higher education. *Int J Adolesc Youth* 2020;25:104–12. <https://doi.org/10.1080/02673843.2019.1596>.
- [86] Choi J. Impact of stress levels on eating behaviors among college students. *Nutrients* 2020;12:1241. <https://doi.org/10.3390/nu12051241>.
- [87] Grimaldi-Puyana M, Fernández-Batanero JM, Fennell C, Sañudo B. Associations of objectively-assessed smartphone use with physical activity, sedentary behavior, mood, and sleep quality in young adults: a cross-sectional study. *Int J Environ Res Publ Health* 2020;17:3499. <https://doi.org/10.3390/ijerph17103499>.
- [88] Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity* 2018;16:643–53. <https://doi.org/10.1038/oby.2007.118>.
- [89] Godos G, Castellano S, Galvano F, Caraci F, Ferri R. Association between diet and sleep quality: a systematic review. *Sleep Med Rev* 2021;57:101430. <https://doi.org/10.1016/j.smrv.2021.101430>.
- [90] Zuraikat FM, Wood RA, Barragán R, St-Onge M-P. Sleep and diet: mounting evidence of a cyclical relationship. *Annu Rev Nutr* 2021;41:309–32. <https://doi.org/10.1146/annurev-nutr-120420-021719>.
- [91] Di Benedetto M, Towt CJ, Jackson ML. A cluster analysis of sleep quality, self-care behaviors, and mental health risk in Australian university students. *Behav Sleep Med* 2020;18:309–20. <https://doi.org/10.1080/15402002.2019.1580194>.
- [92] Dietrich SK, Francis-Jimenez Cm, Knibbs MD, Umali IL, Truglio-Londrigan M. Effectiveness of sleep education programs to improve sleep hygiene and/or sleep quality in college students. *JBI Database Systemat Rev Implement Rep* 2016;14:108–34. <https://doi.org/10.1124/JBISRIR-2016-003088>.
- [93] Veronda AC, Irish LA, Delahanty DL. Effect of smoke exposure on young adults' sleep quality. *Nurs Health Sci* 2020;22:57–63. <https://doi.org/10.1111/nhs.12644>.
- [94] Murray SR, Lyne SJ, Cryan MD, Mullin M, McGrath D, Hayes CB. Not really a smoker? A study on the prevalence of and attitudes to occasional social smoking in a third level institution in Ireland. *Ir J Med Sci* 2021;190:941–8. <https://doi.org/10.1007/s11845-020-02439-5>.
- [95] Gangadi M, Kalpourtzi N, Gavana M, Vantarakis A, Chlouverakis G, Hadjichristodoulou C, et al. Prevalence of tobacco smoking and association with other unhealthy lifestyle risk factors in the general population of Greece: results from the EMENO study. *Tobacco Prev Cessation* 2021;7:61. <https://doi.org/10.18332/tpc/140242>.
- [96] Alkeravi A, Baydarlioglu B, Sauvageot N, Stranges S, Lemmens P, Shivappa N, et al. Smoking status is inversely associated with overall diet quality: findings from the ORISCAV-LUX study. *Clin Nutr* 2017;36:1275–82. <https://doi.org/10.1016/j.clnu.2016.08.013>.
- [97] Cooke R, Beccaria F, Demant J, Fernandes-Jesus M, Fleig L, Negreiros J, et al. Patterns of alcohol consumption and alcohol-related harm among European university students. *Eur J Publ Health* 2019;29:1125–9. <https://doi.org/10.1093/eurpub/ckz067>.
- [98] Carbia C, García-Cabrero R, Cryan JF, Dinan TG. Associations between mental health, alcohol consumption and drinking motives during COVID-19 second lockdown in Ireland. *Alcohol Alcohol* 2021;1–8. <https://doi.org/10.1093/alcalc/agab067>.
- [99] Kilian C, Rehm J, Allebeck P, Braddick F, Gual A, Barták M, et al. Alcohol consumption during the COVID-19 pandemic in Europe: a large-scale cross-sectional study in 21 countries. *Addiction* 2021;116:3369–80. <https://doi.org/10.1111/add.15530>.
- [100] Breslow RA, Guenther PM, Juan W, Graubard BI. Alcoholic beverage consumption, nutrient intakes, and diet quality in the US adult population, 1999–2006. *J Am Diet Assoc* 2010;110:551–62. <https://doi.org/10.1016/j.jada.2009.12.026>.
- [101] Sheehy O'Sullivan E, McCarthy K-M, O'Neill C, Walton J, Bolger L, Bickerdike A. The impact of COVID-19 on the health-related behaviours, mental well-being, and academic engagement of a cohort of undergraduate students in an Irish university setting. *Int J Environ Res Publ Health* 2022;19(23):16096. <https://doi.org/10.3390/ijerph192316096>.
- [102] Gandy J. *Manual of dietetic practice*. 5th ed. Wiley-Blackwell on behalf of the BDA; 2014. ISBN 9780470656228.
- [103] Schoeller DA, Westertep-Plantenga M. *Advances in the assessment of dietary intake*. 1st ed. CRC Press; 2021. ISBN 9781032096520.
- [104] Lesińska-Sawicka M, Pisarek E, Nagórka M. The health behaviours of students from selected countries—a comparative study. *Nurs Rep* 2021;11:404–17. <https://doi.org/10.3390/nursrep11020039>.
- [105] Hammoudeh S, Gadelhaq W, Janahi I. Chapter 2. Prospective cohort studies in medical research. In: Barria RM, editor. *Cohort studies in health Sciences*. 1st ed. Intech Open; 2018. <https://doi.org/10.5772/intechopen.76514> (Accessed January 30th, 2020).
- [106] Kowalski K, Rhodes R, Naylor PJ, Tuokko H, MacDonald S. Direct and indirect measurement of physical activity in older adults: a systematic review of the literature. *Int J Behav Nutr Phys Activ* 2012;9:148. <https://doi.org/10.1186/1479-5868-9-148>.
- [107] Rippin HL, Hutchinson J, Greenwood DC, Jewell J, Breda JJ, Martin A, et al. Inequalities in education and national income are associated with poorer diet: pooled analysis of individual participant data across 12 European countries. *PLoS One* 2020;15(5):e0232447. <https://doi.org/10.1371/journal.pone.0232447>.

**Stephen Doak** is a postgraduate student at the Atlantic Technological University (ATU) exploring dietary and lifestyle behaviors such as stress, sleep, eating attitudes, physical activity, and alcohol consumption, of higher education students.

**John M. Kearney, PhD, RD** is a lecturer in nutrition and Epidemiology and acting chair of the Public Health Nutrition programme at TU Dublin. His research interests include food consumption patterns and health behaviors and their determinants.

**Jacqueline M. McCormack, PhD** is Vice President (Online Development) at the ATU and a Professor of Biomedical Sciences. Her research interests include online learning, nutrition, Immunology and the impact of diet on psychological function.

**Laura Keaver, MPH, RD, RNUTR** is a lecturer in human nutrition and dietetics at ATU. Her research interests include student health, lifestyle interventions, and exploring the role of nutrition in the prevention and management of chronic disease.