

UC Merced

Biogeographia - The Journal of Integrative Biogeography

Title

The new Checklist of the Italian Fauna: Odonata

Permalink

<https://escholarship.org/uc/item/72f1309t>

Journal

Biogeographia - The Journal of Integrative Biogeography, 38(1)

ISSN

1594-7629

Authors

La Porta, Gianandrea
Landi, Federico
Leandri, Fausto
[et al.](#)

Publication Date

2023

DOI

10.21426/B638158781

Supplemental Material

<https://escholarship.org/uc/item/72f1309t#supplemental>

Copyright Information

Copyright 2023 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

The new Checklist of the Italian Fauna: Odonata

GIANANDREA LA PORTA ^{1,2*}, FEDERICO LANDI ¹,
FAUSTO LEANDRI ¹, GIACOMO ASSANDRI ¹

¹ *Società Italiana per lo Studio e la Conservazione delle Libellule ODV,
Via Elce di Sotto, 8, 06123 Perugia (PG), Italy*

² *Department of Chemistry, Biology and Biotechnology (DCBB), University of Perugia,
Via Elce di Sotto, 8, 06123 Perugia (PG), Italy*

* *Corresponding author, email: gianandrea.laporta@unipg.it*

Keywords: Odonata; biodiversity; biogeography; species distribution; checklist

SUMMARY

Seventeen years after the publication of the last checklist of the damselflies and dragonflies found in Italy, an updated list has been compiled. This list reports 95 species belonging to 10 families and 38 genera and includes 2 national endemic and 1 sub-endemic species. Compared to the previous checklist, three species were removed, two subspecies were granted species status, and 10 species were added as new taxa for Italy. The checklist summarizes the current state of the knowledge on the geographical distribution of the Italian species with a regional detail. After the online publication of this dataset on the LifeWatch Italy website in 2021, some minor updates will be included in future releases. The newly discovered species are the result of increased exploration of the national territory combined with some possible range shifts, especially of Libellulidae species of Afro-Asiatic origin. This increased coverage of the country is the result of the efforts of many contributors. It stems from the rapidly growing interest in this zoological group, also enhanced by the activation of a nationwide citizen science project promoted by the Italian Society for the Study and Conservation of Dragonflies (Odonata.it).

INTRODUCTION

Odonata is an insect order composed of approximately 6370 species (Paulson et al. 2022). Dragonflies and damselflies are recognised as valuable ecological indicators in a variety of hydroecosystems and are thus successfully used for: i) the assessment and monitoring of aquatic and terrestrial habitat integrity and ii) the detection and prediction of the biological impact of climate warming (Hassall 2015, Rocha-Ortega et al. 2019) (Fig. 1).

In Europe, 145 species of dragonflies and damselflies are known to occur so far (Boudot and Kalkman 2015, Viganò et al. 2017, López-Estrada et al. 2020). Italy is among the best-known country of Europe with respect to the fauna of Odonata (Boudot and Kalkman 2015). It has been extensively studied from a faunistic point of view since the end of the XIX century (e.g., Ausserer, 1869; Pirota, 1879; Garbini, 1897), although several authors wrote about the Italian dragonflies even earlier (e.g., Vandelli, 1763, Allioni, 1766; Cirillo, 1787).

Checklists and related faunistic occurrence data are key elements in biogeographical and ecological investigations, as they contribute to the identification of priority areas, organization and planning of projects dealing with mapping, inference, and modelling of current or potential species distributions (Gaston and Rodrigues, 2003; Phillips et al. 2006; Possingham et al., 2007; Sandal et al., 2022). Italy has a long history of odonatological checklists. The first ‘modern’ effort was the first volume of the ‘Fauna d’Italia’ series (Conci and Nielsen, 1956), which reported the occurrence of all the Odonata species known at that time for each Italian region, for a total of 80 species for Italy as a whole. Subsequently, other authors provided periodical updates, in particular: Carchini et al. (1985), Carchini and Rota (1986), Utzeri (1994), Utzeri and D’Antonio (2005). The last two checklists were published in the framework of the huge collaborative project of

the ‘Checklist of the Italian Fauna’ (Minelli et al. 1993-1995; Ruffo and Stoch, 2005).

Since the publication of the last Italian Odonata checklist 17 years ago (Utzeri and D’Antonio, 2005), there has been a remarkable increase in the number of valuable faunistic papers on odonates in Italy, even in the traditionally less explored regions of central and southern Italy. This increase, concomitantly reported in other parts of Europe, has been likely triggered by the improved availability of tools for species identification and the presence of internet platforms that allow data, photographs, and observations to be shared, reflecting a rapidly growing interest for this group of insects (Kalkman et al. 2018).

The new checklist (available on the LifeWatch Italy web site, and in a synthetic version in the Supplementary file S1) presents an updated checklist of the damselflies and dragonflies occurring in Italy with a regional geographical detail. New species have been reported in the last 17 years and some taxa have been recently taxonomically revised, making it necessary to update the list of the country to reflect the current state of the knowledge for this order of insects (Supplementary file S2). Thanks to all the researchers and volunteers who have dedicated their time and energy to building an understanding of dragonfly distribution patterns in Italy, the present checklist is based on materials from private and institutional collections and scientific literature, as well as digital biodiversity repositories fed by citizen science projects.

RESULTS

Summary statistics

A total of 95 species belonging to 10 families and 38 genera are listed: Calopterygidae (1 genus, 4 species); Lestidae (3 genera, 9 species); Platycnemididae (1 genus, 1 species); Coenagrionidae (7 genera, 18 species); Aeshnidae (4 genera, 13 species); Gomphidae (6

genera, 8 species); Cordulegastridae (1 genus, 4 species); Corduliidae (2 genera, 6 species); Synthemistidae (1 genus, 1 species); Libellulidae (12 genera, 31 species). 52 species occur in all three Italian biogeographical regions: 68 species live in the Alpine region, 70 in the Continental region and 76 in the Mediterranean region. The Mediterranean region also has the largest number of exclusive species (15 species), followed by the Alpine region (9 exclusive species) and the Continental region (3 exclusive species).

From a chorological point of view, 39% of the species consists of Mediterranean taxa (including species more or less widely spread in the Mediterranean countries), 16% are Asiatic-European species, 12% Sibero-European species, 14% European species, and the remaining 19% belong to other 7 chorotypes. Altogether the 95 species represent about 66% of

the European odonatological fauna. Two species, namely the damselfly *Coenagrion castellani* (formerly known as *C. mercuriale castellanii*) and the dragonfly *Cordulegaster trinacriae*, are endemic of Italy, while *Ischnura genei* is a subendemic species due to its presence in the French island of Corsica. Compared to the previous checklist (Utzeri and D'Antonio, 2005), 3 species (*Ischnura graeelsi*, *Cordulegaster picta*, *Epithea bimaculata*) should not be considered part of the national fauna anymore. Further occurrences of the species in national territory not included in the Supplementary file S2 will be included in future updates of the checklist. It is the case of *Aeshna subarctica* in Veneto, *Cordulegaster bidentata* in Molise, *Somatochlora flavomaculata* in Puglia, and *Trithemis annulata* in Piemonte, just to name a few of the cases of which the authors are currently aware.



Figure 1. Representative Odonata: a) *Coenagrion castellani* – Trevi (PG), 03/06/2022 (foto G. La Porta); b) *Aeshna subarctica* – Altavalle (TN), 09/08/2019 (foto G. Assandri); c) *Oxygastra curtisii* – Calcio (BG), 11/06/2018 (foto F. Leandri); d) *Zygonyx torridus* –S. Caterina Menfi (AG), 14/08/2014 (foto F. Landi).

Data set information

Object name: Checklist of the Italian Fauna: Odonata

Characters encoding: Unicode (UTF-8)

Data set citation: La Porta G., Landi F., Leandri F., Assandri G., 2022. Odonata. In: Bologna M.A., Zapparoli M., Oliverio M., Minelli A., Bonato L., Cianferoni F., Stoch F. (eds.), Checklist of the Italian fauna. Version 1.0. Last update: 2021-05-31.

Format name: xml, Extensible Markup Language, for the online version at LifeWatch.

Format version: 1.0

Distribution: <https://dataportal.lifewatchitaly.eu/view/urn%3Auuid%3Ac1f2ab37-61e4-48e9-b3a9-15bdbf002f9d>

Date of creation: June 15th, 2020

Date of last revision: May 31st, 2021

Date of publication: July 23rd, 2021

Language: English

License of use: Creative Commons Attribution 4.0 International License

Metadata language: English

Metadata managers: Marco Bologna, Lucio Bonato, Fabio Cianferoni, Alessandro Minelli, Marco Oliverio, Fabio Stoch, Marzio Zapparoli & LifeWatch Italy

Management details

Project title: The new Checklist of the Italian Fauna: Odonata

Database manager: Odonata.it; Marco Bologna, Lucio Bonato, Fabio Cianferoni, Alessandro Minelli, Marco Oliverio, Fabio Stoch, Marzio Zapparoli & LifeWatch Italy

Temporal coverage: Anything published until May 2021.

Record basis: Published records in scientific and grey literature and records stored and validated in digital biodiversity repositories fed by citizen science projects.

Funding grants: No funding was specifically available for the project on Odonata; funding for the update of the Checklist of the Italian fauna was obtained from LifeWatch Italy; no funding was specifically available for the update.

Geographic information

General description: The dataset includes records from the national territories of Italy, including the two major islands Sardinia and Sicily, together with archipelagos and minor islands politically under the Italian legislation.

Geographic units: The geographical units within the Italian national territories for terrestrial records refer to the administrative boundaries of the 20 Italian regions, in addition to San Marino Republic and Vatican City (Table 1). Continental Italy was also divided in only two units, namely North (Friuli - Venezia Giulia, Veneto, Trentino - Alto Adige, Lombardia, Valle d'Aosta, Piemonte, Liguria, Emilia Romagna) and South (Toscana, Marche, Umbria, Lazio, Abruzzo, Molise, Campania, Puglia, Basilicata, Calabria), while islands of Sardinia and Sicily were left as independent units. Furthermore, the national territory was also analyzed by considering three European biogeographical regions, as defined for Italy by the European Environment Agency (2017): Alpine, Continental, and Mediterranean.

Bounding box: All areas falling under Italian administration (in addition to San Marino and Vatican City) from 35° 25' to 47° 06' N and from 6° 35' to 19° 20' E (WGS84 reference system) were included.

Sampling design: We did not perform any additional sampling to collect records, but we used only published data and validated sightings coming from georeferenced archives of Odonata.it.

Habitat type: Odonata are amphibiotic hemimetabolous insects, whose nymphs generally live in shallow and slow-flowing waters, whereas adults are good flyers, specialized for dispersal and reproduction. Some species are highly mobile and/or migrant and may disperse over hundreds or even thousands of kilometers. This data paper provides information on dragonflies and damselflies that have been recorded in Italy at fresh flowing or standing waters, as in any terrestrial habitat. Different species inhabit different habitats, such as permanent or temporary waters, unvegetated or with submerged, floating, emergent and waterside plants. Surveys have been conducted across a wide range of habitats such as springs, brooks, streams, rivers, ponds, lakes, ditches, canals, quarries and reservoirs, swamps, peat bogs, marshes, fens, and coastal brackish wetlands. Furthermore, flying adults, dispersing, hunting or maturing, may be encountered in any terrestrial habitat, especially open woods, edges, bushes, meadows and grasslands even far from freshwater sites.

Countries: Italy, without including Corsica, Canton Ticino, and Maltese Archipelago, politically belonging respectively to the French Republic, Swiss Confederation and the Republic of Malta, which have been properly investigated in detail by other authors (Wildermuth et al., 2005, Ebejer, et al., 2008, Roché et al., 2008, Berquier C. 2013, Degabriele 2013, UICN France, 2016, Boudot et al., 2017, Monnerat et al., 2021). San Marino and Vatican City were included even though no odonatological studies were ever carried out due to a lack of relevant sites of odonatological interest.

Quality control for geographic data: We checked that the georeferenced records and the published localities in the papers indeed matched the geographical units used for the checklist at the level of administrative regions.

Literature records

General description: Only published records and publicly available records are included in the dataset. A search through the literature was performed on 15th June 2020.

Literature search methods: We searched through Web of Science, Scopus, and Google Scholar for keywords ('odonata', 'dragonflies', 'damselflies') and ('Italy' or 'Italian'), and individual species by their names.

Literature list: (i) Papers reporting the nine new species found in Italy after checklist of 2005: Bedjanič and Šalamun, (2003); Kunz et al. (2006); Holusa (2008); Festi (2011); Corso et al. (2012a); Corso et al. (2012b); Rattu et al. (2014); Viganò et al. (2017); Corso and Penna (2020); (ii) Papers reporting significant information inserted in species distribution notes: Conci et al. (1944); Capra (1945; 1964); Conci (1948); Roberts (1948); Consiglio (1950; 1953); Nielsen and Conci (1951); Minelli (1966); Bucciarelli (1972; 1978); Bordoni et al. (1980); Mancini et al. (1993); D'Antonio (1995, 1997); Bedjanic and Salamun (1999); Kunz (2006); Boano et al. (2007); Olias et al. (2007); Dijkstra and Matushkina (2009); Fiorenza et al. (2009); Festi (2011); Giugliano and Terzani (2011); Dijkstra et al. (2012, 2013, 2021); Ottonello and Oneto (2013); Terzani et al. (2013); Corso et al. (2014); Riservato et al. (2014); Schneider and Utzeri (2014); Zandigiacomo (2014); Mastropasqua and Liuzzi (2016); Aguzzi et al. (2017); Assandri (2019); Corso (2019); Piretta and Assandri (2019); Bonometto (2020); Assandri et al. (2022); (iii) Other selected papers of local, regional, or national interest published after 2005 (Supplementary file S1).

Taxonomic information

General description: Only records reporting species were included, disregarding records at higher level like genus, family, etc.

Taxonomic coverage: Any species of the order Odonata Fabricius, 1793 of the class Insecta.

Although in the previous checklist (Utzeri and D’Antonio, 2005) 26 taxa were treated at a subspecific level, in the present work subspecies were not included as they require further genetic investigation for a taxonomic resolution (Galimberti et al. 2020) and present doubtful geographic delimitations.

Taxonomic rank: Only species, without subspecies, are reported.

Taxon specialists: Giacomo Assandri, Federico Landi, Fausto Leandri, Gianandrea La Porta

Nomenclature: The adopted nomenclature follows the World Odonata List (Paulson et al. 2022).

Taxonomic remarks: All taxonomic changes that occurred since the publication of the previous checklists (Utzeri and D’Antonio, 2005). Species that were included in the previous checklist and are now considered not valid or for which there is new evidence that they do not occur in Italy were excluded from the dataset.

Quality control for taxonomic data: Taxonomic data were checked and updated to include revision of names, synonyms, delimitation of genera and higher taxa, all conducted through a comparison with the World Odonata List (Paulson et al. 2022) and other published catalogues (Bridges 1994; Banki et al., 2022).

Table 2 - Description of the dataset with specific information relative to definitions and storage type for each field.

Field	Description	Storage type
Phylum	Phylum name, Arthropoda for all records	string
Class	Class name, Insecta for all records	string
Order	Order name, according to Paulson et al. 2022	string
Family	Family name, according to Paulson et al. 2022	string
Subfamily	Subfamily name, according to Bridges (1994)	string
Genus	Valid genus name, according to Paulson et al. 2022	string
Genus authorship	Genus descriptor	string
Species	Valid species name, according to Paulson et al. 2022	string
Species authorship	Species descriptor	string
Fauna Europaea (name)	Species name as reported in FaunaEuropaea	string
Endemic	Species known as endemic for Italy	string
Alien	Species known as alien for Italy (all the Italian species should be considered native)	string
N	Occurrence of the taxon in Northern continental Italy (grouping: Friuli - Venezia Giulia, Veneto, Trentino - Alto Adige, Lombardia, Valle d'Aosta, Piemonte, Liguria, Emilia Romagna)	binary

S	Occurrence of the taxon in Southern continental Italy (grouping: Toscana, Marche, Umbria, Lazio, Abruzzo, Molise, Campania, Puglia, Basilicata, Calabria)	binary
Si	Occurrence of the taxon in Sicily	binary
Sa	Occurrence of the taxon in Sardinia	binary
Ao	Occurrence of the taxon in Valle d'Aosta	binary
Pi	Occurrence of the taxon in Piemonte	binary
Lo	Occurrence of the taxon in Lombardia	binary
VT	Occurrence of the taxon in Trentino - Alto Adige	binary
V	Occurrence of the taxon in Veneto	binary
FVG	Occurrence of the taxon in Friuli - Venezia Giulia	binary
Li	Occurrence of the taxon in Liguria	binary
ER	Occurrence of the taxon in Emilia Romagna	binary
To	Occurrence of the taxon in Toscana	binary
Ma	Occurrence of the taxon in Marche	binary
Um	Occurrence of the taxon in Umbria	binary
La	Occurrence of the taxon in Lazio	binary
Abr	Occurrence of the taxon in Abruzzo	binary
Mo	Occurrence of the taxon in Molise	binary
Cp	Occurrence of the taxon in Campania	binary
Pu	Occurrence of the taxon in Puglia	binary
Bas	Occurrence of the taxon in Basilicata	binary
Cal	Occurrence of the taxon in Calabria	binary
Chorotype	Chorotype according to Vigna Taglianti et al., 1999	string
Taxonomic notes	Relevant taxonomic and systemic notes	string
Distribution notes	Significant notes referring to the regional distribution of each taxon and the relative references	string

DISCUSSION

Taxonomic revisions, evaluations of historical documentary sources and reappraisal of specimens in collections have had little effect on the changes occurred from the previous checklist to the present inventory. Only two species previously considered subspecies have been granted species status (*Calopteryx xanthostoma* and more recently *Coenagrion castellani*, still

listed in Supplementary file S2 as *C. mercuriale*), two species have been reassigned to a different taxonomic position (*Orthetrum anceps* and *Brachythemis leucosticta*), and three species, namely *Ischnura graeelsi*, *Cordulegaster picta*, and *Epitheca bimaculata*, have been removed from the list due to misidentifications of the specimens and lack of clear evidence of past presence in Italy.




Figure 2. The relationship between the species richness and the area of the Italian regions. Linear regression model: Species richness = 50 + 0.53 Area.

Much more impactful have been the changes resulting from the newly discovered species, which are related to: i) the increased exploration of the national territory, ii) the rediscovery of one species thought to be extinct from the country (*Coenagrion ornatum*), iii) the possible recolonization or extension of range from the near-east (*Cordulegaster heros*), iv) the northward shift of the range margins of some species. In fact, seven of the ten new species on the checklist, namely *Orthetrum chrysostigma*, *Sympetrum sinaiticum*, *Diplacodes lefebvrii*, *Pantala flavescens*, *Tramea basilaris*, *Trithemis*

kirby, *Zygonyx torridus* belong to the family of Libellulidae and are thermophilic Afro-Asiatic taxa rapidly expanding northward in most of their range (Polette et al. 2017; Günther 2019; Cerini et al. 2020).

These recent colonizations are likely linked to the dispersal abilities of some species and the climatic suitability of central and southern Italy. However, in some cases, these species are currently known from few or single localities and the long-term persistence of their populations over time should be monitored. These range shifts are also the main reason for

the remarkable increase in diversity of the Mediterranean area, currently the richest of the three Italian bioregions and also the least known and investigated. Fifty-two species are widespread throughout the country, although three of them are currently rare and very localized in the Mediterranean context (*Erythromma najas*, *Sympetrum flaveolum* and *Sympetrum depressiusculum*) and two are found only on the edges of the Alpine area (*Gomphus vulgatissimus* and *Somatochlora meridionalis*).

Considering the individual Italian administrative regions, a statistically significant linear correlation arises between the species richness (S) and the area of the mainland and the Sicily insular system ($R=0.56$; $p<0.05$) (Fig. 2). Sardinia, which is biogeographically distinct because of its geological history, was not considered in the regression and is indicated with a different symbol. The overall rate of increase in species richness per unit area is estimated at 0.53 species/1000 km² (Fig. 2), while the northern regions show a more pronounced increase, as these combine alpine and lowland habitats, and thus have a higher environmental heterogeneity. In contrast, the southern regions are characterized by an increased species turnover and a lower than the expected richness. The situation differs particularly for the island of Sardinia, which is faunistically impoverished compared to the continental regions. The causes for this low species richness can be traced back to its long isolation and its origins from a microplate detached from the Iberian Peninsula (Schmitt et al., 2021). These factors have probably resulted in a low colonization rate of the territory, especially by damselflies. In addition, the current climatic conditions of this region result in a lower availability of wetlands and suitable areas for odonates. Furthermore, it needs to be considered that the odonates of Sardinia, together with the other southern regions, has been less investigated, compared to the northern regions, and thus the rate of false absences is likely to be higher.

The information compiled for country-level checklists - and the present one is no exception - inevitably contains a certain degree of spatial and environmental biases and survey incompleteness. Indeed, the efficiency of biodiversity studies is related to various aspects of survey design and the observer or taxonomist distribution across the territory (Engel, 2021). Effective survey designs are required when the patchiness in the distribution of species richness is high, while some key elements lead to a concentration in species occurrence (e.g., sites accessibility, survey frequency, distance from observer or taxonomist residences). Besides, the availability of information and odonatalogical lists for the localities formerly surveyed can direct field activities towards species-rich sites, but also promote more efficient use of previous efforts. This is the case for northern regions and a few provinces in the rest of Italy for which there is a more consolidated natural history tradition. On the other hand, lack of information and unavailability of collected specimens are factors that lead to the uncoordinated and repeated investigation of a smaller group of localities. Extensive and systematic surveys are indeed expensive and time-consuming. Consequently, better results can be achieved combining the efforts of many contributors. In this perspective, the Italian Society for the Study and Conservation of Dragonflies (Odonata.it) in 2014 launched a nationwide data collection project based on a citizen science (CS) approach. Collecting data through CS activities raises some issues, such as those related to the lack of formal protocols, uneven spatial coverage, erratic fluctuation in the number of volunteers, but also those related to detectability and identifiability of species, population densities, and accessibility of sites. In addition, besides the quantity of collected sightings, the associated quality is an important concern in data reliability. The experience of this project, whose description is beyond the scope of this data-paper, shows that some of these issues can be mitigated. Especially, when activities are conducted through a combination of expert surveys, accurate data validation, and the

creation of a network of active odonatologists. From the start-up phase to the present, the degree of coverage of the national territory on a grid of 10 × 10 km has increased from 72% to 87% and a data set of more than 200,000 field records has been collected. Despite the unevenness in survey effort and research infrastructure, the data collected so far have allowed the reduction of the national Wallacean shortfall (Cardoso et al., 2011), improving the knowledge on the distribution of dragonfly in Italy. On the other hand, no significant progress has been achieved in filling or containing instead the Prestonian shortfall (Cardoso et al., 2011). This impediment, which outlines the limited knowledge on the abundance of species and their changes in space and time, is nowadays one of the most critical challenges in insect conservation and the present checklist hopes to be a trigger for action by the Italian odonatological community.

ACKNOWLEDGEMENTS

We wish to thank the ‘Commissione Scientifica per la Fauna d’Italia’, who pursued the updates for the checklists of the Italian Fauna. We would like to thank D. Baroni, M. Chemollo, I. Chiandetti, C. Christille, A. Festi, A. Galimberti, S. Hardersen, F. Mastropasqua, M. Pettavino, N. Pilon, M. Porciani, G. Soldato, S. Surdo, and the board of Odonata.it. The present work would not have been possible without the efforts and data sharing by the fellowships of the ‘Società Italiana per lo Studio e la Conservazione delle Libellule ODV’ and the many contributors of citizen science projects on odonates in Italy.

REFERENCES

Aguzzi, S., Bogliani, G., Orioli, V. & Pilon, N. (2017) *Nehalennia speciosa* rediscovered in northwestern Italy (Odonata: Coenagrionidae). *Notulae Odonatologicae*, 8 (9), 319–374.

Allioni, C. (1766) *Manipulus insectorum taurinensium* “Mélanges de Philosophie et

Mathématique de la Société royale de Turin”, (per gli anni 1762-65), *Odonata*, III, 7, 185–198.

Assandri, G. (2019) A critical review of the odonate fauna of Trentino: annotated check-list and new relevant data for Italy (Insecta: Odonata). *Fragmenta entomologica*, 51, 75–88. DOI: 10.13133/2284-4880/339

Assandri G., Bazzi G., Festi A., Leandri F. (2022) Distribution, ecology and conservation of *Aeshna caerulea* (Strom, 1793) and *Aeshna subarctica elisabethae* Djakonov, 1922 (Insecta: Odonata) at the southernmost limits of their range. *Aquatic Insect*, in press DOI: 10.1080/01650424.2022.2098341

Ausserer, C., (1869) Neurotteri Tirolesi colla diagnosi di tutti i generi europei. Parte I. Pseudoneurotteri. *Annuario della Società dei Naturalisti in Modena*, 4, 71–156.

Bánki, O., Roskov, Y., Döring, M., Ower, G., Vandepitte, L., Hobern, D., et al. (2022) Catalogue of Life Checklist (Version 2022-01-14). *Catalogue of Life*. DOI: 10.48580/d4tp

Bedjanic, M. & Salamun, A., (1999) Contribution to the knowledge of the Odonate fauna of Sicily, with some additional data from Basilicata, Southern Italy. *Opuscula zoologica fluminensia*, 169, 1–14.

Bedjanič M. & Šalamun A. (2003) Large golden-ringed dragonfly *Cordulegaster heros* Theischinger, 1979, new for the fauna of Italy (Odonata: Cordulegastridae). *Natura Sloveniae*, 5(2), 19–29.

Berquier C., (2013) Plan Régional d’Actions en faveur des Odonates. Région Corse. - 2013-2017. Office de l’Environnement de la Corse – Direction Régionale de l’Environnement, de l’Aménagement et du Logement de Corse, 67 pp.

Boano G., Sindaco R., Riservato E., Fasano S. (2007) Atlante degli odonati del Piemonte e della Valle d’Aosta. *Memorie dell’Associazione Naturalistica Piemontese*, 6, 160 pp.

Bonometto, L. (2020) Le libellule del Cadore. Le specie, gli habitat, le tutele possibili. Parco Naturale Regionale delle Dolomiti d’Ampezzo, Quinto di Treviso, 186 pp.

- Bordoni, A., Carfi S. & Terzani F. (1980) Odonati dell'Alta Val Bormida. IV contributo alla conoscenza degli Odonati italiani. *Redia*, 63, 55–65.
- Boudot, J. P., Grand, D., Wildermuth, H. Monnerat, C., (2017). *Les Libellules de France, Belgique, Luxembourg et Suisse – 2^{ème} Édition*, (Collection Parthénope) Biotope Editions 456 pp. Boudot, J-P. & Kalkman, VJ. (2015) Atlas of the European dragonflies and damselflies. KNNV Publishing, Zeist, the Netherlands.
- Bridges, C. A. (1993) Catalogue of the family-group, genus-group and species-group names of the Odonata of the world (Second Edition). C. A. Bridges, Urbana, Illinois DOI: 10.5962/bhl.title.15291
- Bucciarelli, E. (1972) *L'Aeshna caerulea* (Ström) in Italia e altre interessanti catture nella regione alpina. *Bollettino della Società Entomologica Italiana*, 104, 94–99.
- Bucciarelli, I. (1977) Dati preliminari sul popolamento odonatologico di Calabria, Sicilia e Sardegna (VIII Contributo alla conoscenza degli Odonati). *Annali del Museo Civico di Storia Naturale di Genova*, 81, 374–386.
- Bucciarelli, E. (1978) Odonati della foresta demaniale del Cansiglio (Veneto). *Lavori Società veneziana di Scienze Naturali*, 3, 19–27.
- Capra, F. (1945) Odonati di Liguria. *Annali Museo civico Storia Naturale Genova*, 62, 253–275.
- Capra, F. (1964) Odonati del lago Trasimeno. II. *Rivista di Idrobiologia*, 3(2-3), 173–185.
- Carchini, G., Rota, E. & Utzeri, C. (1985) Lista aggiornata degli odonati italiani e loro distribuzione regionale. *Fragmenta Entomologica*, 18, 91–103.
- Carchini, G. & Rota, E. (1986) Attuali conoscenze sulla distribuzione degli Odonati dell'Italia meridionale. *Biogeographia – The Journal of Integrative Biogeography*, 10, 629–684. DOI: 10.21426/B610110290
- Cardoso, P., Erwin, T.L., Borges, P., New, T. (2011) The seven impediments in invertebrate conservation and how to overcome them. *Biological Conservation*, 144, 2647–2655. DOI: 10.1016/j.biocon.2011.07.024
- Cerini, F., Stellati, L., Luiselli, L. & Vignoli, L. (2020) Long-term shifts in the communities of Odonata: effect of chance or climate change? *North-Western Journal of Zoology*, 16, 1–6.
- Cirillo, D. (1787). *Entomologiae Neapolitanae Specimen primum*. Odonata II, fig. 7, 8 pp.
- Conci, C. (1948) *L'Agrion caerulescens* Fonsc. nell'Italia continentale (Odonata). *Bollettino dei Musei e degli Istituti Biologici Dell'Università di Genova (N.S.)*, 6, 479.
- Conci, C. & Galvani, O. (1944) Alcune interessanti catture di Odonati in Liguria e Venezia Tridentina. *Memorie Società Entomologia Italiana*. Genova, 23, 71–73.
- Conci, C. & Nielsen, C. (1956) *Fauna d'Italia Vol. 1. Odonata*. Calderini, Bologna.
- Consiglio, C. (1950) Cattura del *Gomphus flavipes* (Charp.) a Roma. *Bollettino Società entomologica italiana*, 80, 16.
- Consiglio, C. (1953) Odonati dell'Italia meridionale e degli Abruzzi. *Memorie Società entomologica italiana*, 31, 96–108.
- Corso, A., Janni, O., De Lisio, L., Fracasso, C. (2019) Update to the status of *Lindenia tetraphylla* (Vander Linden, 1825) (Odonata Gomphidae) in Italy, with special reference to the Molise region. *Biodiversity Journal*, 10 (1), 7-12. DOI: 10.31396/Biodiv.Jour.2019.10.1.7.12
- Corso, A., Janni, O., Pavesi, M., Mammut, M., Sciberras, A., Vignoli, M. (2012) Annotated checklist of the dragonflies (Insecta Odonata) of the islands of the Sicilian Channel, including the first records of *Sympetrum sinaiticum* Dumont, 1977 and *Pantala flavescens* (Fabricius, 1798) for Italy. *Biodiversity Journal*, 3 (4), 459–478.
- Corso, A. & Penna, V., (2014) *Orthetrum chrysostigma* (Burmeister, 1839): new for the Italian fauna (Odonata Libellulidae) *Biodiversity Journal*, 11 (2), 359–362. DOI: 10.31396/Biodiv.Jour.2020.11.2.359.362
- Dal Cortivo, M. & Roncen, N. (2019) Nuovi dati sulle Libellule della Provincia di Belluno (Insecta: Odonata). *Bollettino Del Museo Civico di Storia Naturale di Verona*, 43, 5–12.

- D'Antonio, C. (1995). Gli Odonati della Campania (Odonata) Bollettino Società entomologica italiana 127 (2), 103–116.
- D'Antonio, C. (1997) Odonate collection in the department of agricultural entomology and zoology of the University of Naples. Notulae Odonatologicae, 4, 137–152.
- Degabriele, G. (2013) An overview of the dragonflies and damselflies of the Maltese Islands (Central Mediterranean) (Odonata). Bulletin of the entomological Society of Malta Vol. 6, 5–127.
- Dijkstra, K. DB., Achröter, A. & Lewington, R. (2021) Libellule d'Italia e d'Europa, Ricca Editore.
- Dijkstra, K., Bechly, G., Bybee, S., Dow, R., Dumont, H., Fleck, G., et al (2013) The classification and diversity of dragonflies and damselflies (Odonata). In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). Zootaxa, 3703, 36. DOI: 10.11646/zootaxa.3703.1.9
- Dijkstra, K. & Kalkman, V.J. (2012) Phylogeny, classification and taxonomy of European dragonflies and damselflies (Odonata): a review. Organisms Diversity & Evolution, 12, 209–227. DOI: 10.1007/s13127-012-0080-8
- Dijkstra, K. & Matushkina, N. (2009) Kindred spirits: "*Brachythemis leucosticta*", Africa's most familiar dragonfly, consists of two species (Odonata: Libellulidae). International journal of Odonatology, 12 (2), 237–256. DOI: 10.1080/13887890.2009.9748343
- Ebejer, M.J., Degabriele, G. & Scieberras, A. (2008) An Annotated Checklist of Odonata of the Maltese Islands, with evidence for recent influx of species. Libellula, 27 (1/2), 133–145.
- Engel, M.S., Ceriaco, L.M.P., Daniel, G.M., Dellapé, P., Lobl, I., Marinov, M., et al (2021) The taxonomic impediment: a shortage of taxonomists, not the lack of technical approaches. Zoological Journal of the Linnean Society, 193, 381–387. DOI: 10.1093/zoolinnean/zlab072
- Festi, A. (2011) *Aeshna subarctica elisabethae* new to the fauna of Italy (Odonata: Aeshnidae). Libellula, 30 (1-2), 65–76.
- Fiorenza, T. & Pecile, I. (2009) The pygmy damselfly *Nehalennia speciosa* is still part of the odonate fauna of Italy (Insecta, Odonata, Coenagrionidae). Bollettino Museo civico Storia naturale Venezia, 60, 17–27.
- Galimberti, A., Assandri, G., Maggioni, D., Ramazzotti, F., Baroni, D., Bazzi, G., et al. (2020) Italian odonates in the Pandora's box: A comprehensive DNA barcoding inventory shows taxonomic warnings at the Holarctic scale. Molecular Ecology Resources, 21, 183–200. DOI: 10.1101/2020.04.23.056911
- Garbini, A., (1897) Libellulidi del Veronese e delle Provincie limitrofe. Bollettino della Società Entomologia Italiana, 29, 35–135.
- Gaston, K.J. & Rodrigues, A.S.L., (2003) Reserve Selection in Regions with Poor Biological Data. Conservation Biology, 17, 188–195. DOI: 10.1046/J.1523-1739.2003.01268.X
- Giugliano, L. & Terzani, F. (2011). Gli Odonati delle aree umide retrodunali nel Parco Regionale di Migliarino, San Rossore, Massaciuccoli (Toscana settentrionale). Bollettino Società entomologica italiana, 143 (1), 3–13.
- Günther, A. (2019) Erster Freilandnachweis von *Pantala flavescens* in Deutschland (Odonata: Libellulidae). Libellula, 38, 127–136.
- Hassall, C. (2015) Odonata as candidate macroecological barometers for global climate change. Freshwater Science, 34, 1040–1049. DOI: 10.1086/682210
- Holusa, O. (2008) *Trithemis kirbyi* auf Sardinien: Erstnachweis für Europa (Odonata: Libellulidae). Libellula, 27 (1/2), 111.
- Janni, O., Viganò, M., Corso, A. (2020) First records of *Diplacodes lefebvrii* (Rambur, 1842) for Sicily and additional record of *Trithemis kirbyi* Selys, 1891 (Odonata Libellulidae). Biodiversity Journal, 11, 65–68. DOI: 10.31396/Biodiv.Jour.2020.11.1.65.68
- Kalkman, V.J., Boudot, J.-P., Bernard, R., De Knijf, G., Suhling, F., Termaat, T., (2018) Diversity and conservation of European dragonflies and damselflies (Odonata). Hydrobiologia, 811, 269–282. DOI: 10.1007/s10750-017-3495-6

- Kunz, B., Ober S.V., Jödicke, R. (2006) The distribution of *Zygonyx torridus* in the Palaearctic (Odonata: Libellulidae). *Libellula*, 25 (1/2), 89.
- López-Estrada, K. E., Barona Fernández, J., Cardo-Maeso, N., Teruel Montejano, S.; Díaz-Martínez, C. (2020) *Onychogomphus cazuma* sp. nov. from Spain: Molecular and morphological evidence supports the discovery of a new European dragonfly species (Odonata: Gomphidae). *Odonatologica*, 49, 125–154. DOI: 10.5281/ZENODO.3823337
- Mancini, L., Fornasier, F., Jacomini, C., Utzeri, C. (1993) Reperti *Paragomphus genei*, Bollettino Associazione Romana di Entomologia, 47, 121–122.
- Mastropasqua, F. & Liuzzi, C. (2016) New records of *Coenagrion ornatum* in Italy (Odonata: Coenagrionidae) *Fragmenta Entomologica*, 48 (1), 29–31. DOI: 10.13133/2284-4880/167
- Minelli, A., (1966) Odonati del Veneto. *Memorie della Società Entomologica Italiana*, 45, 111–134.
- Minelli, A., Ruffo, S. & La Posta, S. (eds.) (1993-1995) Checklist delle specie della Fauna d'Italia. Ministero dell'Ambiente, Comitato Scientifico per la Fauna d'Italia. Calderini Editore.
- Nielsen, C. & Conci, C. (1951) Note su Odonati Italiani. *Bollettino della Società Entomologica Italiana*, 81, 76–79.
- Olias, M., Weihrauch, F. & Bedjanic, M. (2007) *Lestes parvidens* and *L. viridis* in southeastern Europe: a chorological analysis (Odonata: Lestidae). *Libellula*, 26, 243–247.
- Ottonello, D. & Oneto, F. (2013). *Libellule di Liguria* Estratto dagli Annali del Museo Civico di Storia Naturale "G. Doria", 105, 297–425.
- Paulson, D., Schoor, M., Deliry, C. (2022) World Odonata List. Revision July 2022. <https://www2.pugetsound.edu/academics/academic-resources/slater-museum/biodiversity-resources/dragonflies/world-odonata-list/> - Accessed 06 July 2022
- Phillips, S.J., Anderson, R.P., Schapire, R.E. (2006) Maximum entropy modeling of species geographic distribution. *Ecological Modelling*, 19, 231–259. DOI: 10.1016/j.ecolmodel.2005.03.026
- Piretta, L. & Assandri, G. (2019) First record of the migrant dragonfly *Pantala flavescens* for mainland Italy (Insecta: Odonata). *Fragmenta Entomologica*, 51, 247–250. DOI: 10.13133/2284-4880/376
- Pirotta, R. (1879) *Libellulidi italiani*. *Annali Museo civico di Storia Naturale*, Genova 14, 401–489.
- Polette, P.P., Abbott, C., Gouys, J., Jenard, P., Juliand, P., Darnaud, S. & Boudot, J. (2017) Premières mentions de *Trithemis kirbyi* (Odonata: Libellulidae) en France. *Martinia*, 33, 15–25.
- Possingham, H.P., Grantham, H., Rondinini, C., (2007) How can you conserve species that haven't been found? *Journal of Biogeography*, 34, 758–759. DOI: 10.1111/j.1365-2699.2007.01717.x
- Rattu A., Leo P., Moratin R., Hardersen S. (2014) *Diplacodes lefebvrei* in Sardinia, a new species for the Italian fauna (Odonata: Libellulidae) *Fragmenta Entomologica*, 46, 121–124. DOI: 10.13133/2284-4880/75
- Riservato, E., Fabbri, R., Festi, A., Grieco, C., Hardersen, S., Landi, F., Utzeri, C., Rondinini, C., Battistoni, A., Teofili, C. (compilatori) (2014) Lista Rossa IUCN delle libellule Italiane. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma.
- Riservato, E., Festi, A., Fabbri, R., Grieco, C., Hardersen, S., La Porta, G., Landi, F., Siesa, M.E., Utzeri, C. (2014) Odonata - Atlante delle libellule italiana - preliminare. Società per lo Studio e la Conservazione delle Libellule. Edizioni Belvedere, Latina, 224 pp.
- Roberts, J.E.H. (1948) *Coenagrion castellani*, a new species of dragonfly in Europe (Order Odonata, Zygoptera). *Proceedings of the Royal Entomological Society of London. Series B, Taxonomy* 17 (5/6), 63–66. DOI: 10.1111/j.1365-3113.1948.tb00888.x
- Rocha-Ortega, M., Rodríguez, P., Córdoba-Aguilar, A. (2019) Can dragonfly and damselfly communities be used as bioindicators of land use intensification? *Ecological Indicators*, 107, 105553. DOI: 10.1016/j.ecolind.2019.105553
- Roché, B., Dommanget, J.-L., Grand, D., Papazian, M., (2008) *Atlas des Odonates de Corse*.

- Direction Régionale de l'Environnement, Service de l'Eau et des Milieux Aquatiques (SEMA). Société française d'Odonatologie. Rapport non publié, 128 pp.
- Ruffo, S., & Stoch F. (2005) Checklist and Distribution of the Italian Fauna. Memorie del Museo Civico di Storia Naturale di Verona. 2. Serie. Sezione Scienze della Vita 16, with CD-ROM. ISSN 0392-0097 and ISBN 88-89230-09-6
- Schmitt, T., Fritz, U., Delfino, M., Ulrich, W., Habel, C. (2021) Biogeography of Italy revisited: genetic lineages confirm major phylogeographic patterns and a pre-Pleistocene origin of its biota. *Frontiers in Zoology*, 18, 34. DOI: 10.1186/s12983-021-00418-9
- Schneider, W., & Utzeri, C. (1994) *Gomphus pulchellus* Sel. new to the Italian odonate fauna (Anisoptera: Gomphidae). *Notulae odonatologicae*, 4, 57–76.
- Sandall, E.L., Pinkert, S. & Jetz, W. (2022) Country-level checklists and occurrences for the world's Odonata (dragonflies and damselflies). *Journal of Biogeography*, 49, 1586–1598. DOI: 10.1111/jbi.14457
- Terzani, F., Cianferoni, F. & Rocchi, S. (2013) Aggiornamento e sintesi delle conoscenze odontologiche dell'Arcipelago Toscano (Insecta Odonata). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 37, 175–196.
- Utzeri, C. (1994) Odonata. In: Minelli, A., S. Ruffo & S. La Posta (eds.) (ed.) "Checklist delle specie della fauna italiana", Calderini, Bologna. 35, 1–7.
- Utzeri, C. & D'Antonio, C. (2005) Insecta Odonata. In: Ruffo, S & Stoch, F. (ed.s), Check-list e distribuzione della fauna italiana. Memorie del Museo. civico di Storia Naturale Verona, 2. serie, Sez. Scienze della Vita: 131.
- Vandelli, D. (1763) Saggio d'Istoria Naturale del lago di Como e della Valsassina, Pavia, 232 pp.
- Vigano, M., Janni, O. & Corso, A. (2017) *Tramea basilaris* on Linosa Island, Italy: A new species for Europe and the Western Palaearctic (Odonata: Libellulidae). *Odonatologica*, 46, 55–66. DOI: 10.5281/zenodo.572356
- Vigna Taglianti, A., Audisio, P.A., Biondi, M., Bologna, M., Carpaneto, G.M., De Biase, A., et al. (1999) A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. *Biogeographia*, 20, 31–59. DOI: 10.21426/B6110172
- Wildermuth, H., Gonseth, Y. & Maibach, A. (2005) Odonata - Les Libellules en Suisse. *Fauna Helvetica*, 11, CSCF/SES, Neuchatel.
- Zandigiacomo, P., Chiandetti, I., Fiorenza, T., Nadalon, G., Uboni, C., (2014) Odonati del Friuli Venezia Giulia: secondo aggiornamento della checklist e ulteriori osservazioni. *Gortania, Botanica, Zoologia*, 36, 33–44.

Submitted: 6 October 2022

First decision: 19 November 2022

Accepted: 9 January 2023

*the Guest editors for the special section
"The new Checklist of the Italian Fauna":*

*Marco A. Bologna, Lucio Bonato,
Fabio Cianferoni, Alessandro Minelli,
Marco Oliverio, Fabio Stoch, Marzio Zapparoli*