

Singular periodic solutions to a critical equation in the Heisenberg group

Claudio Afeltra*

Abstract

We construct positive solutions to the equation

$$-\Delta_{\mathbf{H}^n} u = u^{\frac{Q+2}{Q-2}}$$

on the Heisenberg group, singular in the origin, similar to the Fowler solutions of the Yamabe equations on \mathbf{R}^n . These satisfy the homogeneity property $u \circ \delta_T = T^{-\frac{Q-2}{2}} u$ for some T large enough, where $Q = 2n+2$ and δ_T is the natural dilation in \mathbf{H}^n . We use the Lyapunov-Schmidt method applied to a family of approximate solutions built by periodization from the global regular solution classified in [JL].

MSC2010: 35R03, 35H20, 35J20, 35J61.

Keywords: Subelliptic equations, Perturbation methods.

1 Introduction

Let \mathbf{H}^n be the Heisenberg group with its standard pseudohermitian structure. The problem of studying constant *Webster curvature* pseudohermitian structures conformal to the standard one, in the spirit of the classical Riemannian case, is equivalent to find the positive solutions of the equation

$$-\Delta_{\mathbf{H}^n} u = u^{\frac{Q+2}{Q-2}}, \quad (1)$$

where $\Delta_{\mathbf{H}^n}$ is the sublaplacian and $Q = 2n + 2$ is the homogeneous dimension (in Section 2 we will recall the preliminary definitions about the Heisenberg group).

The positive solutions of equation (1) satisfying some integrability hypotheses were classified by Jerison and Lee [JL], and they correspond to conformal factors that transform the standard pseudohermitian structure of \mathbf{H}^n into the push-forward of the pseudohermitian structure of the sphere $\mathbf{S}^{2n+1} \subset \mathbf{C}^{n+1}$ with respect to the Cayley transform, up to translations and dilations. This

*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa (Italy) - claudio.afeltra@sns.it

classification plays an important role in the solution of the *CR Yamabe problem*, see [JL2], [GamYac], [Gam] and [CMY].

In the Euclidean space the analogous equation,

$$-\Delta_{\mathbf{H}^n} u = u^{\frac{n+2}{n-2}}, \quad (2)$$

is well studied, being related to the Yamabe problem, and being analytically interesting due to a lack of compactness.

The Yamabe equation in \mathbf{R}^n also arises when looking for extremals of the critical Sobolev-Gagliardo-Nirenberg inequality. These were classified as “bubble functions” independently by Aubin [Au] and Talenti [Ta].

A complete classification for solutions of (2) (without integrability hypotheses) was given by Caffarelli, Gidas and Spruck [CGS]. In this case also the solutions on $\mathbf{R}^n \setminus \{0\}$ were classified. In addition to the regular ones on the whole space, there is a singular solution corresponding geometrically to the cylindrical metric, and a family of *singular solutions*, the Fowler solutions, which correspond to a family of periodic metrics on the cylinder which are isometric to the Delaunay surfaces (see [MP1] and the references cited therein).

This terminology is in analogy with the structure for axially symmetric constant mean curvature surfaces: in this case Delaunay surfaces bridge the sphere and the cylinder (see [MP2] and the references cited therein). Furthermore, the Fowler solutions have been used as building blocks (see for example [MP1]) for the construction of more general solutions (as well as for the constant mean curvature Delaunay surfaces).

The above classification has been used to study the asymptotic profiles of general singular solution (see [KMPS]), and solutions with singular behavior as the Fowler’s ones arise in the study of blow-up limits for the prescribed scalar curvature problem (see [L1], [L2], [CL]).

The aim of this article is to prove, in analogy with the Euclidean case, the existence of a family of solutions to equation (1) satisfying a periodicity condition with respect to dilations, that is such that $u \circ \delta_T = T^\alpha u$ for some T, α (see Section 2). for the notation). A simple computation shows that it must necessarily hold that $\alpha = -\frac{Q-2}{2}$. The main result of the paper is the following.

Theorem 1.1. *There exists T_0 such that for $T \geq T_0$ there exists a positive solution of the equation (1) on $\mathbf{H}^n \setminus \{0\}$ such that $u \circ \delta_T = T^{-\frac{Q-2}{2}} u$, and T is the smallest period.*

On the Euclidean space the proof of the uniqueness of such solutions relies on a result (in [CGS]), proved by the moving planes method, stating that the positive solutions of equation (1) are radially symmetric. In this way the construction of solutions and their classification is carried out by a standard ODE analysis. This cannot be done on \mathbf{H}^n . We point out that on the Heisenberg group one cannot expect a symmetric solution, because the sublaplacian is not rotationally invariant. We also point out the recent results in [GMM], where solutions with singularities at higher-dimensional sets were constructed with different methods.

Theorem 1.1 is proved by writing equation (1) as the variational equation of the functional

$$\mathcal{J}_T(u) = \int_{\Omega_T} \left(|\nabla_{\mathbf{H}^n} u|^2 - \frac{1}{2^*} |u|^{2^*} \right)$$

on a space of functions satisfying $u \circ \delta_T = T^{-\frac{Q-2}{2}} u$ (the integral is with respect to the Haar measure, see Section 2).

In Section 3 we will find an estimate of the Sobolev constant for periodic solution through a Hardy-Littlewood-Sobolev type theorem for Lorentz spaces. This will be used to carry out the estimates in the subsequent sections.

In Section 4 we will build a family \mathcal{Z}_T of approximate critical points of \mathcal{J}_T by gluing a sequence of suitable dilations of the global regular solution ω_λ . We will show that these solutions are “almost critical” points, in the sense that on \mathcal{Z}_T the differential of the functional \mathcal{J}_T is small.

In Section 5 we will prove that a non degeneracy condition holding for ω_λ can be carried on Ψ_λ .

In the final Section we will prove the existence of the desired solutions through the Lyapunov-Schmidt method, reducing the problem to the orthogonal of the tangent of the curve \mathcal{Z}_T , and therein applying the contraction Theorem.

We believe that the construction should give perspectives for the study of more general singular solutions in the Heisenberg group, in the spirit of the cited results on the Euclidean space.

2 Preliminaries and notation

In this section we recall some basic definitions and facts on the Heisenberg group, widely present in the literature. See, for example, Chapter 10 of [CS].

Let us consider the Heisenberg group $\mathbf{H}^n = \mathbf{C}^n \times \mathbf{R}$, with the convention on the product

$$(z_1, t_1) \cdot (z_2, t_2) = (z_1 + z_2, t_1 + t_2 + 2 \operatorname{Im}(z_1 \cdot \overline{z_2})).$$

Let

$$\begin{aligned} X_i &= T_i = \frac{\partial}{\partial x_i} + 2y_i \frac{\partial}{\partial t}, \\ Y_i &= T_{n+i} = \frac{\partial}{\partial y_i} - 2x_i \frac{\partial}{\partial t}, \\ T &= T_0 = \frac{\partial}{\partial t} \end{aligned}$$

be the standard basis of left invariant vector fields,

$$\nabla_{\mathbf{H}^n} u = \sum_i X_i(u) X_i + Y_i(u) Y_i$$

the subriemannian gradient,

$$\operatorname{div} \left(\sum_i f_i X_i + g_i Y_i \right) = \sum_i X_i(f_i) + Y_i(g_i)$$

the divergence (which coincides with the divergence with respect to a Haar volume form), and

$$\Delta_{\mathbf{H}^n} = \operatorname{div} \circ \nabla_{\mathbf{H}^n} = \sum_i X_i^2 + Y_i^2$$

be the sublaplacian. There exists a constant $C = C(n)$ such that

$$K(x) = \frac{C}{|x|^{Q-2}} \quad (3)$$

is a fundamental solution of the sublaplacian. Let

$$S^1(\mathbf{H}^n) = \{u \in L^2(\mathbf{H}^n) \mid X_i u, Y_i u \in L^2(\mathbf{H}^n)\}.$$

We endow \mathbf{H}^n with the set of dilations

$$\delta_\lambda(z, t) = (\lambda z, \lambda^2 t)$$

and with the homogeneous norm

$$|(z, t)| = (|z|^4 + t^2)^{1/4}.$$

The Lebesgue measure dx is a biinvariant Haar measure on \mathbf{H}^n satisfying

$$(\delta_\lambda)_\# dx = \lambda^Q dx;$$

this is essentially the reason why Q takes the place of the topological dimension n in many analytic questions.

Let us set $B_r = \{|x| < r\}$ and $\Omega_T = \overline{B}_R \setminus B_1$. We define the Hilbert space

$$X_T = \{u \in S^1_{\text{loc}}(\mathbf{H}^n) \mid u \circ \delta_T = T^{-\frac{Q-2}{2}} u\}$$

with the product

$$\langle u, v \rangle = \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v.$$

Let \tilde{X}_T the closed subspace of X_T of the functions of the form $u(|z|, t)$.

It is known that the positive solutions of the equation (1) are

$$\omega_\lambda = \lambda^{(2-Q)/2} \omega \circ \delta_{\lambda^{-1}}$$

and the translates thereof, where

$$\omega(z, t) = c_0 \frac{1}{(t^2 + (1 + |z|^2)^2)^{(Q-2)/4}}.$$

The problem is variational: the solutions in $S^1(\mathbf{H}^n)$ of the equation are the critical points of the functional

$$\mathcal{J}(u) = \int_{\mathbf{H}^n} \left(|\nabla_{\mathbf{H}^n} u|^2 - \frac{1}{2^*} |u|^{2^*} \right).$$

Analogously the solutions of the equation of the equation in X_T are the critical points of the functional

$$\mathcal{J}_T(u) = \int_{\Omega_T} \left(|\nabla_{\mathbf{H}^n} u|^2 - \frac{1}{2^*} |u|^{2^*} \right).$$

It holds that

$$d \mathcal{J}_T(u)[\varphi] = \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \varphi - u |u|^{2^*-2} \varphi$$

and that

$$d^2 \mathcal{J}_T(u)[\varphi, \psi] = \int_{\Omega_T} \nabla_{\mathbf{H}^n} \varphi \cdot \nabla_{\mathbf{H}^n} \psi - (2^* - 1) |u|^{2^*-2} \varphi \psi.$$

We call \mathcal{J}_T'' the operator associated with this bilinear form in the natural way:

$$\langle \mathcal{J}_T''(u)[\varphi], \psi \rangle = d^2 \mathcal{J}_T(u)[\varphi, \psi].$$

Let us notice that, if $u \in X_T$ and $E \subseteq \mathbf{H}^n$ then

$$\int_{\delta_r(E)} |u|^{\frac{2Q}{Q-2}} = \int_E |r^{\frac{Q-2}{2}} u \circ \delta_r|^{\frac{2Q}{Q-2}} \quad (4)$$

and

$$\int_{\delta_r(E)} |\nabla_{\mathbf{H}^n} u|^2 = \int_E |r^{\frac{Q-2}{2}} \nabla_{\mathbf{H}^n} (u \circ \delta_r)|^2. \quad (5)$$

In particular, if $1 \leq r \leq T$ then

$$\begin{aligned} \int_{\delta_r \Omega_T} |u|^{\frac{2Q}{Q-2}} &= \int_{\Omega_T \setminus \Omega_r} |u|^{\frac{2Q}{Q-2}} + \int_{\Omega_r \setminus \Omega_T} |u|^{\frac{2Q}{Q-2}} = \\ &= \int_{\Omega_T \setminus \Omega_r} |u|^{\frac{2Q}{Q-2}} + \int_{\Omega_r} |T^{\frac{Q-2}{2}} u \circ \delta_T|^{\frac{2Q}{Q-2}} = \int_{\Omega_T} |u|^{\frac{2Q}{Q-2}}, \end{aligned} \quad (6)$$

and by induction and inversion one can extend this formula to every value of r . Analogously

$$\int_{\delta_r \Omega_T} |\nabla_{\mathbf{H}^n} u|^2 = \int_{\Omega_T} |\nabla_{\mathbf{H}^n} u|^2, \quad (7)$$

and by polarization

$$\int_{\delta_r \Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v = \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v. \quad (8)$$

The following lemma shows that in integration by parts in X_T boundary terms are null.

Lemma 2.1. *If $u, v \in X_T$ then*

$$\int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v = - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v.$$

Proof. Let us write $v = v_1 + v_2$ with $u_1, u_2 \in X_T$, $\text{supp } u_1 \cap \Omega_T \subset B_{7(T+1)/8} \setminus B_{(T+1)/8}$ and $\text{supp } u_2 \cap \delta_{(T+1)/2}\Omega_T \subset B_{(T+1)^2/8} \setminus B_{3(T+1)/4}$ (this can be carried out through a partition of unity).

Then, using formula (8),

$$\begin{aligned}
\int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v &= \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v_1 + \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v_2 = \\
&= - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v_1 + \int_{\delta_{(T+2)/2}\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} v_2 = \\
&= - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v_1 - \int_{\delta_{(T+2)/2}\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v_2 = \\
&= - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v_1 - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v_2 = - \int_{\Omega_T} \Delta_{\mathbf{H}^n} u \cdot v.
\end{aligned}$$

□

We will need to restrict ourselves to solutions in \tilde{X}_T . In order to do this, we observe that, under the identification $\mathbf{H}^n = \mathbf{R}^{2n} \times \mathbf{R}$, the functional \mathcal{J}_T is invariant by the group of transformations of the form $(z, t) \mapsto (Az, t)$ with $A \in O(\mathbf{R}^{2n}) \cap Sp(\mathbf{R}^{2n})$. In fact it is known that if $A = (a_{ij}) \in Sp(\mathbf{R}^{2n})$ then this transformation is a group automorphism of \mathbf{H}^n (see [Fol], Chapter 1, Section 2), and so it maps the fields T_i into the fields $\sum_j a_{ij} T_j$. So, using the fact that $A \in O(\mathbf{R}^{2n})$, it is easy to verify that \mathcal{J}_T is invariant by this group.

Furthermore, under the canonical identification of \mathbf{R}^{2n} with \mathbf{C}^n , $O(\mathbf{R}^{2n}) \cap Sp(\mathbf{R}^{2n}) = U(\mathbf{C}^n)$ ([Fol], Proposition 4.6).

Since $U(n)$ acts transitively on the unit sphere of \mathbf{C}^n , \tilde{X}_T is the set of the functions in X_T invariant under the transformations of this form, and so, by Palais' criticality principle [Pal], the critical points of the restriction of \mathcal{J}_T to it are critical points in all of X_T .

In the sequel we will need also a particular vector field that plays an important role in \mathbf{H}^n (and more in general in homogeneous groups), the generator of the dilations. It is characterized by the equation

$$\left. \frac{d}{d\lambda} \right|_{\lambda=1} (u \circ \delta_\lambda) = Zu$$

for every $u \in \mathcal{C}^1(\mathbf{H}^n)$. An explicit expression for it is

$$Z = \sum_{i=1}^n x_i \frac{\partial}{\partial x_i} + y_i \frac{\partial}{\partial y_i} + 2t \frac{\partial}{\partial t}.$$

It is easy to verify that

$$\lambda \frac{d}{d\lambda} (u \circ \delta_\lambda) = Z(u \circ \delta_\lambda) = (Zu) \circ \delta_\lambda. \quad (9)$$

Using this formula, it is easy to prove that a function u is homogeneous of degree α if and only if $Zu = \alpha u$ (an extension to \mathbf{H}^n of Euler's theorem).

Furthermore it holds that $[X_i, Z] = X_i$ and $[Y_i, Z] = Y_i$, and so $[\nabla_{\mathbf{H}^n}, Z] = \nabla_{\mathbf{H}^n}$.

2.1 Lorentz spaces

In Section 3, to overcome the non integrability of the functions in X_T in the whole space, will need to use the Lorentz spaces, which we recall briefly.

Given a σ -finite measure space (X, μ) and $1 \leq p < \infty$, $1 \leq q \leq \infty$, the Lorentz quasinorm is defined as

$$\|u\|_{L^{p,q}(X)} = p^{1/q} \left\| \lambda \mu\{|u| > \lambda\}^{1/p} \right\|_{L^q(dt/t)}.$$

Furthermore one defines $\|u\|_{L^{\infty,\infty}(X)} = \|u\|_{L^\infty(X)}$. The Lorentz space $L^{p,q}(X)$ is the set of functions such that this quantity is finite. When $p = q$, $\|u\|_{L^{p,p}} = \|u\|_{L^p}$, while when $q = \infty$, $L^{p,\infty}$ coincides with weak L^p .

We will need the following generalization of the Young inequality, which sometimes is referred in the literature as Young-O'Neil inequality. It can be deduced applying Theorem 2.6 in [ON] (with the corrections in [Yap]) and Theorem 1.2.12, Remark 1.2.11 in [Gr].

Theorem 2.2. *If $1 < p, p_1, p_2 < \infty$, $1 \leq q, q_1, q_2 \leq \infty$ are such that*

$$\frac{1}{p_1} + \frac{1}{p_2} = 1 + \frac{1}{p} \quad \text{and} \quad \frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{q}$$

then there exists C such that for every $f \in L^{p_1, q_1}(\mathbf{H}^n)$, $g \in L^{p_2, q_2}(\mathbf{H}^n)$ it holds

$$\|fg\|_{L^{p,q}(\mathbf{H}^n)} \leq C \|f\|_{L^{p_1, q_1}(\mathbf{H}^n)} \|g\|_{L^{p_2, q_2}(\mathbf{H}^n)}.$$

2.2 Basic definitions on CR geometry

For convenience of the reader, we recall the basic definitions about CR manifolds, also if we will not use them. The reader can find more on the topic in [Bog], [DTom].

A CR manifold is a real smooth manifold M endowed with a subbundle \mathcal{H} of the complexified tangent bundle of M , $T^{\mathbf{C}}M$, such that $\mathcal{H} \cap \overline{\mathcal{H}} = \{0\}$ and $[\mathcal{H}, \mathcal{H}] \subseteq \mathcal{H}$. We will assume M to be of hypersurface type, that is that $\dim M = 2n + 1$ and that $\dim \mathcal{H} = n$. There exists a non-zero real differential form θ that is zero on $\Re(\mathcal{H} \oplus \overline{\mathcal{H}})$; it is unique up to scalar multiple by a function. Such a form is called pseudohermitian structure. On a pseudohermitian manifold, the Levi form is defined as the 2-form on \mathcal{H} $L_\theta(V, W) = -id\theta(V, \overline{W}) = id\theta([V, \overline{W}])$. A CR manifold is said to be pseudoconvex if it admits a positive definite Levi form (this implies every Levi form to be definite).

The Heisenberg group is the simplest pseudoconvex CR manifold, if endowed with the bundle $\mathcal{H} = \text{span}(Z_1, \dots, Z_n)$ with $Z_j = \frac{1}{2}(X_j - iY_j)$.

On a nondegenerate pseudohermitian manifold one can define a connection, the Tanaka-Webster connection. This allows to define curvature operators in an analogous manner as in Riemannian geometry: the pseudohermitian curvature tensor is the curvature of the Tanaka-Webster connection, the Ricci tensor is

$$\text{Ric}(X, Y) = \text{trace}(Z \mapsto R(Z, X)Y),$$

and the Webster scalar curvature is the trace of the Ricci tensor with respect to the Levi form.

Being a pseudohermitian structure defined only up to a conformal factor on a CR manifold, in CR geometry the Yamabe problem is even more natural than in Riemannian geometry. If $\tilde{\theta} = u^{2/n}\theta$, the transformation law of the Webster curvature is

$$\tilde{W} = u^{-1-2/n} \left(\frac{2n+2}{n} \Delta_b u + Wu \right),$$

where Δ_b is the sublaplacian, which can be defined in a similar way as the Heisenberg group. So the Yamabe problem takes to the equation

$$\frac{2n+2}{n} \Delta_b u + Wu = \lambda u^{1+2/n}.$$

Since the Heisenberg group has zero Webster curvature, and since the pseudohermitian sublaplacian coincides with the sublaplacian defined formerly, the Yamabe problem, up to an inessential constant, is equivalent to find positive solution to equation 1.

The solution of this case plays in the solution in the general case the same role that the solution on \mathbf{R}^n plays in the solution of the general Riemannian case.

3 Estimate of the Sobolev constant on X_T

In order to carry out the estimates in the next Sections, we will need an explicit bound on the Sobolev constant on X_T . We will achieve this relating the L^p norm on Ω_T and the $L^{p,\infty}$ norm on the whole space.

Proposition 3.1. *If f is an L^p_{loc} function on $\mathbf{H}^n \setminus \{0\}$ such that $f \circ \delta_T = T^{-\alpha} f$ and $\alpha p = Q$ then*

$$\left(\frac{T^Q - 1}{T^Q} \right)^{1/p} \|u\|_{L^{p,\infty}(\mathbf{H}^n)} \leq C_2 \|u\|_{L^p(\Omega_T)} \leq Q^{1/p} (\log T)^{1/p} \|u\|_{L^{p,\infty}(\mathbf{H}^n)}.$$

Proof. Let us call $f(\lambda) = \mu\{x \in \Omega_T \mid u(x) > \lambda\}$ and $g(t) = \mu\{x \in \mathbf{H}^n \mid u(x) > \lambda\}$. Then it holds that

$$g(\lambda) = \sum_{k \in \mathbf{Z}} T^{Qk} f(\lambda T^{\alpha k}).$$

Therefore for every $\lambda > 0$

$$\begin{aligned}
\|u\|_{L^p(\Omega_T)}^p &= p \int_0^\infty \xi^{p-1} f(\xi) d\xi = p \sum_{k \in \mathbf{Z}} \int_{\lambda T^{\alpha(k-1)}}^{\lambda T^{\alpha k}} \xi^{p-1} f(\xi) d\xi \geq \\
&\geq p \sum_{k \in \mathbf{Z}} f(\lambda T^{\alpha k}) \int_{\lambda T^{\alpha(k-1)}}^{\lambda T^{\alpha k}} \xi^{p-1} d\xi = \frac{T^Q - 1}{T^Q} \lambda^p \sum_{k \in \mathbf{Z}} T^{Qk} f(\lambda T^{\alpha k}) = \\
&= \frac{T^Q - 1}{T^Q} \lambda^p g(\lambda).
\end{aligned}$$

Taking the supremum with respect to λ we get the first inequality.

For the other one, let us pick an integer $N > 0$ and write

$$\begin{aligned}
\|u\|_{L^p(\Omega_T)}^p &= p \int_0^\infty \xi^{p-1} f(\xi) d\xi = p \sum_{k \in \mathbf{Z}} \int_{T^{\alpha k/N}}^{T^{\alpha(k+1)/N}} \xi^{p-1} f(\xi) d\xi \leq \\
&\leq p \sum_{k \in \mathbf{Z}} f(T^{\alpha k/N}) \int_{T^{\alpha k/N}}^{T^{\alpha(k+1)/N}} \xi^{p-1} d\xi = \\
&= \sum_{m=1}^N \sum_{j \in \mathbf{Z}} (T^{\alpha p/N} - 1) T^{\alpha p j} T^{\alpha p m/N} f(T^{\alpha j} T^{\alpha m/N}) = \\
&= (T^{Q/N} - 1) \sum_{m=1}^N T^{Qm/N} \sum_{j \in \mathbf{Z}} T^{Qj} f(T^{\alpha j} T^{\alpha m/N}) = \\
&= (T^{Q/N} - 1) \sum_{m=1}^N T^{Qm/N} g(T^{\alpha m/N}) \leq N (T^{Q/N} - 1) \|u\|_{L^{p,\infty}}^p.
\end{aligned}$$

Taking the limit for $N \rightarrow \infty$ we get the second inequality. \square

Using the Theorem 2.2 we can prove a Sobolev type inequality for weak L^p spaces.

Proposition 3.2. *There exists a constant C such that for every function $u \in L^{2,\infty}(\mathbf{H}^n)$ such that $\nabla u \in L^{2,\infty}(\mathbf{H}^n)$ verifies*

$$\|u\|_{L^{\frac{2Q}{Q-2},\infty}} \leq C \|\nabla u\|_{L^{2,\infty}}.$$

Proof. Let $E = u > 1$, $E^c = \mathbf{H}^n \setminus E$, $u_1 = u \chi_{E^c} + \chi_E$ and $u_2 = (u-1) \chi_E$, so that $u = u_1 + u_2$. It is standard to prove that u_1 and u_2 have weak subriemannian gradient and that $\nabla_{\mathbf{H}} u_1 = (\nabla_{\mathbf{H}} u) \chi_{E^c}$, $\nabla_{\mathbf{H}} u_2 = (\nabla_{\mathbf{H}} u) \chi_E$ (the proof is the same as on \mathbf{R}^n). It is easy to prove that $u_1 \in S^p(\mathbf{H}^n)$ for $p > 2$ and that $u_2 \in S^q(\mathbf{H}^n)$ for $q < 2$. If $\varphi \in \mathcal{C}_c^\infty(\mathbf{H}^n)$ it holds that

$$\varphi(x) = (\varphi * \delta)(x) = (\varphi * (-\Delta_{\mathbf{H}^n} K))(x) =$$

$$= \int_{\mathbf{H}^n} (\nabla_{\mathbf{H}^n} \varphi)(xy^{-1}) * (\nabla_{\mathbf{H}^n} K)(y) dy := (\nabla_{\mathbf{H}^n} \varphi * \nabla_{\mathbf{H}^n} K)(x) \quad (10)$$

Formula (3) implies that $\nabla_{\mathbf{H}^n} K \in L^{\frac{Q}{Q-1}, \infty}$, and so, by Theorem 2.2, the operator $f \mapsto f * \nabla_{\mathbf{H}^n} K$ is bounded from L^p and L^q to some other Lebesgue spaces. Therefore, using the density of \mathcal{C}_c^∞ in $S^p(\mathbf{H}^n)$ for $1 \leq p < \infty$, formula (10) holds almost everywhere for functions in these spaces, and so it holds for u_1 and u_2 . By summing one obtains that

$$u = \nabla_{\mathbf{H}^n} u * \nabla_{\mathbf{H}^n} K.$$

The thesis follows applying Theorem 2.2 once more. \square

We point out that in the proof of the last Proposition the splitting of u in two pieces belonging to some L^p space was necessary because \mathcal{C}_c^∞ functions are not dense in the weak L^p spaces.

Combining Propositions 3.1 and 3.2 we get the following Sobolev theorem for X_T spaces with an explicit constant.

Proposition 3.3. *There exist a constant C independent by T such that for every $u \in X_T$*

$$\|u\|_{L^{\frac{2Q}{Q-2}}(\Omega_T)} \leq C(\log T)^{\frac{Q-2}{2Q}} \left(\frac{T^Q}{T^Q - 1} \right)^{1/2} \|u\|_{X_T}.$$

4 Construction of a family of approximate solutions

In order to apply a perturbative method, we find a family of approximate stationary points of \mathcal{J}_T for T big enough.

The family is the following:

$$\Psi_{\lambda, T} = \sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} = \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \omega_\lambda \circ \delta_{T^k}$$

(we will hide the dependence by T whether not necessary). The series converges uniformly on compact sets, because, if $x \in K$,

$$\begin{aligned} \Psi_\lambda(x) &= \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \omega_\lambda \circ \delta_{T^k} \leq \\ &\leq C_{\lambda, K} \sum_{k \geq 0} T^{\frac{Q-2}{2}k} \frac{1}{T^{k(Q-2)}} + C_{\lambda, K} \sum_{k < 0} T^{\frac{Q-2}{2}k} \leq C_{\lambda, K}. \end{aligned}$$

The subriemannian gradient satisfies

$$|\nabla_{\mathbf{H}^n} \Psi_\lambda(x)| \leq \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} T^k |\nabla_{\mathbf{H}^n} \omega_\lambda| \circ \delta_{T^k} \leq$$

$$\leq C_{\lambda,K} \sum_{k \geq 0} T^{\frac{Q}{2}k} \frac{1}{T^{k(Q-1)}} + C_{\lambda,K} \sum_{k < 0} T^{\frac{Q}{2}k} \leq C_{\lambda,K}$$

and so it converges uniformly on compact sets. The same holds for higher order subriemannian derivatives. $\Psi_\lambda \in X_T$ because

$$\Psi_\lambda \circ \delta_T = \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \omega_\lambda \circ \delta_{T^k} \circ \delta_T = T^{-\frac{Q-2}{2}} \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \omega_\lambda \circ \delta_{T^k} = T^{-\frac{Q-2}{2}} \Psi_\lambda.$$

It holds that

$$\begin{aligned} \Psi_{T\lambda} &= \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \omega_{T\lambda} \circ \delta_{T^k} = \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}k} \frac{1}{(T\lambda)^{\frac{Q-2}{2}}} \omega \circ \delta_{1/T\lambda} \circ \delta_{T^k} = \\ &= \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}(k-1)} \frac{1}{\lambda^{\frac{Q-2}{2}}} \omega \circ \delta_{1/\lambda} \circ \delta_{T^{k-1}} = \sum_{k \in \mathbf{Z}} T^{\frac{Q-2}{2}(k-1)} \omega_\lambda \circ \delta_{T^{k-1}} = \Psi_\lambda. \end{aligned}$$

Therefore the set $\mathcal{X}_T = \{\Psi_\lambda \mid \lambda \in (0, \infty)\}$ is a closed curve in X_T .

Moreover, using formula (9), it can be computed that

$$\begin{aligned} \frac{\partial \Psi_\lambda}{\partial \lambda} &= \frac{\partial}{\partial \lambda} \sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} = \sum_{k \in \mathbf{Z}} \frac{\partial}{\partial \lambda} \left(\lambda^{-\frac{Q-2}{2}} \omega_{1/T^k} \circ \delta_{\lambda^{-1}} \right) = \\ &= \sum_{k \in \mathbf{Z}} \left(-\frac{Q-2}{2} \frac{1}{\lambda} \omega_{\lambda/T^k} - \lambda^{-\frac{Q-2}{2}} \frac{1}{\lambda^2} \lambda Z(\omega_{1/T^k} \circ \delta_{\lambda^{-1}}) \right) = \\ &= \sum_{k \in \mathbf{Z}} \left(-\frac{Q-2}{2} \frac{1}{\lambda} \omega_{\lambda/T^k} - \frac{1}{\lambda} Z(\omega_{\lambda/T^k}) \right) = \\ &= -\frac{Q-2}{2} \frac{1}{\lambda} \Psi_\lambda - \frac{1}{\lambda} Z(\Psi_\lambda). \end{aligned} \tag{11}$$

This implies that the curve \mathcal{X}_T is immersed for T big enough, because if $\frac{\partial \Psi_\lambda}{\partial \lambda}$ was zero then $Z(\Psi_\lambda) = -\frac{Q-2}{2} \Psi_\lambda$ would be zero, and by the aforementioned Euler's theorem Ψ_λ would be homogeneous of degree $-\frac{Q-2}{2}$; but it is clearly not by construction if T is big enough.

We want to prove the following proposition.

Proposition 4.1. *For every ε there exists T_0 , depending only on n , such that if $T \geq T_0$ then $\|\nabla_{\mathbf{H}^n} \mathcal{J}_T\| < \varepsilon$ on \mathcal{X}_T .*

We divide the proof in several lemmas.

First we compute the differential of \mathcal{J}_T in Ψ_λ :

$$\begin{aligned} d\mathcal{J}_T(\Psi_\lambda)[u] &= \int_{\Omega_T} \nabla_{\mathbf{H}^n} \Psi_\lambda \cdot \nabla_{\mathbf{H}^n} u - \Psi_\lambda^{2^*-1} u = \\ &= \int_{\Omega_T} \sum_{k \in \mathbf{Z}} \nabla_{\mathbf{H}^n} \omega_{\lambda/T^k} \cdot \nabla_{\mathbf{H}^n} u - \left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{2^*-1} u = \end{aligned}$$

$$\begin{aligned}
&= \sum_{k \in \mathbf{Z}} \left(\int_{\Omega_T} \nabla_{\mathbf{H}^n} \omega_{\lambda/T^k} \cdot \nabla_{\mathbf{H}^n} u - \omega_{\lambda/T^k}^{\frac{Q+2}{Q-2}} u \right) + \\
&\quad - \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k}^{\frac{Q+2}{Q-2}} \right] u = \\
&\qquad\qquad\qquad := A + B. \tag{12}
\end{aligned}$$

Lemma 4.2. *In the above notation, $A = 0$.*

Proof. We have

$$\begin{aligned}
A &= \sum_{k \in \mathbf{Z}} \int_{\Omega_T} T^{\frac{Q-2}{2}k} (\nabla_{\mathbf{H}^n} \omega_\lambda) \circ \delta_{T^k} \cdot \nabla_{\mathbf{H}^n} u - (T^{\frac{Q-2}{2}k})^{\frac{Q+2}{Q-2}} (\omega_\lambda \circ \delta_{T^k})^{\frac{Q+2}{Q-2}} u = \\
&= \sum_{k \in \mathbf{Z}} \int_{\Omega_T} T^{\frac{Q}{2}k} (\nabla_{\mathbf{H}^n} \omega_\lambda) \circ \delta_{T^k} \cdot \nabla_{\mathbf{H}^n} u - T^{\frac{Q+2}{2}k} (\omega_\lambda \circ \delta_{T^k})^{\frac{Q+2}{Q-2}} u = \\
&= \sum_{k \in \mathbf{Z}} \int_{\delta_{T^k}(\Omega_T)} T^{-kQ} \left[T^{\frac{Q}{2}k} \nabla_{\mathbf{H}^n} \omega_\lambda \cdot (\nabla_{\mathbf{H}^n} u) \circ \delta_{T^{-k}} - T^{\frac{Q+2}{2}k} \omega_\lambda^{\frac{Q+2}{Q-2}} u \circ \delta_{T^{-k}} \right] = \\
&= \sum_{k \in \mathbf{Z}} \int_{\delta_{T^k}(\Omega_T)} T^{-\frac{Q}{2}k} T^k \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} (u \circ \delta_{T^{-k}}) - \omega_\lambda^{\frac{Q+2}{Q-2}} u = \\
&= \sum_{k \in \mathbf{Z}} \int_{\delta_{T^k}(\Omega_T)} \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} u - \omega_\lambda^{\frac{Q+2}{Q-2}} u = \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} u - \omega_\lambda^{\frac{Q+2}{Q-2}} u
\end{aligned}$$

Let us pick a family of smooth functions $\varphi_{\varepsilon,R}$ such that $\varphi_{\varepsilon,R} \equiv 1$ on $B_R \setminus B_{2\varepsilon}$, $\varphi_{\varepsilon,R} \equiv 0$ on B_ε and $\mathbf{H}^n \setminus B_{R+1}$, $|\nabla_{\mathbf{H}^n} \varphi_{\varepsilon,R}| \leq \frac{C}{\varepsilon}$ on $B_{2\varepsilon} \setminus B_\varepsilon$ and $|\nabla_{\mathbf{H}^n} \varphi_{\varepsilon,R}| \leq C$ on $B_{R+1} \setminus B_R$. Then

$$\begin{aligned}
A &= \lim_{\substack{\varepsilon \rightarrow 0 \\ R \rightarrow \infty}} \int_{\mathbf{H}^n} \left(\nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} u - \omega_\lambda^{\frac{Q+2}{Q-2}} u \right) \varphi_{\varepsilon,R} = \\
&= \lim_{\substack{\varepsilon \rightarrow 0 \\ R \rightarrow \infty}} \int_{\mathbf{H}^n} -(\Delta_{\mathbf{H}^n} \omega_\lambda + \omega_\lambda^{\frac{Q+2}{Q-2}}) u \varphi_{\varepsilon,R} - u \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} \varphi_{\varepsilon,R} = \\
&= - \lim_{R \rightarrow \infty} \int_{B_{R+1} \setminus B_R} u \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} \varphi_{\varepsilon,R} - \lim_{\varepsilon \rightarrow 0} \int_{B_{2\varepsilon} \setminus B_\varepsilon} u \nabla_{\mathbf{H}^n} \omega_\lambda \cdot \nabla_{\mathbf{H}^n} \varphi_{\varepsilon,R}.
\end{aligned}$$

If $x \rightarrow \infty$ then $\nabla_{\mathbf{H}^n} \omega_\lambda \lesssim \frac{1}{|x|^{Q-1}}$ and $u \lesssim |x|^{-\frac{Q-2}{2}}$, and so the first limit is zero.

If $x \rightarrow 0$ then $\nabla_{\mathbf{H}^n} \omega_\lambda \lesssim 1$ and $u \lesssim |x|^{-\frac{Q-2}{2}}$, and so also the second limit is zero. Therefore $A = 0$. \square

Now we have to estimate the term B from formula (12).

Lemma 4.3. *In the above notation*

$$|B| \leq C(T) \|u\|_{X_T},$$

where $C(T)$ tends to zero uniformly in λ as T tends to infinity.

Proof.

$$\begin{aligned} |B| &\leq \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k}^{\frac{Q+2}{Q-2}} \right] |u| \leq \\ &\leq \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \omega_{\lambda}^{\frac{Q+2}{Q-2}} \right] |u| \leq \\ &\leq \left\{ \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \omega_{\lambda}^{\frac{Q+2}{Q-2}} \right]^{\frac{2Q}{Q+2}} \right\}^{\frac{Q+2}{2Q}} \|u\|_{L^{\frac{2Q}{Q-2}}(\Omega_T)} \leq \\ &\leq C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \omega_{\lambda}^{\frac{Q+2}{Q-2}} \right]^{\frac{2Q}{Q+2}} \right\}^{\frac{Q+2}{2Q}} = \\ &= C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \cdot \\ &\cdot \left\{ \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} |x|^{\frac{Q-2}{2}} \omega_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - (|x|^{\frac{Q-2}{2}} \omega_{\lambda})^{\frac{Q+2}{Q-2}} \right]^{\frac{2Q}{Q+2}} \frac{dx}{|x|^Q} \right\}^{\frac{Q+2}{2Q}} \end{aligned}$$

by Proposition 3.3 (taking $T \geq T_0 > 1$, since we are going to make a limit for $T \rightarrow \infty$). Let us define $\eta_{\lambda} = |x|^{\frac{Q-2}{2}} \omega_{\lambda}$. Then

$$|B| \leq C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \int_{\Omega_T} \left[\left(\sum_{k \in \mathbf{Z}} \eta_{\lambda/T^k} \right)^{\frac{Q+2}{Q-2}} - \eta_{\lambda}^{\frac{Q+2}{Q-2}} \right]^{\frac{2Q}{Q+2}} \frac{dx}{|x|^Q} \right\}^{\frac{Q+2}{2Q}}.$$

By periodicity we can suppose that $\frac{|x|}{\lambda} \in \left[\frac{1}{\sqrt{T}}, \sqrt{T} \right]$, with $\lambda = \lambda(x)$. The function η_{λ} is bounded and tends to zero for $|x| \rightarrow 0, \infty$. If $k \geq 0$ and T is large enough then η_{λ/T^k} satisfies estimates

$$|\eta_{\lambda/T^k}(x)| \lesssim \left(\frac{\left(\frac{T^k}{\lambda}\right) |x|}{1 + \left(\frac{T^k}{\lambda}\right)^2 |x|^2} \right)^{\frac{Q-2}{2}} \lesssim \left(T^k \frac{|x|}{\lambda} \right)^{-\frac{Q-2}{2}} \leq \left(\frac{1}{T} \right)^{\left(k - \frac{1}{2}\right) \frac{Q-2}{2}}$$

and

$$|\eta_{\lambda/T^{-k}}(x)| \lesssim \left(\frac{\left(\frac{T^{-k}}{\lambda}\right)|x|}{1 + \left(\frac{T^{-k}}{\lambda}\right)^2|x|^2} \right)^{\frac{Q-2}{2}} \lesssim \left(\frac{1}{T^k} \frac{|x|}{\lambda} \right)^{\frac{Q-2}{2}} \leq \left(\frac{1}{T} \right)^{(k-\frac{1}{2})\frac{Q-2}{2}}$$

uniformly in λ . It is easy to verify that, for $\alpha, \beta \geq 1$ the function

$$\frac{[(x+y)^\alpha - x^\alpha]^\beta}{x^{(\alpha-1)\beta}y^\beta + y^{\alpha\beta}}$$

is bounded on $(0, \infty)^2$, and so there exist C such that

$$[(x+y)^\alpha - x^\alpha]^\beta \leq C(x^{(\alpha-1)\beta}y^\beta + y^{\alpha\beta})$$

for $x, y \geq 0$. Taking

$$x = \eta_\lambda, \quad y = \sum_{k \in \mathbf{Z} \setminus \{0\}} \eta_{\lambda/T^k}, \quad \alpha = \frac{Q+2}{Q-2} \quad \text{and} \quad \beta = \frac{2Q}{Q+2}$$

one gets that

$$|B| \leq C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \int_{\Omega_T} \left[\eta_\lambda^{\frac{8Q}{(Q+2)(Q-2)}} \left(\sum_{k \in \mathbf{Z} \setminus \{0\}} \eta_{\lambda/T^k} \right)^{\frac{2Q}{Q+2}} + \left(\sum_{k \in \mathbf{Z} \setminus \{0\}} \eta_{\lambda/T^k} \right)^{\frac{2Q}{Q-2}} \right] \frac{dx}{|x|^Q} \right\}^{\frac{Q+2}{2Q}}.$$

Let

$$\Omega_T^1 = \{x \in \Omega_T \mid \lambda(x) < 1\}$$

and

$$\Omega_T^2 = \{x \in \Omega_T \mid \lambda(x) \geq 1\}.$$

Then

$$|B| \leq C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \left(\int_{\Omega_T^1} + \int_{\Omega_T^2} \right) \left[\eta_\lambda^{\frac{8Q}{(Q+2)(Q-2)}} \left(\sum_{k \in \mathbf{Z} \setminus \{0\}} \eta_{\lambda/T^k} \right)^{\frac{2Q}{Q+2}} + \left(\sum_{k \in \mathbf{Z} \setminus \{0\}} \eta_{\lambda/T^k} \right)^{\frac{2Q}{Q-2}} \right] \frac{dx}{|x|^Q} \right\} \lesssim$$

$$\begin{aligned}
&\lesssim C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \int_{\Omega_T} \left[\left(\frac{1}{T}\right)^{\frac{Q-2}{4} \cdot \frac{2Q}{Q+2}} + \left(\frac{1}{T}\right)^{\frac{Q-2}{4} \cdot \frac{2Q}{Q-2}} \right] \frac{dx}{|x|^Q} \right\}^{\frac{Q+2}{2Q}} \lesssim \\
&\lesssim C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \left(\frac{1}{T}\right)^{\frac{Q(Q-2)}{2(Q+2)}} \int_{\Omega_T} \frac{dx}{|x|^Q} \right\}^{\frac{Q+2}{2Q}} \lesssim \\
&\lesssim C(\log T)^{\frac{Q-2}{2Q}} \|u\|_{X_T} \left\{ \left(\frac{1}{T}\right)^{\frac{Q(Q-2)}{2(Q+2)}} \log T \right\}^{\frac{Q+2}{2Q}} \longrightarrow 0
\end{aligned}$$

uniformly in λ . □

Proof of Proposition 4.1. It follows from the above lemmas. □

5 Non degeneracy of the second differential

In order to verify the non degeneracy of the second differential, we restrict ourselves to the space \tilde{X}_T defined in Section 2 (which contains \mathcal{X}_T). We recall the following result [MU].

Proposition 5.1. *A function $u \in S^1(\mathbf{H}^n)$ is a solution of the following equation:*

$$-\Delta_{\mathbf{H}^n} u = (Q^* - 1)\omega^{Q^* - 2}u \quad (13)$$

if and only if there exist coefficients $\mu, \nu_1, \dots, \nu_{2n} \in \mathbf{R}$ such that

$$u = \mu \frac{\partial \omega_\lambda}{\partial \lambda} \Big|_{\lambda=1} + \sum_{i=0}^{2n} \nu_i T_i(\omega_\lambda).$$

For u to solve (13) is equivalent to being in the kernel of \mathcal{J}'' . Since the operator \mathcal{J}'' is the sum of an isomorphism and a compact operator on $S^1(\mathbf{H}^n)$ (see [MU]) and that it only a negative eigenvalue whose one-dimensional eigenspace is spanned by ω_λ (see [BCD]) there exists a constant C such that if $u \in S^1(\mathbf{H}^n)$ and

$$\int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \frac{\partial \omega_\lambda}{\partial \lambda} = 0, \quad \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} T_i(\omega_\lambda) = 0, \quad \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \omega_\lambda = 0 \quad (14)$$

then

$$d^2 \mathcal{J}(\omega_\lambda)[u, u] \geq C \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n} u|^2. \quad (15)$$

Furthermore, since \mathcal{J}'' is selfadjoint and ω_λ is an eigenfunction,

$$d^2 \mathcal{J}(\omega_\lambda)[\omega_\lambda, u] = 0. \quad (16)$$

We want to use this to prove a similar non degeneracy result for Ψ_λ on Ω_T for T large enough.

In order to do this, we introduce on X_T the norm

$$\|u\|_{T,\mathfrak{H}}^2 = \int_{\Omega_T} \left(|\nabla_{\mathbf{H}^n} u|^2 + \left| \frac{u}{|x|} \right|^2 \right).$$

Thanks to Hardy's inequality in \mathbf{H}^n (see Lemma 2.1 in [BCX], or otherwise apply Hölder inequality for Lorentz spaces), if $u \in S^1(\mathbf{H}^n)$, then, under the aforementioned hypotheses (14),

$$|d^2 \mathcal{J}(\omega_\lambda)[u, u]| \geq C \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n} u|^2 + \left| \frac{u}{|x|} \right|^2.$$

Using this we will prove that, if $u \in \tilde{X}_T$ satisfies

$$\int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} = 0 \quad (17)$$

and

$$\int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \Psi_\lambda = 0, \quad (18)$$

then, given $\varepsilon > 0$, for T large

$$d^2 \mathcal{J}_T(\Psi_\lambda)[u, u] \geq C \int_{\Omega_T} |\nabla_{\mathbf{H}^n} u|^2 + \left| \frac{u}{|x|} \right|^2,$$

$$|d^2 \mathcal{J}_T(\Psi_\lambda)[\Psi_\lambda, \Psi_\lambda]| \geq C \int_{\Omega_T} |\nabla_{\mathbf{H}^n} \Psi_\lambda|^2 + \left| \frac{\Psi_\lambda}{|x|} \right|^2$$

and

$$|d^2 \mathcal{J}_T(\Psi_\lambda)[\Psi_\lambda, u]| < \varepsilon \|\Psi_\lambda\|_{T,\mathfrak{H}} \|u\|_{T,\mathfrak{H}}.$$

This implies that $\mathcal{J}_T''(\Psi_\lambda)$ is invertible orthogonally to $\frac{\partial \Psi_\lambda}{\partial \lambda}$, and that the norm of the inverse is bounded uniformly in T .

Let us take a radial function $\rho = \rho(|x|)$ such that $\rho = 1$ on Ω_T , $\rho = 0$ on $B_{1/2} \cup (\mathbf{H}^n \setminus B_{2T})$, $0 \leq \rho \leq 1$, $|\nabla_{\mathbf{H}^n} \rho| \leq C$ on $B_1 \setminus B_{1/2}$, $|\nabla_{\mathbf{H}^n} \rho| \leq C/T$ on $B_{2T} \setminus B_T$.

By the computations in formula (11) follows that

$$\frac{\partial \omega_\lambda}{\partial \lambda} = -\frac{Q-2}{2} \frac{1}{\lambda} \omega_\lambda - \frac{1}{\lambda} Z(\omega_\lambda).$$

Thanks to formula (11), it can easily be proved that

$$\left| \nabla_{\mathbf{H}^n} \frac{\partial \Psi_{\lambda,T}}{\partial \lambda} \right| \leq \frac{C}{\lambda} \frac{1}{|x|^{\frac{Q}{2}}}. \quad (19)$$

By periodicity with respect to dilations we can suppose the quantity

$$r^Q \int_{B_2 \setminus B_{1/2}} \left(|\nabla_{\mathbf{H}^n} (u \circ \delta_r)|^2 + \left| \frac{u}{|x|} \circ \delta_r \right|^2 \right)$$

to be minimal for $r = 1$. Since there are $\sim \log T$ mutually disjoint annuli in Ω_T of the form $\delta_r\{1/2 \leq |x| \leq 2\}$, by easy computations one gets that

$$\int_{B_2 \setminus B_{1/2}} |\nabla_{\mathbf{H}^n} u|^2 + \left| \frac{u}{|x|} \right|^2 \leq \frac{C}{\log T} \|u\|_{T, \mathfrak{H}}^2,$$

and so, calling $W = (B_{2T} \setminus B_T) \cup (B_1 \setminus B_{1/2})$,

$$\int_W |\nabla_{\mathbf{H}^n} u|^2 + \left| \frac{u}{|x|} \right|^2 \leq \frac{C}{\log T} \|u\|_{T, \mathfrak{H}}^2. \quad (20)$$

Lemma 5.2. *If ρ is a cut-off function as above, for every ε there exists T_0 such that for $T \geq T_0$ if (17) holds then*

$$\left| \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \frac{\partial \Psi_{T, \lambda}}{\partial \lambda} \right| \leq \varepsilon \frac{1}{\lambda} \|u\|_{T, \mathfrak{H}}$$

and

$$\left| \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \Psi_{T, \lambda} \right| \leq \varepsilon \|u\|_{T, \mathfrak{H}}.$$

Proof.

$$\begin{aligned} \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} &= \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} - \int_{\Omega_T} \nabla_{\mathbf{H}^n} u \cdot \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} = \\ &= \int_W \left[(\rho \nabla_{\mathbf{H}^n} u + u \nabla_{\mathbf{H}^n} \rho) \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} \right]. \end{aligned}$$

Thanks to formulas (19) and (20) the first estimate follows by easy computations. The proof of the second one is identical. \square

Lemma 5.3. *For every ε there exists T_0 such that for $T \geq T_0$ if (17) holds then*

$$\int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \frac{\partial \omega_\lambda}{\partial \lambda} \leq \varepsilon \frac{1}{\lambda} \|u\|_{T, \mathfrak{H}}$$

and

$$\int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \nabla_{\mathbf{H}^n} \omega_\lambda \leq \varepsilon \|u\|_{T, \mathfrak{H}}.$$

Proof. Thanks to Lemma 5.2, we can estimate

$$\begin{aligned} \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \lambda \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} - \int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \lambda \nabla_{\mathbf{H}^n} \frac{\partial \omega_\lambda}{\partial \lambda} &\leq \\ &\leq C \|u\|_{T, \mathfrak{H}} \left(\int_{\Omega_T \cup W} \left| \lambda \nabla_{\mathbf{H}^n} \frac{\partial \Psi_\lambda}{\partial \lambda} - \lambda \nabla_{\mathbf{H}^n} \frac{\partial \omega_\lambda}{\partial \lambda} \right|^2 \right)^{1/2}. \end{aligned}$$

This quantity can be estimated almost identically as in the proof of Lemma 4.3. The proof of the second inequality estimate is identical. \square

Lemma 5.4. *For every $\varepsilon > 0$ there exist constants T_0 and C such that for $T \geq T_0$ if (17) and (18) hold then*

$$|d^2 \mathcal{J}(\omega_\lambda)[\rho u, \rho u]| \geq C \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n}(\rho u)|^2 + \left| \frac{\rho u}{|x|} \right|^2,$$

$$|d^2 \mathcal{J}(\omega_\lambda)[\rho \Psi_\lambda, \rho \Psi_\lambda]| \geq C \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n}(\rho \Psi_\lambda)|^2 + \left| \frac{\rho \Psi_\lambda}{|x|} \right|^2$$

and

$$|d^2 \mathcal{J}(\omega_\lambda)[\rho \Psi_\lambda, \rho u]| \leq \varepsilon \|\Psi_\lambda\|_{T, \mathfrak{H}} \|u\|_{T, \mathfrak{H}}.$$

Proof. Since $u \in \tilde{X}_T$, $u\rho$ is invariant with respect to the symmetry $(x, t) \mapsto (-x, t)$, one has

$$\int_{\mathbf{H}^n} \nabla_{\mathbf{H}^n}(\rho u) \cdot \nabla_{\mathbf{H}^n} T_i(\omega_\lambda) = 0.$$

The claim follows by Lemma 5.3, by equations (15) and (16), and elementary linear algebra. \square

Lemma 5.5. *For every $\varepsilon > 0$ there exist constants T_0 and C such that for $T \geq T_0$ if conditions (17) and (18) hold, then*

$$|d^2 \mathcal{J}_T(\Psi_\lambda)[u, u]| \geq C \int_{\Omega_T} |\nabla_{\mathbf{H}^n} u|^2,$$

$$|d^2 \mathcal{J}_T(\Psi_\lambda)[\Psi_\lambda, \Psi_\lambda]| \geq C \int_{\Omega_T} |\nabla_{\mathbf{H}^n} \Psi_\lambda|^2$$

and

$$|d^2 \mathcal{J}_T(\Psi_\lambda)[\Psi_\lambda, u]| < \varepsilon \|\Psi_\lambda\|_{X_T} \|u\|_{X_T}.$$

Proof. By direct computation we find

$$\begin{aligned} & |d^2 \mathcal{J}(\omega_\lambda)[\rho u, \rho u] - d^2 \mathcal{J}_T(\Psi_\lambda)[u, u]| = \\ & = \left| \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n}(\rho u)|^2 - (2^* - 1)|\omega_\lambda|^{2^*-2} \rho^2 u^2 + \right. \\ & \quad \left. - \int_{\Omega_T} |\nabla_{\mathbf{H}^n} u|^2 - (2^* - 1)|\Psi_\lambda|^{2^*-2} u^2 \right| \leq \\ & \leq (2^* - 1) \left| \int_{\Omega_T} \left(|\Psi_\lambda|^{2^*-2} - |\omega_\lambda|^{2^*-2} \right) u^2 \right| + \\ & + (2^* - 1) \left| \left(\int_{B_{2T} \setminus B_T} + \int_{B_1 \setminus B_{1/2}} \right) |\omega_\lambda|^{2^*-2} \rho^2 u^2 \right| + \\ & + 2 \left| \left(\int_{B_{2T} \setminus B_T} + \int_{B_1 \setminus B_{1/2}} \right) (u^2 |\nabla_{\mathbf{H}^n} \rho|^2 + \rho^2 |\nabla_{\mathbf{H}^n} u|^2) \right|. \end{aligned}$$

The first term can be estimated as in Lemma 4.3, the second in a trivial way, and the third has been essentially already estimated, to prove that for every ε there exists T big enough to ensure that the whole sum is bounded by $\varepsilon \|u\|_{X_T}^2$.

Analogously

$$\left| \int_{\mathbf{H}^n} |\nabla_{\mathbf{H}^n}(\rho u)|^2 - \int_{\Omega_T} |\nabla_{\mathbf{H}^n} u|^2 \right| \leq \varepsilon \|u\|_{X_T}^2.$$

This implies the first part of the thesis. The other statements are deduced in an analogous manner. \square

Proposition 5.6. *There exist constants T_0 and C such that for $T \geq T_0$ the operator $\mathcal{J}_T''(\Psi_\lambda)$ is invertible on the orthogonal space of $\frac{\partial \Psi_\lambda}{\partial \lambda}$ in X_T , and $\|\mathcal{J}_T''(\Psi_\lambda)^{-1}\|_{\mathcal{L}(X_T)} \leq C$.*

Proof. It follows from the preceding lemmas and elementary Hilbert space theory. \square

6 Proof of the main Theorem

We have proved that, for T big enough, on the orthogonal in \tilde{X}_T of the tangent of the curve \mathcal{Z}_T the second differential of \mathcal{J}_T is non degenerate, with norm bounded independently by λ and T . Let us call W this orthogonal in the point $\Psi_\lambda \in \mathcal{Z}$ and π the orthogonal projection on W . We remember that our aim is to solve $\nabla_{\mathbf{H}^n} \mathcal{J}_T(u) = 0$. Following the standard reasoning in [AM] we note that this is equivalent to solve

$$\pi \nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda + w) = 0$$

(auxiliary equation) and

$$(I - \pi) \nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda + w) = 0$$

(bifurcation equation) with $w \in W$.

Lemma 6.1. *There exists T_0 such that the auxiliary equation has a unique solution $w_T(\lambda)$; furthermore $\sup_\lambda \|w_T(\lambda)\| \rightarrow 0$ for $T \rightarrow \infty$.*

Proof. Write

$$\nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda + w) = \nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda) + \mathcal{J}_T''[w] + R(\Psi_\lambda, w)$$

with $R(\Psi_\lambda, w) = o(\|w\|)$ and $R(\Psi_\lambda, w) - R(\Psi_\lambda, v) = o(\|w - v\|)$, so that the auxiliary equation becomes

$$\pi \nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda) + \pi \mathcal{J}_T''(\Psi_\lambda)[w] + \pi R(\Psi_\lambda, w) = 0,$$

namely

$$w = -(\pi \mathcal{J}_T''(\Psi_\lambda))^{-1} [\pi \nabla_{\mathbf{H}^n} \mathcal{J}_T(\Psi_\lambda) + \pi R(\Psi_\lambda, w)] := N_\lambda(w).$$

By Propositions 4.1 and 5.6, N is a contraction if T is big enough, and so the auxiliary equation has an unique solution $w = w_T(\lambda)$. Furthermore for every $r > 0$ there exists T big enough such that $B_r(\Psi_\lambda) \cap W$ is mapped into itself by N . So $\sup_\lambda \|w_T(\lambda)\|$ tends to zero for $T \rightarrow \infty$. \square

Proof of Theorem 1.1. Let us consider the function

$$\Phi(\lambda) = \mathcal{J}_T(\Psi_\lambda + w(\lambda)).$$

It is continuous and periodic, so it has a stationary point λ_0 . Following the standard argument of Theorem 2.12 and Remark 2.14 in [AM], with the need for only formal modifications, the fact that $\Phi'(\lambda_0) = \mathcal{J}'_T(\Psi_{\lambda_0} + w(\lambda_0)) \cdot (\frac{\partial \Psi_{\lambda_0}}{\partial \lambda} + w'(\lambda_0))$ implies $u = \Psi_{\lambda_0} + w(\lambda_0)$ to solve the bifurcation equation, and so to be a stationary point of \mathcal{J}_T .

The smoothness of the solution can be proved with the same method of Appendix B in [Str].

Also $\lambda^{(2-Q)/2} u \circ \delta_{\lambda^{-1}}$ is a critical point of \mathcal{J}_T , and by the unicity in the fixed point theorem it must be equal to $\Psi_{\lambda_0 \lambda} + w(\lambda_0 \lambda)$, and so the whole curve $\widetilde{\mathcal{L}}_T = \{\Psi_\lambda + w(\lambda)\}$ consists of critical points of \mathcal{J} .

To prove the positivity, let us notice that from the proof of Proposition 5.6 follows that $\mathcal{J}(\omega_\lambda)$ has Morse index one on $\{\lambda \frac{\partial \omega_\lambda}{\partial \lambda}\}^\perp$. By continuity, the same holds for the orthogonal to the tangent space to $\widetilde{\mathcal{L}}_T$. Since $d\mathcal{J}_T$ is zero on $\widetilde{\mathcal{L}}_T$, the tangent of $\widetilde{\mathcal{L}}_T$ is in the kernel of \mathcal{J}_T'' . So the Morse index of \mathcal{J}_T on \widetilde{X}_T is one.

By a slight adaptation of the proof of Proposition 3.2 in [BCD] the set $\{u \neq 0\}$ has at most one connected component modulo δ_T , and so u does not change sign. By construction it is evident that it must be weakly positive (and even if it was not, it would be enough to change sign). The strict positivity follows from Bony's maximum principle (see [Bon]).

The last assertion follows by construction. \square

References

- [AM] Ambrosetti, Antonio; Malchiodi, Andrea, *Perturbation methods and semi-linear elliptic problems on \mathbf{R}^n* . Progress in Mathematics, 240. Birkhäuser Verlag, Basel, 2006. xii+183 pp.
- [Au] Aubin, Thierry *Problèmes isopérimétriques et espaces de Sobolev*. J. Differential Geometry 11 (1976), no. 4, 573–598.
- [BCX] H. Bahouri, J. Y. Chemin, and C. J. Xu. *Trace and trace lifting theorems in weighted sobolev spaces*. Journal of the Institute of Mathematics of Jussieu, 4(4):509–552, 2005
- [BCD] Birindelli, I., Capuzzo Dolcetta I., *Morse index and Liouville property for superlinear elliptic equations on the Heisenberg group*. , 2000, Ricerche di matematica, vol. 49, pp. 1-15

- [Bog] Boggess, Albert *CR manifolds and the tangential Cauchy-Riemann complex* Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1991. xviii+364 pp.
- [Bon] Jean-Michel Bony, *Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés*, Ann. Inst. Fourier (Grenoble) 19 (1969), no. fasc. 1, 277–304 xii (French, with English summary).
- [CGS] Caffarelli, Luis A.; Gidas, Basilis; Spruck, Joel, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*. Comm. Pure Appl. Math. 42 (1989), no. 3, 271–297.
- [CL] Chen, Chiun-Chuan; Lin, Chang-Shou, *Estimate of the conformal scalar curvature equation via the method of moving planes. II*. J. Differential Geom. 49 (1998), no. 1, 115–178.
- [CS] Chen, So-Chin; Shaw, Mei-Chi, *Partial differential equations in several complex variables*, AMS/IP Studies in Advanced Mathematics, 19. American Mathematical Society, Providence, RI; International Press, Boston, MA, 2001. xii+380 pp.
- [CMY] Cheng, Jih-Hsin; Malchiodi, Andrea; Yang, Paul, *A positive mass theorem in three dimensional Cauchy-Riemann geometry*. Adv. Math. 308 (2017), 276–347.
- [DTom] Dragomir, Sorin; Tomassini, Giuseppe *Differential geometry and analysis on CR manifolds*, Progress in Mathematics, 246. Birkhäuser Boston, Inc., Boston, MA, 2006. xvi+487 pp.
- [Fol] Folland, Gerald B., *Harmonic analysis in phase space.*, Annals of Mathematics Studies, 122. Princeton University Press, Princeton, NJ, 1989. x+277 pp.
- [Gam] Gamara, Najoua, *The CR Yamabe conjecture—the case $n=1$* . J. Eur. Math. Soc. (JEMS) 3 (2001), no. 2, 105–137.
- [GamYac] Gamara, Najoua; Yacoub, Ridha; *CR Yamabe conjecture—the conformally flat case*. Pacific J. Math. 201 (2001), no. 1, 121–175.
- [Gr] Grafakos, Loukas, *Classical Fourier analysis. Second edition.*, Graduate Texts in Mathematics, 249. Springer, New York, 2008. xvi+489 pp.
- [GMM] Guidi, Chiara; Maalaoui, Ali; Martino, Vittorio; *Singular CR structures of constant Webster curvature and applications*, preprint.
- [JL] Jerison, David; Lee, John M., *Extremals for the Sobolev inequality on the Heisenberg group and the CR Yamabe problem*. J. Amer. Math. Soc. 1 (1988), no. 1, 1–13.

- [JL2] Jerison, David; Lee, John M.; *The Yamabe problem on CR manifolds.*, J. Differential Geom. 25 (1987), no. 2, 167–197.
- [KMPS] Korevaar, Nick; Mazzeo, Rafe; Pacard, Frank; Schoen, Richard, *Refined asymptotics for constant scalar curvature metrics with isolated singularities.* Invent. Math. 135 (1999), no. 2, 233–272.
- [L1] Li, Yan Yan *Prescribing scalar curvature on S_n and related problems. I.* J. Differential Equations 120 (1995), no. 2, 319–410.
- [L2] Li, Yanyan *Prescribing scalar curvature on S_n and related problems. II. Existence and compactness.* Comm. Pure Appl. Math. 49 (1996), no. 6, 541–597.
- [MU] Malchiodi, Andrea; Uguzzoni, Francesco, *A perturbation result for the Webster scalar curvature problem on the CR sphere.* J. Math. Pures Appl. (9) 81 (2002), no. 10, 983–997.
- [MP1] Mazzeo, Rafe; Pacard, Frank, *Constant scalar curvature metrics with isolated singularities.* Duke Math. J. 99 (1999), no. 3, 353–418.
- [MP2] Mazzeo, Rafe; Pacard, Frank, *Constant mean curvature surfaces with Delaunay ends.* Comm. Anal. Geom. 9 (2001), no. 1, 169–237.
- [ON] O’Neil, Richard, *Convolution operators and $L(p,q)$ spaces,* Duke Math. J. 30 (1963), 129–142.
- [Pal] Palais, Richard S. *The principle of symmetric criticality.* Comm. Math. Phys. 69 (1979), no. 1, 19–30.
- [Str] M. Struwe, *Variational Methods,* Ergeb. der Math. u. Grenzgeb., Vol. 34, Berlin, Springer, 1996.
- [Ta] Talenti, Giorgio, *Best constant in Sobolev inequality.*, Ann. Mat. Pura Appl. (4) 110 (1976), 353–372.
- [Yap] Yap, Leonard Y. H. *Some remarks on convolution operators and $L(p,q)$ spaces,* Duke Math. J. 36 (1969), 647–658.