



Metagenomic Survey of Tomato Rhizosphere Microbiome Using the Shotgun Approach

 Olubukola Oluranti Babalola,^a Afeez Adesina Adedayo,^a Ayomide Emmanuel Fadiji^a

^aFood Security and Safety Niche, Faculty of Natural and Agricultural Sciences, North-West University, Mmabatho, South Africa

Olubukola Oluranti Babalola is the main investigator. Afeez Adesina Adedayo and Ayomide Emmanuel Fadiji are co-investigators. All authors contributed equally to this work.

ABSTRACT Food sustainability, e.g., fruit and vegetables, is a major agricultural problem that requires monitoring. Rhizosphere microbiomes' abundance and functionality are essential in promoting tomato plants' growth and health. We selected farms in South Africa's North West Province and present the metagenomes of their tomato rhizospheres and associated functional potentials.

The North West Province is a semiarid region with high temperatures. The soil in this region is populated by important microorganisms, with essential characteristics that promote the planting of tomatoes (1). Tomatoes produce annual crops of about 600,000 tonnes and find their way to South Africa through Europe and South American countries like Peru and Ecuador (2). The availability of these fruits in South Africa will enhance food production for human consumption because of their richness in essential vitamins and carotenoids (3). Three main cultivars of tomatoes are predominant in South Africa, namely, round or fresh tomatoes, Roma tomatoes, and cherry tomatoes, contributing about 24% of the total vegetable production in South Africa (4). Tomatoes are cultivated in open fields under irrigation in the following South African provinces: Eastern Cape, Western Cape, northern Kwazulu-Natal, North West, Mpumalanga, and Limpopo.

We collected the soil samples used in this study from the root region of Roma tomato plants at the experimental farm of the North-West University, Mafikeng (26°019'36.9"S, 26°053'19.0"E; 25°47'19.1"S, 25°37'05.1"E; 25°47'17.0"S, 25°37'03.2"E; altitude, 159 km). The site has a 450-mm annual rainfall record and regional temperatures ranging from 25°C to 37°C (5). The experimental soil samples were collected from the rhizosphere of healthy and diseased tomato plants and the bulk soil. Bulk soil, which served as the control soil, was collected from a natural grassland with no tomato plantation, 20 m from the tomato plantation field. We took rhizosphere soil from the Roma tomato plants from three different areas of a site. Fifteen samples were put in separate sterile polyethylene bags, kept in a cold box at –4°C, and transported to the laboratory. All collected soil samples were then stored at a temperature of –20°C before extraction of DNA for shotgun metagenomic sequencing.

From the stored rhizosphere soil, 5 g from each sample was measured using a calibrated scale. DNA was extracted using the NucleoSpin Soil kit (Macherey-Nagel, Germany). The quality of the extracted DNA was assessed using a NanoDrop spectrophotometer.

The libraries were prepared with 50 ng DNA using a Nextra DNA Flex kit, undergoing fragmentation and the ligation of adapter sequences. The final concentrations of the libraries were measured using the Qubit double-stranded DNA (dsDNA) HS assay kit (Life Technologies), and the mean lengths of the DNA fragments were ascertained using a 2100 Bioanalyzer (Agilent Technologies). The libraries were then monitored, combined at 0.6 nM, and sequenced using a NovaSeq 6000 system (Illumina) with 300 cycles.

SolexaQA v1.6 was used to conduct the quality control (QC) of raw data, reduce low-quality reads, and remove replicate data (6). Duplicate read inferred sequencing error estimation (DRISSE) enables us to assess the error of sequenced samples

Editor John J. Dennehy, Queens College CUNY

Copyright © 2022 Babalola et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Olubukola Oluranti Babalola, olubukola.babalola@nwu.ac.za.

The authors declare no conflict of interest.

Received 22 November 2021

Accepted 23 December 2021

Published 3 February 2022

TABLE 1 Sequence reads for the rhizosphere soil samples analyzed

| Sample ^a | Data before QC | | | | Data after processing | | | | Data after QC | | | | Data after alignment | | | |
|---------------------|-----------------|-----------------------|---------------------|---------------------------|--|---------------------------------|-------------------------------|---------------|-----------------------|---------------------|---------------------------|----------------------------------|--------------------------------|--|--|--|
| | Size (bp) | No. of sequence reads | Mean GC content (%) | Mean sequence length (bp) | No. of artificial duplicate read sequences | No. of known proteins predicted | No. of RNA features predicted | Size (bp) | No. of sequence reads | Mean GC content (%) | Mean sequence length (bp) | No. of known proteins identified | No. of RNA features identified | | | |
| HT21 (HR) | 2,152,004,650.3 | 13,739,258.3 | 64 ± 10 | 154 ± 32 | 836,435 | 7,850,484.3 | 28,853 | 765,041,235.3 | 12,665,143.7 | 63 ± 9 | 155 ± 33 | 3,985,525.7 | 5,962.7 | | | |
| HT21 (DR) | 1,409,528,303.3 | 19,765,082 | 64 ± 10 | 155 ± 32 | 757,641.7 | 4,208,873.7 | 30,027 | 1,352,124,415 | 11,966,279.7 | 64 ± 9 | 156 ± 39 | 4,178,206.3 | 7,656.7 | | | |
| HT21 (BR) | 1,477,197,003 | 138,145,859.3 | 65 ± 9 | 155 ± 32 | 374,748 | 11,448,097.3 | 25,322.7 | 1,385,218,693 | 5,247,459 | 64 ± 9 | 156 ± 34 | 2,515,439.7 | 5,439.3 | | | |

^a HR, healthy rhizosphere; DR, diseased rhizosphere; BR, bulk rhizosphere.




FIG 1 Abundant phyla obtained according to the taxonomic system.

caused by artificial replicated sequenced data (7). We employed the default settings of the MG-RAST v4.0.3 server to perform analytical processing downstream (8, 9) (Table 1).

The domain or kingdoms obtained according to the taxonomic system are *Bacteria*, *Eukaryota*, and *Archaea*. The most abundant phyla belonged to the *Bacteria* domain; *Proteobacteria* (38.8 to 54%) and *Actinobacteria* (25.4 to 35.5%) were the most abundant, and others, such as *Acidobacteria* (2.3 to 5.2%), *Bacteroidetes* (3.0 to 3.9%), *Planctomycetes* (2.6 to 3.4%), *Verrucomicrobia* (2.2 to 2.3%), and *Firmicutes* (1.7 to 2.4%), were also significant. Moreover, reads for fungi (*Ascomycota* and *Basidiomycota*) and archaea (*Thaumarchaeota* and *Euryarchaeota*) were also identified but at <1% relative abundance (Fig. 1).

Functional annotation after mapping with SEED subsystems (10) revealed the presence of the following important attributes: carbohydrates (13.2 to 14.8%), clustering-based systems (12.7 to 12.8%), amino acids and derivatives (10.1 to 10.3%), protein metabolism (8.2 to 8.3%), DNA metabolism (4.3 to 4.6%), cell wall and capsule (3.4 to 3.6%), RNA metabolism (3.3 to 3.5%), and stress response (2.5 to 2.7%).

Data availability. The metagenomes of rhizosphere soil sequence reads were submitted to the NCBI with BioProject accession number [PRJNA766489](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA766489) and Sequence Read Archive (SRA) accession numbers [SRX12366062](https://www.ncbi.nlm.nih.gov/sra/SRX12366062), [SRX12366063](https://www.ncbi.nlm.nih.gov/sra/SRX12366063), and [SRX12366064](https://www.ncbi.nlm.nih.gov/sra/SRX12366064) (healthy), [SRX12366065](https://www.ncbi.nlm.nih.gov/sra/SRX12366065), [SRX12366066](https://www.ncbi.nlm.nih.gov/sra/SRX12366066), and [SRX12366067](https://www.ncbi.nlm.nih.gov/sra/SRX12366067) (diseased), and [SRX12366068](https://www.ncbi.nlm.nih.gov/sra/SRX12366068), [SRX12366069](https://www.ncbi.nlm.nih.gov/sra/SRX12366069), and [SRX12366070](https://www.ncbi.nlm.nih.gov/sra/SRX12366070) (bulk).

ACKNOWLEDGMENTS

O.O.B. acknowledges the National Research Foundation (South Africa) grants (UID123634 and UID132595) that support research in our laboratory.

REFERENCES

- Babalola OO, Akinola SA, Ayangbenro AS. 2020. Shotgun metagenomic survey of maize soil rhizobiome. *Microbiol Resour Announc* 9:e00860-20. <https://doi.org/10.1128/MRA.00860-20>.
- Manzanero-Medina GI, Vásquez-Dávila MA, Lustre-Sánchez H, Pérez-Herrera A. 2020. Ethnobotany of food plants (quelites) sold in two traditional markets of Oaxaca, Mexico. *S Afr J Bot* 130:215–223. <https://doi.org/10.1016/j.sajb.2020.01.002>.
- Llauradó Maury G, Méndez Rodríguez D, Hendrix S, Escalona Arranz JC, Fung Boix Y, Pacheco AO, García Díaz J, Morris-Quevedo HJ, Ferrer Dubois A, Aleman EI, Beenaerts N, Méndez-Santos IE, Orberá Ratón T, Cos P, Cuypers A. 2020. Antioxidants in plants: a valorization potential emphasizing the need for the conservation of plant biodiversity in Cuba. *Antioxidants* 9:1048. <https://doi.org/10.3390/antiox9111048>.
- Bontempo L, van Leeuwen KA, Paolini M, Holst Laursen K, Micheloni C, Prenzler PD, Ryan D, Camin F. 2020. Bulk and compound-specific stable isotope ratio analysis for authenticity testing of organically grown tomatoes. *Food Chem* 318:126426. <https://doi.org/10.1016/j.foodchem.2020.126426>.
- Babalola OO, Enebe MC. 2020. Metagenomes of maize rhizosphere samples after different fertilization treatments at Molelwane Farm, located in North-West Province, South Africa. *Microbiol Resour Announc* 9:e00937-20. <https://doi.org/10.1128/MRA.00937-20>.
- Cox MP, Peterson DA, Biggs PJ. 2010. SolexaQA: at-a-glance quality assessment of Illumina second-generation sequencing data. *BMC Bioinformatics* 11:485. <https://doi.org/10.1186/1471-2105-11-485>.
- Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, McGill SK, Dougherty MK. 2019. High-throughput amplicon sequencing of the full-

- length 16S rRNA gene with single-nucleotide resolution. *Nucleic Acids Res* 47:e103. <https://doi.org/10.1093/nar/gkz569>.
8. Meyer F, Paarmann D, D'Souza M, Olson R, Glass E, Kubal M, Paczian T, Rodriguez A, Stevens R, Wilke A, Wilkening J, Edwards R. 2008. The metagenomics RAST server: a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9:386. <https://doi.org/10.1186/1471-2105-9-386>.
 9. Enagbonma BJ, Aremu BR, Babalola OO. 2019. Profiling the functional diversity of termite mound soil bacteria as revealed by shotgun sequencing. *Genes* 10:637. <https://doi.org/10.3390/genes10090637>.
 10. Mitra S, Rupek P, Richter DC, Urich T, Gilbert JA, Meyer F, Wilke A, Huson DH. 2011. Functional analysis of metagenomes and metatranscriptomes using SEED and KEGG. *BMC Bioinformatics* 12(Suppl 1):S21. <https://doi.org/10.1186/1471-2105-12-S1-S21>.