

The Jackson Laboratory

The Mouseion at the JAXlibrary

Faculty Research 2023

Faculty Research

12-1-2022

Overview of triple negative breast cancer prognostic signatures in the context of data science-driven clinico-genomics research.

Enrico Capobianco

Follow this and additional works at: <https://mouseion.jax.org/stfb2023>



Overview of triple negative breast cancer prognostic signatures in the context of data science-driven clinico-genomics research

Enrico Capobianco

The Jackson Laboratory, CT, USA

Correspondence to: Enrico Capobianco. The Jackson Laboratory, 10 Discovery Drive Farmington, CT 06032, USA. Email: enrico.capobianco@jax.org.

Comment on: Chen C, Lin CJ, Li SY, *et al.* Identification of a novel signature with prognostic value in triple negative breast cancer through clinico-transcriptomic analysis. *Ann Transl Med* 2022;10:1095.

Keywords: Triple-negative breast cancer (TNBC); genomics; prognostic signatures; data science

Submitted Nov 03, 2022. Accepted for publication Dec 15, 2022.

doi: 10.21037/atm-22-5477

View this article at: <https://dx.doi.org/10.21037/atm-22-5477>

Background

Triple-negative breast cancer (TNBC) is a highly aggressive tumor with inherent heterogeneity leading to a variety of outcomes, including early relapse, metastatic spread, and poor survival. Chen *et al.*'s (1) proposed a prognostic signature to improve risk stratification in TNBC patients obtained by combining expressions from The Cancer Genome Atlas (TCGA), with 1,090 primary tumors and 113 normal, and RNA-seq data FUSCCTNBC (Sequence Read Archive, SRP15). As the samples were classified according to subtypes (HR+/HER2-, HR+/HER2+, HR-/HER2+, TNBC, and normal samples with status of ER, PR, and HER2), the differentially expressed gene (DEG) profiles were computed from pairwise comparisons between TNBC and the other subtypes.

In general, signatures may inherently determine stratifications based on groups of patients sharing similar tumor characteristics and biological properties. Ideally, a signature should be linked to some novel mechanism, a pathway or a biological process or a drug mechanism. Algorithmic or semi-algorithmic identifications of (non-redundant) groups call for validation at both biological and clinical levels. For instance, prognostic validity may be established because of its association with overall survival. One usually relies on the 'big data evidence' that omics are expected to support. However, the reliability assigned to these and other types of signatures is a controversial matter. First, many cancer signatures are not fully justified. Signatures may fail to show biological value with reference

to the specific cancer they address. This is typically proven algorithmically, by enabling gene signature replacement and showing that surrogate genes have measurable prognostic prediction power not affecting the overall signature, but with opposite biological interpretation compared to the replaced genes with respect to mechanisms. Therefore, the problem becomes lack of provable causality regarding outcome prediction, depending on the usable surrogate gene space.

Second, reproducibility implies the ability to assess the robustness of a signature relatively to specific methods or platforms. A demonstration for association with outcomes should be generally valid across independent groups and replicate experiments, and with no role played by factors excluded from the model generating the signature, including noise. Gene expression signatures (GES) connecting phenotypes to mRNA suffer from quite limited reproducibility. Therefore, one possible strategy is to combine significantly enhanced GES across datasets and experiments covering various phenotypes. Another strategy is to correlate gene expressions with other measurements to increase the signature's prediction power, for instance through co-expression networks.

The strategy followed by Chen *et al.* [2022] to obtain the prognostic GES involves multiple steps (*Figure 1A*). Chen *et al.*' results are now contextualized (*Figures 1B,2*) with reference to a selected non-comprehensive list of recent signature results retrieved from PubMed (query: "TNBC prognostic signatures", 164 results).




Figure 1 Overview of Chen *et al.*'s work and contextualization of their signature. (A) Basic sketch of the approach in Chen *et al.* The results have highlighted a multi-year (1-to-5) prognostic 6-gene signature integrated to a clinical scoring system utilizing pathological stage information through a stably performing nomogram and resulting in an improved concordance index. (B) Biomedical contexts selected from recent literature and defining TNBC prognostic signatures. Note that the highlighted composite signatures have data science as a driver. TNBC, triple-negative breast cancer.

Contextualization

NACT and EMT

TNBC patients respond better to neoadjuvant therapy compared to other BC subtypes, though some patients still have a poor prognosis (2). A study represented the transcriptional portrait associated with neoadjuvant chemotherapy (NACT) resistance and found candidate genes as prognostic markers (3). In Blaye *et al.*'s study (4), an immunologic transcriptomic signature was found using 115 post-NACT samples to predict relapse in TNBC patients by profiling 770 TME-related genes using the NanoString PanCancer IO360 panel. The 8 genes linked to immunity (*BLK*, *GZMM*, *CXCR6*, *LILRA1*, *SPIB*, *CCL4*, *CXCR4*, *SLAMF7*) were predictive of intra-cohort relapse (validated externally in KMplot and METABRIC), indicating that lack of immune activation after NACT is associated with a high risk of distant relapse. Another study (5) revealed a 6-epithelial-mesenchymal transition (EMT) gene signature (*LUM*, *SFRP4*, *COL6A3*, *MMP2*, *CXCL12*, and *HTRA1*) at diagnosis potentially predictive for metastasis in TNBC patients who did not achieve pathological complete response to NACT and in patients treated with surgery in combination with adjuvant therapy. The EMT pathway was significantly enriched in pre- and post-NACT patients with metastasis after NACT. The 6-EMT gene signature was validated in a GEO dataset (HR = 0.36,

P = 0.0008, 95% CI: 0.200–0.658) and in another GEO dataset of TNBC patients who received surgery followed by adjuvant chemotherapy (HR = 0.46, 95% CI: 0.225–0.937). Other results (6) informing on cell plasticity relatively to EMT were centered on a signature of circulating tumor cells (CTCs) from 32 patients aimed to identify markers associated with TNBC outcome and aggressiveness, especially tissue inhibitor of metalloproteinases 1 (*TIMP1*) and androgen receptor (AR), whose blockade was further analyzed *in vitro* and *in vivo* in relation to proliferation and dissemination.

Immunotherapy & hypoxia-related

Immunotherapy (ImT) research and TNBC prognosis present signature gaps. A hypoxia-immune-based cross-cohort classifier predictive of prognosis was developed and validated in (7) to potentially guide TNBC ImT. In study of Yang *et al.* (8), six gene cross-cohort prognostic hypoxia-immune related signatures were identified to stratify patients into high-/low-risk groups, with the former group showing poorer prognosis, and correlation with hypoxia status in tumor microenvironment (TME). Interestingly, there were relatively less activated immune cells and lower expression levels in immune checkpoint inhibitors (ICI). In study of Sun *et al.* (9), a robust predictive 3-gene signature (*PEKL*, *ALDOA*, *PGK1*) associated with the glycolytic process was

Immunity Prognostic Signatures

NACT & relapse		Immune Infiltration		Immune Checkpoint Inhibitors		TME	Hypoxia
		Network-driven		Low risk	OS favorable	Poor prognosis	
BLK		RMND5A	TOP2A	PD-1	PDCD1	PPF1A4	PEKL
GZMM		ZNF829	CCNA2	PD-L1	PDCD1LG2	TNFRSF1B	ALDOA
CXCR6		KDM5B	PCNA	PD-L2	KIR3DL2	ARHGAP9	PGK1
LILRA1		NCBP2	MSH2	CTLA4		ZNF831	
SPIB		GPI	CDK6			CTLA4	
CCL4		BGN				BLK	
CXCR4		CCND2				ANKRD22	
SLAMF7		PLBD1				CLEC4E	
		ZYG11A					
		IL17					
Macrophage and Resistance		EMT		Paracancer		Bioinfo-driven	
Network-driven			With CTC			Poor OS	
CD79A		LUM	AR	ABCC9 (cg06951626)		CENPW	
CXCL13		SFRP4	TIMP1	NKAPL (cg18675097, cg01031101, cg17384889)		HORMAD1	
IGLL5		COL6A3		TMEM132C (cg03530754)		APOD	
LHFPL2		MMP2				PIP	
PLEKHF1		CXCL12				ZNF703	
		HTRA1					
DRFS		DFS		PFI			
		Poor, GEO, network-driven					
CD1B		IL18R1	FAM83B	FAM83B			
CD53		CD53	KITLG	EXO1			
CT45A1		TRIM	CFD	S100B			
GTF3C1		Jaw1	RBM24	TYMS			
IL11RA		LTB		CFD			
IL1RN		PTPRCAP					
LRRN3							
MAPK1							
NEFL							
PRKCE							
PTPRC							
SPACA3							
TNFSF11							

DFS= disease-free survival
 PFI= progression-free interval
 DRFS=Distant recurrence-free survival

Chen et al.
HCAR1
ORM2
TMCC2
ERICH3
KRT83
FABP7

Methylation Prognostic Signatures

m6A modification-hypoxia	m6A	m6A-related lncRNA
	Favorable Unfavorable	
PIM2	METTL14	ALKBH5
PET117		SAMD12-AS1
SMARCA5		BVES-AS1
TAF9		LINC00593
ABCB10		MIR205HG
MKP1		LINC00571
		ANKRD10-IT1
		CIRBP-AS1
		SUCLG2-AS1
		BLACAT1
		HOXB-AS1

ncRNA Prognostic Signatures

miRNA	lncRNA immune-related	
Relapse predictive	OS	Distant metastases-free
miR-139-5p	hsa-miR-342-3p	hsa-miR-29c
miR-10b-5p	hsa-miR-195	hsa-miR-342-3p
miR-486-5p	hsa-miR-155	hsa-let-7c
miR-455-3p	hsa-miR-497	hsa-let-7b
miR-107		hsa-miR-497
miR-146b-5p		
miR-324-5p		
miR-20a-5p		
		RP11-890B15.3
		RP11-1024P17.1
		MF12-AS1
		RP11-180N14.1

Figure 2 Prognostic signatures associated to multiple domains and macro-categorized.

identified based on 48 hypoxia-related prognostic genes and a risk score model was set as a prognostic biomarker for TNBC. Future ImT developments will involve ICI toward TNBC treatment, as in Islam *et al.*'s study (10) (PI3K, MEK, PARP, EGFR, VEGF, and AR). Immune prognostic signatures were investigated (11) based on 66 prognostic genes, showing predictive power of the response to ICI (PD-1, PD-L1, PD-L2 and CTLA4 gene expressions higher in the low-risk group). ICI expression (and clinical) profiles of TNBC samples from TCGA and METABRIC were used to build 1-, 2-, 3-year OS-related GES (PDCD1, PDCD1LG2 and KIR3DL2, with receiver operating characteristic (ROC) curves showing AUC values of 0.925, 0.822 and 0.835, respectively) from potentially favorable OS predictors (*IDO1, CD274, PDCD1LG2, PDCD1, CTLA4, ICOS, KIR3DL2, HLA-B, HLA-F, LGA3*) (12).

Studies of underlying mechanisms will include the TME and the immune infiltration (ImI). In study of Qin *et al.* (13), the TNBC TME landscape was elucidated (GEO, N=107, training: METABRIC, N=299, validation cohorts). Two distinct TME clusters were identified, one characterized by low ImI with poor prognosis, the other with high ImI and better survival probability. These clusters showed differential expression (37 upregulated and 778 downregulated genes) delivering then a final 8-gene TME signature (*PPF1A4, TNFRSF1B, ARHGAP9, ZNF831, CTLA4, BLK, ANKRD22* and *CLEC4E*) in which *TNFRSF1B* and *BLK* were risk factors for poor prognosis in TNBC patients. In (14), from the study of 1,145 patients a tumor immune risk score (TIRS) was obtained with 8 biomarkers stratifying patients in high- and low-risk groups. Significant survival and ImI pattern differences were found at both transcriptome and protein levels indicating 4 different tumor immune microenvironment types (TIMT): one associated with the best prognosis and immune status, another with the opposite effect, the other two associated with highly unstable genomes and stem-cell-related characteristics along with high stromal scores.

Bioinformatics composite signatures

TNBC prognostic signatures may involve hub genes and pathways (15), without establishing genes as drivers of network causality and directionality (16). For example, in (17) a 5-gene prognostic signature (*CD79A, CXCL13, IGLL5, LHFPL2, and PLEKHF1*) was found based on co-expressed hub genes associated with macrophages and related to gemcitabine resistance in TNBC patients. Then,

a survival-related 5-gene hub signature (*TOP2A, CCNA2, PCNA, MSH2, CDK6*) was combined with ImI analysis to show relevance for immune monitoring across GEO cancer data. Tumor-infiltrating lymphocytes (TILs) are a robust prognostic factor for improved patient survival, with T cells being predominant, but without resolving the uncertainty about the relationship with patient prognosis (18).

The real predictive power of cancer-related immune prognostic markers was investigated in (19) through a bioinformatic approach identifying a 6-gene signature (*IL18R1, CD53, TRIM, JAW1, LTB, and PTPRCAP*) positively correlated with disease-free survival (DFS) in TNBC patients and showing significant immune expression profiles in cancer cells and in intra- and peri-TILs. In Kim *et al.*'s study (20), it was identified the associations between gene expression and distant recurrence-free survival (DRFS) from a 13-gene expression profile (*CD1B, CD53, CT45A1, GTF3C1, IL11RA, IL1RN, LRRN3, MAPK1, NEFL, PRKCE, PTPRC, SPACA3* and *TNFSF11*) associated with patient prognosis ($P < 0.05$) (AUC =0.923). Combined with stage, it was predictive of distant recurrence of early TNBC, with 3 genetic risk groups classifying DRFS rate significantly even intra-stage ($P < 0.001$) and informing on treatment with new possible ImT targets. Finally, cell differentiation together with grade- and tumor mutational burden-related DEGs were identified using integrated single-cell and bulk RNA-seq data analysis, delivering a composite 10-gene prognostic signature (*RMND5A, ZNF829, KDM5B, NCBP2, GPI, BGN, CCND2, PLBD1, ZYG11A, IL17RD*), closely related to tumor ImI (21).

Epigenetics

Despite its relevance, aberrant DNA methylation is still underutilized in prognostic models for TNBC. For instance, the N⁶-methyladenosine (m⁶A) modification is relevant in cancer development, but not enough evidenced in TNBC. Based on the TCGA transcriptome profiles of the 13 m⁶A methylation regulators measured from 98 TNBC tumor and normal tissue samples, two expression levels were emphasized, specifically ALKBH5 (unfavorable prognostic factor, HR =3.327, $P = 0.006$) and METTL14 (favorable, HR =0.425, $P = 0.009$), both contributing to improve the accuracy of a prognostic model of risk prediction when combined with TNM stage (AUC =0.791) (22). Aberrant DNA methylation was also investigated in study of Zhang *et al.* (23) from TCGA paracancer samples (PCS). Differential methylation-DEG correlation revealed

1,525 DEGs and 150 differentially methylated genes between TNBC and PCS, with a signature including prognostic markers such as ABCC9 (cg06951626), NKAPL (cg18675097, cg01031101, and cg17384889), and TMEM132C (cg03530754).

In Gao *et al.*'s study (24), methylation and expression data from TCGA were combined, and 743 differentially methylated sites (DMS) were identified corresponding to 332 genes, with 357 hypermethylated sites and 386 hypomethylated sites, of which 103 were prognosis related. A LASSO-driven 5-DMSs prognostic signature was found to classify TNBC patients with significant survival difference (log-rank $P=4.97E-03$), then validated in GSE78754 (HR =2.42, 95% CI: 1.27–4.59, log-rank $P=0.0055$) and verified for DFS in GSE141441 (HR =2.09, 95% CI: 1.28–3.44, log-rank $P=0.0027$). Two DMS were related to high risk (cg21234506 and cg21580376; HR >1), and three DMS were protective (cg15724876, cg17887364, and cg19419246; HR <1). Interestingly, a recent study (25) on TME of TNBC associated m⁶A modification and hypoxia status, identifying 26 genes related to both regulation types and characterizing two clusters, one being with significantly worse prognosis. A 6-gene prognostic signature was identified (*PIM2*, *PET117*, *SMARCA5*, *TAF9*, *ABCB10*, *MKP1*) among the m⁶A modification-hypoxia genes to evaluate risk and predict ImT response of patients.

Network- and machine learning (ML)-driven signatures

Key DEGs (*CENPW*, *HORMAD1*, *APOD*, *PIP*, and *ZNF703*) were found as candidate prognostic markers associated with poor OS through integrated bioinformatics performed on GEO microarray TNBC data (26). Network analysis focused on 147 co-DEGs in TNBC vs non-cancerous tissue samples, identifying a 15-gene signature inclusive of *BUB1* and *CENPF* significantly correlated with OS, while *BUB1*, *CCNA2*, and *PACC1* showed significant poor DFS (27). In study of Liu *et al.* (28), 105 heterogeneous DEGs were identified between TNBC and other subtypes. Two prognostic signatures were significant: a 4-gene (*FAM83B*, *KITLG*, *CFD* and *RBM24*) in disease-free interval, and a 5-gene (*FAM83B*, *EXO1*, *S100B*, *TYMS* and *CFD*) in progression-free interval. The multivariate Cox regression models showed predictive performance by time-dependent ROC analysis and survival analysis of TNBC subtypes (mesenchymal stem-like and mesenchymal) emphasized the *FAM83B* prognostic role.

In study of Liu *et al.* (29), bioinformatics and network

analyses explored the role of alternative splicing events (ASE) events and their correlation with TNBC prognostic DEGs, delivering ASE profiles, prognostic interaction networks, and splice factor-AS interaction networks. Finally, the use of ML was central to (30) where first a Recursive Feature Elimination (RFE) algorithm identified signatures (20, 25, 30, 35, 40, 45, and 50 genes) differentiating TNBC from other subtypes, then XGBoost was found the best performer for 45 genes (mixed signatures), of which 34 genes differentially regulated, 4 specifically relevant for distant metastasis free survival, and 2 potentially prognostic (*POU2AF1* and *S100B*) associated with MAPK, PI3-AkT, Wnt, TGF- β , and other signal transduction pathways involved in metastasis. In study of Gong *et al.* (31), 151 TNBC patients obtained from the TCGA SpliceSeq database showed relevance for the Exon Skip (ES) type of AS events, more robust in predicting performance in TNBC prognosis. The ES AS signature confers a strong oncogenic phenotypic enrichment in a high-risk group (cell cycle and SUMOylating pathways of tumorigenesis), and a low-risk group (programmed cell death and metabolism pathways).

Non-coding RNAs

Many ncRNAs are likely dysregulated in TNBC, some being promising biomarkers and prognostic too. For instance, an 8-miRNAs signature (miR-139-5p, miR-10b-5p, miR-486-5p, miR-455-3p, miR-107, miR-146b-5p, miR-324-5p and miR-20a-5p) supported an accurate predictive model of relapse in TNBC patients highly correlated with prognosis (AUC of 0.80), independently validated twice (AUCs of 0.89 and 0.90) (32). In study of Zaka *et al.* (33), Cox analysis was employed to identify 25 miRNAs associated with prognosis with both risky and protective outcomes. Some miRNAs were associated with OS (hsa-miR-342-3p, hsa-miR-195, hsa-miR-155, and hsa-miR-497) and others with distant metastases-free survival (hsa-miR-29c, hsa-miR-342-3p, hsa-let-7c, hsa-let-7b, and hsa-miR-497). Based on both TCGA and GEO, and supported by ImmPort relatively to immune-related mRNAs, of 62 identified immune-related lncRNAs a signature of 4 (RP11-890B15.3, RP11-1024P17.1, MFI2-AS1 and RP11-180N14.1) showed regulation power mediated by miRNAs and prognostic value stratifying TNBC patients into high- and low-risk groups. The high-risk group had unfavorable outcomes together with significant immune response to tumor cell and high infiltrating abundance of regulatory T cell (34). In study

of Wu *et al.* (35), a 10-m6A-related lncRNA signature predictive of the prognostic risk in TNBC (SAM12-AS1, BVES-AS1, LINC00593, MIR205HG, LINC00571, ANKRD10-IT1, CIRBP-AS1, SUCLG2-AS1, BLACAT1, and HOXB-AS1) was found, establishing a prognostic score risk model (AUC of 0.997 and 0.864 in TCGA and GSE76250 datasets, respectively) and showing patients with lower score associated to better OS.

Radiomics & integrated signatures

A TNBC radiogenomic study (N=202) generated quantitative radiomic features (n=860) from contrast-enhanced magnetic resonance images (CE-MRI). Notably, a radiomic feature capturing peritumoral heterogeneity (PH) was an established prognostic factor for RFS (P=0.01) and OS (P=0.004). Combined with matching transcriptomic and metabolomic data, the PH was associated with immune suppression and upregulated fatty acid synthesis (36). A role for radiomics with reference images to achieve consistent assessments is linked to the assessment of stromal TILs to be incorporated into clinical practice scores based on the reproducibility of multiple factors central to the heterogeneity in lymphocyte distribution (slide-related issues; outside tumor boundary; minimal assessable stroma; inflammatory cells etc.) (37). In study of Wu *et al.* (38), 17 dynamic CE-MRI features and TILs were combined to characterize tumor and parenchyma in a TGCA cohort (n=126). A LASSO imaging signature for TILs was found and a prediction model built with imaging signature and molecular features, resulting in 4 imaging features significantly associated with TILs: tumor volume, cluster shade of signal enhancement ratio (SER), mean SER of tumor-surrounding background parenchymal enhancement (BPE), and proportion of BPE. The imaging signature during validation allowed the predicted TILs to separate TNBC patients into two groups with distinct RFS (log-rank P=0.042), with TNBC showing no/minimal TILs associated with a worse prognosis.

To conclude, TNBC research offers room for further improvement. Especially longitudinal studies based on patients with long or more prolonged survival life periods (5–10 years) and involving a multitude of data/marker types including imaging, will be valuable, bringing future challenges for integrative data science approaches.

Acknowledgments

Funding: The author acknowledges support from JAX

Computational Sciences, JAX Cancer Center (JAXCC) and NCI CCSG (P30CA034196) and support from grant NSF 19-500, DMS 1918925/1922843.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Annals of Translational Medicine*. The article did not undergo external peer review.

Conflicts of Interest: The author has completed the ICMJE uniform disclosure form (available at <https://atm.amegrouops.com/article/view/10.21037/atm-22-5477/coif>). The author has no conflicts of interest to declare.

Ethical Statement: The author is accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Chen C, Lin CJ, Li SY, et al. Identification of a novel signature with prognostic value in triple-negative breast cancer through clinico-transcriptomic analysis. *Ann Transl Med* 2022;10:1095.
2. Sporikova Z, Koudelakova V, Trojanec R, et al. Genetic Markers in Triple-Negative Breast Cancer. *Clin Breast Cancer* 2018;18:e841-50.
3. Vishnubalaji R, Alajez NM. Transcriptional landscape associated with TNBC resistance to neoadjuvant chemotherapy revealed by single-cell RNA-seq. *Mol Ther Oncolytics* 2021;23:151-62.
4. Blaye C, Darbo É, Debled M, et al. An immunological signature to predict outcome in patients with triple-negative breast cancer with residual disease after neoadjuvant chemotherapy. *ESMO Open* 2022;7:100502.
5. Wei LY, Zhang XJ, Wang L, et al. A Six-Epithelial-

- Mesenchymal Transition Gene Signature May Predict Metastasis of Triple-Negative Breast Cancer. *Onco Targets Ther* 2020;13:6497-509.
6. Abreu M, Cabezas-Sainz P, Pereira-Veiga T, et al. Looking for a Better Characterization of Triple-Negative Breast Cancer by Means of Circulating Tumor Cells. *J Clin Med* 2020;9:353.
 7. Zheng S, Zou Y, Liang JY, et al. Identification and validation of a combined hypoxia and immune index for triple-negative breast cancer. *Mol Oncol* 2020;14:2814-33.
 8. Yang X, Weng X, Yang Y, et al. A combined hypoxia and immune gene signature for predicting survival and risk stratification in triple-negative breast cancer. *Aging (Albany NY)* 2021;13:19486-509.
 9. Sun X, Luo H, Han C, et al. Identification of a Hypoxia-Related Molecular Classification and Hypoxic Tumor Microenvironment Signature for Predicting the Prognosis of Patients with Triple-Negative Breast Cancer. *Front Oncol* 2021;11:700062.
 10. Islam R, Lam KW. Recent progress in small molecule agents for the targeted therapy of 454 triple-negative breast cancer. *Eur J Med Chem* 2020;207:112812.
 11. Wang C, Feng G, Zhu J, et al. Developing an immune signature for triple-negative breast cancer to predict prognosis and immune checkpoint inhibitor response. *Future Oncol* 2022;18:1055-66.
 12. Liu J, Ling Y, Su N, et al. A novel immune checkpoint-related gene signature for predicting overall survival and immune status in triple-negative breast cancer. *Transl Cancer Res* 2022;11:181-92.
 13. Qin Y, Deng J, Zhang L, et al. Tumor microenvironment characterization in triple-negative breast cancer identifies prognostic gene signature. *Aging (Albany NY)* 2021;13:5485-505.
 14. Liu C, Li Y, Xing X, et al. Immunogenomic landscape analyses of immune molecule signature-based risk panel for patients with triple-negative breast cancer. *Mol Ther Nucleic Acids* 2022;28:670-84.
 15. Ma J, Chen C, Liu S, et al. Identification of a five genes prognosis signature for triple-negative breast cancer using multi-omics methods and bioinformatics analysis. *Cancer Gene Ther* 2022;29:1578-89.
 16. Dill CD, Dammer EB, Griffen TL, et al. A network approach reveals driver genes associated with survival of patients with triple-negative breast cancer. *iScience* 2021;24:102451.
 17. Su P, Peng Z, Xu B, et al. Establishment and validation of an individualized macrophage-related gene signature to predict overall survival in patients with triple negative breast cancer. *PeerJ* 2021;9:e12383.
 18. Savas P, Virassamy B, Ye C, et al. Single-cell profiling of breast cancer T cells reveals a tissue-resident memory subset associated with improved prognosis. *Nat Med* 2018;24:986-93.
 19. Marchetti P, Antonov A, Anemona L, et al. New immunological potential markers for triple negative breast cancer: IL18R1, CD53, TRIM, Jaw1, LTB, PTPRCAP. *Discov Oncol* 2021;12:6.
 20. Kim JY, Jung HH, Sohn I, et al. Prognostication of a 13-immune-related-gene signature in patients with early triple-negative breast cancer. *Breast Cancer Res Treat* 2020;184:325-34.
 21. Wang X, Chen H. Prognosis Prediction Through an Integrated Analysis of Single-Cell and Bulk RNA-Sequencing Data in Triple-Negative Breast Cancer. *Front Genet* 2022;13:928175.
 22. Wang S, Zou X, Chen Y, et al. Effect of N6-Methyladenosine Regulators on Progression and Prognosis of Triple-Negative Breast Cancer. *Front Genet* 2020;11:580036.
 23. Zhang X, Kang X, Jin L, et al. ABCC9, NKAPL, and TMEM132C are potential diagnostic and prognostic markers in triple-negative breast cancer. *Cell Biol Int* 2020;44:2002-10.
 24. Gao Y, Wang X, Li S, et al. Identification of a DNA Methylation-Based Prognostic Signature for Patients with Triple-Negative Breast Cancer. *Med Sci Monit* 2021;27:e930025.
 25. Shen X, Zhong J, He J, et al. Identification of m6A modification patterns and development of m6A-hypoxia prognostic signature to characterize tumor microenvironment in triple-negative breast cancer. *Front Immunol* 2022;13:978092.
 26. Bissanum R, Kamolphiwong R, Navakanitworakul R, et al. Integrated bioinformatic analysis of potential biomarkers of poor prognosis in triple-negative breast cancer. *Transl Cancer Res* 2022;11:3039-49.
 27. Chen DL, Cai JH, Wang CCN. Identification of Key Prognostic Genes of Triple Negative Breast Cancer by LASSO-Based Machine Learning and Bioinformatics Analysis. *Genes (Basel)* 2022;13:902.
 28. Liu Y, Teng L, Fu S, et al. Highly heterogeneous-related genes of triple-negative breast cancer: potential diagnostic and prognostic biomarkers. *BMC Cancer* 2021;21:644.
 29. Liu Q, Wang X, Kong X, et al. Prognostic Alternative mRNA Splicing Signature and a Novel Biomarker

- in Triple-Negative Breast Cancer. *DNA Cell Biol* 2020;39:1051-63.
30. Thalor A, Kumar Joon H, Singh G, et al. Machine learning assisted analysis of breast cancer gene expression profiles reveals novel potential prognostic biomarkers for triple-negative breast cancer. *Comput Struct Biotechnol J* 2022;20:1618-31.
 31. Gong S, Song Z, Spezia-Lindner D, et al. Novel Insights Into Triple-Negative Breast Cancer Prognosis by Comprehensive Characterization of Aberrant Alternative Splicing. *Front Genet* 2020;11:534.
 32. Hong HC, Chuang CH, Huang WC, et al. A panel of eight microRNAs is a good predictive parameter for triple-negative breast cancer relapse. *Theranostics* 2020;10:8771-89.
 33. Zaka M, Sutton CW, Peng Y, et al. Model-Based Integration Analysis Revealed Presence of Novel Prognostic miRNA Targets and Important Cancer Driver Genes in Triple-Negative Breast Cancers. *Cancers (Basel)* 2020;12:632.
 34. Li YX, Wang SM, Li CQ. Four-lncRNA immune prognostic signature for triple-negative breast cancer Running title: Immune lncRNAs predict prognosis of TNBC. *Math Biosci Eng* 2021;18:3939-56.
 35. Wu J, Cai Y, Zhao G, et al. A ten N6-methyladenosine-related long non-coding RNAs signature predicts prognosis of triple-negative breast cancer. *J Clin Lab Anal* 2021;35:e23779.
 36. Jiang L, You C, Xiao Y, et al. Radiogenomic analysis reveals tumor heterogeneity of triple-negative breast cancer. *Cell Rep Med* 2022;3:100694.
 37. Kos Z, Roblin E, Kim RS, et al. Pitfalls in assessing stromal tumor infiltrating lymphocytes (sTILs) in breast cancer. *NPJ Breast Cancer* 2020;6:17.
 38. Wu J, Li X, Teng X, et al. Magnetic resonance imaging and molecular features associated with tumor-infiltrating lymphocytes in breast cancer. *Breast Cancer Res* 2018;20:101.

Cite this article as: Capobianco E. Overview of triple negative breast cancer prognostic signatures in the context of data science-driven clinico-genomics research. *Ann Transl Med* 2022;10(24):1300. doi: 10.21037/atm-22-5477