

QCSP on Reflexive Tournaments*

BENOÎT LAROSE, LACIM, Université du Québec a Montréal, Canada

BARNABY MARTIN, Department of Computer Science, Durham University, UK

PETAR MARKOVIĆ, Department of Mathematics and Informatics, University of Novi Sad, Serbia

DANIËL PAULUSMA, Department of Computer Science, Durham University, UK

SIANI SMITH, Department of Computer Science, Durham University, UK

STANISLAV ŽIVNÝ[†], Department of Computer Science, University of Oxford, UK

We give a complexity dichotomy for the Quantified Constraint Satisfaction Problem QCSP(H) when H is a reflexive tournament. It is well-known that reflexive tournaments can be split into a sequence of strongly connected components H_1, \dots, H_n so that there exists an edge from every vertex of H_i to every vertex of H_j if and only if $i < j$. We prove that if H has both its initial and final strongly connected component (possibly equal) of size 1, then QCSP(H) is in NL and otherwise QCSP(H) is NP-hard.

CCS Concepts: • **Theory of computation** → *Design and analysis of algorithms; Logic; Computational complexity and cryptography.*

Additional Key Words and Phrases: quantified constraints, constraint satisfaction, graph theorem, logic, computational complexity

ACM Reference Format:

Benoît Larose, Barnaby Martin, Petar Marković, Daniël Paulusma, Siani Smith, and Stanislav Živný. 2021. QCSP on Reflexive Tournaments. *ACM Trans. Comput. Logic* 1, 1, Article 1 (January 2021), 22 pages. <https://doi.org/10.1145/3508069>

1 INTRODUCTION

The *Quantified Constraint Satisfaction Problem* QCSP(B), for a fixed *template* (structure) B, is a popular generalisation of the *Constraint Satisfaction Problem* CSP(B). In the latter, one asks if a primitive positive sentence (the existential quantification of a conjunction of atoms) φ is true

*A conference version of this article will appear under the same name at the 29th Annual European Symposium on Algorithms (ESA 2021).

[†]Stanislav Živný was supported by a Royal Society University Research Fellowship. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 714532). The paper reflects only the authors' views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein.

Authors' addresses: Benoît Larose, blarose@lacim.ca, LACIM, Université du Québec a Montréal, Montréal, Canada; Barnaby Martin, barnabymartin@gmail.com, Department of Computer Science, Durham University, Durham, UK; Petar Marković, pera@dm.uns.ac.rs, Department of Mathematics and Informatics, University of Novi Sad, Novi Sad, Serbia; Daniël Paulusma, daniel.paulusma@durham.ac.uk, Department of Computer Science, Durham University, Durham, UK; Siani Smith, siani.smith@durham.ac.uk, Department of Computer Science, Durham University, Durham, UK; Stanislav Živný, standa.zivny@cs.ox.ac.uk, Department of Computer Science, University of Oxford, Oxford, UK.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2021 Association for Computing Machinery.

1529-3785/2021/1-ART1 \$15.00
<https://doi.org/10.1145/3508069>

50 on B , while in the former this sentence may also have universal quantification¹. Much of the
 51 theoretical research into (finite-domain²) CSPs has been in respect of a complexity classification
 52 project [5, 11], recently completed by [4, 22, 24], in which it is shown that all such problems are
 53 either in P or NP -complete. Various methods, including combinatorial (graph-theoretic), logical
 54 and universal-algebraic were brought to bear on this classification project, with many remarkable
 55 consequences.

56 Complexity classifications for QCSPs appear to be harder than for CSPs. Indeed, a classification
 57 for QCSPs will give a fortiori a classification for CSPs (if $B \uplus K_1$ is the disjoint union of B with
 58 an isolated element, then $QCSP(B \uplus K_1)$ and $CSP(B)$ are polynomial-time many-one equivalent).
 59 Just as $CSP(B)$ is always in NP , so $QCSP(B)$ is always in $Pspace$. However, no polychotomy has
 60 been conjectured for the complexities of $QCSP(B)$, though, until recently, only the complexities P ,
 61 NP -complete and $Pspace$ -complete were known. Recent work [25] has shown that this complexity
 62 landscape is considerably richer, and that dichotomies of the form P versus NP -hard (using Turing
 63 reductions) might be the sensible place to be looking for classifications.

64 $CSP(B)$ may equivalently be seen as the *homomorphism* problem which takes as input a structure
 65 A and asks if there is a homomorphism from A to B . The *surjective CSP*, $SCSP(B)$, is a cousin of
 66 $CSP(B)$ in which one requires that this homomorphism from A to B be surjective. From the logical
 67 perspective this translates to the stipulation that all elements of B be used as witnesses to the
 68 (existential) variables of the primitive positive input φ . The surjective CSP appears in the literature
 69 under a variety of names, including *surjective homomorphism* [2], *surjective colouring* [12, 15] and
 70 *vertex compaction* [19, 20]. $CSP(B)$ and $SCSP(B)$ have various other cousins: see the survey [2] or,
 71 in the specific context of reflexive tournaments, [15]. The only one we will dwell on here is the
 72 *retraction* problem $CSP^c(B)$ which can be defined in various ways but, in keeping with the present
 73 narrative, we could define logically as allowing atoms of the form $v = b$ in the input sentence φ
 74 where b is some element of B (the superscript c indicates that constants are allowed). It has only
 75 recently been shown that there exists a B so that $SCSP(B)$ is in P while $CSP^c(B)$ is NP -complete
 76 [23]. It is still not known whether such an example exists among the (partially reflexive) graphs.

77 It is well-known that the binary *cousin* relation is not transitive, so let us ask the question
 78 as to whether the surjective CSP and QCSP are themselves cousins? The algebraic operations
 79 pertaining to the CSP are *polymorphisms* and for QCSP these become *surjective* polymorphisms.
 80 On the other hand, a natural use of universal quantification in the QCSP might be to ensure some
 81 kind of surjective map (at least under some evaluation of many universally quantified variables).
 82 So it is that there may appear to be some relationship between the problems. Yet, there are known
 83 irreflexive graphs H for which $QCSP(H)$ is in NL , while $SCSP(H)$ is NP -complete (take the 6-
 84 cycle [18, 20]). On the other hand, one can find a 3-element B whose relations are preserved by a
 85 *semilattice-without-unit* operation such that both $CSP^c(B)$ and $SCSP(B)$ are in P but $QCSP(B)$ is
 86 $Pspace$ -complete. We are not aware of examples like this among graphs and it is perfectly possible
 87 that for (partially reflexive) graphs H , $SCSP(H)$ being in P implies that $QCSP(H)$ is in P .

88 Tournaments, both irreflexive and reflexive (and sometimes in between), have played a strong
 89 role as a testbed for conjectures and a habitat for classifications, for relatives of the CSP both
 90 complexity-theoretic [1, 10, 15] and algebraic [14, 21]. Looking at Table 1 one can see the last
 91 unresolved case is precisely QCSP on reflexive tournaments. This is the case we address in this
 92 paper. For irreflexive tournaments H , $QCSP(H)$ is in P if and only if $SCSP(H)$ in P , but for reflexive

93
 94 ¹Typically, primitive positive logic also possesses equality, but these can be propagated out by substitution, or removed in
 95 the case $x = x$. In the presence of universal quantification, any atom $x = y$ whose innermost variable is universal is false
 96 (unless x and y are the same variable). Other instances of equality may be propagated out as before. It follows that the
 97 complexity of $QCSP(B)$ is not affected by the presence or absence of equality, up to logarithmic space reducibility.

98 ²All structures considered in this article are finite.

99 tournaments this is not the case. When H is a reflexive tournament, we prove that $\text{QCSP}(H)$ is in
 100 NL if H has both initial and final strongly connected components trivial, and is NP-hard otherwise.
 101 In contrast to the proof from [10] and like the proof of [15], we will henceforth work largely
 102 combinatorially rather than algebraically. Note that we do not investigate beyond NP-hard, so our
 103 dichotomy cannot be compared directly to the trichotomy of [10] for irreflexive tournaments which
 104 distinguishes between P, NP-complete and Pspace-complete.

	QCSP	CSP	Surjective CSP	Retraction
irreflexive tournaments	trichotomy [10]	dichotomy [1]	dichotomy [1]	dichotomy [1]
reflexive tournaments	this paper	all trivial	dichotomy [15]	dichotomy [14]

111 Table 1. Our result in a wider context. The results for irreflexive tournaments were all proved in the more
 112 general setting of irreflexive semicomplete digraphs in the papers cited.

113
 114
 115
 116 In Section 3 we prove the NP-hard cases of our dichotomy. Our proof method follows that from
 117 [15], while adapting the ideas of [8] in order to make what was developed for Surjective CSP
 118 applicable to QCSP. The QCSP is not naturally a combinatorial problem but can be seen thusly
 119 when viewed in a certain way. We indeed closely mirror [15] with [8] in the strongly connected
 120 case. For the not strongly connected case, the adaptation from the strongly connected case was
 121 straightforward for the Surjective CSP in [15]. However, the straightforward method does not work
 122 for the QCSP. Instead, we seek a direct argument that essentially sees us extending the method
 123 from [15].

124 In Section 4 we prove the NL cases of our dichotomy. Here, we use ideas originally developed in
 125 (the conference version of) [8] and first used in the wild in [17]. Thus, we do not introduce new
 126 proof techniques as such but rather weave our proof through the reasonably intricate synthesis
 127 of different known techniques. In Section 5 we state our dichotomy and give some directions for
 128 future work.

129 2 PRELIMINARIES

130
 131 For an integer $k \geq 1$, we write $[k] := \{1, \dots, k\}$. A vertex $u \in V(G)$ in a digraph G is *backwards-*
 132 *adjacent* to another vertex $v \in V$ if $(u, v) \in E$. It is *forwards-adjacent* to another vertex $v \in V$ if
 133 $(v, u) \in E$. If a vertex u has a self-loop (u, u) , then u is *reflexive*; otherwise u is *irreflexive*. A digraph
 134 G is *reflexive* or *irreflexive* if all its vertices are reflexive or irreflexive, respectively.

135 The *directed path* on k vertices is the digraph with vertices u_0, \dots, u_{k-1} and edges (u_i, u_{i+1}) for
 136 $i = 0, \dots, k - 2$. By adding the edge (u_{k-1}, u_0) , we obtain the *directed cycle* on k vertices. A digraph
 137 G is *strongly connected* if for all $u, v \in V(G)$ there is a directed path in $E(G)$ from u to v . A *double*
 138 *edge* in a digraph G consists in a pair of distinct vertices $u, v \in V(G)$, so that (u, v) and (v, u) belong
 139 to $E(G)$. A digraph G is *semicomplete* if for every two distinct vertices u and v , at least one of (u, v) ,
 140 (v, u) belongs to $E(G)$. A semicomplete digraph G is a *tournament* if for every two distinct vertices
 141 u and v , exactly one of (u, v) , (v, u) belongs to $E(G)$. A reflexive tournament G is *transitive* if for
 142 every three vertices u, v, w with (u, v) , $(v, w) \in E(G)$, also (u, w) belongs to $E(G)$. A digraph F is a
 143 *subgraph* of a digraph G if $V(F) \subseteq V(G)$ and $E(F) \subseteq E(G)$. It is *induced* if $E(F)$ coincides with $E(G)$
 144 restricted to pairs containing only vertices of $V(F)$. A *subtournament* is an induced subgraph of a
 145 tournament. It is well known that a reflexive tournament H can be split into a sequence of strongly
 146 connected components H_1, \dots, H_n for some integer $n \geq 1$ so that there exists an edge from every
 147

vertex of H_i to every vertex of H_j if and only if $i < j$. We will use the notation $H_1 \Rightarrow \dots \Rightarrow H_n$ for H and we refer to H_1 and H_n as the *initial* and *final* components of H , respectively.

A *homomorphism* from a digraph G to a digraph H is a function $f : V(G) \rightarrow V(H)$ such that for all $u, v \in V(G)$ with $(u, v) \in E(G)$ we have $(f(u), f(v)) \in E(H)$. We say that f is (*vertex*)-*surjective* if for every vertex $x \in V(H)$ there exists a vertex $u \in V(G)$ with $f(u) = x$. A digraph H' is a *homomorphic image* of a digraph H if there is a surjective homomorphism from H to H' that is also *edge-surjective*, that is, for all $(x', y') \in E(H')$ there exists an $(x, y) \in E(H)$ with $x' = h(x)$ and $y' = h(y)$.

The problem H-RETRACTION takes as input a graph G of which H is an induced subgraph and asks whether there is a homomorphism from G to H that is the identity on H . This definition is polynomial-time many-one equivalent to the one we suggested in the introduction (see e.g. [2]). The *quantified constraint satisfaction problem* QCSP(H) takes as input a sentence $\varphi := \forall x_1 \exists y_1 \dots \forall x_n \exists y_n \Phi(x_1, y_1, \dots, x_n, y_n)$, where Φ is a conjunction of positive atomic (binary edge) relations. This is a yes-instance to the problem just in case $H \models \varphi$.

The *canonical query* of G (from [13]) is a primitive positive sentence φ_G that has the property that, for all H , G has a homomorphism to H iff $H \models \varphi_G$. It is built by mapping edges (x, y) from $E(G)$ to atoms $E(x, y)$ is an existentially quantified conjunctive query.

The *direct product* of two digraphs G and H , denoted $G \times H$, is the digraph on vertex set $V(G) \times V(H)$ with edges $((x, y), (x', y'))$ if and only if $(x, x') \in E(G)$ and $(y, y') \in E(H)$. We denote the direct product of k copies of G by G^k . A k -ary *polymorphism* of G is a homomorphism f from G^k to G ; if $k = 1$, then f is also called an *endomorphism*. A k -ary polymorphism f is *essentially unary* if there exists a unary operation g and $i \in [k]$ so that $f(x_1, \dots, x_k) = g(x_i)$ for every $(x_1, \dots, x_k) \in G^k$. Let us say that a k -ary polymorphism f is *uniformly z* for some $z \in V(G)$ if $f(x_1, \dots, x_k) = z$ for every $(x_1, \dots, x_k) \in V(G^k)$. We need the following two lemmas.

LEMMA 2.1. *Let H be a reflexive tournament and f be a k -ary polymorphism of H . If $f(x_1, \dots, x_k) = z$ for every $x \in V(H)$, then f is uniformly equal to z .*

PROOF. Consider some tuple (x_1, \dots, x_k) which has m distinct vertices. We proceed by induction on m , where the base case $m = 1$ is given as an assumption. Suppose we have the result for m vertices and let (x_1, \dots, x_k) have $m + 1$ distinct entries. For simplicity (and w.l.o.g.) we will consider this reordered and without duplicates as $(y_1, \dots, y_m, y_{m+1})$. Suppose f maps (x_1, \dots, x_k) to z' . Assume $(y_m, y_{m+1}) \in E(H)$ (the case (y_{m+1}, y_m) is symmetric). Then consider the tuples (y_1, \dots, y_m, y_m) and $(y_1, \dots, y_{m+1}, y_{m+1})$. By the inductive hypothesis, f maps each of these (when reordered and padded appropriately with duplicates) to z . Furthermore, we have co-ordinatewise edges from (y_1, \dots, y_m, y_m) to $(y_1, \dots, y_m, y_{m+1})$ and from $(y_1, \dots, y_m, y_{m+1})$ to $(y_1, \dots, y_{m+1}, y_{m+1})$. Since we deduce by the definition of polymorphism that both (z, z') , $(z', z) \in E(H)$, it follows that $z' = z$. Thus, f maps also $(y_1, \dots, y_m, y_{m+1})$ (when reordered and padded appropriately with duplicates) to z . That is, $f(x_1, \dots, x_k) = z$. \square

LEMMA 2.2. *Let H be the reflexive tournament $H_1 \Rightarrow \dots \Rightarrow H_i \Rightarrow \dots \Rightarrow H_n$. If f is a k -ary surjective polymorphism of H , then f preserves each of $V(H_1), \dots, V(H_n)$; that is, for every i and every tuple of k vertices $x_1, \dots, x_k \in V(H_i)$, $f(x_1, \dots, x_k) \in V(H_i)$.*

PROOF. Suppose f maps some tuple (x_1, \dots, x_m) from $V(H_i)$ to $y \in V(H_\ell)$. Let (x'_1, \dots, x'_m) be any tuple from $V(H_i)$. Since H_i is strongly connected, $f(x'_1, \dots, x'_m) \in V(H_\ell)$. It follows that if $\ell \neq i$, e.g. w.l.o.g. $\ell < i$, then some component $\ell' \geq i$ can not be in the range of f . \square

The relevance of this lemma is in its sequent corollary, which follows according to Proposition 3.15 of [3].

197 COROLLARY 2.3. *Let H be the reflexive tournament $H_1 \Rightarrow \dots \Rightarrow H_i \Rightarrow \dots \Rightarrow H_n$. Each subset of*
 198 *the domain $V(H_i)$ is definable by a QCSP instance in one free variable.*

199 An endomorphism e of a digraph G is a *constant map* if there exists a vertex $v \in V(G)$ such that
 200 $e(u) = v$ for every $u \in V(G)$, and e is the *identity* if $e(u) = u$ for every $u \in G$. An *automorphism* is a
 201 bijective endomorphism whose inverse is a homomorphism. An endomorphism is *trivial* if it is
 202 either an automorphism or a constant map; otherwise it is *non-trivial*. A digraph is *endo-trivial* if
 203 all of its endomorphisms are trivial. An endomorphism e of a digraph G *fixes* a subset $S \subseteq V(G)$
 204 if $e(S) = S$, that is, $e(x) \in S$ for every $x \in S$, and e fixes an induced subgraph F of G if it is the
 205 identity on $V(F)$. It fixes an induced subgraph F *up to automorphism* if $e(F)$ is an automorphic copy
 206 of F . An endomorphism e of G is a *retraction* of G if e is the identity on $e(V(G))$. A digraph is
 207 *retract-trivial* if all of its retractions are the identity or constant maps. Note that endo-triviality
 208 implies retract-triviality, but the reverse implication is not necessarily true (see [15]). However, on
 209 reflexive tournaments both concepts do coincide [15].

210 We need a series of results from [15]. The third one follows from the well-known fact that every
 211 strongly connected tournament has a directed Hamilton cycle [6].

212 LEMMA 2.4 ([15]). *A reflexive tournament is endo-trivial if and only if it is retract-trivial.*

213 LEMMA 2.5 ([15]). *Let H be an endo-trivial reflexive digraph with at least three vertices. Then every*
 214 *polymorphism of H is essentially unary.*

215 LEMMA 2.6 ([15]). *If H is an endo-trivial reflexive tournament, then H contains a directed Hamilton*
 216 *cycle.*

217 LEMMA 2.7 ([15]). *If H is an endo-trivial reflexive tournament, then every homomorphic image of H*
 218 *of size $1 < n < |V(H)|$ has a double edge.*

219 COROLLARY 2.8. *If H is an endo-trivial reflexive digraph on at least three vertices, then QCSP(H) is*
 220 *NP-hard (in fact it is even Pspace-complete).*

221 PROOF. This follows from Lemma 2.5 and [3]. □

222 3 THE PROOF OF THE NP-HARD CASES OF THE DICHOTOMY

223 We commence with the NP-hard cases of the dichotomy. The simpler NL cases will follow, in
 224 Section 4. In this section, the central results will appear as Corollaries 3.9 and 3.15. However, each
 225 of these proceeds via an induction where there are two base cases and two inductive (general) cases.
 226 Thus, there are eight principal propositions. Propositions 3.3, 3.5, 3.7 and 3.8 lead to Corollary 3.9
 227 and Propositions 3.11, 3.12, 3.13 and 3.14 lead to Corollary 3.15. The base cases are the simplest to
 228 understand and are given in the most detail. The principal propositions commence in Section 3.2.
 229 Before this we introduce our construction with some supporting lemmas.

230 3.1 The NP-Hardness Gadget

231 We introduce the gadget Cyl_m^* from [15] drawn in Figure 1. Take m disjoint copies of the (reflexive)
 232 directed m -cycle DC_m^* arranged in a cylindrical fashion so that there is an edge from i in the j th copy
 233 to i in the $(j + 1)$ th copy (drawn in red), and an edge from i in the $(j + 1)$ th copy to $(i + 1) \bmod m$
 234 in the j th copy (drawn in green). We consider DC_m^* to have vertices $\{1, \dots, m\}$. Recall that every
 235 strongly connected (reflexive) tournament on m vertices has a Hamilton Cycle HC_m . We label the
 236 vertices of HC_m as $1, \dots, m$ in order to attach it to the gadget Cyl_m^* .³

237 ³The superscripted * indicates that the corresponding graph is reflexive. This notation is inherited from [15]. It is not
 238 significant since we could safely assume every graph we work with is reflexive as the template is a reflexive tournament.
 239




Fig. 1. The gadget Cyl_m^* in the case $m := 4$ (self-loops are not drawn). We usually visualise the right-hand copy of DC_4^* as the “bottom” copy and then we talk about vertices “above” and “below” according to the red arrows.

The following lemma follows from induction on the copies of DC_m^* , since a reflexive tournament has no double edges.

LEMMA 3.1 ([15]). *In any homomorphism h from Cyl_m^* , with bottom cycle DC_m^* , to a reflexive tournament, if $|h(\text{DC}_m^*)| = 1$, then $|h(\text{Cyl}_m^*)| = 1$.*

We will use another property, denoted (\dagger) , of Cyl_m^* , which is that the retractions from Cyl_m^* to its bottom copy of DC_m^* , once propagated through the intermediate copies, induce on the top copy precisely the set of automorphisms of DC_m^* . That is, the top copy of DC_m^* is mapped isomorphically to the bottom copy, and all such isomorphisms may be realised. The reason is that in such a retraction, the $(j + 1)$ th copy may either map under the identity to the j th copy, or rotate one edge of the cycle clockwise, and Cyl_m^* consists of sufficiently many (namely m) copies of DC_m^* . Now let H be a reflexive tournament that contains a subtournament H_0 on m vertices that is endo-trivial. By Lemma 2.6, we find that H_0 contains at least one directed Hamilton cycle HC_0 . Define $\text{Spill}_m(H[H_0, \text{HC}_0])$ as follows. Begin with H and add a copy of the gadget Cyl_m^* , where the bottom copy of DC_m^* is identified with HC_0 , to build a digraph $F(H_0, \text{HC}_0)$. Now ask, for some $y \in V(H)$ whether there is a retraction r of $F(H_0, \text{HC}_0)$ to H so that some vertex x (not dependent on y) in the top copy of DC_m^* in Cyl_m^* is such that $r(x) = y$. Such vertices y comprise the set $\text{Spill}_m(H[H_0, \text{HC}_0])$.

Remark 1. If x belongs to some copy of DC_m^* that is not the top copy, we can find a vertex x' in the top copy of DC_m^* and a retraction r' from $F(H_0, \text{HC}_0)$ to H with $r'(x') = r(x) = y$, namely by letting r' map the vertices of higher copies of DC_m^* to the image of their corresponding vertex in the copy that contains x . In particular this implies that $\text{Spill}_m(H[H_0, \text{HC}_0])$ contains $V(H_0)$.

We note that the set $\text{Spill}_m(H[H_0, \text{HC}_0])$ is potentially dependent on which Hamilton cycle in H_0 is chosen. We now recall that $\text{Spill}_m(H[H_0, \text{HC}_0]) = V(H)$ if H retracts to H_0 .

LEMMA 3.2 ([15]). *If H is a reflexive tournament that retracts to a subtournament H_0 with Hamilton cycle HC_0 , then $\text{Spill}_m(H[H_0, \text{HC}_0]) = V(H)$.*

We now review a variant of a construction from [8]. Let G be a graph containing H where $|V(H)|$ is of size n . Consider all possible functions $\lambda : [n] \rightarrow V(H)$ (let us write $\lambda \in V(H)^{[n]}$ of cardinality N). For some such λ , let $\mathcal{G}(\lambda)$ be the graph G enriched with constants c_1, \dots, c_n where these are interpreted over $V(H)$ according to λ in the natural way (acting on the subscripts). We use calligraphic notation to remind the reader the signature has changed from $\{E\}$ to $\{E, c_1, \dots, c_n\}$

but we will still treat these structures as graphs. If we write $G(\lambda)$ without calligraphic notation we mean we look at only the $\{E\}$ -reduct, that is, we drop the constants. Of course, $G(\lambda)$ will always be G .

Let $\mathcal{G} = \bigotimes_{\lambda \in V(H)^{[n]}} \mathcal{G}(\lambda)$. That is, the vertices of \mathcal{G} are N -tuples over $V(G)$ and there is an edge between two such vertices (x_1, \dots, x_N) and (y_1, \dots, y_N) if and only if $(x_1, y_1), \dots, (x_N, y_N) \in E(G)$. Finally, the constants c_i are interpreted as (x_1, \dots, x_N) so that $\lambda_1(c_i) = x_1, \dots, \lambda_N(c_i) = x_N$. An important induced substructure of \mathcal{G} is $\{(x, \dots, x) : x \in V(G)\}$. It is a copy of G called the *diagonal* copy and will play an important role in the sequel. To comprehend better the construction of \mathcal{G} from the sundry $\mathcal{G}(\lambda)$, confer on Figure 2.

The final ingredient of our fundamental construction involves taking some structure \mathcal{G} and making its canonical query with all vertices other than those corresponding to c_1, \dots, c_n becoming existentially quantified variables (as usual in this construction). We then turn the c_1, \dots, c_n to variables y_1, \dots, y_n to make $\varphi_{\mathcal{G}}(y_1, \dots, y_n)$. Let \mathcal{H} come from the given construction in which $G = H$. It is proved in [8] that $H' \models \forall y_1, \dots, y_n \varphi_{\mathcal{H}}(y_1, \dots, y_n)$ if and only if $\text{QCSP}(H) \subseteq \text{QCSP}(H')$ (here we identify $\text{QCSP}(H)$ with the set of sentences that form its yes-instances). By way of a side note, let us consider a k -ary relation R over H with tuples $(x_1^1, \dots, x_k^1), \dots, (x_1^r, \dots, x_k^r)$. For $i \in [r]$, let λ_i map (c_1, \dots, c_k) to (x_1^i, \dots, x_k^i) . Let $\mathcal{H} = \bigotimes_{\lambda \in \{\lambda_1, \dots, \lambda_r\}} \mathcal{H}(\lambda)$. Then $\varphi_{\mathcal{H}}(y_1, \dots, y_n)$ is the closure of R under the polymorphisms of H .

3.2 The strongly connected case: Two Base Cases

Recall that if H is a (reflexive) endo-trivial tournament, then $\text{QCSP}(H)$ is NP-hard due to Lemma 2.5 combined with the results from [3]. Indeed, Theorem 5.2 in [3] states that any H with more than one element, such that all surjective polymorphisms of H are essentially unary, satisfies that $\text{QCSP}(H)$ is Pspace-complete. However H may not be endo-trivial. We will now show how to deal with the case where H is not endo-trivial but retracts to an endo-trivial subtournament. For doing this we use the NP-hardness gadget, but we need to distinguish between two different cases.

PROPOSITION 3.3 (BASE CASE I). *Let H be a reflexive tournament that retracts to an endo-trivial subtournament H_0 with Hamilton cycle HC_0 . Assume that H retracts to H_0^d for every isomorphic copy $H_0^d = i(H_0)$ of H_0 in H with $\text{Spill}_m(H[H_0^d, i(\text{HC}_0)]) = V(H)$. Then H_0 -RETRACTION can be polynomially reduced to $\text{QCSP}(H)$.*

PROOF. Let m be the size of $|V(H_0)|$ and n be the size of $|V(H)|$. Let G be an instance of H_0 -RETRACTION. We build an instance φ of $\text{QCSP}(H)$ in the following fashion. First, take a copy of H together with G and build G' by identifying these on the copy of H_0 that they both possess as an induced subgraph. Now, consider all possible functions $\lambda : [n] \rightarrow V(H)$. For some such λ , let $\mathcal{G}'(\lambda)$ be the graph enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H)$ according to λ in the natural way (acting on the subscripts).

Let $\mathcal{G}' = \bigotimes_{\lambda \in V(H)^{[n]}} \mathcal{G}'(\lambda)$. Let G'^d, H^d and H_0^d be the diagonal copies of G', H and H_0 in \mathcal{G}' . Let \mathcal{H} be the subgraph of \mathcal{G}' induced by $V(H) \times \dots \times V(H)$. Note that the constants c_1, \dots, c_n live in \mathcal{H} . Now build \mathcal{G}'' from \mathcal{G}' by augmenting a new copy of Cyl_m^* for every vertex $v \in V(\mathcal{H}) \setminus V(H_0^d)$. Vertex v is to be identified with any vertex in the top copy of DC_m^* in Cyl_m^* and the bottom copy of DC_m^* is to be identified with HC_0 in H_0^d according to the identity function. (Thus, in each case, the new vertices are the middle cycles of Cyl_m^* and all but one of the vertices in the top cycle of Cyl_m^* .)

Finally, build φ from the canonical query of \mathcal{G}'' where we additionally turn the constants c_1, \dots, c_n to outermost universal variables. The size of φ is doubly exponential in n (the size of H) but this is constant, so still polynomial in the size of G .

We claim that G retracts to H_0 if and only if $\varphi \in \text{QCSP}(H)$.




Fig. 2. Illustrations of direct product with constants.

First suppose that G retracts to H_0 . Let λ be some assignment of the universal variables of φ to H . To prove $\varphi \in \text{QCSP}(H)$ it suffices to prove that there is a homomorphism from \mathcal{G}'' to H that extends λ . Then for this it suffices to prove that there is a homomorphism h from \mathcal{G}' that extends λ . Let us explain why. Because H retracts to H_0 , we have $\text{Spill}_m(H[H_0, HC_0]) = V(H)$ due to Lemma 3.2. Hence, if $h(x) = y$ for two vertices $x \in V(\mathcal{H}) \setminus V(H_0^d)$ and $y \in V(H)$, we can always find a retraction of the graph $F(H_0, HC_0)$ to H that maps x to y , and we mimic this retraction on the corresponding subgraph in \mathcal{G}'' . The crucial observation is that this can be done independently for each vertex in $V(\mathcal{H}) \setminus V(H_0^d)$, as two vertices of different copies of Cyl_m^* are only adjacent if they both belong to \mathcal{H} .

Henceforth let us consider the homomorphic image of \mathcal{G}' that is $\mathcal{G}'(\lambda)$. To prove $\varphi \in \text{QCSP}(H)$ it suffices to prove that there is a homomorphism from $G'(\lambda)$ to H that extends λ . Note that it will be sufficient to prove that G' retracts to H . Let h be the natural retraction from G' to H that extends the known retraction from G to H_0 . We are done.




Fig. 3. An interesting tournament H on six vertices (self-loops are not drawn). This tournament does not retract to the DC_3^* on the left-hand side, yet $\text{Spill}_3(H[DC_3^*, DC_3]) = V(H)$.

Suppose now $\varphi \in \text{QCSP}(H)$. Choose some surjection for λ , the assignment of the universal variables of φ to H . Recall $N = |V(H)^{[n]}|$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H)$ induces a surjective homomorphism s from \mathcal{G}' to H which contains within it a surjective homomorphism s' from $\mathcal{H} = H^N$ to H . Consider the diagonal copy of $H_0^d \subset H^d \subset G'^d$ in \mathcal{G}' . By abuse of notation we will also consider each of s and s' acting just on the diagonal. If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}' , we have $|s'(H^d)| = 1$. Indeed, this was the property we noted in Lemma 3.1. By Lemma 2.1, this would mean s' is uniformly mapping \mathcal{H} to one vertex, which is impossible as s' is surjective. Now we will work exclusively in the diagonal copy G'^d . As $1 < |s'(H_0^d)| < m$ is not possible either due to Lemma 2.7, we find that $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy of itself in H which we will call $H'_0 = i(H_0^d)$ for some isomorphism i .

We claim that $\text{Spill}_m(H[H'_0, i(HC_0^d)]) = V(H)$. In order to see this, consider a vertex $y \in V(H)$. As s' is surjective, there exists a vertex $x \in V(\mathcal{H})$ with $s'(x) = y$. By construction, x belongs to some top copy of DC_m^* in Cyl_m^* in $F(H_0, HC_0)$. We can extend i^{-1} to an isomorphism from the copy of Cyl_m^* (which has $i(HC_0^d)$ as its bottom cycle) in the graph $F(H'_0, i(HC_0^d))$ to the copy of Cyl_m^* (which has HC_0^d as its bottom cycle) in the graph $F(H_0, HC_0)$. We define a mapping r^* from $F(H'_0, i(HC_0^d))$ to H by $r^*(u) = s' \circ i^{-1}(u)$ if u is on the copy of Cyl_m^* in $F(H'_0, i(HC_0^d))$ and $r^*(u) = u$ otherwise. We observe that $r^*(u) = u$ if $u \in V(H'_0)$ as s' coincides with i on H_0 . As H_0^d separates the other vertices of the copy of Cyl_m^* from $V(H^d) \setminus V(H_0^d)$, in the sense that removing H_0^d would disconnect them, this means that r^* is a retraction from $F(H'_0, i(HC_0^d))$ to H . We find that r^* maps $i(x)$ to $s' \circ i^{-1}(i(x)) = s'(x) = y$. Moreover, as x is in the top copy of DC_m^* in $F(H_0, HC_0)$, we conclude that y always belongs to $\text{Spill}_m(H[H'_0, i(HC_0^d)])$.

As $\text{Spill}_m(H[H'_0, i(HC_0^d)]) = V(H)$, we find, by assumption of the lemma, that there exists a retraction r from H to H'_0 . Now, recalling that we can view s' acting just on the diagonal copy H^d of H , $i^{-1} \circ r \circ s'$ is the desired retraction of G to H_0 . \square

We now need to deal with the situation in which we have an isomorphic copy $H'_0 = i(H_0)$ of H_0 in H with $\text{Spill}_m(H[H'_0, i(HC_0)]) = V(H)$, such that H does not retract to H'_0 (see Figure 3 for an example). We cannot deal with this case in a direct manner and first show another base case. For this we need the following lemma and an extension of endo-triviality that we discuss afterwards.

LEMMA 3.4 ([15]). *Let H be a reflexive tournament, containing a subtournament H_0 so that any endomorphism of H that fixes H_0 as a graph is an automorphism. Then any endomorphism of H that maps H_0 to an isomorphic copy $H'_0 = i(H_0)$ of itself is an automorphism of H .*

Let H_0 be an induced subgraph of a digraph H . We say that the pair (H, H_0) is *endo-trivial* if all endomorphisms of H that fix H_0 are automorphisms.

PROPOSITION 3.5 (BASE CASE II). *Let H be a reflexive tournament with a subtournament H_0 with Hamilton cycle HC_0 so that (H, H_0) and H_0 are endo-trivial and $\text{Spill}_m(H[H_0, HC_0]) = V(H)$. Then H-RETRACTION can be polynomially reduced to QCSP(H).*

PROOF. Let G be an instance of H-RETRACTION. Let m be the size of $|V(H_0)|$ and n be the size of $|V(H)|$. We build an instance φ of QCSP(H) in the following fashion. Consider all possible functions $\lambda : [n] \rightarrow V(H)$. For some such λ , let $\mathcal{G}(\lambda)$ be the graph enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H)$ according to λ in the natural way (acting on the subscripts).

Let $\mathcal{G} = \bigotimes_{\lambda \in V(H)^{[n]}} \mathcal{G}(\lambda)$. Let G^d, H^d and H_0^d be the diagonal copies of G, H and H_0 in \mathcal{G} . Let \mathcal{H} be the subgraph of \mathcal{G} induced by $V(H) \times \dots \times V(H)$. Note that the constants c_1, \dots, c_n live in \mathcal{H} . Now build \mathcal{G}' from \mathcal{G} by augmenting a new copy of Cyl_m^* for every vertex $v \in V(\mathcal{H}) \setminus V(H_0^d)$. Vertex v is to be identified with any vertex in the top copy of DC_m^* in Cyl_m^* and the bottom copy of DC_m^* is to be identified with HC_0 in H_0^d according to the identity function.

Finally, build φ from the canonical query of \mathcal{G}' where we additionally turn the constants c_1, \dots, c_n to outermost universal variables.

First suppose that G retracts to H by r . Let λ be some assignment of the universal variables of φ to H . To prove $\varphi \in \text{QCSP}(H)$ it suffices to prove that there is a homomorphism from \mathcal{G}' to H that extends λ and for this it suffices to prove that there is a homomorphism from \mathcal{G} that extends λ . This is always possible since we have $\text{Spill}_m(H[H_0, HC_0]) = V(H)$ by assumption.

Henceforth let us consider the homomorphic image of \mathcal{G} that is $\mathcal{G}(\lambda)$. To prove $\varphi \in \text{QCSP}(H)$ it suffices to prove that there is a homomorphism from $G(\lambda)$ to H that extends λ . Note that it will be sufficient to prove that G retracts to H . Well this was our original assumption so we are done.

Suppose now $\varphi \in \text{QCSP}(H)$. Choose some surjection for λ , the assignment of the universal variables of φ to H . Recall $N = |V(H)^{[n]}|$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H)$ induces a surjective homomorphism s from \mathcal{G}' to H which contains within it a surjective homomorphism s' from $\mathcal{H} = H^N$ to H . Consider the diagonal copy of $H_0^d \subset H^d \subset G^d$ in $(G)^N$. By abuse of notation we will also consider each of s and s' acting just on the diagonal. If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}' , we have $|s'(H^d)| = 1$. By Lemma 2.1, this would mean s' is uniformly mapping \mathcal{H} to one vertex, which is impossible as s' is surjective. Now we will work exclusively on the diagonal copy G^d . As $1 < |s'(H_0^d)| < m$ is not possible either due to Lemma 2.7, we find that $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy of itself in H which we will call $H'_0 = i(H_0^d)$ for some isomorphism i .

As (H, H_0) is endo-trivial, Lemma 3.4 tells us that the restriction of s' to H^d is an automorphism of H^d , which we call α . The required retraction from G to H is now given by $\alpha^{-1} \circ s'$. \square

3.3 The strongly connected case: Generalising the Base Cases

We now generalise the two base cases to more general cases via some recursive procedure. Afterwards we will show how to combine these two cases to complete our proof. We will first need a slightly generalised version of Lemma 3.4, which nonetheless has virtually the same proof. For completeness of this article we provide this proof from [15].

LEMMA 3.6 ([15]). *Let $H_2 \supset H_1 \supset H_0$ be a sequence of strongly connected reflexive tournaments, each one a subtournament of the one before. Suppose that any endomorphism of H_1 that fixes H_0 is an automorphism. Then any endomorphism h of H_2 that maps H_0 to an isomorphic copy $H'_0 = i(H_0)$ of itself also gives an isomorphic copy of H_1 in $h(H_1)$.*

will be sufficient to prove that G' retracts to H_{k+1} . Let h be the natural retraction from G' to H_{k+1} that extends the known retraction from G to H_k . We are done.

Suppose now $\varphi \in \text{QCSP}(H_{k+1})$. Choose some surjection for λ , the assignment of the universal variables of φ to H_{k+1} . Let $N = |V(H_{k+1})|^{[n]}$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H_{k+1})$ induces a surjective homomorphism s from \mathcal{G}' to H_{k+1} which contains within it a surjective homomorphism s' from $\mathcal{H} = H_{k+1}^N$ to H_{k+1} . Consider the diagonal copy of $H_0^d \subset \dots \subset H_k^d \subset H_{k+1}^d \subset G'^d$ in \mathcal{G}' . By abuse of notation we will also consider each of s and s' acting just on the diagonal. If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}'' , we could follow the chain of spills to deduce that $|s'(H_{k+1}^d)| = 1$, which is not possible by Lemma 2.1. Moreover, $1 < |s'(H_0^d)| < |V(H_0^d)|$ is impossible due to Lemma 2.7. Now we will work exclusively on the diagonal copy G'^d .

Thus, $|s'(H_0^d)| = |V(H_0^d)|$ and indeed s' maps H_0^d to an isomorphic copy of itself in H_{k+1} which we will call $H'_0 = i(H_0^d)$. We now apply Lemma 3.6 as well as our assumed endo-trivialities to derive that s' in fact maps H_k^d by the isomorphism i to a copy of itself in H_{k+1} which we will call H'_k . Since s' is surjective, we can deduce that $\text{Spill}_{a_k}(H_{k+1}[H'_k, i(\text{HC}_k^d)]) = V(H_{k+1})$ in the same way as in the proof of Proposition 3.3. and so there exists a retraction r from H_{k+1} to H'_k . Now $i^{-1} \circ r \circ s'$ gives the desired retraction of G to H_k . \square

PROPOSITION 3.8 (GENERAL CASE II). *Let $H_0, H_1, \dots, H_k, H_{k+1}$ be reflexive tournaments, the first $k+1$ of which have Hamilton cycles $\text{HC}_0, \text{HC}_1, \dots, \text{HC}_k$, respectively, so that $H_0 \subseteq H_1 \subseteq \dots \subseteq H_k \subseteq H_{k+1}$. Suppose that $H_0, (H_1, H_0), \dots, (H_k, H_{k-1}), (H_{k+1}, H_k)$ are endo-trivial and that*

$$\begin{aligned} \text{Spill}_{a_0}(H_1[H_0, \text{HC}_0]) &= V(H_1) \\ \text{Spill}_{a_1}(H_2[H_1, \text{HC}_1]) &= V(H_2) \\ &\vdots \\ \text{Spill}_{a_{k-1}}(H_k[H_{k-1}, \text{HC}_{k-1}]) &= V(H_k) \\ \text{Spill}_{a_k}(H_{k+1}[H_k, \text{HC}_k]) &= V(H_{k+1}) \end{aligned}$$

Then H_{k+1} -RETRACTION can be polynomially reduced to $\text{QCSP}(H_{k+1})$.

PROOF. Let $n = a_{k+1} = |V(H_{k+1})|$ and let a_k, \dots, a_0 be the cardinalities of $|V(H_k)|, \dots, |V(H_0)|$, respectively. Let G be an instance of H_{k+1} -RETRACTION. We build an instance φ of $\text{QCSP}(H_{k+1})$ in the following fashion. Consider all possible functions $\lambda : [n] \rightarrow V(H_{k+1})$. For some such λ , let $\mathcal{G}(\lambda)$ be the graph enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H_{k+1})$ according to λ in the natural way (acting on the subscripts).

Let $\mathcal{G} = \bigotimes_{\lambda \in V(H_{k+1})^{[n]}} \mathcal{G}(\lambda)$. Let $G^d, H_{k+1}^d, H_k^d, \dots, H_0^d$ be the diagonal copies of $G, H_{k+1}, H_k, \dots, H_0$ in \mathcal{G} . Let \mathcal{H}_{k+1} be the subgraph of \mathcal{G} induced by $V(H_{k+1}) \times \dots \times V(H_{k+1})$. Note that the constants c_1, \dots, c_n live in \mathcal{H}_{k+1} .

Build \mathcal{G}' from \mathcal{G} by first augmenting a new copy of $\text{Cyl}_{a_k}^*$ for every vertex $v \in V(\mathcal{H}_{k+1}) \setminus V(H_k^d)$. Vertex v is to be identified with any vertex in the top copy of DC_{a_k} in $\text{Cyl}_{a_k}^*$ and the bottom copy of DC_{a_k} is to be identified with HC_k in H_k^d according to the identity function. Now, for each $i \in [k]$, and $v \in V(H_i^d) \setminus V(H_{i-1}^d)$, we add a copy of $\text{Cyl}_{a_{i-1}}^*$, where v is identified with any vertex in the top copy of $\text{DC}_{a_{i-1}}^*$ in $\text{Cyl}_{a_{i-1}}^*$ and the bottom copy of $\text{DC}_{a_{i-1}}^*$ is to be identified with H_{i-1}^d according to the identity map of $\text{DC}_{a_{i-1}}^*$ to HC_{i-1}^d .

Finally, build φ from the canonical query of \mathcal{G}' where we additionally turn the constants c_1, \dots, c_n to outermost universal variables.

First suppose that G retracts to H_{k+1} . Let h be a retraction from G to H_{k+1} . Let λ be some assignment of the universal variables of φ to H_{k+1} . To prove $\varphi \in \text{QCSP}(H_{k+1})$ it suffices to prove that there is a homomorphism from \mathcal{G}' to H_{k+1} that extends λ and for this it suffices to prove that

there is a homomorphism from \mathcal{G} that extends λ . The extension of the latter to the former will always be possible due to the spill assumptions.

Henceforth let us consider the homomorphic image of \mathcal{G} that is $\mathcal{G}(\lambda)$. To prove $\varphi \in \text{QCSP}(H_{k+1})$ it suffices to prove that there is a homomorphism from $\mathcal{G}(\lambda)$ to H_{k+1} that extends λ . Note that it will be sufficient to prove that G retracts to H_{k+1} . Well this was our original assumption so we are done.

Suppose now $\varphi \in \text{QCSP}(H_{k+1})$. Choose some surjection for λ , the assignment of the universal variables of φ to H_{k+1} . Let $N = |V(H_{k+1})|^{[n]}$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H_{k+1})$ induces a surjective homomorphism s from \mathcal{G} to H_{k+1} which contains within it a surjective homomorphism s' from $\mathcal{H}_{k+1} = H_{k+1}^N$ to H_{k+1} . Consider the diagonal copy of $H_0^d \subset H_1^d \subset \dots \subset H_{k+1}^d$ in \mathcal{G} . By abuse of notation we will also consider each of s and s' acting just on the diagonal. If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}' , we have $|s'(H^d)| = 1$. Now we follow the chain of spills to deduce that $|s'(\mathcal{H}_{k+1})| = 1$, a contradiction. We now apply Lemma 3.6 as well as our assumed endo-trivialities to derive that s' in fact maps H_k^d by the isomorphism i to a copy of itself in H_{k+1} , which we will call H'_k . Now we can deduce, via Lemma 3.4, that $s'(H_{k+1}^d)$ is an automorphism of H_{k+1} , which we call α . The required retraction from G to H_{k+1} is now given by $\alpha^{-1} \circ s'$. \square

COROLLARY 3.9. *Let H be a non-trivial strongly connected reflexive tournament. Then $\text{QCSP}(H)$ is NP-hard.*

PROOF. As H is a strongly connected reflexive tournament, which has more than one vertex by our assumption, H is not transitive. Note that H -RETRACTION is NP-complete (see Section 4.5 in [15], using results from [5, 14, 16]). Thus, if H is endo-trivial, the result follows from Proposition 3.3 (note that we could also have used Corollary 2.8).

Suppose H is not endo-trivial. Then, by Lemma 2.4, H is not retract-trivial either. This means that H has a non-trivial retraction to some subtournament H_0 . We may assume that H_0 is endo-trivial, as otherwise we will repeat the argument until we find a retraction from H to an endo-trivial (and consequently strongly connected) subtournament.

Suppose that H retracts to all isomorphic copies $H'_0 = i(H_0)$ of H_0 within it, except possibly those for which $\text{Spill}_m(H[H'_0, i(\text{HC}_0)]) \neq V(H)$. Then the result follows from Proposition 3.3. So there is a copy $H'_0 = i(H_0)$ to which H does not retract for which $\text{Spill}_m(H[H'_0, i(\text{HC}_0)]) = V(H)$. If (H, H'_0) is endo-trivial, the result follows from Proposition 3.5. Thus we assume (H, H'_0) is not endo-trivial and we deduce the existence of $H'_0 \subset H_1 \subset H$ (H_1 is strictly between H and H'_0) so that (H_1, H'_0) and H'_0 are endo-trivial and H retracts to H_1 . Now we are ready to break out. Either H retracts to all isomorphic copies of $H'_1 = i(H_1)$ in H , except possibly for those so that $\text{Spill}_m(H[H'_1, i(\text{HC}_1)]) \neq V(H)$, and we apply Proposition 3.7, or there exists a copy H'_1 , with $\text{Spill}_m(H[H'_1, i(\text{HC}_1)]) = V(H)$, to which it does not retract. If (H, H'_1) is endo-trivial, the result follows from Proposition 3.8. Otherwise we iterate the method, which will terminate because our structures are getting strictly smaller. \square

3.4 An initial strongly connected component that is non-trivial

Let H^+ denote any reflexive tournament that has an initial strongly connected component H that is non-trivial (not of size 1). Let Cyl_m^{**} be Cyl_m^* but with a pendant out-edge hanging from the top-most cycle. This edge is directed to the vertex x . Thus, Cyl_m^{**} contains one additional vertex to Cyl_m^* and this has an incoming edge from some vertex in the top-most cycle DC_m^* (it does not matter which one). Cyl_m^{**} is drawn in Figure 4.

	Strongly connected	Initial component strongly connected
Graph	H	H^+
Gadget	Cyl_m^*	Cyl_m^{**+}
Subgraph (strongly connected)	H_0	H_0
Hamilton cycle	HC_0	HC_0
Spill	$Spill_m(H[H_0, HC_0])$	$Spill_m^+(H^+[H_0, HC_0])$

Table 2. Mapping notation from the strongly connected case to the case in which there is an initial strongly connected component that is non-trivial.




Fig. 4. The gadget Cyl_m^{**+} in the case $m := 4$ (self-loops are not drawn). We usually visualise the right-hand copy of DC_4^* as the “bottom” copy and then we talk about vertices “above” and “below” according to the red arrows. The vertex x is depicted at the left-hand extremity.

Define $Spill_m^+$ as $Spill_m$ but with respect to Cyl_m^{**+} instead of Cyl_m^* . At this point we risk confusion with our overburdened notation. Let us address in Table 2 how our notation maps from the strongly connected case to that in which there is an initial strongly connected component that is non-trivial.

Note that Lemma 3.1, with Cyl_m^* replaced by Cyl_m^{**+} , does not hold.

LEMMA 3.10. *Let H^+ be some reflexive tournament that has an initial strongly connected component H that is non-trivial and contains endo-trivial H_0 with Hamilton cycle HC_0 . Suppose $Spill_m^+(H[H_0, HC_0]) = V(H)$, then $Spill_m^+(H^+[H_0, HC_0]) = V(H^+)$.*

PROOF. We only need to argue for the $x \in H^+ \setminus H$. In this case, we may evaluate all the cycles in Cyl_m^{**+} onto HC_0 with each vertex mapping to the one directly beneath it. This works as x is forward-adjacent from every vertex in HC_0 . \square

The condition of endo-triviality of H_0 was not used in the proof of Lemma 3.10.

PROPOSITION 3.11 (BASE CASE A-I). *Let H^+ be some reflexive tournament that has an initial strongly connected component H that is non-trivial and contains endo-trivial H_0 with Hamilton cycle HC_0 . Assume that H retracts to H'_0 for every isomorphic copy $H'_0 = i(H_0)$ of H_0 in H with $Spill_m^+(H[H'_0, i(HC_0)]) = V(H)$. Then H_0 -RETRACTION can be polynomially reduced to QCSP(H^+).*

PROOF. Let m be the size of $|V(H_0)|$ and n be the size of $|V(H)|$. Let G be an instance of H_0 -RETRACTION. We build an instance φ of QCSP(H^+) in the following fashion. First, take a copy of H

687 together with G and build G' by identifying these on the copy of H_0 that they both possess as an
 688 induced subgraph.

689 Now, consider all possible functions $\lambda : [n] \rightarrow V(H)$. For some such λ , let $\mathcal{G}'(\lambda)$ be the graph
 690 enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H)$ according
 691 to λ in the natural way (acting on the subscripts).

692 Let $\mathcal{G}' = \bigotimes_{\lambda \in V(H)^{[n]}} \mathcal{G}'(\lambda)$. Let G'^d, H^d and H_0^d be the diagonal copies of G', H and H_0 in \mathcal{G}' . Let
 693 \mathcal{H} be the subgraph of \mathcal{G}' induced by $V(H) \times \dots \times V(H)$. Note that the constants c_1, \dots, c_n live in
 694 \mathcal{H} . Now build \mathcal{G}'' from \mathcal{G}' by augmenting a new copy of Cyl_m^{*+} for every vertex $v \in V(\mathcal{H}) \setminus V(H_0^d)$.
 695 Vertex v is to be identified with the vertex x that is at the end of the out-edge pendant on the top
 696 copy of DC_m^* in Cyl_m^{*+} and the bottom copy of DC_m^* is to be identified with HC_0 in H_0^d according to
 697 the identity function. Call these *the Cyl_m^{*+} of the second stage*.

698 Now build \mathcal{G}''' by adding an edge from each vertex c_i to a new vertex d_i (for each $i \in [n]$). Now
 699 add a copy of Cyl_m^{*+} for every vertex $v \in \{d_1, \dots, d_n\}$. Vertex v is to be identified with the vertex x
 700 that is at the end of the out-edge pendant on the top copy of DC_m^* in Cyl_m^{*+} and the bottom copy
 701 of DC_m^* is to be identified with HC_0 in H_0^d according to the identity function. Call these *the Cyl_m^{*+}*
 702 *of the third stage*.

703 Finally, build φ from the canonical query of \mathcal{G}''' , where we additionally turn the vertices d_1, \dots, d_n
 704 to outermost universal variables z_1, \dots, z_n . Then existentially quantify all remaining constants and
 705 vertices innermost. Finally, restrict all except the universal variables to be in $V(H)$, appealing to
 706 the definition guaranteed by Corollary 2.3.

707 We claim that G retracts to H_0 if and only if $\varphi \in \text{QCSP}(H^+)$.

708 First suppose that G retracts to H_0 by r . Let λ' be some assignment of the universal variables
 709 z_1, \dots, z_n of φ to H^+ and choose y_1, \dots, y_n backwards-adjacent to these in H , mapped by λ . To prove
 710 $\varphi \in \text{QCSP}(H^+)$ it suffices to prove that there is a homomorphism from \mathcal{G}'' to H^+ that extends λ
 711 and for this it suffices to prove that there is a homomorphism h from \mathcal{G}' to H that extends λ . Let
 712 us explain why. Because H retracts to H_0 , we have $\text{Spill}_m(H[H_0, \text{HC}_0]) = V(H)$ due to Lemma 3.2
 713 which implies the weaker $\text{Spill}_m^+(H[H_0, \text{HC}_0]) = V(H)$. For the Cyl_m^{*+} of the second stage, the
 714 weaker statement suffices, but for the Cyl_m^{*+} of the third stage, the stronger statement is needed.

715 Henceforth let us consider the homomorphic image of \mathcal{G}' that is $\mathcal{G}'(\lambda)$. To prove $\varphi \in \text{QCSP}(H^+)$
 716 it suffices to prove that there is a homomorphism from $G'(\lambda)$ to H that extends λ . Note that it will
 717 be sufficient to prove that G' retracts to H . Let h be the natural retraction from G' to H that extends
 718 the known retraction r from G to H_0 . We are done.

719 Suppose now $\varphi \in \text{QCSP}(H^+)$. Choose some surjection for λ' mapping z_1, \dots, z_n to H . Choose
 720 some y_1, \dots, y_n backwards-adjacent to these and let this be the map λ . Note that it is not possible
 721 for all y_1, \dots, y_n to be evaluated as a single vertex as the initial strongly connected component is
 722 non-trivial.

723 The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H)$ induces a non-trivial
 724 homomorphism s from \mathcal{G}'' to H which contains within it a non-trivial homomorphism s' from
 725 $\mathcal{H} = H^N$ to H . Consider the diagonal copy of $H_0^d \subset H^d \subset G'^d$ in \mathcal{G}' . By abuse of notation we will
 726 also consider each of s and s' acting just on the diagonal.

727 If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}'' , we have that $s'(H^d)$ is an in-star (that is, a single terminal
 728 vertex receiving an edge from potentially numerous initial vertices), but this is not possible as H^d
 729 is strongly connected. As $1 < |s'(H_0^d)| < m$ is not possible either due to Lemma 2.7, we find that
 730 $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy of itself in H which we will call $H'_0 = i(H_0^d)$ for some
 731 isomorphism i .

732 We claim that $\text{Spill}_m^+(H[H'_0, i(\text{HC}_0^d)]) = V(H)$. Since λ' is surjective on H^+ , this is enforced
 733 explicitly by the Cyl_m^{*+} of the third stage. As $\text{Spill}_m^+(H[H'_0, i(\text{HC}_0^d)]) = V(H)$, we find, by assumption
 734

of the lemma, that there exists a retraction r from H^d to H'_0 . Now $i^{-1} \circ r \circ s'$ is the desired retraction of G to H_0 . \square

PROPOSITION 3.12 (BASE CASE A-II). *Let H^+ be some reflexive tournament that has an initial strongly connected component H that is non-trivial and contains H_0 with Hamilton cycle HC_0 so that (H, H_0) and H_0 are endo-trivial and $\text{Spill}_m^+(H[H_0, HC_0]) = V(H)$. Then H-RETRACTION can be polynomially reduced to QCSP(H^+).*

PROOF. Let m be the size of $|V(H_0)|$ and n be the size of $|V(H)|$. Let G be an instance of H-RETRACTION. We build an instance φ of QCSP(H^+) in the following fashion. Consider all possible functions $\lambda : [n] \rightarrow V(H)$. For some such λ , let $\mathcal{G}(\lambda)$ be the graph enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H)$ according to λ in the natural way (acting on the subscripts).

Let $\mathcal{G} = \bigotimes_{\lambda \in V(H)^{[n]}} \mathcal{G}(\lambda)$. Let G^d, H^d and H_0^d be the diagonal copies of G, H and H_0 in \mathcal{G} . Let \mathcal{H} be the subgraph of \mathcal{G} induced by $V(H) \times \dots \times V(H)$. Note that the constants c_1, \dots, c_n live in \mathcal{H} . Now build \mathcal{G}' from \mathcal{G} by augmenting a new copy of Cyl_m^{*+} for every vertex $v \in V(\mathcal{H}) \setminus V(H_0^d)$. Vertex v is to be identified with the vertex x that is at the end of the out-edge pendant on the top copy of DC_m^* in Cyl_m^{*+} and the bottom copy of DC_m^* is to be identified with HC_0 in H_0^d according to the identity function.

Now build \mathcal{G}'' by adding an edge from each vertex c_i to a new vertex d_i (for each $i \in [n]$).

Finally, build φ from the canonical query of \mathcal{G}'' , where we additionally turn the vertices d_1, \dots, d_n to outermost universal variables z_1, \dots, z_n . Then existentially quantify all remaining constants and vertices innermost. Finally, restrict all except the universal variables to be in $V(H)$.

First suppose that G retracts to H by r . Let λ' be some assignment of the universal variables z_1, \dots, z_n of φ to H^+ and choose y_1, \dots, y_n backwards-adjacent to these in H , mapped by λ .

To prove $\varphi \in \text{QCSP}(H^+)$ it suffices to prove that there is a homomorphism from \mathcal{G}' to H^+ that extends λ and for this it suffices to prove that there is a homomorphism h from \mathcal{G} to H that extends λ . Let us explain why. By assumption, we have $\text{Spill}_m^+(H[H_0, HC_0]) = V(H)$.

Henceforth let us consider the homomorphic image of \mathcal{G} that is $\mathcal{G}(\lambda)$. To prove $\varphi \in \text{QCSP}(H^+)$ it suffices to prove that there is a homomorphism from $G(\lambda)$ to H that extends λ . Note that it will be sufficient to prove that G retracts to H . We are done.

Suppose now $\varphi \in \text{QCSP}(H^+)$. Choose some surjection for λ' mapping z_1, \dots, z_n to H . Choose some y_1, \dots, y_n backwards-adjacent to these (and therefore in H) and let this be the map λ . Note that it is not possible for all y_1, \dots, y_n to be evaluated as a single vertex as H is strongly connected. Recall $N = |V(H)^{[n]}|$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H)$ induces a non-trivial homomorphism s from \mathcal{G}' to H which contains within it a non-trivial homomorphism s' from $\mathcal{H} = H^N$ to H . Consider the diagonal copy of $H_0^d \subset H^d \subset G^d$ in \mathcal{G} . By abuse of notation we will also consider each of s and s' acting just on the diagonal. If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}'' with the Cyl_m^{*+} , we have $s'(H^d)$ is an in-star, but this is not possible as H^d is strongly connected. As $1 < |s'(H_0^d)| < m$ is not possible either due to Lemma 2.7, we find that $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy of itself in H which we will call $H'_0 = i(H_0^d)$ for some isomorphism i .

As (H, H_0) is endo-trivial, Lemma 3.4 tells us that the restriction of s' to H^d is an automorphism of H^d , which we call α . The required retraction from G to H is now given by $\alpha^{-1} \circ s'$. \square

It remains to generalise these base cases.

PROPOSITION 3.13 (GENERAL CASE A-I). *Let H_{k+1}^+ be some reflexive tournament that has an initial strongly connected component H_{k+1} . Let $H_0, H_1, \dots, H_k, H_{k+1}$ be reflexive tournaments, the first k of which have Hamilton cycles HC_0, HC_1, \dots, HC_k , respectively, so that $H_0 \subseteq H_1 \subseteq \dots \subseteq H_k \subseteq H_{k+1}$.*

785 Assume that $H_0, (H_1, H_0), \dots, (H_k, H_{k-1})$ are endo-trivial and that

$$\begin{aligned}
 786 \quad & \text{Spill}_{a_0}^+(H_1[H_0, HC_0]) &= & V(H_1) \\
 787 \quad & \text{Spill}_{a_1}^+(H_2[H_1, HC_1]) &= & V(H_2) \\
 788 \quad & \vdots & & \vdots \\
 789 \quad & \vdots & & \vdots \\
 790 \quad & \text{Spill}_{a_{k-1}}^+(H_k[H_{k-1}, HC_{k-1}]) &= & V(H_k). \\
 791
 \end{aligned}$$

792 Moreover, assume that H_{k+1} retracts to H_k and also to every isomorphic copy $H'_k = i(H_k)$ of H_k in H_{k+1}
 793 with $\text{Spill}_{a_k}^+(H_{k+1}[H'_k, i(HC_k)]) = V(H_{k+1})$. Then H_k -RETRACTION can be polynomially reduced to
 794 $\text{QCSP}(H_{k+1}^+)$.
 795

796 PROOF. Let $n = a_{k+1} = |V(H_{k+1})|$ and let a_k, \dots, a_0 be the cardinalities of $|V(H_k)|, \dots, |V(H_0)|$,
 797 respectively. Let G be an instance of H_k -RETRACTION. We will build an instance φ of $\text{QCSP}(H_{k+1}^+)$
 798 in the following fashion. First, take a copy of H_{k+1} together with G and build G' by identifying
 799 these on the copy of H_k that they both possess as an induced subgraph.

800 Consider all possible functions $\lambda : [n] \rightarrow V(H_{k+1})$. For some such λ , let $\mathcal{G}'(\lambda)$ be the graph en-
 801 riched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H_{k+1})$ according
 802 to λ in the natural way (acting on the subscripts).

803 Let $\mathcal{G}' = \bigotimes_{\lambda \in V(H_{k+1})^{[n]}} \mathcal{G}'(\lambda)$. Let G'^d, H_{k+1}^d and H_k^d etc. be the diagonal copies of G', H_{k+1} and
 804 H_k in \mathcal{G}' . Let \mathcal{H}_{k+1} be the subgraph of \mathcal{G}' induced by $V(H_{k+1}) \times \dots \times V(H_{k+1})$. Note that the
 805 constants c_1, \dots, c_n live in \mathcal{H}_{k+1} .

806 Now build \mathcal{G}'' from \mathcal{G}' by augmenting a new copy of $\text{Cyl}_{a_k}^{**}$ for every vertex $v \in V(\mathcal{H}_{k+1}) \setminus V(H_k^d)$.
 807 Vertex v is to be identified with the vertex x that is at the end of the out-edge pendant on the top
 808 copy of DC_{a_k} in $\text{Cyl}_{a_k}^{**}$ and the bottom copy of DC_{a_k} is to be identified with HC_k in H_k^d according
 809 to the identity function. Call these *the $\text{Cyl}_{a_k}^{**}$ of the second stage*. Then, for each $i \in [k]$, and
 810 $v \in V(H_i^d) \setminus V(H_{i-1}^d)$, add a copy of $\text{Cyl}_{a_{i-1}}^{**}$, where v is identified with the vertex x that is at the
 811 end of the out-edge pendant on the top copy of $\text{DC}_{a_{i-1}}^*$ in $\text{Cyl}_{a_{i-1}}^{**}$ and the bottom copy of $\text{DC}_{a_{i-1}}^*$ is
 812 to be identified with H_{i-1} according to the identity map of $\text{DC}_{a_{i-1}}^*$ to HC_{i-1} .

813 Now build \mathcal{G}''' by adding an edge from each vertex c_i to a new vertex d_i (for each $i \in [n]$). Now
 814 add a copy of $\text{Cyl}_{a_k}^{**}$ for every vertex $v \in \{d_1, \dots, d_n\}$. Vertex v is to be identified with the vertex x
 815 that is at the end of the out-edge pendant on the top copy of DC_{a_k} in $\text{Cyl}_{a_k}^{**}$ and the bottom copy
 816 of DC_{a_k} is to be identified with HC_k in H_k^d according to the identity function. Call these *the $\text{Cyl}_{a_k}^{**}$*
 817 *of the third stage*.
 818

819 Finally, build φ from the canonical query of \mathcal{G}''' , where we additionally turn the vertices d_1, \dots, d_n
 820 to outermost universal variables z_1, \dots, z_n . Then existentially quantify all remaining constants and
 821 vertices innermost. Finally, restrict all except the universal variables to be in $V(H)$.

822 First suppose that G retracts to H_k by r . Let λ' be some assignment of the universal variables
 823 z_1, \dots, z_n of φ to H_{k+1}^+ and choose y_1, \dots, y_n backwards-adjacent to these in H_{k+1} , mapped by λ . To
 824 prove $\varphi \in \text{QCSP}(H_{k+1}^+)$ it suffices to prove that there is a homomorphism from \mathcal{G}'' to H_{k+1}^+ that
 825 extends λ and for this it suffices to prove that there is a homomorphism h from \mathcal{G}' that extends
 826 λ . Let us explain why. Because H_{k+1} retracts to H_k , we have $\text{Spill}_{a_k}^+(H_{k+1}[H_k, \text{HC}_k]) = V(H_{k+1})$
 827 due to Lemma 3.2 which implies the weaker $\text{Spill}_{a_k}^+(H_{k+1}[H_k, \text{HC}_k]) = V(H_{k+1})$. For the $\text{Cyl}_{a_k}^{**}$ of
 828 the second stage, the weaker statement suffices, but for the $\text{Cyl}_{a_k}^{**}$ of the third stage, the stronger
 829 statement is needed. We continue mapping now the various copies of $\text{Cyl}_{a_{i-1}}^{**}$ in G'' in any suitable
 830 fashion, which will always exist due to our assumptions.

831 Henceforth let us consider the homomorphic image of \mathcal{G}' that is $\mathcal{G}'(\lambda)$. To prove $\varphi \in \text{QCSP}(H_{k+1}^+)$
 832 it suffices to prove that there is a homomorphism from $G'(\lambda)$ to H_{k+1} that extends λ . Note that it
 833

will be sufficient to prove that G' retracts to H_{k+1} . Let h be the natural retraction from G' to H_{k+1} that extends the known retraction r from G to H_k . We are done.

Suppose now $\varphi \in \text{QCSP}(H_{k+1}^+)$. Choose some surjection for λ , the assignment of the universal variables of φ to H_{k+1} . Choose some y_1, \dots, y_n backwards-adjacent to these (and therefore in H_{k+1}) and let this be the map λ . Note that it is not possible for all y_1, \dots, y_n to be evaluated as a single vertex as H_{k+1} is strongly connected. Let $N = |V(H_{k+1})|^{[n]}$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H_{k+1}^+)$ induces a non-trivial homomorphism s from \mathcal{G}' to H_{k+1} which contains within it a non-trivial homomorphism s' from $\mathcal{H} = H_{k+1}^N$ to H_{k+1} . Consider the diagonal copy of $H_0^d \subset \dots \subset H_k^d \subset H_{k+1}^d \subset G'^d$ in \mathcal{G}' . By abuse of notation we will also consider each of s and s' acting just on the diagonal.

If $|s'(H_0^d)| = 1$, by construction of \mathcal{G}'' , we have that $s'(H_1^d)$ is either an in-star or a loop, but the former is not possible as H_1^d is strongly connected. Iterating this argument we find that $|s'(H_{k+1}^d)| = 1$, but this would mean s' is uniformly mapping \mathcal{H}_{k+1} to one vertex, which is impossible as s' is non-trivial. As $1 < |s'(H_0^d)| < m$ is not possible either due to Lemma 2.7, we find that $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy of itself in H which we will call $H'_0 = i(H_0^d)$ for some isomorphism i .

We now apply Lemma 3.6 as well as our assumed endo-trivialities to derive that s' in fact maps H_k^d by the isomorphism i to a copy of itself in H_{k+1} which we will call H'_k .

We claim that $\text{Spill}_{a_k}^+(H_{k+1}[H'_{k+1}, i(\text{HC}_{a_k}^d)]) = V(H_{k+1})$. Since λ' is surjective on H_{k+1}^+ , this is enforced explicitly by the $\text{Cyl}_{a_k}^{*+}$ of the third stage. Thus, there exists a retraction r from H_{k+1} to H'_k . Now $i^{-1} \circ r \circ s'$ gives the desired retraction of G to H_k . \square

PROPOSITION 3.14 (GENERAL CASE A-II). *Let H_{k+1}^+ be some reflexive tournament that has an initial strongly connected component H_{k+1} that is non-trivial. Let $H_0, H_1, \dots, H_k, H_{k+1}$ be reflexive tournaments, the first $k+1$ of which have Hamilton cycles $\text{HC}_0, \text{HC}_1, \dots, \text{HC}_k$, respectively, so that $H_0 \subseteq H_1 \subseteq \dots \subseteq H_k \subseteq H_{k+1}$. Suppose that $H_0, (H_1, H_0), \dots, (H_k, H_{k-1}), (H_{k+1}, H_k)$ are endo-trivial and that*

$$\begin{aligned} \text{Spill}_{a_0}^+(H_1[H_0, \text{HC}_0]) &= V(H_1) \\ \text{Spill}_{a_1}^+(H_2[H_1, \text{HC}_1]) &= V(H_2) \\ &\vdots \\ \text{Spill}_{a_{k-1}}^+(H_k[H_{k-1}, \text{HC}_{k-1}]) &= V(H_k) \\ \text{Spill}_{a_k}^+(H_{k+1}[H_k, \text{HC}_k]) &= V(H_{k+1}) \end{aligned}$$

Then H_{k+1} -RETRACTION can be polynomially reduced to $\text{QCSP}(H_{k+1}^+)$.

PROOF. Let $n = a_{k+1} = |V(H_{k+1})|$ and let a_k, \dots, a_0 be the cardinalities of $|V(H_k)|, \dots, |V(H_0)|$, respectively. Let G be an instance of H_{k+1} -RETRACTION. We build an instance φ of $\text{QCSP}(H_{k+1}^+)$ in the following fashion. Consider all possible functions $\lambda : [n] \rightarrow V(H_{k+1})$. For some such λ , let $\mathcal{G}(\lambda)$ be the graph enriched with constants c_1, \dots, c_n where these are interpreted over some subset of $V(H_{k+1})$ according to λ in the natural way (acting on the subscripts).

Let $\mathcal{G} = \bigotimes_{\lambda \in V(H_{k+1})^{[n]}} \mathcal{G}(\lambda)$. Let $G^d, H_{k+1}^d, H_k^d, \dots, H_0^d$ be the diagonal copies of $G, H_{k+1}, H_k, \dots, H_0$ in \mathcal{G} . Let \mathcal{H}_{k+1} be the subgraph of \mathcal{G} induced by $V(H_{k+1}) \times \dots \times V(H_{k+1})$. Note that the constants c_1, \dots, c_n live in \mathcal{H}_{k+1} .

Now build \mathcal{G}' from \mathcal{G} by the following procedure. For each $i \in [k+1]$, and $v \in V(H_i^d) \setminus V(H_{i-1}^d)$, add a copy of $\text{Cyl}_{a_{i-1}}^{*+}$, where v is identified with the vertex x that is at the end of the out-edge pendant on the top copy of $\text{DC}_{a_{i-1}}^*$ in $\text{Cyl}_{a_{i-1}}^{*+}$ and the bottom copy of $\text{DC}_{a_{i-1}}^*$ is to be identified with H_{i-1} according to the identity map of $\text{DC}_{a_{i-1}}^*$ to HC_{i-1} .

Now build \mathcal{G}'' by adding an edge from each vertex c_i to a new vertex d_i (for each $i \in [n]$).

883 Finally, build φ from the canonical query of \mathcal{G}'' , where we additionally turn the vertices d_1, \dots, d_n
 884 to outermost universal variables z_1, \dots, z_n . Then existentially quantify all remaining constants and
 885 vertices innermost. Finally, restrict all except the universal variables to be in $V(H_{k+1})$.

886 First suppose that G retracts to H_{k+1} by r . Let λ' be some assignment of the universal variables
 887 z_1, \dots, z_n of φ to H_{k+1}^+ and choose y_1, \dots, y_n backwards-adjacent to these in H_{k+1} , mapped by λ . To
 888 prove $\varphi \in \text{QCSP}(H_{k+1}^+)$ it suffices to prove that there is a homomorphism from \mathcal{G}' to H_{k+1}^+ that
 889 extends λ and for this it suffices to prove that there is a homomorphism h from \mathcal{G} that extends λ .
 890 The extension of the latter to the former will always be possible due to the spill assumptions.

891 Henceforth let us consider the homomorphic image of \mathcal{G} that is $\mathcal{G}(\lambda)$. To prove $\varphi \in \text{QCSP}(H_{k+1}^+)$
 892 it suffices to prove that there is a homomorphism from $\mathcal{G}(\lambda)$ to H_{k+1} that extends λ . Note that it
 893 will be sufficient to prove that G retracts to H_{k+1} . Well this was our original assumption so we are
 894 done.

895 Suppose now $\varphi \in \text{QCSP}(H_{k+1}^+)$. Choose some surjection for λ' mapping z_1, \dots, z_n to H_{k+1} . Choose
 896 some y_1, \dots, y_n backwards-adjacent to these (and therefore in H_{k+1}) and let this be the map λ . Note
 897 that it is not possible for all y_1, \dots, y_n to be evaluated as a single vertex as H_{k+1} is strongly connected.
 898 Recall $N = |V(H)^{[n]}|$. The evaluation of the existential variables that witness $\varphi \in \text{QCSP}(H_{k+1}^+)$
 899 induces a non-trivial homomorphism s from \mathcal{G} to H_{k+1} which contains within it a non-trivial
 900 homomorphism s' from $\mathcal{H}_{k+1} = H_{k+1}^N$ to H_{k+1} . Consider the diagonal copy of $H_0^d \subset H_1^d \subset \dots \subset H_{k+1}^d$
 901 in \mathcal{G} . By abuse of notation we will also consider each of s and s' acting just on the diagonal.

902 If $|s'(H_0^d)| = 1$ we deduce that $s'(H_1^d)$ is either an in-star or a loop, but the former is not possible as
 903 H_1^d is strongly connected. Iterating this argument we find that $|s'(H_{k+1}^d)| = 1$, but this would mean
 904 s' is uniformly mapping to one vertex, which is impossible as s' is non-trivial. As $1 < |s'(H_0^d)| < m$
 905 is not possible either due to Lemma 2.7, we find that $|s'(H_0^d)| = m$, and indeed s' maps H_0^d to a copy
 906 of itself in H which we will call $H'_0 = i(H_0^d)$ for some isomorphism i .

907 We now apply Lemma 3.6 as well as our assumed endo-trivialities to derive that s' in fact maps
 908 H_k^d by the isomorphism i to a copy of itself in H_{k+1} , which we will call H'_k . Now we can deduce,
 909 via Lemma 3.4, that $h(H_{k+1}^d)$ is an automorphism of H_{k+1} , which we call α . The required retraction
 910 from G to H_{k+1} is now given by $\alpha^{-1} \circ s'$. \square

912 The proof of the following is exactly as that for Corollary 3.9 modulo Spill becoming Spill⁺.

913
 914 **COROLLARY 3.15.** *Let H be a reflexive tournament with an initial strongly connected component*
 915 *that is non-trivial. Then $\text{QCSP}(H)$ is NP-hard.*

916 4 THE PROOF OF THE NL CASES OF THE DICHOTOMY

917 A particular role in the tractable part of our dichotomy will be played by TT_2^* , the reflexive transitive
 918 2-tournament, which has vertex set $\{0, 1\}$ and edge set $\{(0, 0), (0, 1), (1, 1)\}$.

919
 920 **LEMMA 4.1.** *Let $H = H_1 \Rightarrow \dots \Rightarrow H_n$ be a reflexive tournament on $m+2$ vertices with $V(H_1) = \{s\}$*
 921 *and $V(H_n) = \{t\}$. Then there exists a surjective homomorphism from $(\text{TT}_2^*)^m$ to H .*

922
 923 **PROOF.** Build a surjective homomorphism f from $(\text{TT}_2^*)^m$ to H in the following fashion. Let \bar{x}_i
 924 be the m -tuple which has 1 in the i th position and 0 in all other positions. For $i \in [m]$, let f map \bar{x}_i
 925 to i . Let f map $(0, \dots, 0)$ to s and everything remaining to t .

926 By construction, f is surjective. To see that f is a homomorphism, let $((y_1, \dots, y_m), (z_1, \dots, z_m)) \in$
 927 $E((\text{TT}_2^*)^m)$, which is the case exactly when $y_i \leq z_i$ for all $i \in [m]$. Let $f(y_1, \dots, y_m) = u$ and
 928 $f(z_1, \dots, z_m) = v$. First suppose that y_1, \dots, y_m are all 0. Then $u = s$. As s has an out-edge to
 929 every vertex of H , we find that $(u, v) \in E(H)$. Now suppose that y_1, \dots, y_m contains a single 1. If
 930 $(y_1, \dots, y_m) = (z_1, \dots, z_m)$, then $u = v$. As H is reflexive, we find that $(u, v) \in H$. If $(y_1, \dots, y_m) \neq$
 931

(z_1, \dots, z_m), then $v = t$. As t has an in-edge from every vertex of H , we find that $(u, v) \in E(H)$. Finally suppose that y_1, \dots, y_m contains more than one 1. Then $u = v = t$. As H is reflexive, we find that $(u, v) \in E(H)$. \square

We also need the following lemma, which follows from combining some known results.

LEMMA 4.2. *If H is a transitive reflexive tournament then $\text{QCSP}(H)$ is in NL.*

PROOF. It is noted in [15] that H has the ternary median operation as a polymorphism. It follows from well-known results (e.g. in [7, 9]) that $\text{QCSP}(H)$ is in NL. Specifically, one can apply Theorem 5.16 from [7] to reduce $\text{QCSP}(H)$ to an ensemble of instances of $\text{CSP}(H)$, which may also reference constants, each of which can be solved in NL by Corollary 4 from [9]. Each of these instances may be solved independently and the ensemble is polynomial in number, hence the whole procedure can be accomplished in NL. \square

The other tractable cases are more interesting.

We are now ready to prove the main result of this section.

THEOREM 4.3. *Let $H = H_1 \Rightarrow \dots \Rightarrow H_n$ be a reflexive tournament. If $|V(H_1)| = |V(H_n)| = 1$, then $\text{QCSP}(H)$ is in NL.*

PROOF. Let $|V(H)| = m + 2$ for some $m \geq 0$. By Lemma 4.1, there exists a surjective homomorphism from $(\text{TT}_2^*)^m$ to H . There exists also a surjective homomorphism from H to TT_2^* ; we map s to 0 and all other vertices of H to 1. It follows from Theorem 3.4 in [8] that $\text{QCSP}(H) = \text{QCSP}(\text{TT}_2^*)$ meaning we may consider the latter problem. We note that TT_2^* is a transitive reflexive tournament. Hence, we may apply Lemma 4.2. \square

5 FINAL RESULT AND REMARKS

We are now in a position to prove our main dichotomy theorem.

THEOREM 5.1. *Let $H = H_1 \Rightarrow \dots \Rightarrow H_n$ be a reflexive tournament. If $|V(H_1)| = |V(H_n)| = 1$, then $\text{QCSP}(H)$ is in NL; otherwise it is NP-hard.*

PROOF. The NL case follow from Theorem 4.3. The NP-hard cases follow from Corollary 3.9 and Corollary 3.15, bearing in mind the case with a non-trivial final strongly connected component is dual to the case with a non-trivial initial strongly connected component (map edges (x, y) to (y, x)). \square

Theorem 5.1 resolved the open case in Table 1. It is difficult to position this result in the overall classification program for finite-domain QCSPs save to say that our methods are tailored, indeed specialised, to reflexive tournaments. It is not clear that they can be applied easily to different or wider classes (in this vein we return to mixed-type tournaments below). Since complexities outside of P, NP-complete and Pspace-complete were discovered for QCSPs in [25], for example co-NP-complete, DP-complete and Θ_2^P , the whole classification task has been thrown wide open. Classes such as that of reflexive tournaments might provide comfort, as it is doubtful such monstrous complexities could be found here. Though, we cannot be sure, with our lacuna between NP-hard and Pspace-complete.

Recall that the results for the irreflexive tournaments in this table were all proven in a more general setting, namely for irreflexive semicomplete graphs. One natural direction for future research is to determine a complexity dichotomy for QCSP and SCSP for reflexive semicomplete graphs. We leave this as an interesting open direction.

The task of promoting our NP-hardness results to Pspace-complete, while using the same method, seems to require corresponding Pspace-hardness results for reflexive tournaments with constants.

981 If $\text{QCSP}^c(H)$ were Pspace-complete, for H a non-trivial reflexive strongly connected tournament,
 982 then likely our NP-hardness results, for the similar class of graphs, would easily rise to Pspace-
 983 complete. The cases that are not strongly connected require additional arguments, and perhaps
 984 even a different method.

985 Mixed-type tournaments, where some vertices are reflexive and others irreflexive, are well-
 986 understood algebraically [21]. Indeed, from this paper there follows a complexity dichotomy for
 987 $\text{CSP}^c(H)$ where H is a mixed-type tournament. Furthermore, $\text{CSP}(H)$ is either trivial or H is an
 988 irreflexive tournament, so the complexity dichotomy for $\text{CSP}(H)$ is also known. Though many of
 989 our supporting lemmas hold for mixed-type tournaments, some do not. For example, Lemma 2.1
 990 fails for the transitive 2-tournament TT_2 in which one vertex is a self-loop and the other is not. To
 991 extend our classification to mixed-type tournaments thus requires still some work.

992

993 ACKNOWLEDGEMENTS

994 We are grateful to several referees for careful reading of the paper and good advice.
 995

996

997 REFERENCES

- 998 [1] Jørgen Bang-Jensen, Pavol Hell, and Gary MacGillivray. The complexity of colouring by semicomplete digraphs. *SIAM*
 999 *Journal on Discrete Mathematics*, 1(3):281–298, 1988.
- 1000 [2] Manuel Bodirsky, Jan Kára, and Barnaby Martin. The complexity of surjective homomorphism problems - a survey.
 1001 *Discrete Applied Mathematics*, 160(12):1680–1690, 2012.
- 1002 [3] Ferdinand Börner, Andrei A. Bulatov, Hubie Chen, Peter Jeavons, and Andrei A. Krokhin. The complexity of constraint
 1003 satisfaction games and QCSP. *Inf. Comput.*, 207(9):923–944, 2009.
- 1004 [4] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In *58th IEEE Annual Symposium on Foundations of*
 1005 *Computer Science, FOCS 2017, Berkeley, CA, USA, October 15–17, 2017*, pages 319–330, 2017.
- 1006 [5] Andrei A. Bulatov, Peter Jeavons, and Andrei A. Krokhin. Classifying the complexity of constraints using finite
 1007 algebras. *SIAM Journal on Computing*, 34(3):720–742, 2005.
- 1008 [6] Paul Camion. Chemins et circuits hamiltoniens de graphes complets. *Comptes Rendus de l'Académie des Sciences Paris*,
 1009 249:2151–2152, 1959.
- 1010 [7] Hubie Chen. The complexity of quantified constraint satisfaction: Collapsibility, sink algebras, and the three-element
 1011 case. *SIAM J. Comput.*, 37(5):1674–1701, 2008. doi:<http://dx.doi.org/10.1137/060668572>.
- 1012 [8] Hubie Chen, Florent R. Madelaine, and Barnaby Martin. Quantified constraints and containment problems. *Logical*
 1013 *Methods in Computer Science*, 11(3), 2015. Extended abstract appeared at LICS 2008. This journal version incorporates
 1014 principal part of CP 2012 *Containment, Equivalence and Coreness from CSP to QCSP and Beyond*. URL: [http://dx.doi.org/10.2168/LMCS-11\(3:9\)2015](http://dx.doi.org/10.2168/LMCS-11(3:9)2015), doi:10.2168/LMCS-11(3:9)2015.
- 1015 [9] Víctor Dalmau and Andrei A. Krokhin. Majority constraints have bounded pathwidth duality. *European Journal of*
 1016 *Combinatorics*, 29(4):821–837, 2008.
- 1017 [10] Petar Dapic, Petar Markovic, and Barnaby Martin. Quantified constraint satisfaction problem on semicomplete
 1018 digraphs. *ACM Trans. Comput. Log.*, 18(1):2:1–2:47, 2017. doi:10.1145/3007899.
- 1019 [11] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction:
 1020 A study through datalog and group theory. *SIAM Journal on Computing*, 28(1):57–104, 1998.
- 1021 [12] Petr A. Golovach, Daniël Paulusma, and Jian Song. Computing vertex-surjective homomorphisms to partially reflexive
 1022 trees. *Theoretical Computer Science*, 457:86–100, 2012.
- 1023 [13] P. G. Kolaitis and M. Y. Vardi. *Finite Model Theory and Its Applications (Texts in Theoretical Computer Science. An EATCS*
 1024 *Series)*, A logical Approach to Constraint Satisfaction. Springer-Verlag New York, Inc., 2005.
- 1025 [14] Benoit Larose. Taylor operations on finite reflexive structures. *International Journal of Mathematics and Computer*
 1026 *Science*, 1(1):1–26, 2006.
- 1027 [15] Benoit Larose, Barnaby Martin, and Daniël Paulusma. Surjective H-colouring over reflexive digraphs. *TOCT*, 11(1):3:1–
 1028 3:21, 2019. doi:10.1145/3282431.
- 1029 [16] Benoit Larose and László Zádori. Finite posets and topological spaces in locally finite varieties. *Algebra Universalis*,
 52(2):119–136, 2005.
- [17] Barnaby Martin. QCSP on partially reflexive forests. In *Principles and Practice of Constraint Programming - CP*
 2011 - 17th International Conference, CP 2011, Perugia, Italy, September 12–16, 2011. *Proceedings*, pages 546–560, 2011.
 doi:10.1007/978-3-642-23786-7_42.

- 1030 [18] Barnaby Martin and Florent R. Madelaine. Towards a trichotomy for quantified H -coloring. In *Logical Approaches to*
1031 *Computational Barriers, Second Conference on Computability in Europe, CiE 2006, Swansea, UK, June 30-July 5, 2006,*
1032 *Proceedings*, pages 342–352, 2006. doi:10.1007/11780342_36.
- 1033 [19] Narayan Vikas. Algorithms for partition of some class of graphs under compaction and vertex-compaction. *Algorith-*
1034 *mica*, 67(2):180–206, 2013.
- 1035 [20] Narayan Vikas. Computational complexity of graph partition under vertex-compaction to an irreflexive hexagon.
1036 In *42nd International Symposium on Mathematical Foundations of Computer Science, MFCS 2017, August 21-25, 2017 -*
Aalborg, Denmark, pages 69:1–69:14, 2017.
- 1037 [21] Alexander Wires. Dichotomy for finite tournaments of mixed-type. *Discrete Mathematics*, 338(12):2523–2538, 2015.
1038 doi:10.1016/j.disc.2015.06.024.
- 1039 [22] Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In *58th IEEE Annual Symposium on Foundations of Computer*
1040 *Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017*, pages 331–342, 2017.
- 1041 [23] Dmitriy Zhuk. No-rainbow problem and the surjective constraint satisfaction problem, 2020. To appear LICS 2021.
1042 arXiv:2003.11764.
- 1043 [24] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. *J. ACM*, 67(5):30:1–30:78, 2020. doi:10.1145/3402029.
- 1044 [25] Dmitriy Zhuk and Barnaby Martin. QCSP monsters and the demise of the Chen Conjecture. In Konstantin Makarychev,
1045 Yury Makarychev, Madhur Tulsiani, Gautam Kamath, and Julia Chuzhoy, editors, *Proceedings of the 52nd Annual ACM*
SIGACT Symposium on Theory of Computing, STOC 2020, Chicago, IL, USA, June 22-26, 2020, pages 91–104. ACM, 2020.
doi:10.1145/3357713.3384232.

1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078