



PHARMACOLOGY 2019

15–17 December | Edinburgh

SUBMIT AN ABSTRACT

- Participate in the UK's leading pharmacology event
- Share your research with over 1,200 attendees
- Apply for awards and attendance bursaries
- Have your work published in the British Journal of Pharmacology or the British Journal of Clinical Pharmacology



**SUBMIT
NOW**



Deadline to submit
9 September




BRITISH
PHARMACOLOGICAL
SOCIETY



@BritPharmSoc #Pharmacology2019

RESEARCH PAPER

Endothelial TRPV4 channels modulate vascular tone by Ca²⁺-induced Ca²⁺ release at inositol 1,4,5-trisphosphate receptors

Helen R. Heathcote¹ | Matthew D. Lee¹ | Xun Zhang¹ | Christopher D. Saunter² | Calum Wilson¹ | John G. McCarron¹ 

¹Strathclyde Institute of Pharmacy and Biomedical Science, University of Strathclyde, Glasgow, UK

²Centre for Advanced Instrumentation, Biophysical Sciences Institute, Department of Physics, Durham University, Durham, UK

Correspondence

John G. McCarron, Strathclyde Institute of Pharmacy and Biomedical Science, University of Strathclyde, 161 Cathedral Street, Glasgow G4 0RE, UK.
Email: john.mccarron@strath.ac.uk

Funding information

British Heart Foundation, Grant/Award Numbers: PG/16/54/32230 and PG16/82/32439; Wellcome Trust, Grant/Award Numbers: 202924/Z/16/Z and 204682/Z/16/Z

Background and Purpose: The TRPV4 ion channels are Ca²⁺ permeable, non-selective cation channels that mediate large, but highly localized, Ca²⁺ signals in the endothelium. The mechanisms that permit highly localized Ca²⁺ changes to evoke cell-wide activity are incompletely understood. Here, we tested the hypothesis that TRPV4-mediated Ca²⁺ influx activates Ca²⁺ release from internal Ca²⁺ stores to generate widespread effects.

Experimental Approach: Ca²⁺ signals in large numbers (~100) of endothelial cells in intact arteries were imaged and analysed separately.

Key Results: Responses to the TRPV4 channel agonist GSK1016790A were heterogeneous across the endothelium. In activated cells, Ca²⁺ responses comprised localized Ca²⁺ changes leading to slow, persistent, global increases in Ca²⁺ followed by large propagating Ca²⁺ waves that moved within and between cells. To examine the mechanisms underlying each component, we developed methods to separate slow persistent Ca²⁺ rise from the propagating Ca²⁺ waves in each cell. TRPV4-mediated Ca²⁺ entry was required for the slow persistent global rise and propagating Ca²⁺ signals. The propagating waves were inhibited by depleting internal Ca²⁺ stores, inhibiting PLC or blocking IP₃ receptors. Ca²⁺ release from stores was tightly controlled by TRPV4-mediated Ca²⁺ influx and ceased when influx was terminated. Furthermore, Ca²⁺ release from internal stores was essential for TRPV4-mediated control of vascular tone.

Conclusions and Implications: Ca²⁺ influx via TRPV4 channels is amplified by Ca²⁺-induced Ca²⁺ release acting at IP₃ receptors to generate propagating Ca²⁺ waves and provide a large-scale endothelial communication system. TRPV4-mediated control of vascular tone requires Ca²⁺ release from the internal store.

Abbreviations: 2-APB, 2-aminoethoxydiphenyl borate; CPA, cyclopiazonic acid; F, fluorescence intensity; F₀, baseline fluorescence intensity; GSK, GSK1016790A; HC067, HC067047; IP₃, inositol 1,4,5-trisphosphate; IP₃R, inositol 1,4,5-trisphosphate receptor; RuR, ruthenium red; RyRs, ryanodine receptors; SNP, sodium nitroprusside

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. British Journal of Pharmacology published by John Wiley & Sons Ltd on behalf of British Pharmacological Society.

1 | INTRODUCTION

The endothelium plays a critical role in the regulation of numerous vascular processes such as maintaining vascular tone, regulating the passage of macromolecules and oxygen to tissues, modulating immune responses, initiating angiogenesis, and controlling vascular remodelling. The control by the endothelium of each process is mediated by the generation of various signalling molecules that include NO, prostacyclin, von Willebrand factor, tissue plasminogen activator, and endothelial-derived hyperpolarizing and contracting factors (Taylor et al., 2003; Whorton, Willis, Kent, & Young, 1984). The generation of each of these signalling molecules is controlled tightly by changes in the cytoplasmic Ca^{2+} concentration in the endothelium. Central therefore to an understanding of endothelial function is an appreciation of the control of intracellular Ca^{2+} concentrations.

There are two major sources of Ca^{2+} for the endothelium: the extracellular space and the intracellular Ca^{2+} store. Ca^{2+} influx from the extracellular space is mediated mainly by channels present on the plasmalemmal membrane. In many cell types, such as smooth muscle and cardiomyocytes, the influx pathways are well characterized, and **voltage-dependent Ca^{2+} channels** are major contributors (Catterall, 2011; Nelson, Patlak, Worley, & Standen, 1990). In endothelial cells, however, understanding of the Ca^{2+} influx pathways is incomplete (Nilius, Droogmans, & Wondergem, 2003). Several members of the transient receptor potential (TRP) superfamily of cation channels are present in endothelial cells (Hofmann, Schaefer, Schultz, & Gudermann, 2002; Nilius, Droogmans, & Wondergem, 2003). Of these, the **TRPV4** channel has attracted substantial interest because of the channels' relatively high Ca^{2+} permeability (Filosa, Yao, & Rath, 2013; Watanabe et al., 2002). TRPV4 channels are expressed widely in endothelial cells (Mendoza et al., 2010; Sullivan, Francis, Pitts, Taylor, & Earley, 2012; Watanabe et al., 2002). TRPV4 channels were initially demonstrated to be activated by hypotonicity-induced cell swelling (Liedtke et al., 2000; Mizuno, Matsumoto, Imai, & Suzuki, 2003). Later studies revealed that the channel is also activated by various other stimuli including mechanical stimulation (Gao, Wu, & O'Neil, 2003; Kohler et al., 2006; Liedtke, Tobin, Bargmann, & Friedman, 2003), moderate heat, endogenous chemicals such as **anandamide**, **arachidonic acid**, and its epoxyeicosatrienoic acid metabolites, as well as by a number of exogenous chemical ligands (Nilius & Voets, 2013; Vriens et al., 2004; Watanabe et al., 2003) or UVB (Moore et al., 2013; Yin et al., 2008). In the endothelium, by regulating the cytoplasmic Ca^{2+} concentration, TRPV4 channels may contribute to the response to arachidonic acid metabolites, physical forces, and agonists (Bagher et al., 2012; Freichel et al., 2001; Hartmannsgruber et al., 2007; Kohler et al., 2006; Marrelli, O'Neil, Brown, & Bryan, 2007; Mendoza et al., 2010; Watanabe et al., 2002; Zhang et al., 2009; Zhang, Papadopoulos, & Hamel, 2013).

While large in amplitude, the Ca^{2+} rise evoked by activation of endothelial TRPV4 channels is reported to be highly localized and confined to within a few micrometres of the channels (Sonkusare et al., 2012; Sonkusare et al., 2014). These highly localized Ca^{2+} signals

What is already known

- TRPV4 channels are key Ca^{2+} permeable channels that control endothelial function.
- TRPV4 channels generate large but highly localized Ca^{2+} signals.

What this study adds

- Ca^{2+} -induced Ca^{2+} release from IP_3 -sensitive internal store amplifies local Ca^{2+} signals from TRPV4 channel activity.
- Ca^{2+} -induced Ca^{2+} release at IP_3 receptors mediates control of vascular contractility by endothelial TRPV4 channels.

What is the clinical significance

- Endothelial TRPV4 channels are involved in several disease conditions such as cancer progression and hypertension.
- Endothelial TRPV4-mediated Ca^{2+} -induced Ca^{2+} release at IP_3 receptors offers a new target for drug development.

evoke endothelial and smooth muscle hyperpolarization and vascular relaxation (F. Gao & Wang, 2010; Kohler et al., 2006; Ma et al., 2013; Sonkusare et al., 2012; D. X. Zhang et al., 2009). The highly localized nature of the Ca^{2+} signal provides tight spatial control of the cellular effectors activated, for example, **Ca^{2+} -activated K^+ channels** (Gao & Wang, 2010; Ma et al., 2013; Sonkusare et al., 2012). However, more widespread Ca^{2+} rises throughout the cytoplasm have been implicated in the regulation of endothelial structure, maintenance of the normal orientation of endothelial cells, the control and selectively of endothelial permeability, and the production of anti-thrombotic factors. TRPV4 channels have been proposed to play a role in each of these processes (Noren et al., 2016; Phuong et al., 2017; Thodeti et al., 2009; Thoppil et al., 2016). The question arises as to how activation of TRPV4 channels may lead to Ca^{2+} -dependent events throughout the cytoplasm if the increase in Ca^{2+} concentration arising from TRPV4 channel activity remains localized to within a few microns of the channel.

The second major source of Ca^{2+} in endothelial cells is the internal Ca^{2+} store. Ca^{2+} release from the internal store may occur via **ryanodine** receptors (**RyRs**) or IP_3 receptors (**IP_3 Rs**). While RyRs may be expressed in endothelial cells (Moccia, Berra-Romani, & Tanzi, 2012; Mumtaz, Burdyga, Borisova, Wray, & Burdyga, 2011; Rusko, Wang, & Vanbreemen, 1995), the functional role of the channels in the endothelium is not clear given that RyR activators such as **caffeine** do not increase cytoplasmic Ca^{2+} in endothelial cells (Borisova, Wray, Eisner, & Burdyga, 2009; Wilson et al., 2019; Wilson, Lee, & McCarron, 2016). Ca^{2+} release from the internal store in endothelial cells may be mediated mainly by IP_3 Rs (Mumtaz et al., 2011; Wilson

et al., 2019). Ca^{2+} release via IP_3R occurs in response to physical forces and circulating vasoactive substances and contributes to the control of several vascular activities such as vasorelaxation, endothelial permeability, and production of anti-thrombotic factors (Sun, Geyer, & Komarova, 2017).

IP_3Rs are themselves regulated by Ca^{2+} , in addition to IP_3 . Ca^{2+} released via IP_3Rs may induce a positive feedback Ca^{2+} -induced Ca^{2+} release process at neighbouring IP_3Rs . Indeed, Ca^{2+} signals at IP_3Rs begin as a localized Ca^{2+} blip at a single receptor ($\sim 1 \mu\text{m}$ spread) which expand by Ca^{2+} -induced Ca^{2+} release opening neighbouring IP_3Rs generating a Ca^{2+} puff ($\sim 4 \mu\text{m}$ spread). Puffs may further propagate via Ca^{2+} -induced Ca^{2+} release to form a transient global Ca^{2+} elevation throughout the cell or a Ca^{2+} "wave" (Berridge, 1997; Bootman, Niggli, Berridge, & Lipp, 1997). These waves may move through all or part of the cell (Bootman, Berridge, & Lipp, 1997). In some conditions that are not fully understood, IP_3 -evoked Ca^{2+} waves may also move between cells to provide a signalling system capable of coordinating the activity of many cells (Leybaert & Sanderson, 2012).

While Ca^{2+} influx and release are activated separately, they are not independent: Ca^{2+} influx can trigger release, and release can trigger influx (McCarron, Chalmers, Bradley, Macmillan, & Muir, 2006). In cardiomyocytes, Ca^{2+} influx via voltage-dependent Ca^{2+} channels may activate RyRs to evoke a large rise in Ca^{2+} , inducing cell contraction. In smooth muscle cells, TRPV4 channel activity may lead to Ca^{2+} -induced Ca^{2+} release acting at RyRs (Earley, Heppner, Nelson, & Brayden, 2005). In astrocytes, Ca^{2+} -induced Ca^{2+} release occurs in response to Ca^{2+} influx via TRPV4 channels but at IP_3Rs rather than at RyRs. This process amplifies and propagates the Ca^{2+} signal arising from TRPV4-mediated Ca^{2+} influx (Dunn, Hill-Eubanks, Liedtke, & Nelson, 2013). TRPV4-mediated Ca^{2+} influx may also lead to recruitment of IP_3Rs and Ca^{2+} -induced Ca^{2+} release in murine-derived cultured neuronal cells (Shen et al., 2018). These observations raise the possibility of TRPV4-mediated Ca^{2+} influx being able to generate a more global increase in intracellular Ca^{2+} via Ca^{2+} -induced Ca^{2+} release in endothelial cells.

To explore this possibility, we examined the role that the internal store plays in regulating alterations in intracellular Ca^{2+} evoked by activation of TRPV4 channels. Here, we report that the IP_3 -sensitive Ca^{2+} store is critical for the TRPV4-evoked global Ca^{2+} rise in endothelial cells in intact blood vessels. Ca^{2+} influx generates Ca^{2+} -induced Ca^{2+} release at IP_3Rs to evoke Ca^{2+} waves in the vascular endothelium. IP_3R -dependent Ca^{2+} waves are required for the endothelium-dependent vascular smooth muscle cell relaxation in response to activation of TRPV4 channels.

2 | METHODS

2.1 | Animals

All animal care and experimental procedures complied with the relevant UK Home Office Regulations, (Schedule 1 of the Animals [Scientific Procedures] Act 1986, UK) and were approved by the

University of Strathclyde Animal Welfare and Ethical Review Body. Animal studies are reported in compliance with the ARRIVE guidelines (Kilkenny et al., 2010) and with the recommendations made by the *British Journal of Pharmacology*. The Strathclyde Biological Protection Unit is a conventional unit which undertakes FELASA quarterly health monitoring. Male Sprague–Dawley rats (10–12 week old; 250–300 g), from an in-house colony, were used for the study. Sprague–Dawley rats are a widely used experimental model with a wealth of background information to aid interpretation of results. The animals were housed three per cage, and the cage type was North Kent Plastic model RC2F with nesting material "Sizzle Nest." A 12:12 light dark cycle was used with a temperature range of 19–23°C (set point 21°C) and humidity levels between 45% and 65%. Animals had free access to fresh water and SDS diet RM1 (rodent maintenance). The enrichment in the cages was aspen wood chew sticks and hanging huts.

All experiments used first- or second-order mesenteric arteries isolated from rats killed by either CO_2 or by injection with pentobarbital; no differences in the results were observed with either method. Controls and experimental treatments were carried out in the same tissue, so blinding and randomization were not used.

2.2 | High-resolution imaging of endothelial Ca^{2+} signalling

Arteries (diameter of 200–230 μm) were isolated from the mesenteric bed, cleaned, then cut and pinned flat *en face* (endothelial side facing up) on a Sylgard block. The endothelium was then preferentially loaded with Cal-520/AM (5 μM ; with 0.02% Pluronic F-127) in physiological saline solution (PSS) at 37°C for 30 min (Wilson, Lee, & McCarron, 2016; Wilson, Saunter, Girkin, & McCarron, 2016). Following incubation, arteries were gently washed with PSS, and the Sylgard blocks were placed face down on a custom flow chamber. Arteries were then continuously perfused with PSS at 1.5 $\text{ml}\cdot\text{min}^{-1}$ using a syringe pump. Endothelial Ca^{2+} activity was stimulated by swapping the PSS syringe for one containing **ACh** (100 nM) or **GSK1016790A** (GSK, 20 nM). GSK1016790A is a selective TRPV4 channel agonist and evokes Ca^{2+} influx in wild type but not TRPV4 channel knockout mice (Mannaa et al., 2018; Sonkusare et al., 2012).

In experiments examining the effects of various pharmacological agents on stimulated endothelial Ca^{2+} signalling, drugs were added to the perfusate and remained thereafter. Images were acquired at 10 Hz using an inverted fluorescence microscope (Eclipse TE300, Nikon, Tokyo, Japan) equipped with a 40 \times objective (S Fluor, Nikon, Tokyo, Japan, NA = 1.3) and an electron-multiplying charge-coupled device camera (iXon Life; Andor Technology Limited, Belfast, Northern Ireland, UK) or on an upright epi-fluorescence microscope (FN-1, Nikon, Tokyo, Japan) equipped with a 60 \times objective (CFI Fluor, Nikon, Tokyo, Japan, NA = 1.0). Cal520/AM was excited at 488 nm using a monochromator (Polychrome IV, TILL Photonics,

Graefelfing, Germany) or an LED illumination system (PE-300Ultra, CoolLED, Andover, UK).

In some experiments, the endothelium was loaded with Cal-520 (5 μM) and a membrane permeant, caged IP_3 (5 μM), and the Ca^{2+} response to local photolysis of caged IP_3 examined. In these experiments, the endothelium was imaged as above, and a xenon flash lamp (Rapp Optoelectronic, Germany) was used to uncage IP_3 (Olson, Chalmers, & McCarron, 2012; Olson, Sandison, Chalmers, & McCarron, 2012; Wilson et al., 2019).

2.3 | Extraction and analysis of endothelial Ca^{2+} signals

Endothelial Ca^{2+} signals were extracted automatically from fluorescence recordings using custom written Python software (RRID: SCR_001658; Lee et al., 2018; Wilson, Lee, & McCarron, 2016; Wilson, Saunter, et al., 2016). In brief, cellular regions of interest (ROIs) were first generated from average-intensity projections of each time series recording. Intensity projections were sharpened (un-sharp mask filter) and thresholded, generating ROIs that encompassed the majority of each cell's area. To facilitate comparisons in paired experiments, for example, where Ca^{2+} activity was recorded before and after pharmacological inhibition, ROIs were aligned and tracked across separate image acquisitions. Only cells that remained within the field of view for all recordings were included.

Cellular Ca^{2+} signals were extracted by averaging fluorescence intensity within each of the ROIs, for each frame of the image stack. Raw fluorescence signals (F) were expressed as fractional changes in fluorescence intensity (F/F_0) by dividing each intensity value by the average intensity of a 100-frame period exhibiting the least activity/noise (F_0 , red in Figure S1B). The F_0 period was determined automatically by calculating, in series, the derivative of the signal, the rolling (100-frame) SD of the derivative, and the rolling (100-frame) summation of the rolling SD. The minimum value of the rolling summation indicates the centre of the portion of the signal with the least activity/noise.

The Ca^{2+} response to activation of TRPV4 channels contained two main components: a "slow" persistent Ca^{2+} elevation rise in the baseline Ca^{2+} levels and fast intracellular Ca^{2+} waves. To determine the underlying mechanisms, we isolated the two components from each F/F_0 trace. The slow persistent Ca^{2+} elevation (baseline) component of each signal was extracted by applying an asymmetric least squares (ALS) smoothing function (Figure S1C–E; Eilers & Boelens, 2005). Signalling metrics describing this slow persistent Ca^{2+} elevation (amplitude; time of activation, 10–90% rise time) were then calculated from a fitted sigmoidal curve (Figure S1C–E). Cells were considered to exhibit a persistent Ca^{2+} elevation if the slow F/F_0 component rose by more than 10-fold the SD of the baseline noise. The propagating Ca^{2+} wave (fast) component of each signal was extracted by dividing each F/F_0 trace by the ALS-smoothed signal. This procedure effectively flattens the baseline and removes any slow drift from the signal (Figure S1C–E). Peaks in each Ca^{2+} signal arising from the waves were

then detected automatically, using a zero-crossing detector on derivative F/F_0 traces (Lee et al., 2018; Wilson, Lee, & McCarron, 2016). Various signal metrics (number of peaks, peak amplitudes, peak durations, 10–90% rise time, and 90–10% fall time) were extracted from the corresponding ALS-smoothed F/F_0 trace. Cells were considered to exhibit Ca^{2+} wave activity if the F/F_0 component exhibited at least one event with an amplitude more than 10-fold the SD of the baseline noise.

In some experiments, Ca^{2+} signals were calibrated using $[\text{Ca}^{2+}] = K_D \frac{F - F_{\min}}{F_{\max} - F}$, where $K_D = 320$ nM. F_{\max} was determined at the end of the experiment by the addition of ionomycin (1 μM) and F_{\min} the signal in the absence of light to the camera.

2.4 | Measurement of vascular reactivity

To assess the endothelium dependence of contractile and dilation responses, vascular reactivity to various agonists were assessed in *en face* artery preparations. This technique permits simultaneous visualization of the endothelium and assessment of functional responses (contraction and dilation) and was used so that we could confirm the removal of the endothelium as required.

Arteries of 200–230 μm diameter were loaded with Cal-520/AM as above, except that they were pinned to the bottom of a custom, Sylgard-coated flow chamber. This flow chamber was designed to facilitate the perfusion of solutions (1.5 $\text{ml}\cdot\text{min}^{-1}$) when mounted on an upright microscope (FN-1, Nikon, Tokyo, Japan). Handling and pinning of the arteries were restricted to the outermost corners, leaving the central portion of the vessels with intact endothelium and able to contract/dilate freely.

Following incubation, arteries were gently washed with PSS and left to equilibrate for 30 min in PSS (continuously perfused 1.5 $\text{ml}\cdot\text{min}^{-1}$ using a syringe pump). Contraction was stimulated by swapping the PSS for one containing phenylephrine. The concentration of phenylephrine was titrated to contract arteries to ~20% of resting diameter. Relaxation was stimulated by changing the phenylephrine-containing syringe for one containing phenylephrine (at the same concentration) and either ACh (100 nM), GSK1016790A (GSK, 20 nM), or sodium nitroprusside (SNP, 100 μM). Contractile and dilator responses were assessed before and after (paired) various treatments in the same artery, as described in the text. After treatment, antagonists were continuously present in the PSS. The removal of the endothelium was achieved by gentle mechanical disruption with a human hair that was superglued onto a pipette tip. Endothelium removal was confirmed by the absence of endothelial cells stained with the Ca^{2+} indicator Cal-520/AM. After the removal of the endothelium, arteries were left to equilibrate for 60 min.

Images were acquired at 10 or 5 Hz (consistent within each experimental protocol) using an upright fluorescence microscope equipped with a 16 \times objective lens (0.8 NA; Nikon, Tokyo, Japan) and large

format (1,024 × 1,024; 13- μ m pixels) back-illuminated electron-multiplying charge-coupled device camera (iXon 888; Andor, Belfast, UK) and stored for offline analysis. An edge-detection algorithm (Lawton et al., 2019) was used to track the width of arteries in image recordings. The algorithm extracts an intensity profile along a scanline orientated perpendicular to the longitudinal axis of the vessel. Each intensity profile was smoothed using a 251-point, fifth-order Savitzky–Golay filter, and the first-order derivative calculated. The edges of the artery correspond to the maxima and minima of the first-order derivative, which were identified and tracked using a zero-crossing detector. The width of the artery equates to the unfolded circumference of the intact vessel and was converted to the equivalent diameter. To control for variation in the resting diameters, summary contractile response data are expressed as the percentage reduction from resting diameter, while relaxation responses are expressed as the percentage increase in diameter compared to resting diameter (Figure S2).

2.5 | Matching arterial tone across experimental conditions

Following the removal of the endothelium, arteries were significantly more sensitive to phenylephrine. The concentration of phenylephrine was titrated to achieve a level of tone that was comparable to that achieved in the presence of a functional endothelium (Figure S2A). In the presence of **cyclopiazonic acid** (CPA), Phenylephrine-induced contractions were transient, and it was not possible to achieve a stable contraction. Therefore, assessing the effect of store depletion on TRPV4-induced dilation was not feasible. Instead, vessels were pretreated with GSK and CPA, and the magnitude of the transient contraction induced by phenylephrine was assessed. Furthermore, after depletion of Ca^{2+} stores using CPA, a significantly higher concentration of phenylephrine was required to generate comparable levels of vascular tone (Figure S2B).

2.6 | Pressure myography

Diameter measurements of pressurized arteries were performed using the VasoTracker open source pressure myograph system (system (RRID:SCR_017233; Lawton et al., 2019). Mesenteric arteries were dissected free from surrounding tissue, cleaned of connective tissue, and mounted on similar sized glass pipettes in a VasoTracker pressure myograph chamber. Arteries were pressurized to 70 mmHg, checked for leaks, and left to equilibrate for 30 min in pre-warmed (37°C) circulating PSS, with intraluminal flow ($\sim 100 \mu\text{l}\cdot\text{min}^{-1}$) established by a 10 cmH_2O pressure gradient.

Artery diameter was monitored using an sCMOS camera (DCC1545M, Thorlabs, New Jersey, USA) and recorded by VasoTracker acquisition software. Arteries were contracted using phenylephrine ($\sim 500 \text{ nM}$) to $\sim 80\%$ of resting diameter, which was added to the perfusion solution. All other drugs (e.g., ACh and

GSK) were applied intraluminally. Summary relaxation data are expressed as the percentage increase in diameter compared to resting diameter.

2.7 | Materials

Cal-520/AM was obtained from Abcam (Cambridge, MA, USA). Caged-IP₃ (caged-IP₃ 4,5-dimethoxy-2-nitrobenzyl) was obtained from SicheM (Germany). Pluronic F-127 was obtained from Invitrogen (Carlsbad, CA, USA). U73343 was obtained from Tocris (St Louis, MO, USA). All other drugs and chemicals were obtained from Sigma (St Louis, MO, USA). GSK, **HC067047**, CPA, **U73122**, U73343, **2-aminoethoxydiphenyl borate** (2-APB), and ionomycin were dissolved in DMSO and diluted to working concentration in PSS such that the total volume of DMSO was less than or equal to 0.1%. All other drug stocks were dissolved in water. The PSS consisted of the following (mM): 145 NaCl, 4.7 KCl, 2.0 MOPS, 1.2 NaH_2PO_4 , 5.0 glucose, 0.02 EDTA, 1.17 MgCl_2 , and 2.0 CaCl_2 (adjusted to pH 7.4 with NaOH). In experiments using a Ca^{2+} -free PSS, Ca^{2+} was substituted with Mg^{2+} on an equimolar basis, and EGTA (1 mM) was included. All solutions were freshly prepared each day.

2.8 | Data and statistical analysis

The data and statistical analysis in this study comply with the recommendations of the British Journal of Pharmacology on experimental design and analysis in pharmacology. The n values shown are the numbers of biological replicates (number of animals). For each replicate, a single field of endothelial cells (Ca^{2+} imaging experiments) or a single artery segment (vascular reactivity studies) was studied. Summary data are presented graphically as paired mean responses, or as the grand mean \pm SEM of n biological replicates. Two-tailed, paired t test (for paired observations), independent two-sample t tests or repeated measures one-way ANOVA with multiple comparisons were used as indicated in the text. All statistical analyses were performed using GraphPad Prism version 6.0 (RRID:SCR_002798; GraphPad Software, La Jolla, CA, USA). A P value of $< .05$ was considered statistically significant.

2.9 | Nomenclature of targets and ligands

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Harding et al., 2018), and are permanently archived in the Concise Guide to PHARMACOLOGY 2017/18 (Alexander, Christopoulos et al., 2017; Alexander, Fabbro et al., 2017; Alexander, Peters et al., 2017; Alexander, Striessnig et al., 2017).

3 | RESULTS

3.1 | Characteristics of ACh- and GSK-evoked Ca^{2+} signals

To determine if activation of TRPV4 channels in endothelial cells evoked Ca^{2+} release from the internal store, endothelial cells were loaded with the fluorescent Ca^{2+} indicator Cal520/AM and activated with ACh (100 nM) or the TRPV4 channel agonist GSK (20 nM).

Ca^{2+} activity was then visualized in the fields of ~100 endothelial cells. Cellular responses were analysed individually (Figures 1 and S1).

Ca^{2+} responses evoked by ACh and GSK were clearly different (Figure 1). ACh evoked a rapid elevation in intracellular Ca^{2+} which was followed by asynchronous oscillations across the field of view (Figure 1; Mumtaz et al., 2011). Activation of TRPV4 channels resulted in heterogeneous Ca^{2+} responses (Figure 1c, Movie S1, and Figure S3) that consisted of several distinct phases (Figures 1 and 2).




FIGURE 1 ACh and GSK1016790A stimulate different Ca^{2+} signalling patterns. (a, b) Ca^{2+} dynamics visualized in a single *en face* endothelial preparation stimulated with ACh (a), and GSK1016790A (GSK, b). Panels show: (i) composite images illustrating endothelial Ca^{2+} activity over the time course of the experiment, scale bars = 50 μm ; (ii) whole-cell Ca^{2+} traces (F/F_0) for all cells shown in the corresponding panel in (i) ranked and coloured according to the magnitude of the signal (dark blue, low; dark red, high); (iii) a representative Ca^{2+} trace from a single cell; (iv) a rolling summation (30 s) of the number of Ca^{2+} events (peaks) across the-field-of-view; and (v) a rastergram display of Ca^{2+} activity, where each dot represents the occurrence of a peak in the Ca^{2+} response from each cell. (c) Representative images showing endothelial cells that were most sensitive (first 25% of endothelial cells to respond) to application of either ACh (green) or GSK (red). On average, 32.6% of cells that responded to one agonist also responded to the other (indicated by yellow in right most panel; $n = 3$). Scale bars = 20 μm

The response to GSK was, initially, small localized Ca^{2+} spikes which led to a slowly increasing global cytoplasmic Ca^{2+} concentration (Figures 2b and S4). The GSK-evoked, global Ca^{2+} elevation eventually plateaued, but the time taken to plateau was significantly longer when compared to ACh (Figure 1; 55.9 ± 6.4 s for GSK; 16.7 ± 4.7 s for ACh; $n = 6$). Finally, large propagating Ca^{2+} waves developed (Figure 2b,c and Movies S1 and S2). These waves propagated within and between cells at a constant velocity of ~ 5 to $15 \mu\text{m}\cdot\text{s}^{-1}$ (Figure 2b,c, Movie S2, and Figure S5). GSK-evoked intracellular Ca^{2+} waves

were significantly lower in frequency than those for ACh (0.05 ± 0.01 Hz for GSK; 0.22 ± 0.03 Hz for ACh; $n = 6$).

These results suggest there are at least major two components of the Ca^{2+} signal arising from activation of TRPV4 channels: (a) initial local signals which lead to a slow global rise in Ca^{2+} and (b) large fast propagating Ca^{2+} waves.

We next sought to examine the mechanisms that give rise to the GSK-evoked Ca^{2+} signals. We first confirmed the reproducibility of GSK-evoked responses (20 nM) in the same preparation so that a




FIGURE 2 Activation of TRPV4 channels stimulates local and propagating Ca^{2+} events. (a) Representative Ca^{2+} image (left), cell outlines (middle), and $\Delta F/F_0$ maximum intensity projection (right) illustrating the endothelial Ca^{2+} response to activation of TRPV4 channels with GSK1016790A (GSK; 20 nM). The $\Delta F/F_0$ image shows an increase in Ca^{2+} across the field of endothelial cells. (b) GSK stimulates local Ca^{2+} events and propagating Ca^{2+} waves. Panels show: (i) Ca^{2+} traces illustrating the effect of GSK on whole-cell Ca^{2+} levels from cells indicated by the coloured outlines in (a); (ii) line scans (kymographs) showing GSK-evoked local Ca^{2+} events (1) and propagating Ca^{2+} waves (2). Kymographs in (ii) correspond to the times indicated by dashed lines in (i) and were generated from $F/F_{\Delta t}$ movies, where $F_{\Delta t}$ is the running (20 s) baseline. (c) Ca^{2+} waves propagate across multiple cells. Panels show: (i) time series of the active front (determined by sequential subtraction) of a multicellular Ca^{2+} wave that spreads across four endothelial cells; (ii, iii) polar coordinate system (ii), example $F/F_{\Delta t}$ radial line scans (iii, 20° integration angle), and polar plot (iv) showing the angular dependence of propagation distance/velocity for the depicted Ca^{2+} event. Data in (c) are shown in Movie S2. All scale bars = 20 μm

paired experimental design could be used to examine the effects of blockers. There was no significant difference in any of the Ca^{2+} signal parameters measured (number of peaks, peak amplitudes, peak durations, 10–90% rise time, and 90–10% fall time) on repeated applications of GSK (Figure S3).

3.2 | Ca^{2+} influx via TRPV4 channels is required to initiate endothelial Ca^{2+} signals

To examine the mechanisms underlying each component of the GSK-evoked Ca^{2+} signal, we developed methods to extract the slow global rise in Ca^{2+} and the propagating Ca^{2+} waves (fast) component from raw Ca^{2+} signals in each cell (Figure S1).

To determine if each component is dependent on Ca^{2+} entry, we first performed experiments using a Ca^{2+} -free PSS (Figure 3a,b). The removal of extracellular Ca^{2+} significantly decreased the percentage of cells exhibiting “slow” global Ca^{2+} elevations and “fast” propagating waves in response to GSK (Figure 3c). The velocity of residual Ca^{2+}

waves was similar to those occurring in the presence of external Ca^{2+} (Figure 3d; $10.0 \pm 1.1 \mu\text{m}\cdot\text{s}^{-1}$ for control; $8.8 \pm 0.7 \mu\text{m}\cdot\text{s}^{-1}$ for Ca^{2+} -free; $n = 5$). However, removal of external Ca^{2+} significantly reduced the amplitude (Figure 3e; $0.28 \pm 0.11 \Delta\text{F}/\text{F}_0$ for control; $0.02 \pm 0.10 \Delta\text{F}/\text{F}_0$ for Ca^{2+} -free; $n = 6$) and the frequency of occurrence (Figure 3f; 4.1 ± 0.6 peaks per cell for control; 1.7 ± 0.1 peaks per cell for Ca^{2+} -free; $n = 6$) of these residual propagating Ca^{2+} waves.

These results suggest that Ca^{2+} influx across the plasma membrane is essential for the slow global increase in Ca^{2+} and for the propagating Ca^{2+} waves evoked by GSK.

To test whether or not the slow global Ca^{2+} rise and propagating Ca^{2+} waves evoked by GSK arose from the activation of TRPV4 channels, we pre-incubated endothelial cells with the non-selective TRPV channel antagonist, ruthenium red (RuR; $5 \mu\text{M}$, 20 min; Figure 4a,b). RuR significantly reduced the percentage of cells responding to GSK with slow global Ca^{2+} elevations and propagating Ca^{2+} waves (Figure 4c). Pretreatment with RuR reduced the amplitude (Figure 4e; $0.33 \pm 0.06 \Delta\text{F}/\text{F}_0$ for control; $0.07 \pm 0.01 \Delta\text{F}/\text{F}_0$ for RuR; $n = 6$) and the frequency of occurrence (Figure 4f; 13.4 ± 3.3 peaks per cell for




FIGURE 3 The removal of extracellular Ca^{2+} abolishes GSK-evoked Ca^{2+} signalling. (a) Composite images showing GSK1016790A (GSK)-evoked Ca^{2+} activity in a single field of native mesenteric artery endothelial cells before (left, control) and after (right, Ca^{2+} -free) the removal of external Ca^{2+} . Images show basal Cal-520/AM fluorescence (grey) with Ca^{2+} activity overlaid (red). Scale bars = $50 \mu\text{m}$. (b) GSK-evoked (20 nM; grey box) cellular Ca^{2+} signals (F/F_0) in the presence (left) and absence (right) of extracellular Ca^{2+} . F/F_0 signals (top) were decomposed into fast (middle) and slow (bottom) components. (c) Summary bar chart showing the percentage of cells that exhibited propagating Ca^{2+} waves (fast) and slow global Ca^{2+} elevations (slow). (d–f) Paired summary data showing the effect of external Ca^{2+} inhibition on Ca^{2+} wave propagation velocity (d), peak amplitude ($\Delta\text{F}/\text{F}_0$, e), and the number of peaks per cell (f). The left plots in (e and f) are scatter plots showing the mean Ca^{2+} event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a and b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. $*P < .05$, significantly different as indicated; paired Student's t test ($n = 6$)




FIGURE 4 Inhibition of TRPV channels with ruthenium red inhibits propagating Ca²⁺ waves and slow global Ca²⁺ elevations evoked by GSK1016790A. (a) Composite images showing GSK1016790A (GSK)-evoked (20 nM) Ca²⁺ activity in the absence (left) and presence (right) of the broad-spectrum TRPV channel antagonist ruthenium red (RuR; 5 μM, 20 min) in the same native mesenteric artery endothelial cells. Images show basal Cal-520/AM fluorescence (grey) with Ca²⁺ activity overlaid (red). Scale bars = 50 μm. (b) Raw F/F₀ (top), fast (middle), and slow (bottom) components of the GSK-evoked (20 nM; grey box) Ca²⁺ signal in the absence (left) and presence (right) of RuR. (c) Summary bar chart illustrating the effect of RuR on the percentage of cells exhibiting propagating Ca²⁺ waves and slow global Ca²⁺ signal components. (d–f) Paired summary data showing the effect of TRPV channel inhibition on Ca²⁺ wave propagation velocity (d), peak amplitude (ΔF/F₀, e), and the number of peaks per cell (f). The left plots in (e–f) are scatter plots showing the mean Ca²⁺ event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a) and (b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. **P* < .05, significantly different as indicated; paired Student's *t* test (*n* = 6)

control; 3.2 ± 0.1 peaks per cell for RuR; *n* = 6) of propagating Ca²⁺ waves but had no effect on the velocity of propagating waves (Figure 4d; $11.8 \pm 0.7 \mu\text{m}\cdot\text{s}^{-1}$ for control; $11.6 \pm 1.7 \mu\text{m}\cdot\text{s}^{-1}$ for RuR; *n* = 5). A similar reduction in GSK-evoked Ca²⁺ signalling was observed using the selective TRPV4 channel antagonist, HC067047 (HC067; 10 μM, 20 min; Figure 5a,b). HC067 reduced the percentage of cells exhibiting slow, global elevations and the percentage exhibiting fast, propagating Ca²⁺ waves (Figure 5c). HC067 reduced the amplitude (Figure 5e; $0.29 \pm 0.08 \Delta\text{F}/\text{F}_0$ for control; $0.08 \pm 0.02 \Delta\text{F}/\text{F}_0$ for HC067; *n* = 6) and the frequency (Figure 5f; 5.5 ± 0.8 peaks per cell for control; 0.7 ± 0.2 peaks per cell for HC067; *n* = 6) of propagating Ca²⁺ waves. The velocity of residual Ca²⁺ waves was unaffected by HC067 (Figure 5d; $13.1 \pm 0.8 \mu\text{m}\cdot\text{s}^{-1}$ for control; $13.6 \pm 0.7 \mu\text{m}\cdot\text{s}^{-1}$ for HC067; *n* = 5).

Together, these results suggest that Ca²⁺ influx via TRPV4 channels is required to induce both the slow global rise component and the propagating Ca²⁺ waves induced by GSK.

3.3 | GSK-induced Ca²⁺ waves require a replete internal Ca²⁺ store, IP₃ synthesis, and IP₃ receptor activation

Having confirmed a role for TRPV4-mediated Ca²⁺ influx in GSK-stimulated Ca²⁺ signals, we next investigated the contribution of the internal Ca²⁺ store using the sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase (SERCA) inhibitor, CPA (Figure 6a,b). Depletion of the internal store by CPA decreased the percentage of cells exhibiting the slow global rise component of the GSK-evoked Ca²⁺ signal and the percentage of cells in which propagating Ca²⁺ waves occurred (Figure 6c). CPA significantly inhibited the wave propagation speed (Figure 6d; $8.6 \pm 0.4 \mu\text{m}\cdot\text{s}^{-1}$ for control; $3.6 \pm 0.5 \mu\text{m}\cdot\text{s}^{-1}$ for CPA; *n* = 6), amplitude (Figure 6e; $0.48 \pm 0.12 \Delta\text{F}/\text{F}_0$ for control; $0.17 \pm 0.04 \Delta\text{F}/\text{F}_0$ for CPA; *n* = 6), and frequency (Figure 6f; 4.3 ± 0.4 peaks per cell for control; 1.5 ± 0.2 peaks per cell for CPA; *n* = 6) of Ca²⁺ waves. The amplitude of the slow global Ca²⁺ rise




FIGURE 5 Inhibition of TRPV4 channels with HC067047 inhibits both propagating Ca^{2+} waves and slow global Ca^{2+} elevations evoked by GSK1016790A. (a) Composite images showing GSK1016790A (GSK)-evoked (20 nM) Ca^{2+} activity in the absence (left) and presence (right) of the broad-spectrum TRPV channel antagonist HC067047 (HC067; 10 μM , 20 min) in the same native mesenteric artery endothelial cells. Images show basal Cal-520/AM fluorescence (grey) with Ca^{2+} activity overlaid (red). Scale bars = 50 μm . (b) Raw F/F_0 (top), fast (middle), and slow (bottom) components of the GSK-evoked (grey box) Ca^{2+} signal in the absence (left) and presence (right) of HC067. (c) Summary bar chart illustrating the effect of HC067 on the percentage of cells exhibiting propagating Ca^{2+} waves and slow global Ca^{2+} signal components. (d–f) Paired summary data showing the effect of TRPV4 channel inhibition on Ca^{2+} wave propagation velocity (d), peak amplitude ($\Delta F/F_0$, e), and the number of peaks per cell (f). The left plots in (e–f) are scatter plots showing the mean Ca^{2+} event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a) and (b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. * $P < .05$, significantly different as indicated; paired Student's *t* test ($n = 6$)

initially appeared to be reduced by CPA (Figure 6b). However, this apparent decrease appears to occur in large part because of the increase in baseline Ca^{2+} evoked by CPA and the non-linear relationship between Ca^{2+} concentration and F/F_0 value. When the signals were calibrated (see Section 2), the mean Ca^{2+} concentration increase evoked by GSK was 123 ± 75 nM, but after CPA, GSK evoked a Ca^{2+} rise of 93 ± 44 nM ($n = 3$), showing no significant effect of CPA. CPA itself evoked an increase in Ca^{2+} concentration from a resting value of 176 ± 52 to 335 ± 58 nM ($n = 3$). The elevated Ca^{2+} may account for the reduced number of cells activated by GSK (Watanabe et al., 2003). These results suggest that the internal store is required for the propagating Ca^{2+} waves.

To explore the role of Ca^{2+} release from the internal store, we examined the contribution of IP_3 using pharmacological inhibition of phospholipase C (PLC; Figure 7a,b). The PLC inhibitor, U73122, significantly reduced the percentage of cells activated by GSK, both for the

fast propagating Ca^{2+} waves and for the slow persistent Ca^{2+} elevation (Figure 7c). Although the mean amplitude of propagating Ca^{2+} waves was not significantly different in the presence of U73122 ($1.32 \pm 0.25 \Delta F/F_0$ for control; 0.67 ± 0.14 for U73122; $n = 5$), there was a significant reduction in the propagation velocity ($10.2 \pm 0.4 \mu\text{m}\cdot\text{s}^{-1}$ for control; $5.5 \pm 0.7 \mu\text{m}\cdot\text{s}^{-1}$ for U73122; $n = 5$) and frequency of those events (Figure 7d–f).

U73343, the inactive analogue of the PLC inhibitor U73122, had no effect on GSK-induced Ca^{2+} signals (Figure S6). The percentage of cells exhibiting slow and propagating Ca^{2+} wave components of the GSK-evoked signal were unaltered. Pretreatment with U73343 did not significantly alter the propagation velocity ($9.4 \pm 0.4 \mu\text{m}\cdot\text{s}^{-1}$ for control; $7.7 \pm 0.3 \mu\text{m}\cdot\text{s}^{-1}$ for U73343; $n = 5$), amplitude ($1.34 \pm 0.26 \Delta F/F_0$ for control; 1.34 ± 0.14 for U73343; $n = 5$), nor the frequency (3.5 ± 0.3 peaks/cell for control; 4.3 ± 0.6 peaks/cell for U73343; $n = 5$) of propagating Ca^{2+} waves. Moreover, although




FIGURE 6 Depletion of the internal Ca²⁺ store modifies the Ca²⁺ signal induced by activation of TRPV4 channels. (a) Composite images showing GSK1016709A (GSK)-evoked Ca²⁺ activity in the absence (left) and presence (right) of cyclopiazonic acid (CPA; 10 μM, 5 min) in the same native mesenteric artery endothelial cells. Images show basal Cal-520/AM fluorescence (grey) with Ca²⁺ activity overlaid (red). Scale bars = 50 μm. (b) GSK-evoked (20 nM; grey box) cellular Ca²⁺ signals in the absence and presence of CPA (left and right respectively). Original F/F₀ traces (top) were decomposed into fast (middle) and slow (bottom) components. CPA evoked a substantial increase in basal Ca²⁺ (see Section 3). In control and in the presence of CPA, F/F₀ was normalized to the Ca²⁺ change before GSK so that the resting values in each case were 1. CPA caused a substantial increase in resting Ca²⁺ (see text). (c) Summary bar chart illustrating the effect of CPA on the percentage of cells exhibiting propagating Ca²⁺ waves and slow global Ca²⁺ elevations to GSK. (d–f) Paired summary data showing the effect of SERCA inhibition on Ca²⁺ wave propagation velocity (d), peak amplitude (ΔF/F₀, e), and the number of peaks per cell (f). The left plots in (e–f) are scatter plots showing the mean Ca²⁺ event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a and b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. **P* < .05, significantly different as indicated; paired Student's *t* test (*n* = 6)

U73122 inhibits the internal store Ca²⁺ pump in colonic smooth muscle cells (Macmillan & McCarron, 2010), the blocker did not reduce IP₃-evoked Ca²⁺ release in endothelial cells (Figure S7).

These results suggest that the GSK-evoked slow global rise in Ca²⁺ and propagating Ca²⁺ waves each have contributions from PLC activation.

To investigate any role of IP₃-sensitive Ca²⁺ stores in propagating Ca²⁺ waves induced by activation of TRPV4 channels, we examined the effects of the IP₃R antagonist, 2-APB, on GSK-evoked endothelial Ca²⁺ signalling (Figure 8). 2-APB significantly reduced the percentage of cells exhibiting propagating Ca²⁺ waves but had no effect on the percentage of cells exhibiting slow global elevations in Ca²⁺ in response to GSK (Figure 8c). 2-APB had no effect on the amplitude (0.67 ± 0.09 ΔF/F₀ for control; 0.60 ± 0.15 ΔF/F₀ for 2-APB; *n* = 6) but significantly reduced propagation velocity (11.0 ± 0.8 μm·s⁻¹ for control; 2.0 ± 0.4 μm·s⁻¹ for 2-APB; *n* = 5) and frequency (3.1 ± 0.3

peaks per cell for control; 0.9 ± 0.3 peaks per cell for 2-APB; *n* = 6) of fast Ca²⁺ waves (Figure 8d–f).

To crosscheck the contribution of IP₃R in the propagating Ca²⁺ waves, we used caffeine—a potent inhibitor of IP₃R (Ehrlich, Kaftan, Bezprozvannaya, & Bezprozvanny, 1994; Parker & Ivorra, 1991; Saleem, Tovey, Molinski, & Taylor, 2014). Caffeine, which does not evoke Ca²⁺ release in the endothelial cells under study and inhibits Ca²⁺ release evoked by IP₃ (Wilson et al., 2019), also blocked GSK-evoked propagating Ca²⁺ waves (Figure S8). Interestingly, caffeine also reduced the slow global Ca²⁺ rise suggesting an effect of caffeine also on TRPV4 channels.

While Ca²⁺ influx via TRPV4 channels triggered large propagating Ca²⁺ waves from the internal Ca²⁺ store, the release and influx did not become an uncontrolled self-regenerative process but remained under the control of Ca²⁺ influx. When activation of TRPV4 channels ceases, by washout of GSK, the propagating Ca²⁺ waves stop (Figure S9 and Movie S3), and Ca²⁺ returned towards resting values.




FIGURE 7 PLC inhibition reduces the slow global Ca²⁺ rises and propagating Ca²⁺ waves of the GSK-evoked Ca²⁺ response. (a) Composite images illustrating GSK1016790A (GSK)-evoked (20 nM) Ca²⁺ activity in the absence (left) and presence (right) of the PLC antagonist U73122 (2 μM, 10 min) in a single field of native mesenteric artery endothelial cells. Images show basal Cal-520/AM fluorescence (grey) with Ca²⁺ activity overlaid (red). Scale bars = 50 μm. (b) GSK-evoked F/F₀ (top) Ca²⁺ signals, and corresponding fast (middle) and slow (bottom) signal components in the absence (left) and presence (right) of U73122. (c) Summary data showing the effect of U73122 on the percentage of endothelial cells exhibiting propagating Ca²⁺ waves (fast) and global Ca²⁺ elevations (slow). (d–f) Paired summary data showing the effect of PLC inhibition on Ca²⁺ wave propagation velocity (d), peak amplitude (ΔF/F₀) (e), and the number of peaks per cell (f). The left plots in (e–f) are scatter plots showing the mean Ca²⁺ event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a and b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. **P* < .05, significantly different as indicated; paired Student's *t* test (*n* = 5)

This result demonstrates that TRPV4 channels maintain control of Ca²⁺ release from the internal Ca²⁺ store and the propagation of Ca²⁺ waves both within and between cells.

3.4 | The functional effects of TRPV4 channels in controlling vascular tone

Collectively, our results suggest that endothelial TRPV4-mediated Ca²⁺ influx activates Ca²⁺-induced Ca²⁺ release at IP₃Rs. To investigate any physiological contribution of this process, we examined vascular tone in mesenteric arteries (Figure 8) with and without a functionally intact endothelial layer.

In intact, pressurized arteries, ACh and GSK each relaxed arteries and reversed phenylephrine-mediated contractions (Figure S10; *n* = 5). Relaxations to GSK were reversed by the selective TRPV4 channel antagonist, HC067. However, the dilations to ACh were preserved after blockade of TRPV4 channels with HC067 (Figure S10; *n* = 5). These results suggest that these channels do not contribute to ACh-evoked

relaxations in rat mesenteric arteries (see also Hartmannsgruber et al., 2007; Kohler et al., 2006; Wilson, Lee, & McCarron, 2016).

We next investigated the contribution of the internal Ca²⁺ store to TRPV4-mediated relaxation evoked by GSK. In *en face* preparations (225 ± 4 μm resting diameter; *n* = 10), ACh and GSK each relaxed phenylephrine-constricted arteries back to pre-constricted levels (Figure 9 and Figure S2). In these same arteries, the removal of the endothelium increased the sensitivity to phenylephrine and significantly reduced relaxation to ACh (to 26 ± 7%; *n* = 5 vs. control with endothelium). The residual ACh-evoked response may have arisen from the incomplete removal of the endothelium. The functional removal of the endothelium also prevented GSK-evoked relaxations. Indeed, after the removal of the endothelium, GSK caused an increase in tone (Figure 9e; *n* = 5). Endothelium removal did not prevent relaxation to the endothelium-independent vasodilator SNP (Figure 9).

These results demonstrate that vascular relaxation to GSK occurs via an endothelium-dependent mechanism and not by acting directly on the underlying smooth muscle.




FIGURE 8 Inhibiting IP₃ receptors prevents propagating Ca²⁺ waves but not slow global Ca²⁺ elevations induced by activation of TRPV4 channels. (a) Composite images showing Ca²⁺ activity evoked by GSK1016790A (GSK, 20 nM) in the absence (left) and presence (right) of the IP₃R antagonist, 2-APB (500 μM, 10 min) in a single field of native mesenteric artery endothelial cells. Images show basal Cal-520/AM fluorescence (grey) with Ca²⁺ activity overlaid (red). Scale bars = 50 μm. (b) GSK-evoked (grey box) cellular Ca²⁺ signals (F/F₀; top) were decomposed into fast (middle) and slow (bottom) components. (c) Summary data illustrating the percentage of cells exhibiting propagating Ca²⁺ waves and slow global Ca²⁺ increases. (d–f) Paired summary data showing the effect of IP₃R inhibition on Ca²⁺ wave propagation velocity (d), peak amplitude (ΔF/F₀, e), and the number of peaks per cell (f). The left plots in (e–f) are scatter plots showing the mean Ca²⁺ event amplitude, or oscillation frequency, within each cell visualized in the experiment shown in panels (a and b). Individual data points are coloured (from blue, low to red, high) according to the density (i.e., occurrence) of particular values. **P* < .05, significantly different as indicated; paired Student's *t* test (*n* = 6)

3.5 | Store dependence of the TRPV4 channel response

We next examined the contribution of Ca²⁺-induced Ca²⁺ release at IP₃Rs in TRPV4 mediated relaxation. To do this, we assessed the effect of GSK on vascular reactivity before and after depletion of internal Ca²⁺ stores.

Depletion of the internal Ca²⁺ store using CPA (6 μM) prevents stable contractions such that it was not possible to reliably assess vasodilator responses in these arteries. Instead, we examined whether pretreatment with GSK was capable of modulating PE-evoked contraction (Figure 10). Pretreatment with GSK significantly attenuated PE-induced contractions (Figure 10a, c).

After Ca²⁺ store depletion using CPA, a higher concentration of phenylephrine (3.6 ± 0.2 μM for control; 12.8 ± 2.2 μM for CPA; *n* = 5) was required to achieve a level of contraction (20 ± 2%; *n* = 5) equivalent to control conditions (Figure S2). After store depletion with CPA, pretreatment with GSK no longer attenuated

phenylephrine-induced contraction (4 ± 2% contraction for GSK; 27 ± 6% contraction for GSK + CPA; *n* = 5; *P* < .05). Rather, GSK tended to enhance the contraction to phenylephrine (Figure 10b, c). Together, these results demonstrate that depletion of internal Ca²⁺ stores reduces the inhibitory effect of GSK pretreatment on phenylephrine-induced contraction.

4 | DISCUSSION

TRPV4 channels are emerging as key plasmalemmal channels that mediate Ca²⁺ influx and physiological functions in the endothelium (Fiorio Pla et al., 2012; Gao & Wang, 2010; Gifford et al., 2014; Ma et al., 2013; Sayed et al., 2010; Schierling et al., 2011; Sonkusare et al., 2012; Thodeti et al., 2009; Troidl et al., 2009). We have shown that the Ca²⁺ increase in endothelial cells, arising from Ca²⁺ entry via TRPV4 channels, may be amplified by a Ca²⁺-induced Ca²⁺ release-like




FIGURE 9 ACh and GSK1016790A both cause endothelium-dependent vasodilation. (a and b) The vasodilator effect of ACh (a; 100 nM) and GSK1016790A (b; GSK; 20 nM) on phenylephrine (PE)-contracted arteries before (+ endothelium) and after the mechanical removal of the endothelium (- endothelium). After the removal of the endothelium, sodium nitroprusside (SNP) was used to test endothelium-independent vasodilation. Panels show: (i) images of an *en face* artery, illustrating the vessel's responses to phenylephrine and either ACh or GSK (in the continued presence of phenylephrine); (ii) full time course of the experiments shown in the corresponding panel (i); (iii) Summary data (mean \pm SEM, $n = 5$) illustrating the effect of the endothelium removal on vasodilator responses to the indicated agonist. Relaxations are expressed as % change from PE-induced tone. A negative value indicates an enhancement of phenylephrine-induced contraction. Coloured arrows in (ii) indicate the time at which relaxation to each agonist was assessed. All scale bars = 100 μm . * $P < .05$, significantly different as indicated; repeated measures one-way ANOVA with multiple comparison

mechanism acting at IP₃R_s to generate propagating Ca²⁺ waves. When IP₃R-mediated Ca²⁺ release was inhibited, TRPV4-induced increases in intracellular [Ca²⁺] and TRPV4-mediated control of vascular tone was

suppressed. These results highlight the significance of endothelial TRPV4-mediated Ca²⁺-induced Ca²⁺ release at IP₃R_s in controlling vascular contractility (Figure 11).




FIGURE 10 Ca^{2+} store depletion prevents TRPV4-mediated opposition of vascular tone. (a and b) Effect of GSK1016790A (GSK, 20 nM) on phenylephrine (PE)-induced contraction in vessels with an intact internal Ca^{2+} store (a) or after depletion of the internal Ca^{2+} store (b) using cyclopiazonic acid (CPA; 6 μM ; bottom trace). Upper panels show images of *en face* arteries, pretreated with the indicated drug, before and after exposure to phenylephrine. Corresponding diameter traces are shown in the lower panels. Data in (i) and (ii) show responses from the same artery. Pretreatment with CPA and GSK consistently delayed the onset but increased the magnitude of the response to phenylephrine. The reason for the delay is not clear. All scale bars = 100 μm . (c) Summary data (mean \pm SEM, $n = 5$) showing the magnitude of phenylephrine-induced contraction (% resting diameter) under each of the conditions. Diameter measured at the times indicated by the red arrows on diameter traces. * $P < .05$, significantly different as indicated; repeated measures one-way ANOVA with multiple comparisons

In this study, we analysed individual TRPV4-mediated Ca^{2+} responses in large numbers of endothelial cells in intact blood vessels. Two components, a slow global increase and propagating Ca^{2+} waves, contributed to the Ca^{2+} response to activation of TRPV4 channels. The response to activation of TRPV4 channels was also heterogeneous among cells. Some cells were highly responsive, while other cells responded weakly or not at all (see also Aird, 2012; Huang,

Chu, Chen, & Jen, 2000; Lee et al., 2018; Marie & Beny, 2002; McCarron, Lee, & Wilson, 2017; McCarron et al., 2019; Wilson, et al., 2016). We developed methods to automatically extract and separate the slow global increases and propagating Ca^{2+} waves occurring in each endothelial cell. The changes in Ca^{2+} concentration triggered by activation of TRPV4 channels consisted of different phases which operated sequentially. First, rapid localized Ca^{2+} changes occurred




FIGURE 11 Model of TRPV4-mediated control of vascular tone. Activation of TRPV4 channels (1) induces Ca^{2+} influx which results in a persistent elevation in cytoplasmic Ca^{2+} concentration (2). Basal (Hardie, Gu, Martin, Sweeney, & Raghu, 2004; Running Deer, Hurley, & Yarfitz, 1995; Willars, Nahorski, & Challiss, 1998) or Ca^{2+} -activated PLC (γ , δ , or β) activity decreases PIP_2 and generates IP_3 (3). Ca^{2+} derived from TRPV4 channel activity together with IP_3 (McCarron, Chalmers, MacMillan, & Olson, 2010) act at IP_3 receptors on the internal Ca^{2+} store to induce Ca^{2+} release (4). The released Ca^{2+} triggers propagating Ca^{2+} waves (5) by activating neighbouring IP_3 receptors that ultimately result in vasodilation (6). The entire process remains under the control of Ca^{2+} influx and terminates when TRPV4 channel activity ceases. This channel activity is suppressed by PIP_2 (7). ACh, via the **M3 receptor**, also activates PLC to generate IP_3 and activate Ca^{2+} release from the store. The characteristics of M_3 - and TRPV4-mediated Ca^{2+} release differ significantly (see text)

which led to a slow persistent global elevation in Ca^{2+} throughout the cytoplasm. Then, large transient increases as a result of propagating Ca^{2+} waves occurred. The large transient increases as a result of propagating Ca^{2+} waves dominated the Ca^{2+} changes occurring in the endothelium after a short time (~ 30 s).

Each of the Ca^{2+} changes occurring as a result of activation of TRPV4 channels was critically dependent on Ca^{2+} influx across the plasma membrane. The Ca^{2+} changes were abolished after the removal of external Ca^{2+} or by the inhibition of TRPV4 channels, with RuR or HC067047. These results suggest that Ca^{2+} influx via TRPV4 channels generates the persistent global Ca^{2+} rise. Inhibiting IP_3 -mediated Ca^{2+} release by depleting the internal Ca^{2+} store, or inhibiting IP_3 Rs, had little effect on the slow persistent global elevation in Ca^{2+} but prevented the large transient increases arising from propagating Ca^{2+} waves. This result suggests that Ca^{2+} entry via TRPV4 channels triggers activation of IP_3 Rs which amplifies the initial Ca^{2+} rise and initiates regenerative release of Ca^{2+} in the form of propagating waves. In support of the active nature of the Ca^{2+} waves, the velocity of wave progression away from the release site was relatively constant within and between cells (~ 5 to $15 \mu\text{m}\cdot\text{s}^{-1}$). If diffusion alone was responsible for wave progression, velocity and amplitude would decline with distance from the release site. The mechanisms contributing to TRPV4-mediated activation of Ca^{2+} release and propagation may include direct activation of IP_3 Rs (in a Ca^{2+} -induced Ca^{2+} release like process) or indirect Ca^{2+} -dependent activation of PLC. The partial inhibition of the propagating Ca^{2+} waves by the PLC-blocker, U73122, is consistent with being a Ca^{2+} -induced Ca^{2+} release-like process at IP_3 Rs. Basal PLC

activity has been reported for many PLC isoforms in biochemical experiments (Running Deer et al., 1995; Willars et al., 1998) and in living cells (Hardie et al., 2004; Willars et al., 1998) which may create a background level of IP_3 for Ca^{2+} -induced Ca^{2+} release to occur.

Interestingly, blocking of PLC by U73122 also reduced the slow global Ca^{2+} response (i.e., TRPV4-mediated Ca^{2+} influx). Previous work has shown that TRPV4 channels are inhibited by the phospholipid, **phosphatidylinositol 4,5-bisphosphate** (PIP_2 ; Harraz, Longden, Hill-Eubanks, & Nelson, 2018). PLC-mediated depletion of PIP_2 results in disinhibition of TRPV4 channels and increased Ca^{2+} influx (Harraz et al., 2018). Thus, blocking of PLC by U73122 might be expected to decrease TRPV4 channel activity.

To block IP_3 Rs, we used both 2-APB and caffeine. 2-APB is a broad-spectrum inhibitor that, although effective in blocking IP_3 Rs in native endothelial cells, may inhibit Ca^{2+} entry pathways (see Peppiatt et al., 2003; Trebak, Bird, McKay, & Putney, 2002; Voets et al., 2001; Wilson, Lee, & McCarron, 2016). However, in the present experimental conditions, 2-APB did not reduce the slow global Ca^{2+} rise evoked by activation of TRPV4 channels suggesting that it is not inhibiting influx in the present experiments. Caffeine, while often used as a RyR activator, is a potent inhibitor of IP_3 Rs in the same concentration range used to activate RyR (Ehrlich et al., 1994; Parker & Ivorra, 1991; Saleem et al., 2014). Caffeine does not evoke Ca^{2+} release in endothelial cells (Wilson, Saunter, Girkin, & McCarron, 2015) and inhibits Ca^{2+} release induced by photolysis of caged- IP_3 in the endothelium (Wilson et al., 2019). That caffeine was effective in inhibiting propagating Ca^{2+}

waves supports the conclusion that IP₃Rs contribute to TRPV4-mediated Ca²⁺ responses.

RuR and HC067047 were each used to block TRPV4 channels. RuR is an effective TRPV4 channel antagonist but also has effects unrelated to this channel. For example, RuR is an antagonist of channels such as the mitochondrial uniporter, the RyR, voltage-dependent Ca²⁺ channels, and other TRP channels. However, the effects of RuR on these other channels are unlikely to explain the present findings. RuR is not membrane permeant and will not have access to the cytoplasm, so the effects on the mitochondrial uniporter or RyR are unlikely to be of significance in the present study. While RuR may have effects on voltage-sensitive Ca²⁺ channels (Cibulsky & Sather, 1999), these channels do not appear to play a major role in endothelial Ca²⁺ signalling. RuR may also inhibit other TRP channels, for example, TRPC and TRPA. However, those channels are not activated by the agonist used in the present study—GSK1016790A—suggesting that channels other than TRPV4 are unlikely contributors. The present findings with RuR are also supported by the use of the more selective TRPV4 channel antagonist HC067047 which also blocked the response to GSK1016790A. Together, the findings point to TRPV4 channels as being the major target for RuR in the present study.

The increase in cytoplasmic Ca²⁺ concentration evoked by TRPV4-mediated Ca²⁺-induced Ca²⁺ release at IP₃Rs is large and regenerative, as demonstrated by the long distance propagation. This process arises (at least in part) because IP₃Rs are activated by Ca²⁺. TRPV4 channels are also activated by Ca²⁺ (Strotmann, Schultz, & Plant, 2003). The Ca²⁺-dependent processes operating at IP₃Rs and TRPV4 channels are positive feedback mechanisms that should self-reinforce and reach a maximum in all-or-none fashion. However, TRPV4-mediated Ca²⁺-induced Ca²⁺ release at IP₃Rs does not become uncontrolled and, while regenerative, remains tightly regulated by Ca²⁺ influx. The question arises as to why maximum responses do not occur each time TRPV4-mediated influx begins. The answer may lie in IP₃Rs and TRPV4 channels being regulated to prevent uncontrolled influx and release (Foskett, White, Cheung, & Mak, 2007; Nilius, Watanabe, & Vriens, 2003). Two observations support this conclusion. First, when TRPV4-mediated Ca²⁺ influx ceases, the release of Ca²⁺ from IP₃Rs stops. Thus, Ca²⁺ release is tightly coupled to influx. Second, coordinated activity of IP₃Rs is required for propagating waves to occur. The propagation of Ca²⁺ waves requires sequential opening and then closing of IP₃Rs. Thus, after activation, the IP₃R must become refractory to ensure a directional propagation of the signal. The decline in Ca²⁺ occurring at the back of the wave presumably follows from a functional compartmentalization of the endoplasmic reticulum which renders the site of IP₃-mediated Ca²⁺ release refractory to the phosphoinositide. The functional compartmentalization may arise from a lowered local Ca²⁺ concentration in the lumen of the endoplasmic reticulum, or increased Ca²⁺ concentrations could deactivate the receptors (Adkins & Taylor, 1999; McCarron, MacMillan, Bradley, Chalmers, & Muir, 2004; Oancea & Meyer, 1996). These observations highlight a system that is activated by Ca²⁺ influx, but is regulated to prevent an uncontrolled positive feedback processes dominating the TRPV4-mediated increases in cytoplasmic Ca²⁺ concentrations.

As well as propagating at a constant velocity within cells, TRPV4-mediated Ca²⁺ waves could move seamlessly, at the same velocity, across cell boundaries into neighbouring cells as intercellular Ca²⁺ waves (Movie S2). These waves were abolished by IP₃R blockers or removal of GSK. The mechanisms of intercellular wave propagation are not entirely clear, but IP₃Rs and (since the process stops when GSK is removed) TRPV4 channel activities are both required for these intercellular waves to occur.

In other tissues, intercellular Ca²⁺ waves may be a fundamental mechanism for coordinating multicellular responses (Leybaert & Sanderson, 2012). In the endothelium of small arteries of the mouse cremaster muscle, rapidly spreading intercellular Ca²⁺ waves propagate at a velocity of ~116 μm·s⁻¹ for distances up to ~1 mm (Tallini et al., 2007). These intercellular waves are associated with vasodilation and are hypothesized to regulate blood flow to the parenchyma by inducing upstream dilation of arterioles (Tallini et al., 2007). The intercellular waves observed in the present study had a slower velocity (~5 to 15 μm·s⁻¹) than those in mouse cremaster arterioles, suggesting a different physiological control mechanism.

In previous studies, when endothelial IP₃R-mediated signalling was eliminated by depletion of the internal Ca²⁺ store, activation of TRPV4 channels induced a large but highly localized (few square micrometres) increase in Ca²⁺ concentrations (Sonkusare et al., 2012). These localized Ca²⁺ signals were reported to activate endothelial Ca²⁺-activated K⁺ channels (Sonkusare et al., 2012) and lead to endothelial hyperpolarization and vasodilation (Sonkusare et al., 2012). However, activation of TRPV4 channels is known to also control the orientation of endothelial cells, regulate endothelial permeability, and modulate the production of antithrombotic factors, each of which may require a more global [Ca²⁺] increase throughout the cytoplasm (Noren et al., 2016; Phuong et al., 2017; Thodeti et al., 2009; Thoppil et al., 2016). In the present study, IP₃R-mediated Ca²⁺ release generated propagating Ca²⁺ waves and provided a mechanism by which TRPV4 channel activity, in the presence of a functioning Ca²⁺ store, may generate large rises in Ca²⁺ throughout the cytoplasm. These propagating waves are critical for TRPV4-mediated control of vascular tone. When Ca²⁺ release from the internal store is inhibited, TRPV4-mediated endothelial control of tone is abolished (Figure 10). This result suggests that control of vascular tone by TRPV4 channels is mediated by Ca²⁺-induced Ca²⁺ release via IP₃Rs. It is tempting to also speculate that the more general rise in Ca²⁺ throughout the cells, which occurs as a result of the propagating IP₃R mediated Ca²⁺ waves, will facilitate TRPV4-mediated control of endothelial permeability and the production of antithrombotic factors (Noren et al., 2016; Phuong et al., 2017; Thodeti et al., 2009; Thoppil et al., 2016).

ACh evoked rapid asynchronous Ca²⁺ waves in neighbouring endothelial cells. ACh-evoked changes in intracellular Ca²⁺ concentration are linked closely to the IP₃-sensitive Ca²⁺ store, but they do not involve TRPV4 channels in rat arteries (Hartmannsgruber et al., 2007; see also Kohler et al., 2006; Wilson, Lee, & McCarron, 2016; Figure 10). While the response to the activation by ACh and activation of TRPV4 channels each involved IP₃, the Ca²⁺ signals generated had very different characteristics. ACh-evoked Ca²⁺ increases did not

appear to coordinate between cells. Activation of TRPV4 channels evoked a slowly increasing baseline and large propagating Ca^{2+} waves.

Endothelial TRPV4 channels are involved in several cardiovascular control mechanisms, including inhibiting vasoconstriction in small skeletal muscle arteries in response to increases in temperature (Gifford et al., 2014), regulating myogenic tone (Bagher et al., 2012), remodeling of the cytoskeleton and reorientation of the endothelial cells in response to mechanical forces (Thodeti et al., 2009), collateral vessel growth (Sayed et al., 2010; Schierling et al., 2011; Troidl et al., 2009), and arachidonic acid-induced endothelial cell migration required for angiogenesis (Fiorio Pla et al., 2012). The present results show that when the internal IP_3 sensitive store is intact, TRPV4 channel activity evokes IP_3 R signalling to generate Ca^{2+} waves that propagate within and between cells, and through the Ca^{2+} store, TRPV4 channels modulate vascular contractility. The results demonstrate a link between TRPV4 channel activity and Ca^{2+} -induced Ca^{2+} release at IP_3 R in endothelial cells and offer a new target for drug development.

ACKNOWLEDGEMENTS

This work was funded by the Wellcome Trust (202924/Z/16/Z and 204682/Z/16/Z) and the British Heart Foundation (PG/16/54/32230 and PG16/82/32439), whose support is gratefully acknowledged. The authors would like to thank Margaret MacDonald for her excellent technical support.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

H.R.H., M.D.L., C.W., and J.G.M. developed the concept. H.R.H., M.D.L., X.Z., and C.W. performed the experiments. C.W. and C.D.S. wrote the analysis software. H.R.H., M.D.L., X.Z., and C.W. analysed the data. J.G.M. and H.R.H. drafted the manuscript. J.G.M., H.R.H., C.W., C.D.S., and M.D.L. edited the manuscript. J.G.M., M.D.L., C.W., and X.Z. revised the manuscript. C.W., C.D.S., and J.G.M. sourced funding. All authors approved the final version of the manuscript.

DECLARATION OF TRANSPARENCY AND SCIENTIFIC RIGOUR

This Declaration acknowledges that this paper adheres to the principles for transparent reporting and scientific rigour of preclinical research as stated in the *BJP* guidelines for [Design & Analysis](#) and [Animal Experimentation](#), and as recommended by funding agencies, publishers and other organisations engaged with supporting research

DATA AVAILABILITY

All data underpinning this study is available from the authors upon reasonable request.

ORCID

John G. McCarron  <https://orcid.org/0000-0002-3302-3984>

REFERENCES

- Adkins, C. E., & Taylor, C. W. (1999). Lateral inhibition of inositol 1,4,5-trisphosphate receptors by cytosolic Ca^{2+} . *Current Biology*, 9, 1115–1118. [https://doi.org/10.1016/S0960-9822\(99\)80481-3](https://doi.org/10.1016/S0960-9822(99)80481-3)
- Aird, W. C. (2012). Endothelial cell heterogeneity. *Cold Spring Harbor Perspectives in Medicine*, 2(1), a006429. <https://doi.org/10.1101/cshperspect.a006429>
- Alexander, S. P. H., Christopoulos, A., Davenport, A. P., Kelly, E., Marrion, N. V., Peters, J. A., ... CGTP Collaborators. (2017). The Concise Guide to PHARMACOLOGY 2017/18: G protein-coupled receptors. *British Journal of Pharmacology*, 174, S17–S129. <https://doi.org/10.1111/bph.13878>
- Alexander, S. P. H., Fabbro, D., Kelly, E., Marrion, N. V., Peters, J. A., Faccenda, E., ... CGTP Collaborators. (2017). The Concise Guide to PHARMACOLOGY 2017/18: Enzymes. *British Journal of Pharmacology*, 174, S272–S359. <https://doi.org/10.1111/bph.13877>
- Alexander, S. P. H., Peters, J. A., Kelly, E., Marrion, N. V., Faccenda, E., Harding, S. D., ... CGTP Collaborators. (2017). The Concise Guide to PHARMACOLOGY 2017/18: Ligand-gated ion channels. *British Journal of Pharmacology*, 174, S130–S159. <https://doi.org/10.1111/bph.13879>
- Alexander, S. P. H., Striessnig, J., Kelly, E., Marrion, N. V., Peters, J. A., Faccenda, E., ... CGTP Collaborators. (2017). The Concise Guide to PHARMACOLOGY 2017/18: Voltage-gated ion channels. *British Journal of Pharmacology*, 174, S160–S194. <https://doi.org/10.1111/bph.13884>
- Bagher, P., Beleznai, T., Kansui, Y., Mitchell, R., Garland, C. J., & Dora, K. A. (2012). Low intravascular pressure activates endothelial cell TRPV4 channels, local Ca^{2+} events, and IKCa channels, reducing arteriolar tone. *Proceedings of the National Academy of Sciences of the United States of America*, 109(44), 18174–18179. <https://doi.org/10.1073/pnas.1211946109>
- Berridge, M. J. (1997). Elementary and global aspects of calcium signalling. *The Journal of Experimental Biology*, 200(Pt 2), 315–319.
- Bootman, M., Niggli, E., Berridge, M., & Lipp, P. (1997). Imaging the hierarchical Ca^{2+} signalling system in HeLa cells. *The Journal of Physiology*, 499, 307–314. <https://doi.org/10.1113/jphysiol.1997.sp021928>
- Bootman, M. D., Berridge, M. J., & Lipp, P. (1997). Cooking with calcium: The recipes for composing global signals from elementary events. *Cell*, 91, 367–373. [https://doi.org/10.1016/S0092-8674\(00\)80420-1](https://doi.org/10.1016/S0092-8674(00)80420-1)
- Borisova, L., Wray, S., Eisner, D. A., & Burdya, T. (2009). How structure, Ca signals, and cellular communications underlie function in precapillary arterioles. *Circulation Research*, 105(8), 803–810. <https://doi.org/10.1161/CIRCRESAHA.109.202960>
- Catterall, W. A. (2011). Voltage-gated calcium channels. *Cold Spring Harbor Perspectives in Biology*, 3(8), a003947. <https://doi.org/10.1101/cshperspect.a003947>
- Cibulsky, S. M., & Sather, W. A. (1999). Block by ruthenium red of cloned neuronal voltage-gated calcium channels. *The Journal of Pharmacology and Experimental Therapeutics*, 289(3), 1447–1453.
- Dunn, K. M., Hill-Eubanks, D. C., Liedtke, W. B., & Nelson, M. T. (2013). TRPV4 channels stimulate Ca^{2+} -induced Ca^{2+} release in astrocytic endfeet and amplify neurovascular coupling responses. *Proceedings of the National Academy of Sciences of the United States of America*, 110(15), 6157–6162. <https://doi.org/10.1073/pnas.1216514110>
- Earley, S., Heppner, T. J., Nelson, M. T., & Brayden, J. E. (2005). TRPV4 forms a novel Ca^{2+} signaling complex with ryanodine receptors and

- BKCa channels. *Circulation Research*, 97(12), 1270–1279. <https://doi.org/10.1161/01.RES.0000194321.60300.d6>
- Ehrlich, B. E., Kaftan, E., Bezprozvannaya, S., & Bezprozvanny, I. (1994). The pharmacology of intracellular Ca^{2+} -release channels. *Trends in Pharmacological Sciences*, 15(5), 145–149. [https://doi.org/10.1016/0165-6147\(94\)90074-4](https://doi.org/10.1016/0165-6147(94)90074-4)
- Eilers, P. H. C., & Boelens, H. F. M. (2005). Baseline correction with asymmetric least squares smoothing. *Leiden University Medical Centre Report*.
- Filosa, J. A., Yao, X., & Rath, G. (2013). TRPV4 and the regulation of vascular tone. *Journal of Cardiovascular Pharmacology*, 61(2), 113–119. <https://doi.org/10.1097/FJC.0b013e318279ba42>
- Fiorio Pla, A., Ong, H. L., Cheng, K. T., Brossa, A., Bussolati, B., Lockwich, T., ... Ambudkar, I. S. (2012). TRPV4 mediates tumor-derived endothelial cell migration via arachidonic acid-activated actin remodeling. *Oncogene*, 31(2), 200–212. <https://doi.org/10.1038/onc.2011.231>
- Foskett, J. K., White, C., Cheung, K. H., & Mak, D. O. (2007). Inositol trisphosphate receptor Ca^{2+} release channels. *Physiological Reviews*, 87(2), 593–658. <https://doi.org/10.1152/physrev.00035.2006>
- Freichel, M., Suh, S. H., Pfeifer, A., Schweig, U., Trost, C., Weissgerber, P., ... Nilius, B. (2001). Lack of an endothelial store-operated Ca^{2+} current impairs agonist-dependent vasorelaxation in TRPV4^{-/-} mice. *Nature Cell Biology*, 3(2), 121–127. <https://doi.org/10.1038/35055019>
- Gao, F., & Wang, D. H. (2010). Hypotension induced by activation of the transient receptor potential vanilloid 4 channels: Role of Ca^{2+} -activated K^+ channels and sensory nerves. *Journal of Hypertension*, 28(1), 102–110. <https://doi.org/10.1097/HJH.0b013e328332b865>
- Gao, X., Wu, L., & O'Neil, R. G. (2003). Temperature-modulated diversity of TRPV4 channel gating: Activation by physical stresses and phorbol ester derivatives through protein kinase C-dependent and -independent pathways. *The Journal of Biological Chemistry*, 278(29), 27129–27137. <https://doi.org/10.1074/jbc.M302517200>
- Gifford, J. R., Ives, S. J., Park, S. Y., Andtbacka, R. H., Hynstrom, J. R., Mueller, M. T., ... Richardson, R. S. (2014). α 1- and α 2-adrenergic responsiveness in human skeletal muscle feed arteries: The role of TRPV ion channels in heat-induced sympatholysis. *American Journal of Physiology. Heart and Circulatory Physiology*, 307(9), H1288–H1297. <https://doi.org/10.1152/ajpheart.00068.2014>
- Hardie, R. C., Gu, Y., Martin, F., Sweeney, S. T., & Raghu, P. (2004). In vivo light-induced and basal phospholipase C activity in Drosophila photoreceptors measured with genetically targeted phosphatidylinositol 4,5-bisphosphate-sensitive ion channels (Kir2.1). *The Journal of Biological Chemistry*, 279(46), 47773–47782. <https://doi.org/10.1074/jbc.M407525200>
- Harding, S. D., Sharman, J. L., Faccenda, E., Southan, C., Pawson, A. J., Ireland, S., ... NC-IUPHAR. (2018). The IUPHAR/BPS guide to pharmacology in 2018: Updates and expansion to encompass the new guide to immunopharmacology. *Nucleic Acids Research*, 46(D1), D1091–D1106. <https://doi.org/10.1093/nar/gkx1121>
- Harraz, O. F., Longden, T. A., Hill-Eubanks, D., & Nelson, M. T. (2018). PIP2 depletion promotes TRPV4 channel activity in mouse brain capillary endothelial cells. *eLife*, 7. <https://doi.org/10.7554/eLife.38689>
- Hartmannsgruber, V., Heyken, W. T., Kacik, M., Kaistha, A., Grgic, I., Harteneck, C., ... Köhler, R. (2007). Arterial response to shear stress critically depends on endothelial TRPV4 expression. *PLoS ONE*, 2(9), e827. <https://doi.org/10.1371/journal.pone.0000827>
- Hofmann, T., Schaefer, M., Schultz, G., & Gudermann, T. (2002). Subunit composition of mammalian transient receptor potential channels in living cells. *Proceedings of the National Academy of Sciences of the United States of America*, 99(11), 7461–7466. <https://doi.org/10.1073/pnas.102596199>
- Huang, T. Y., Chu, T. F., Chen, H. I., & Jen, C. J. (2000). Heterogeneity of $[\text{Ca}^{2+}]_i$ signaling in intact rat aortic endothelium. *The FASEB Journal*, 14(5), 797–804. <https://doi.org/10.1096/fasebj.14.5.797>
- Kohler, R., Heyken, W. T., Heinau, P., Schubert, R., Si, H., Kacik, M., ... Hoyer, J. (2006). Evidence for a functional role of endothelial transient receptor potential V4 in shear stress-induced vasodilatation. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 26(7), 1495–1502. <https://doi.org/10.1161/01.ATV.0000225698.36212.6a>
- Kilkenny, C., Browne, W., Cuthill, I. C., Emerson, M., & Altman, D. G. (2010). Animal research: Reporting in vivo experiments: The ARRIVE guidelines. *British Journal of Pharmacology*, 160, 1577–1579.
- Lawton, P. F., Lee, M. D., Saunter, C. D., Girkin, J. M., McCarron, J. G., & Wilson, C. (2019). VasoTracker, a low-cost and open source pressure myograph system for vascular physiology has been approved for production and accepted for publication in Frontiers in Physiology, section Vascular Physiology. *Frontiers in Physiology*, 10. <https://doi.org/10.3389/fphys.2019.00099>
- Lee, M. D., Wilson, C., Saunter, C. D., Kennedy, C., Girkin, J. M., & McCarron, J. G. (2018). Spatially structured cell populations process multiple sensory signals in parallel in intact vascular endothelium. *Science Signaling*, 11(561), eaar4411. <https://doi.org/10.1126/scisignal.aar4411>
- Leybaert, L., & Sanderson, M. J. (2012). Intercellular Ca^{2+} waves: Mechanisms and function. *Physiological Reviews*, 92(3), 1359–1392. <https://doi.org/10.1152/physrev.00029.2011>
- Liedtke, W., Choe, Y., Marti-Renom, M. A., Bell, A. M., Denis, C. S., Sali, A., ... Heller, S. (2000). Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell*, 103(3), 525–535. [https://doi.org/10.1016/S0092-8674\(00\)00143-4](https://doi.org/10.1016/S0092-8674(00)00143-4)
- Liedtke, W., Tobin, D. M., Bargmann, C. I., & Friedman, J. M. (2003). Mammalian TRPV4 (VR-OAC) directs behavioral responses to osmotic and mechanical stimuli in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the United States of America*, 100(Suppl 2), 14531–14536. <https://doi.org/10.1073/pnas.2235619100>
- Ma, X., Du, J., Zhang, P., Deng, J., Liu, J., Lam, F. F., ... Yao, X. (2013). Functional role of TRPV4-KCa2.3 signaling in vascular endothelial cells in normal and streptozotocin-induced diabetic rats. *Hypertension*, 62(1), 134–139. <https://doi.org/10.1161/HYPERTENSIONAHA.113.01500>
- Macmillan, D., & McCarron, J. G. (2010). The phospholipase C inhibitor U-73122 inhibits Ca^{2+} release from the intracellular sarcoplasmic reticulum Ca^{2+} store by inhibiting Ca^{2+} pumps in smooth muscle. *British Journal of Pharmacology*, 160(6), 1295–1301. <https://doi.org/10.1111/j.1476-5381.2010.00771.x>
- Mannaa, M., Marko, L., Balogh, A., Vigolo, E., N'Diaye, G., Kassmann, M., ... Gollasch, M. (2018). Transient receptor potential vanilloid 4 channel deficiency aggravates tubular damage after acute renal ischaemia reperfusion. *Scientific Reports*, 8(1), 4878. <https://doi.org/10.1038/s41598-018-23165-0>
- Marie, I., & Beny, J. L. (2002). Calcium imaging of murine thoracic aorta endothelium by confocal microscopy reveals inhomogeneous distribution of endothelial cells responding to vasodilator agents. *Journal of Vascular Research*, 39(3), 260–267. <https://doi.org/10.1159/000063691>
- Marrelli, S. P., O'Neil, R. G., Brown, R. C., & Bryan, R. M. Jr. (2007). PLA2 and TRPV4 channels regulate endothelial calcium in cerebral arteries. *American Journal of Physiology. Heart and Circulatory Physiology*, 292(3), H1390–H1397. <https://doi.org/10.1152/ajpheart.01006.2006>
- McCarron, J. G., Chalmers, S., Bradley, K. N., Macmillan, D., & Muir, T. C. (2006). Ca^{2+} microdomains in smooth muscle. *Cell Calcium*, 40, 461–493. <https://doi.org/10.1016/j.ceca.2006.08.010>

- McCarron, J. G., Chalmers, S., MacMillan, D., & Olson, M. L. (2010). Agonist-evoked Ca^{2+} wave progression requires Ca^{2+} and IP_3 . *Journal of Cell Physiology*, 244, 334–344.
- McCarron, J. G., Lee, M. D., & Wilson, C. (2017). The endothelium solves problems that endothelial cells do not know exist. *Trends in Pharmacological Sciences*, 38(4), 322–338. <https://doi.org/10.1016/j.tips.2017.01.008>
- McCarron, J. G., MacMillan, D., Bradley, K. N., Chalmers, S., & Muir, T. C. (2004). Origin and mechanisms of Ca^{2+} waves in smooth muscle as revealed by localized photolysis of caged inositol 1,4,5-trisphosphate. *The Journal of Biological Chemistry*, 279, 8417–8427. <https://doi.org/10.1074/jbc.M311797200>
- McCarron, J. G., Wilson, C., Heathcote, H. R., Zhang, X., Buckley, C., & Lee, M. D. (2019). Heterogeneity and emergent behaviour in the vascular endothelium. *Current Opinion in Pharmacology*, 45, 23–32. <https://doi.org/10.1016/j.coph.2019.03.008>
- Mendoza, S. A., Fang, J., Gutterman, D. D., Wilcox, D. A., Bubolz, A. H., Li, R., ... Zhang, D. X. (2010). TRPV4-mediated endothelial Ca^{2+} influx and vasodilation in response to shear stress. *American Journal of Physiology. Heart and Circulatory Physiology*, 298(2), H466–H476. <https://doi.org/10.1152/ajpheart.00854.2009>
- Mizuno, A., Matsumoto, N., Imai, M., & Suzuki, M. (2003). Impaired osmotic sensation in mice lacking TRPV4. *American Journal of Physiology. Cell Physiology*, 285(1), C96–C101. <https://doi.org/10.1152/ajpcell.00559.2002>
- Moccia, F., Berra-Romani, R., & Tanzi, F. (2012). Update on vascular endothelial Ca^{2+} signalling: A tale of ion channels, pumps and transporters. *World Journal of Biological Chemistry*, 3(7), 127–158. <https://doi.org/10.4331/wjbc.v3.i7.127>
- Moore, C., Cevikbas, F., Pasolli, H. A., Chen, Y., Kong, W., Kempkes, C., ... Liedtke, W. B. (2013). UVB radiation generates sunburn pain and affects skin by activating epidermal TRPV4 ion channels and triggering endothelin-1 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, 110(34), E3225–E3234. <https://doi.org/10.1073/pnas.1312933110>
- Mumtaz, S., Burdya, G., Borisova, L., Wray, S., & Burdya, T. (2011). The mechanism of agonist induced Ca^{2+} signalling in intact endothelial cells studied confocally in in situ arteries. *Cell Calcium*, 49(1), 66–77. <https://doi.org/10.1016/j.ceca.2010.11.010>
- Nelson, M. T., Patlak, J. B., Worley, J. F., & Standen, N. B. (1990). Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. *The American Journal of Physiology*, 259(1 Pt 1), C3–C18. <https://doi.org/10.1152/ajpcell.1990.259.1.C3>
- Nilius, B., Droogmans, G., & Wondergem, R. (2003). Transient receptor potential channels in endothelium: Solving the calcium entry puzzle? *Endothelium: Journal of Endothelial Cell Research*, 10(1), 5–15. <https://doi.org/10.1080/10623320303356>
- Nilius, B., & Voets, T. (2013). The puzzle of TRPV4 channelopathies. *EMBO Reports*, 14(2), 152–163. <https://doi.org/10.1038/embor.2012.219>
- Nilius, B., Watanabe, H., & Vriens, J. (2003). The TRPV4 channel: Structure–function relationship and promiscuous gating behaviour. *Pflügers Archiv*, 446(3), 298–303. <https://doi.org/10.1007/s00424-003-1028-9>
- Noren, D. P., Chou, W. H., Lee, S. H., Qutub, A. A., Warmflash, A., Wagner, D. S., ... Levchenko, A. (2016). Endothelial cells decode VEGF-mediated Ca^{2+} signaling patterns to produce distinct functional responses. *Science Signaling*, 9(416), ra20. <https://doi.org/10.1126/scisignal.aad3188>
- Oancea, E., & Meyer, T. (1996). Reversible desensitization of inositol trisphosphate-induced calcium release provides a mechanism for repetitive calcium spikes. *The Journal of Biological Chemistry*, 271, 17253–17260. <https://doi.org/10.1074/jbc.271.29.17253>
- Olson, M. L., Chalmers, S., & McCarron, J. G. (2012). Mitochondrial organization and Ca^{2+} uptake. *Biochemical Society Transactions*, 40(1), 158–167. <https://doi.org/10.1042/BST20110705>
- Olson, M. L., Sandison, M. E., Chalmers, S., & McCarron, J. G. (2012). Microdomains of muscarinic acetylcholine and $\text{Ins}(1,4,5)\text{P}_3$ receptors create ' $\text{Ins}(1,4,5)\text{P}_3$ junctions' and sites of Ca^{2+} wave initiation in smooth muscle. *Journal of Cell Science*, 125(Pt 22), 5315–5328. <https://doi.org/10.1242/jcs.105163>
- Parker, I., & Ivorra, I. (1991). Caffeine inhibits inositol trisphosphate-mediated liberation of intracellular calcium in *Xenopus* oocytes. *The Journal of Physiology*, 433, 229–240. <https://doi.org/10.1113/jphysiol.1991.sp018423>
- Peppiatt, C. M., Collins, T. J., Mackenzie, L., Conway, S. J., Holmes, A. B., Bootman, M. D., ... Roderick, H. L. (2003). 2-Aminoethoxydiphenyl borate (2-APB) antagonises inositol 1,4,5-trisphosphate-induced calcium release, inhibits calcium pumps and has a use-dependent and slowly reversible action on store-operated calcium entry channels. *Cell Calcium*, 34(1), 97–108. [https://doi.org/10.1016/S0143-4160\(03\)00026-5](https://doi.org/10.1016/S0143-4160(03)00026-5)
- Phuong, T. T. T., Redmon, S. N., Yarishkin, O., Winter, J. M., Li, D. Y., & Krizaj, D. (2017). Calcium influx through TRPV4 channels modulates the adherens contacts between retinal microvascular endothelial cells. *The Journal of Physiology*, 595(22), 6869–6885. <https://doi.org/10.1113/JP275052>
- Running Deer, J. L., Hurley, J. B., & Yarfitz, S. L. (1995). G protein control of *Drosophila* photoreceptor phospholipase C. *The Journal of Biological Chemistry*, 270(21), 12623–12628. <https://doi.org/10.1074/jbc.270.21.12623>
- Rusko, J., Wang, X., & Vanbreemen, C. (1995). Regenerative caffeine-induced responses in native rabbit aortic endothelial-cells. *British Journal of Pharmacology*, 115(5), 811–821. <https://doi.org/10.1111/j.1476-5381.1995.tb15005.x>
- Saleem, H., Tovey, S. C., Molinski, T. F., & Taylor, C. W. (2014). Interactions of antagonists with subtypes of inositol 1,4,5-trisphosphate (IP_3) receptor. *British Journal of Pharmacology*, 171(13), 3298–3312. <https://doi.org/10.1111/bph.12685>
- Sayed, A., Schierling, W., Troidl, K., Ruding, I., Nelson, K., Apfelbeck, H., ... Schmitz-Rixen, T. (2010). Exercise linked to transient increase in expression and activity of cation channels in newly formed hind-limb collaterals. *European Journal of Vascular and Endovascular Surgery*, 40(1), 81–87. <https://doi.org/10.1016/j.ejvs.2010.02.014>
- Schierling, W., Troidl, K., Apfelbeck, H., Troidl, C., Kasprzak, P. M., Schaper, W., & Schmitz-Rixen, T. (2011). Cerebral arteriogenesis is enhanced by pharmacological as well as fluid-shear-stress activation of the *Trpv4* calcium channel. *European Journal of Vascular and Endovascular Surgery*, 41(5), 589–596. <https://doi.org/10.1016/j.ejvs.2010.11.034>
- Shen, J., Tu, L., Chen, D., Tan, T., Wang, Y., & Wang, S. (2018). TRPV4 channels stimulate Ca^{2+} -induced Ca^{2+} release in mouse neurons and trigger endoplasmic reticulum stress after intracerebral hemorrhage. *Brain Research Bulletin*, 146, 143–152.
- Sonkusare, S. K., Bonev, A. D., Ledoux, J., Liedtke, W., Kotlikoff, M. I., Heppner, T. J., ... Nelson, M. T. (2012). Elementary Ca^{2+} signals through endothelial TRPV4 channels regulate vascular function. *Science*, 336(6081), 597–601. <https://doi.org/10.1126/science.1216283>
- Sonkusare, S. K., Dalsgaard, T., Bonev, A. D., Hill-Eubanks, D. C., Kotlikoff, M. I., Scott, J. D., ... Nelson, M. T. (2014). AKAP150-dependent cooperative TRPV4 channel gating is central to endothelium-dependent vasodilation and is disrupted in hypertension. *Science Signaling*, 7(333), ra66. <https://doi.org/10.1126/scisignal.2005052>
- Strotmann, R., Schultz, G., & Plant, T. D. (2003). Ca^{2+} -dependent potentiation of the nonselective cation channel TRPV4 is mediated by a C-terminal calmodulin binding site. *The Journal of Biological Chemistry*, 278(29), 26541–26549. <https://doi.org/10.1074/jbc.M302590200>

- Sullivan, M. N., Francis, M., Pitts, N. L., Taylor, M. S., & Earley, S. (2012). Optical recording reveals novel properties of GSK1016790A-induced vanilloid transient receptor potential channel TRPV4 activity in primary human endothelial cells. *Molecular Pharmacology*, *82*(3), 464–472. <https://doi.org/10.1124/mol.112.078584>
- Sun, M. Y., Geyer, M., & Komarova, Y. A. (2017). IP3 receptor signaling and endothelial barrier function. *Cellular and Molecular Life Sciences*, *74*(22), 4189–4207. <https://doi.org/10.1007/s00018-017-2624-8>
- Tallini, Y. N., Brekke, J. F., Shui, B., Doran, R., Hwang, S. M., Nakai, J., ... Kotlikoff, M. I. (2007). Propagated endothelial Ca²⁺ waves and arteriolar dilation in vivo: measurements in Cx40BAC GCaMP2 transgenic mice. *Circulation Research*, *101*(12), 1300–1309. <https://doi.org/10.1161/CIRCRESAHA.107.149484>
- Taylor, M. S., Bonev, A. D., Gross, T. P., Eckman, D. M., Brayden, J. E., Bond, C. T., ... Nelson, M. T. (2003). Altered expression of small-conductance Ca²⁺-activated K⁺ (SK3) channels modulates arterial tone and blood pressure. *Circulation Research*, *93*(2), 124–131. <https://doi.org/10.1161/01.RES.0000081980.63146.69>
- Thodeti, C. K., Matthews, B., Ravi, A., Mammoto, A., Ghosh, K., Bracha, A. L., & Ingber, D. E. (2009). TRPV4 channels mediate cyclic strain-induced endothelial cell reorientation through integrin-to-integrin signaling. *Circulation Research*, *104*(9), 1123–1130. <https://doi.org/10.1161/CIRCRESAHA.108.192930>
- Thoppil, R. J., Cappelli, H. C., Adapala, R. K., Kanugula, A. K., Paruchuri, S., & Thodeti, C. K. (2016). TRPV4 channels regulate tumor angiogenesis via modulation of Rho/Rho kinase pathway. *Oncotarget*, *7*(18), 25849–25861. <https://doi.org/10.18632/oncotarget.8405>
- Trebak, M., Bird, G. S., McKay, R. R., & Putney, J. W. Jr. (2002). Comparison of human TRPC3 channels in receptor-activated and store-operated modes. Differential sensitivity to channel blockers suggests fundamental differences in channel composition. *Journal of Biological Chemistry*, *277*(24), 21617–21623. <https://doi.org/10.1074/jbc.M202549200>
- Troidl, C., Troidl, K., Schierling, W., Cai, W. J., Nef, H., Mollmann, H., ... Schaper, W. (2009). Trpv4 induces collateral vessel growth during regeneration of the arterial circulation. *Journal of Cellular and Molecular Medicine*, *13*(8B), 2613–2621. <https://doi.org/10.1111/j.1582-4934.2008.00579.x>
- Voets, T., Prenen, J., Fleig, A., Vennekens, R., Watanabe, H., Hoenderop, J. G., ... Nilius, B. (2001). CaT1 and the calcium release-activated calcium channel manifest distinct pore properties. *The Journal of Biological Chemistry*, *276*(51), 47767–47770. <https://doi.org/10.1074/jbc.C100607200>
- Vriens, J., Watanabe, H., Janssens, A., Droogmans, G., Voets, T., & Nilius, B. (2004). Cell swelling, heat, and chemical agonists use distinct pathways for the activation of the cation channel TRPV4. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(1), 396–401. <https://doi.org/10.1073/pnas.0303329101>
- Watanabe, H., Davis, J. B., Smart, D., Jerman, J. C., Smith, G. D., Hayes, P., ... Nilius, B. (2002). Activation of TRPV4 channels (hVRL-2/mTRP12) by phorbol derivatives. *The Journal of Biological Chemistry*, *277*(16), 13569–13577. <https://doi.org/10.1074/jbc.M200062200>
- Watanabe, H., Vriens, J., Janssens, A., Wondergem, R., Droogmans, G., & Nilius, B. (2003). Modulation of TRPV4 gating by intra- and extracellular Ca²⁺. *Cell Calcium*, *33*(5–6), 489–495. [https://doi.org/10.1016/S0143-4160\(03\)00064-2](https://doi.org/10.1016/S0143-4160(03)00064-2)
- Watanabe, H., Vriens, J., Prenen, J., Droogmans, G., Voets, T., & Nilius, B. (2003). Anandamide and arachidonic acid use epoxyeicosatrienoic acids to activate TRPV4 channels. *Nature*, *424*(6947), 434–438. <https://doi.org/10.1038/nature01807>
- Whorton, A. R., Willis, C. E., Kent, R. S., & Young, S. L. (1984). The role of calcium in the regulation of prostacyclin synthesis by porcine aortic endothelial cells. *Lipids*, *19*(1), 17–24. <https://doi.org/10.1007/BF02534603>
- Willars, G. B., Nahorski, S. R., & Challiss, R. A. (1998). Differential regulation of muscarinic acetylcholine receptor-sensitive polyphosphoinositide pools and consequences for signaling in human neuroblastoma cells. *The Journal of Biological Chemistry*, *273*(9), 5037–5046. <https://doi.org/10.1074/jbc.273.9.5037>
- Wilson, C., Lee, M., & McCarron, J. G. (2016). Acetylcholine released by endothelial cells facilitates flow-mediated dilatation. *The Journal of Physiology*, *594*, 7267–7307. <https://doi.org/10.1113/JP272927>
- Wilson, C., Lee, M. D., Heathcote, H. R., Zhang, X., Buckley, C., Girkin, J. M., ... McCarron, J. G. (2019). Mitochondrial ATP production provides long-range control of endothelial inositol trisphosphate-evoked calcium signaling. *The Journal of Biological Chemistry*, *294*(3), 737–758. <https://doi.org/10.1074/jbc.RA118.005913>
- Wilson, C., Saunter, C. D., Girkin, J. M., & McCarron, J. G. (2015). Pressure-dependent regulation of Ca²⁺ signaling in the vascular endothelium. *The Journal of Physiology*, *593*, 5231–5253. <https://doi.org/10.1113/JP271157>
- Wilson, C., Saunter, C. D., Girkin, J. M., & McCarron, J. G. (2016). Clusters of specialized detector cells provide sensitive and high fidelity receptor signaling in intact endothelium. *The FASEB Journal*, *30*, 2000–2013. <https://doi.org/10.1096/fj.201500090>
- Yin, J., Hoffmann, J., Kaestle, S. M., Neye, N., Wang, L., Baeurle, J., ... Kuebler, W. M. (2008). Negative-feedback loop attenuates hydrostatic lung edema via a cGMP-dependent regulation of transient receptor potential vanilloid 4. *Circulation Research*, *102*(8), 966–974. <https://doi.org/10.1161/CIRCRESAHA.107.168724>
- Zhang, D. X., Mendoza, S. A., Bubolz, A. H., Mizuno, A., Ge, Z. D., Li, R., ... Gutterman, D. D. (2009). Transient receptor potential vanilloid type 4-deficient mice exhibit impaired endothelium-dependent relaxation induced by acetylcholine in vitro and in vivo. *Hypertension*, *53*(3), 532–538. <https://doi.org/10.1161/HYPERTENSIONAHA.108.127100>
- Zhang, L., Papadopoulos, P., & Hamel, E. (2013). Endothelial TRPV4 channels mediate dilation of cerebral arteries: Impairment and recovery in cerebrovascular pathologies related to Alzheimer's disease. *British Journal of Pharmacology*, *170*(3), 661–670. <https://doi.org/10.1111/bph.12315>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Heathcote HR, Lee MD, Zhang X, Saunter CD, Wilson C, McCarron JG. Endothelial TRPV4 channels modulate vascular tone by Ca²⁺-induced Ca²⁺ release at inositol 1,4,5-trisphosphate receptors. *Br J Pharmacol*. 2019;1–21. <https://doi.org/10.1111/bph.14762>