

Nernst Advection and the Field-Generating Thermal Instability Revisited

J. J. Bissell^{1,2}†

¹Department of Mathematical Sciences, University of Durham, DH1 3LE, England

²Blackett Laboratory, Imperial College London, SW7 2BZ, England

(Received ?; revised ?; accepted ?. - To be entered by editorial office)

It is widely held that the Nernst effect can drive instability in un-magnetised laser-plasmas by laterally compressing seed \mathbf{B} -fields arising from the field-generating thermal instability [Tidman & Shanny, *Phys. Fluids*, **12**:1207 (1974)]. Indeed, for wavelike perturbations, differential compression by the Nernst mechanism is thought to be most pronounced in the limit of low wave-number $k \rightarrow 0$, and is considered particularly important given that it can ostensibly lead to instability when the more usual field-generating mechanism is stable. However, as part of a recent article [Bissell *et al.*, *New J. Phys.*, **15**:025017 (2013)] we noted some irregularities to the Nernst mechanism which obscure its operation. For example, by taking characteristic density and temperature length-scales l_n and l_T respectively, we observed that consistent analytical treatment of the instability requires $kl_{n,T} \gg 1$, preventing the peak-growth limit $k \rightarrow 0$. Furthermore, the Nernst term—which compresses magnetic field perturbations—does not couple to a corresponding term acting on thermal perturbations, and as such does not describe an unstable feedback mechanism. In this article we probe the origin of such ambiguities more formally, and in so doing argue (contrary to reports existing elsewhere in the literature) that the Nernst effect does not drive instability in un-magnetised conditions, at least not in the fashion typically cited.

PACS codes: Authors should not enter PACS codes directly on the manuscript, as these must be chosen during the online submission process and will then be added during the typesetting process (see <http://www.aip.org/pacs/> for the full list of PACS codes)

1. Introduction

It has long been known that otherwise un-magnetised plasmas can self-generate large magnetic fields ($\sim 100\text{T}$) through a variety of mechanisms (Stamper *et al.* 1971; Pert 1977; Raven *et al.* 1978; Haines 1986a; Thomas *et al.* 2009; Li *et al.* 2013). These fields strongly affect electron transport by suppressing the cross-field thermal conductivity (Braginskii 1965) and are thus key to understanding a range of laser-plasma interactions, including ongoing efforts to achieve controlled inertial confinement fusion (Glenzer *et al.* 1999; Lindl *et al.* 2004; Nilson *et al.* 2006; Froula *et al.* 2007; Li *et al.* 2007a,b; Schurtz *et al.* 2007; Froula *et al.* 2009; Li *et al.* 2009, 2013). Of special importance in such contexts is the role transport effects might play in driving instabilities, especially given that such instabilities are themselves often candidate mechanisms for producing the self-generated field (Weibel 1959; Tidman & Shanny 1974; Bol'shov *et al.* 1974; Ogasawara *et al.* 1980; Haines 1981; Bissell *et al.* 2010, 2012; Gao *et al.* 2012; Manuel *et al.* 2013).

† Email address for correspondence: john.bissell@durham.ac.uk

As part of a recent article (Bissell *et al.* 2013) we discussed how super-Gaussian transport effects arising from strong inverse bremsstrahlung (I.B.) heating can suppress growth-rates of one such candidate, the *field-generating thermal instability*, which was first reported in 1974 (Tidman & Shanny 1974; Bol’shov *et al.* 1974), and remains an important phenomena in laser-plasma interactions (see, for example, experimental studies of coronal plasmas by Manuel *et al.* (2013) reported earlier this year). It is widely held that this instability may be driven by two mechanisms which—denoting the electron temperature and density as T_e and n_e respectively, and taking $\mathbf{b} = \mathbf{B}/|\mathbf{B}|$ as a unit vector in the direction of the magnetic flux density \mathbf{B} —may be summarised as follows. In the first case (i) feedback acts between $\nabla T_e \times \nabla n_e$ generated field and the consequent cross-gradient Righi-Leduc heat-flow $\mathbf{q}_\perp \propto \mathbf{b} \times \nabla T_e$ (Haines 1986a; Braginskii 1965; Tidman & Shanny 1974; Bol’shov *et al.* 1974). In the second (ii) it is commonly understood that lateral advection of the field with the diffusive heat-flow $\mathbf{q}_\perp \propto \nabla T_e$ *via* the Nernst effect (Nishiguchi *et al.* 1985; Bissell *et al.* 2013), i.e., with velocity $\mathbf{v}_N \approx \mathbf{q}_\perp / (\frac{3}{2} n_e T_e)$, can lead to exponential compression of the perturbation in \mathbf{B} (Brownell 1979; Hirao & Ogasawara 1981). In the absence of hydrodynamic effects, case (i) requires that zeroth-order temperature and density gradients be parallel, i.e., $l_T l_n > 0$, where in an x -coordinate aligned geometry the length scales l_T and l_n may be defined

$$\frac{1}{l_T} = \frac{1}{T_0} \frac{\partial T_0}{\partial x} \quad \text{and} \quad \frac{1}{l_n} = \frac{1}{n_0} \frac{\partial n_0}{\partial x}, \quad (1.1)$$

respectively, with the subscripts ‘0’ denoting zeroth-order profiles (see §3). This feature of case (i) makes the mechanism in case (ii) particularly important, since Nernst advection may both *contribute* to instability when $l_T l_n > 0$ holds, but also (ostensibly) *drive* instability when the parallel gradient condition fails, that is, if $l_n l_T < 0$ (Brownell 1979; Hirao & Ogasawara 1981).

Our original comments (Bissell *et al.* 2013) on the field-generating thermal instability focused on how super-Gaussian modifications to electron transport can suppress classically predicated instability growth-rates (by as much as $\sim 80\%$ under both $l_T l_n > 0$ and $l_T l_n < 0$ conditions). For this reason, our analysis followed in the tradition of previous work in its treatment of the Nernst advection terms (Brownell 1979; Hirao & Ogasawara 1981); however, we noted a curious feature of the Nernst mechanism in that it seems to predict peak instability growth-rates as the perturbation wave-number k vanishes. When trying to develop a mathematically consistent picture of the instability, this feature is problematic because analytical treatments typically assume some local conditions $l_{n,T} k \gg 1$ for the unstable modes, precluding $k \rightarrow 0$ (Tidman & Shanny 1974; Bol’shov *et al.* 1974; Ogasawara *et al.* 1980; Brownell 1979; Hirao & Ogasawara 1981). Furthermore, the Nernst term (which compresses the field perturbations) does not couple to a corresponding term acting on thermal perturbations, meaning that mechanism (ii) does not account for unstable feedback in the usual way.

Given the need for clarity when establishing the stability of laser-plasma configurations, this short article revisits the impact of the Nernst effect on the field-generating thermal instability, placing special emphasis on resolving the ambiguities described above. By reviewing the basic instability theory (§2 and §3) we identify various inconsistencies of treatment which imply that Nernst advection does not drive instability under the linear assumptions usually made (Brownell 1979; Hirao & Ogasawara 1981), and consequently (in the absence of hydrodynamic motion, and a more advanced treatment of lateral effects) that laser-plasmas should in fact be stable to the field-generating thermal instability mechanism whenever $l_n l_T < 0$ (§4). In the light of these observations, we also comment on the role Nernst plays in driving instability more generally (§5 and §6).

2. Governing Equations

Since we are primarily interested in Nernst advection effects we neglect hydrodynamics, and the governing equations are simply the thermal energy equation and Faraday's law,

$$\frac{3}{2}n_e\frac{\partial T_e}{\partial t} + \nabla \cdot \mathbf{q} - \mathbf{E} \cdot \mathbf{j} = \dot{U}_L \quad \text{and} \quad \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} \quad (2.1)$$

respectively, where \dot{U}_L describes the rate-of-change of thermal energy $U = \frac{3}{2}n_eT_e$ due to laser heating, and $\mathbf{j} = (\nabla \times \mathbf{B})/\mu_0$ is the current (with μ_0 as the permeability of free-space), while the electric field \mathbf{E} and total heat-flow \mathbf{q} are given by Braginskii (1965):

$$en_e\mathbf{E} = -\nabla P_e + \mathbf{j} \times \mathbf{B} + \frac{m_e}{e\tau_B}\underline{\underline{\alpha}}^c \cdot \mathbf{j} - n_e\underline{\underline{\beta}}^c \cdot \nabla T_e \quad (2.2)$$

$$\text{and} \quad \mathbf{q} = -\frac{n_e\tau_B T_e}{m_e}\underline{\underline{\kappa}}^c \cdot \nabla T_e - \underline{\underline{\psi}}' \cdot \mathbf{j} \frac{T_e}{e}. \quad (2.3)$$

Here the usual notation applies (Bissell *et al.* 2013), so that e is the elementary electronic charge, $P_e = n_eT_e$ is the isotropic pressure, m_e is the electronic mass, and the Braginskii collision time $\tau_B = c_B\tau_T$ is proportional to the thermal collision time τ_T by the constant factor $c_B = 3\sqrt{\pi}/4$. By quasi-neutrality, the ion number density may be written in terms of the atomic number Z as $n_i \approx n_e/Z$, so that with permittivity of free space ϵ_0 , and Coulomb logarithm $\log \Lambda_{ei} \approx 8$, τ_T is itself given by $\tau_T = (4\pi v_T^3)/(n_i[Ze^2/\epsilon_0 m_e]^2 \log \Lambda_{ei})$, where $v_T = (2T_e/m_e)^{1/2}$ is the mean thermal velocity and defines a thermal mean-free-path $\lambda_T = v_T\tau_T$. As usual in such treatments, we shall a geometry for which gradients and fluxes are perpendicular to \mathbf{B} , i.e, scalar ϕ and vector \mathbf{A} quantities obey $\mathbf{B} \cdot \nabla \phi = \mathbf{B} \cdot \mathbf{A} = 0$. Under these conditions, the resistivity $\underline{\underline{\alpha}}^c$, conductivity $\underline{\underline{\kappa}}^c$, and thermoelectric tensors $\underline{\underline{\beta}}^c$ and $\underline{\underline{\psi}}' = \underline{\underline{\beta}}^c + [5/2]\mathbf{I}$ (here \mathbf{I} is the identity tensor) may be written in the general form

$$\underline{\underline{\eta}} \cdot \mathbf{s} = \eta_{\perp} \mathbf{s} + \eta_{\wedge} \mathbf{b} \times \mathbf{s}, \quad \text{with} \quad \eta \in \{\alpha, \beta, \kappa\} \quad \text{and} \quad \mathbf{s} \in \{\mathbf{j}, \nabla T_e\}, \quad (2.4)$$

where the components η_{\perp} and η_{\wedge} are dimensionless functions of both Z and the Hall parameter $\chi = \omega_L\tau_B$, with $\omega_L = (e|\mathbf{B}|/m_e)$ as the electron Larmor frequency (Braginskii 1965; Epperlein 1984; Epperlein & Haines 1986). Note that for our un-magnetised conditions the ' \perp ' components are constants, that is, $\eta_{\perp}(0) = \eta_{\parallel}$, while convention dictates that equation 2.4 takes the slightly different form $\underline{\underline{\alpha}} \cdot \mathbf{j} = \alpha_{\perp} \mathbf{j} - \alpha_{\wedge} \mathbf{b} \times \mathbf{j}$ for the resistivity.

3. Review of the Basic Linear Theory

Let us begin our analysis by reviewing the basic linear theory. As usual, we assume that the plasma is initially un-magnetised, so that in zeroth-order our governing equations—the thermal energy equation and induction equation (Faraday's Law and Ohm's Law combined)—are satisfied by solutions $\mathbf{B} = \mathbf{B}_0 = \mathbf{0}$, $T_e = T_0(x, t)$ and $n_e = n_0(x, t)$, where gradients in the latter two quantities exist in the x -direction only, and define length-scales l_n and l_T according to equations 1.1. In addition, we note that for laser-plasmas one may take

$$\Lambda = \frac{\lambda_T}{\delta} \gg 1, \quad \text{where} \quad \delta = \frac{c}{\omega_{pe}} \quad \text{and} \quad \omega_{pe} = \left(\frac{n_e e^2}{\epsilon_0 m_e} \right)^{1/2} \quad (3.1)$$

are the collisionless-skin-depth and plasma frequency respectively, and $c = 1/\sqrt{\mu_0\epsilon_0}$ is the speed of light *in vacuo*.

To the zeroth-order solutions we add small wavelike perturbations with wavenumber

k , frequency γ , and periodicity $\propto (iky + \gamma t)$, that is,

$$T_e = T_0 + \delta T \exp(iky + \gamma t) \quad \text{and} \quad \mathbf{B} = \delta B \exp(iky + \gamma t) \hat{\mathbf{z}}, \quad (3.2)$$

with $\delta T \ll T_0$ and δB as some complex amplitudes. Hence, after substituting these perturbed forms into the governing system, subtracting zeroth-order solutions, and neglecting second-order perturbed terms or higher, the linearised energy and induction equations become

$$\Gamma + D_T K^2 - Q = +iC_E \left(\frac{e\lambda_T^2}{\tau_T} \frac{\delta B}{\delta T} \right) K \quad (3.3)$$

and

$$\Gamma + D_R K^2 - N = -iC_I \left(\frac{\tau_T}{e\lambda_T^2} \frac{\delta T}{\delta B} \right) K \quad (3.4)$$

respectively. Notice here that we have adopted the dimensionless notation

$$\Gamma = \gamma\tau_T, \quad K = \lambda_T k, \quad L_{T,n} = \frac{l_{T,n}}{\lambda_T}, \quad D_T = \frac{c_B}{3} \kappa_{\parallel}^c, \quad D_R = \frac{\alpha_{\parallel}^c}{c_B \Lambda^2}, \quad (3.5)$$

where D_T and D_R are the thermal and resistive diffusion coefficients respectively (Bissell *et al.* 2012). The remaining terms arise from: i) differential thermal diffusion Q lateral to the perturbation (down zeroth-order temperature gradients); ii) differential Nernst advection N lateral to the perturbation; iii) divergent Righi-Leduc heat-flow C_E ; and iv) magnetic field generation C_I by the $\nabla T_e \times \nabla n_0$ mechanism; these are defined by

$$Q = \frac{5}{2} D_T \frac{\lambda_T^2}{n_0 \tau_T T_0} \frac{\partial}{\partial x} \left(n_0 \tau_T \frac{\partial T_0}{\partial x} \right) \sim D_T / L_{T,n}^2, \quad C_E = \frac{c_B^2}{6L_T} \left(\frac{\partial \kappa_{\Lambda}^c}{\partial \chi} \right), \quad (3.6)$$

$$N = \frac{c_B}{2} \frac{\partial \beta_{\Lambda}^c}{\partial \chi} \frac{\lambda_T^2}{\tau_T T_0} \frac{\partial}{\partial x} \left(\tau_T \frac{\partial T_0}{\partial x} \right), \quad \text{and} \quad C_I = \frac{1}{L_n}. \quad (3.7)$$

At this stage in the usual analysis one assumes the local condition $KL_{n,T} \gg 1$ (Brownell 1979; Hirao & Ogasawara 1981), so that the lateral diffusion term $Q \sim D_T / L_{T,n}^2$ may be neglected in equation 3.3 when compared to $D_T K^2$, i.e.,

$$[D_T K^2 - Q] = D_T K^2 [1 - O(1/K^2 L_{T,n}^2)] \approx D_T K^2. \quad (3.8)$$

Thus, after eliminating terms in $(\delta T / \delta B)$ from equations 3.3 and 3.4, one obtains the dispersion relation (Bissell *et al.* 2013; Brownell 1979; Hirao & Ogasawara 1981)

$$\Gamma_{\pm} = \frac{1}{2} \left\{ -[(D_T + D_R)K^2 - N] \pm \sqrt{[(D_T + D_R)K^2 - N]^2 + 4D_T K^2 [D_R(K_G^2 - K^2) + N]} \right\}, \quad (3.9)$$

where K_G is the source term describing coupling between the Righi-Leduc heat-flow C_E and $\nabla T_e \times \nabla n_e$ field generation C_I as described by Tidman & Shanny (1974) and Bissell *et al.* (2012), *viz*

$$K_G^2 = \frac{C_E C_I}{D_T D_R} = \frac{c_B^2 \Lambda^2}{2L_T L_n \alpha_{\parallel}^c \kappa_{\parallel}^c} \left(\frac{\partial \kappa_{\Lambda}^c}{\partial \chi} \right). \quad (3.10)$$

Equation 3.9 is the dispersion relation quoted by the author in our earlier context (Bissell *et al.* 2013) where we observed that in the absence of Nernst advection ($N \rightarrow 0$) the cut-off wave-number becomes K_G , so that instability requires $L_T L_n > 0$ (Tidman &

Shanny 1974; Brownell 1979; Hirao & Ogasawara 1981). In terms of plasma stability, this result makes the contribution from the advection term N particularly important, since it ostensibly predicts instability when $L_T L_n < 0$ provided $N > 0$ (Brownell 1979; Hirao & Ogasawara 1981). As both Brownell (1979) and Hirao & Ogasawara (1981) describe, equation 3.9 suggests that lateral compression of the field by the advection term should be especially effective in the limit of long-wavelength perturbations $K \rightarrow 0$ (when damping effects, such as thermal diffusion, are minimised), in which case the peak growth-rate becomes $\Gamma = N$. Of course, the value $K = 0$ is not a physically meaningful one to take for the peak wave-number; however, as we noted in our original context (Bissell *et al.* 2013), the apparent mathematical inconsistency between the local assumption $K L_{n,T} \gg 1$ made in the analysis, and the (ideal) conclusion that instability peaks when $K = 0$, invites us to treat the basic theory with some caution. What is more, Nernst compression by N does not describe coupling between the energy and induction equations, and so fails to provide a mechanism for unstable feedback. For these reasons we stated that the Nernst term cannot be understood as driving instability *proper*, though it may lead to exponential compression of the local field. In what follows we consider the source of such ambiguities in more formal detail by re-examining the basic linear theory, and reflecting further on the physical basis of the Nernst effect. Ultimately, we conclude by taking the opposing view that Nernst will not drive instability in un-magnetised conditions, at least not without a more sophisticated treatment of lateral effects.

4. Inconsistencies in the Basic Linear Theory

In essence, the ambiguities described in the previous section arise from miss-application of the local approximation; however, it is often as instructive to consider the shortcomings of physical arguments as their advantages, and to this end it is worth discussing such miss-application in relative detail. Indeed, by probing related problems in the basic linear theory (Brownell 1979; Hirao & Ogasawara 1981), one can gain insight into both the meaning of various analytical steps, and the physical processes involved. Here three issues are considered: first, physical interpretation (§4.1); second, ordering of terms (§4.2); and third, lateral effects (§4.3).

4.1. Physical Interpretation

Recall that the Nernst effect describes advection of the magnetic field with the diffusive heat-flow \mathbf{q}_\perp , i.e., the heat-flow representing thermal diffusion down temperature gradients (Nishiguchi *et al.* 1985; Bissell *et al.* 2013). Indeed, one may write the velocity \mathbf{v}_N of Nernst advection as

$$\mathbf{v}_N = \frac{2A_N \mathbf{q}_\perp}{3D_T P_e} \approx \frac{\mathbf{q}_\perp}{3n_e T_e / 2}, \quad \text{where} \quad \mathbf{q}_\perp = -\frac{3}{2} n_e D_T \frac{\lambda_T^2}{\tau_T} \nabla T_e \quad (4.1)$$

is the diffusive heat-flow, A_N is a dimensionless *Nernst advection coefficient* defined by

$$A_N = \frac{c_B}{2\chi} \beta_\wedge^c \approx D_T = \frac{c_B}{3} \kappa_\perp^c, \quad (4.2)$$

and the approximate equality follows by considering values for the transport coefficients over a range of χ (Haines 1986b; Bissell *et al.* 2012). Physically, therefore, our term N in the linearised induction equation 3.4 arises from differential advection of the field perturbation by heat-flow electrons moving down the zeroth-order temperature gradient, i.e., those electrons which account for lateral diffusive transport of the temperature perturbation Q in the linearised energy equation 3.3. It seems curious that we should retain N

(lateral \mathbf{B} -field advection) in the dispersion relation 3.9, whilst simultaneously arguing for the exclusion of the thermal term $Q \sim D_T/L_{T,n}^2$ responsible for its physical origin.

4.2. Ordering of Terms

That such a contradiction arises may be understood by reviewing the local approximation $KL_{n,T} \gg 1$; as we saw in equation 3.8, should this condition hold it is legitimate to neglect terms in $Q \sim D_T/L_{n,T}$ when compared to those in D_T in the linearised energy equation 3.3. However, closer inspection of the Nernst term reveals that the local approximation also applies to N . Indeed, for our un-magnetised conditions the cross-field thermoelectric coefficient is directly proportional to the Hall parameter, i.e., $\beta_\lambda^c \propto \chi$ (Braginskii 1965), so that

$$\frac{\partial \beta_\lambda^c}{\partial \chi} = \frac{\beta_\lambda^c}{\chi} \quad \Rightarrow \quad N = A_N \frac{\lambda_T^2}{\tau_T T_0} \frac{\partial}{\partial x} \left(\tau_T \frac{\partial T_0}{\partial x} \right) \sim A_N / L_{T,n}^2, \quad (4.3)$$

and thus, since $D_R \approx A_N$, we have (cf. equation 3.8 and 3.9)

$$[D_T K^2 - N] = D_T K^2 [1 - O(1/K^2 L_{T,n}^2)] \approx D_T K^2. \quad (4.4)$$

Consistent application of the local condition $KL_{n,T} \gg 1$ to neglect Q in equation 3.3 therefore demands that we neglect N in the dispersion relation (equation 3.9). That we cannot do so at the stage of linearising the induction equation 3.4 is a consequence of the relatively small value of the resistive diffusion coefficient D_R when compared to the thermal diffusion coefficient D_T (by the factor $1/\Lambda^2$, cf. equations 3.1 and 3.5).

4.3. Lateral Effects

Our discussion in the previous two sections demonstrates that (as a consequence of their common physical basis) the term Q arising from the divergence of the heat-flow down zeroth-order gradients has the same functional form as N . Furthermore, equations 3.8 and 4.4 establish that *both* Q and N should be properly neglected in the theory provided that the local condition $KL_{n,T} \gg 1$ holds. It is natural to wonder, therefore, whether these terms should be retained in the analysis under conditions for which the local approximation $KL_{n,T} \gg 1$ does not apply, in which case we might expect Q to modify the growth-rate alongside N by acting as an additional source of instability.

To address this supposition, let us consider more carefully the role played by the local condition in the process of linearisation. Strictly speaking, the proper form for the perturbations applied in §3 should postulate some x -dependence, and this is especially true when we are interested in differential advection or diffusion, because any x -dependence will have consequences for the lateral compression (or rarefaction) of the perturbations normal to the wave-vector. To account for such lateral effects, therefore, one must set $\delta T \equiv \delta T(x)$ and $\delta B \equiv \delta B(x)$, and in this case equations 3.3 and 3.4 acquire new terms. For example, after neglecting terms of order $(\delta T)^2$ and higher, the thermal diffusion component to equation 3.3 undergoes the transformation

$$D_T K^2 \left[1 - \frac{Q}{D_T K^2} \right] \rightarrow D_T K^2 \left[1 - \frac{Q}{D_T K^2} + O\left(\frac{\lambda_T^2}{K^2 \delta T} \frac{d^2 \delta T}{dx^2} \right) + O\left(\frac{\lambda_T}{K^2 L_T \delta T} \frac{d \delta T}{dx} \right) \right]. \quad (4.5)$$

Thus, our perturbation equations 3.3 and 3.4 become coupled second-order ordinary differential equations reminiscent of those encountered in problems involving hydrodynamic stability (Chandrasekhar 1961). Indeed, although the final two differential terms describe gradients in small quantities, because the $\nabla T_e \times \nabla n_0$ field generating mechanism responsible for driving initial growth of the instability operates over the density length-scale l_n ,

one expects both $(d\delta T/dx)/\delta T \sim 1/l_n$ and $(d^2\delta T/dx^2)/\delta T \sim 1/l_n^2$. The differential terms will thus be of a similar order to the lateral heat-flow term in $Q/D_T \sim 1/L_{T,n}^2$. In this way, we see that if the local condition does not hold, then one is obliged to retain not only Q , but also the differential terms describing gradients in δT , and the eigenvalue problem becomes one requiring solution of a non-linear system of coupled ordinary differential equations with eigenfunction solutions δT and δB dependent on appropriate boundary conditions (cf. Chandrasekhar (1961)). In general such a problem will be non-trivial.

On the other hand, should the local approximation hold, then its utility is now clear. Provided $KL_{n,T} \gg 1$ we can effect a ‘secondary linearisation process’ whereby we discard the differential terms in equation 4.5 (and similar), and thence solve for the growth rate Γ algebraically; in so doing, however, consistency requires us to surrender terms in both Q and N (as described in §4.2). To address the supposition made at the beginning of this section directly, therefore, we conclude that while it is possible for laterally divergent transport to drive instability when the local approximation does not apply, proper treatment of such effects requires a more sophisticated analysis than that considered here; it is not sufficient simply to add terms in Q and N to the dispersion relation 3.9. Indeed, substantial further investigation inclusive of numerical simulations is needed to better determine the role played by lateral effects in de-stabilising laser plasmas under such conditions.

5. Super-Gaussian Transport and Suppression of Instability

Although our argument here has focused on classical (Braginskii) transport effects, it remains for us to comment briefly on how the loss of the advection mechanism N affects those results stated in our earlier context (Bissell *et al.* 2013) where the primary concern was super-Gaussian transport phenomena arising from strong inverse-bremsstrahlung heating. One obvious consequence is stabilisation of the plasma to the field-generating instability when $l_T l_n < 0$. Nevertheless, the reduction of both $\nabla T_e \times \nabla n_e$ field generation and the Righi-Leduc heat-flow when super-Gaussian transport applies, means that our original predictions (Bissell *et al.* 2013), i.e., heavy suppression of instability growth-rates due to I.B. (by as much as $\sim 80\%$), remain valid for $l_T l_n > 0$. Of course, if hydrodynamic motion is included in the analysis, then instability can prevail without Nernst for $l_T l_n < 0$ (Ogasawara *et al.* 1980); however, in this case we would continue to expect significant growth-rate suppression by I.B. heating because super-Gaussian transport phenomena are unaffected by hydrodynamic flow (Bissell *et al.* 2013).

6. Conclusion

The usual treatment of Nernst advection effects (Brownell 1979; Hirao & Ogasawara 1981) on the field-generating thermal instability (Tidman & Shanny 1974) leads to ambiguities in the dispersion relation for the growth of unstable modes. For example, and as we noted in a recent article (Bissell *et al.* 2013), lateral compression by the Nernst effect ostensibly yields peak-growth rates which correspond to vanishing wavenumber $k \rightarrow 0$, violating the local approximation $kl_{n,T} \gg 1$ (§1 and §3). In this short article we have sought to determine the source of such ambiguities more formally, both by re-examining the basic analysis, and returning to the physical meaning of the Nernst effect (§2 and §3). In particular, we argued that consistent (and necessary) application of the local approximation requires the Nernst advection term to be omitted from the dispersion relation 3.9 (§3 and §4), meaning that for un-magnetised conditions Nernst cannot drive instability in the fashion commonly cited (Brownell 1979; Hirao & Ogasawara 1981). One

consequence of such an interpretation is that—in the absence of significant hydrodynamic motion (Ogasawara *et al.* 1980)—un-magnetised laser-plasmas should be stable to the field-generating instability whenever zeroth-order temperature and density gradients are anti-parallel, i.e., $l_n l_T < 0$ (§5).

Naturally, we do not go so far as to state that Nernst advection can never drive instability, since it is possible that further investigation of lateral effects might lead to a compressive mechanism precluded by the approach taken here; though such an investigation would require solving a more complex eigenvalue problem formulated in terms of second-order differential equations, in combination with a thorough numerical investigation, and is therefore left as future research (§4.3). Indeed, theoretical and computational work reported elsewhere has shown that the Nernst effect is expected to drive a related *field-compressing magneto-thermal instability* in laser-plasmas under sufficiently magnetised conditions (Bissell *et al.* 2010, 2012).

The author would like to thank two anonymous referees for insightful comments on the original manuscript which have helped to clarify its argument. JJB is supported by a Leverhulme Trust Grant (*Tipping Points Project*, University of Durham).

REFERENCES

- BISSELL, J. J., RIDGERS, C. P. & KINGHAM, R. J. 2010 Field Compressing Magnetothermal Instability in Laser Plasmas. *Phys. Rev. Lett.* **105**(17):175001.
- BISSELL, J. J., KINGHAM, R. J. & RIDGERS, C. P. 2012 Magnetothermal instability in laser plasmas including hydrodynamic effects. *Physics of Plasmas*. **19**(5):052107.
- BISSELL, J. J., RIDGERS, C. P. & KINGHAM, R. J. 2013 Super-Gaussian transport theory and the field-generating thermal instability in laser-plasmas. *New J. Phys.* **15**:025017.
- BOL'SHOV, L. A., DREIZIN, Y. A. & DYKHNE, A. M. 1974 Spontaneous magnetization of electronic thermal conductivity in a laser plasma. *JETP Letters*. **19**(5):168–170.
- BRAGINSKII, S. I. 1965 Transport processes in a plasma. *Reviews of Plasma Physics*. **1**:205.
- BROWNELL, J. H. 1979 Magnetic Field Generation in Plasma Implosions due to Nernst Refrigeration. *Comments on Plasma Physics and Controlled Fusion*. **4**(5):131–138.
- CHANDRASEKHAR, S. 1961 (Dover Edition 1981) *Hydrodynamic and Hydromagnetic Stability*.
- EPPERLEIN, E. M. 1984 The accuracy of Braginskii's transport coefficients for a Lorentz plasma. *Journal of Physics D: Applied Physics*. **17**(9):1823–1827.
- EPPERLEIN, E. M. & HAINES, M. G. 1986 Plasma transport coefficients in a magnetic field by direct numerical simulation of the Fokker-Planck equation. *Physics of Fluids*. **29**(4):1029–1041.
- FROULA, D. H., ROSS, J. S., POLLOCK, B. B., DAVIS, P., JAMES, A. N., DIVOL, L., EDWARDS, M. J., OFFENBERGER, A. A., PRICE, D., TOWN, R. P. J., TYNAN, G. R., & GLENZER, S. H. 2007 Quenching of the Nonlocal Electron Heat Transport by Large External Magnetic Fields in a Laser-Produced Plasma Measured with Imaging Thomson Scattering. *Phys. Rev. Lett.* **98**:135001.
- FROULA, D. H., DIVOL, L., DAVIS, P., PALASTRO, J. P., MICHEL, P., LEURENT, V., GLENZER, S. H., POLLOCK, B. B. & TYNAN, G. 2009 Magnetically controlled plasma waveguide for laser wakefield acceleration. *Plasma Physics and Controlled Fusion*. **51**(2):024009.
- GAO, L., NILSON, P. M., IGUMENSHEV, I. V., HU, S. X., DAVIES, J. R., STOECKL, C., HAINES, M. G., FROULA, D. H., BETTI, R. & MEYERHOFER, D. D. 2012 Magnetic Field Generation by the Rayleigh-Taylor Instability in Laser-Driven Planar Plastic Targets. *Phys. Rev. Lett.*, **109**:115001.
- GLENZER, S. H., ALLEY, W. E., ESTABROOK, K. G., DE GROOT, J. S., HAINES, M. G., HAMMER, J. H., JADAUD, J. P., MACGOWAN, B. J., MOODY, J. D., ROZMUS, W., SUTER, L. J., WEILAND, T. L. & WILLIAMS, E. A. 1999 Thomson scattering from laser plasmas. *Physics of Plasmas*. **6**(5):2117–2128.
- HAINES, M. G. 1981 Thermal Instability and Magnetic Field Generation by Large Heat Flow in a Plasma, Especially Under Laser-Fusion Conditions. *Phys. Rev. Lett.* **74**:917.

- HAINES, M. G. 1986a Magnetic field generation in laser fusion and hot-electron transport. *Canadian Journal of Physics*. **64**:912–918.
- HAINES, M. G. 1986b Heat Flux Effects in Ohms Law. *Plasma Physics and Controlled Fusion*. **28**(11):1705–1716.
- HIRAO, A. & OGASAWARA, M. 1981 Magnetic Field Generating Thermal Instability Including the Nernst Effect. *Journal of the Physical Society of Japan*. **50**(2):668–672.
- LI, C. K., SÉGUIN, F. H., FRENJE, J. A., RYGG, J. R. & PETRASSO, R. D. 2007 Observation of the Decay Dynamics and Instabilities of Megagauss Field Structures in Laser-Produced Plasmas. *Phys. Rev. Lett.* **99**(1):015001.
- LI, C. K., SÉGUIN, F. H., FRENJE, J. A., RYGG, J. R. & PETRASSO, R. D. 2007 Observation of Megagauss-Field Topology Changes due to Magnetic Reconnection in Laser-Produced Plasmas *Phys. Rev. Lett.*, **99**(5):055001.
- LI, C. K., FRENJE, J. A., PETRASSO, R. D. SÉGUIN, AMENDT, P. A., LANDEN, O. L., TOWN, R. P. J., BETTI, R., KNAUER, J. P., MEYERHOFER, D. D. & SOURES, J. M. 2009 Pressure-driven, resistive magnetohydrodynamic interchange instabilities in laser-produced high-energy-density plasmas. *Phys. Rev. E*. **80**(1):016407.
- LI, C. K., ZYLSTRA, A. B., FRENJE, J. A., SÉGUIN, F. H., SINENIAN, N., PETRASSO, R. D., AMENDT, P. A., BIONTA, R., FRIEDRICH, S., COLLINS, G. W. DEWALD, E., DÖPPNER, T., GLENZER, S. H., HICKS, D. G., LANDEN, O. L., KILKENNY, J. D., MACKINNON, A. J., MEEZAN, N., RALPH, J., RYGG, J. R., KLINE, J. & KYRALA, G. 2013 Observation of strong electromagnetic fields around laser-entrance holes of ignition-scale hohlraums in inertial-confinement fusion experiments at the National Ignition Facility. *New J. Phys.* **15**:025040.
- LINDL, J. D., AMENDT, P., BERGER, R. L., GLENDINNING, S. G., GLENZER, S. H., HAAN, S. W., KAUFFMAN, R. L., LANDEN, O. L. & SUTER, L. J. 2004 The physics basis for ignition using indirect-drive targets on the National Ignition Facility *Physics of Plasmas*. **11**(2):339–491.
- NILSON, P. M., WILLINGALE, L., KALUZA, M. C., KAMPERIDIS, C. MINARDI, S., WEI, M. S., FERNANDES, P., NOTLEY, M., BANDYOPADHYAY, S., SHERLOCK, M., KINGHAM, R. J., TATARAKIS, M., NAJMUDIN, Z., ROZMUS, W., EVANS, R. G., HAINES, M. G., DANGOR, A. E., & KRUSHELNICK, K. 2006 Magnetic Reconnection and Plasma Dynamics in Two-Beam Laser-Solid Interactions *Phys. Rev. Lett.* **97**(25):255001.
- MANUEL, M. J. E., LI, C. K., SÉGUIN, F. H., SINENIAN, N., FRENJE, J. A., CASEY, D. T., PETRASSO, R. D., HAGER, J. D., BETTI, R. HU, S. X., DELETTREZ, J., & MEYERHOFER, D. D. 2013 Instability-driven electromagnetic fields in coronal plasmas. *Physics of Plasmas*, **20**:056301.
- NISHIGUCHI, A., YABE, T. & HAINES, M. G. 1985 Nernst effect in laser-produced plasmas. *Physics of Fluids*. **28**(12):3683–3690.
- OGASAWARA, M., HIRAO, A. & OHKUBO, H. 1980 Hydrodynamic effects on field-generating thermal instability in laser-heated plasma. *Journal of the Physical Society of Japan*. **49**(1):322–326,
- PERT, G. J. 1977 Self-generated magnetic fields in plasmas. *J. Plasma Phys.* **18**:227–241.
- RAVEN, A., WILLI, O., & RUMSBY, P. T. 1978 Megagauss Magnetic Field Profiles in Laser-produced Plasmas. *Phys. Rev. Lett.* **41**:554.
- SCHURTZ, G., GARY, S., HULIN, S., CHENAIS-POPOVICS, C., GAUTHIER, J. C., THAIS, F., BREIL, J., DURUT, F., FEUGEAS, J. L., MAIRE, P. H., NICOLAÏ, P., PEYRUSSE, O., REVERDIN, C., SOULLIÉ, G., TIKHONCHUK, V., VILLETTE, B. & FOURMENT, C. 2007 Revisiting Nonlocal Electron-Energy Transport in Inertial-Fusion Conditions. *Phys. Rev. Lett.* **98**:095002.
- STAMPER, J. A., PAPADOPOULOS, K., SUDAN, R. N., DEAN, S. O., MCLEAN, E. A. & DAWSON, J. M. 1971 Spontaneous Magnetic Fields in Laser-Produced Plasmas. *Phys. Rev. Lett.* **26**:1012.
- THOMAS, A. G. R., KINGHAM, R. J. & RIDGERS, C. P. 2009 Rapid self-magnetization of laser speckles in plasmas by nonlinear anisotropic instability. *New J. Phys.* **11**:033001.
- TIDMAN, D. A. & SHANNY, R. A. 1974 Field-generating thermal instability in laser-heated plasma. *Physics of Fluids*. **17**(6):1207–1210.
- WEIBEL, E. S. 1959 Spontaneously Growing Transverse Waves in a Plasma Due to an Anisotropic Velocity Distribution. *Phys. Rev. Lett.* **2**:89.