

Review

## The Metal-Dependent Regulators FurA and FurB from *Mycobacterium Tuberculosis*

Debora Lucarelli <sup>1,#</sup>, Michael L. Vasil <sup>2</sup>, Wolfram Meyer-Klaucke <sup>1</sup> and Ehmke Pohl <sup>3,\*</sup>

<sup>1</sup> European Molecular Biology Laboratory, Hamburg Outstation, Notkestr. 85, D-22603 Hamburg, Germany. E-Mails: Debora.Lucarelli@cancer.org.uk (D. L.); wolfram@embl-hamburg.de (W. M-K.)

<sup>2</sup> Department of Microbiology, University of Colorado, Denver Anschutz Medical Campus, Aurora, CO 80045, USA. E-Mail: Mike.Vasil@UCHSC.edu (M. V.)

<sup>3</sup> Department of Chemistry & School of Biological and Biomedical Sciences, Durham University, Durham DH1 3LE, UK

# Current address: Cancer Research UK, Clare Hall Laboratories, London Research Institute, Blanche Lane, South Mimms, Potters Bar, Herts EN6 3LD, UK

\* Author to whom correspondence should be addressed; E-Mail: ehmke.pohl@durham.ac.uk; Tel. +44-191-3343619; Fax: +44-191-3844737

Received: 9 May 2008; in revised form: 29 July 2008 / Accepted: 30 July 2008 / Published: 28 August 2008

---

**Abstract:** The ferric uptake regulators (Fur) form a large family of bacterial metal-activated DNA-binding proteins that control a diverse set of genes at the transcriptional level. *Mycobacterium tuberculosis*, the causative agent of tuberculosis, expresses two members of the Fur family, designated FurA and FurB. Although both belong to the same family, they share only approximately 25% sequence identity and as a consequence, they differ significantly in some of their key biological functions. FurA appears to be a specialized iron-dependent regulator that controls the *katG* gene, which encodes for a catalase-peroxidase involved in the response of *M. tuberculosis* to oxidative stress. KatG is also the key mycobacterial enzyme responsible for the activation of the first-line tuberculosis drug Isoniazid. FurB in contrast requires Zn<sup>2+</sup> rather than Fe<sup>2+</sup>, to bind to its target sequence in regulated genes, which include those involved in Zn<sup>2+</sup>-homeostasis. Recent biochemical, crystallographic and spectroscopic data have now shed light on the activation and metal discrimination mechanisms in this protein family.

**Keywords:** Metal uptake, regulator, ferric, zinc, *Mycobacterium tuberculosis*.

---

## 1. Introduction

*Mycobacterium tuberculosis*, a Gram-positive bacteria and a widespread human pathogen, resides mainly in alveolar macrophages within the lungs of infected individuals. *M. tuberculosis* has infected up to one third of the world's population but only a small proportion manifests the disease, tuberculosis (TB). TB is most often associated with poor socio-economic conditions and/or co-infections such as HIV. In 2006, it was estimated that there were approximately 9.2 million new cases of TB, with a death toll of more than 1.5 million people [1]. Even though efficacious short-course chemotherapy (DOTS) and the Bacille Guerin-Calmette (BGC) vaccine are now available for a large proportion of the populations of developed countries, the threat remains, as the incidence of multi-drug resistance is steadily increasing in *M. tuberculosis* [2].

Like all human pathogens, *M. tuberculosis* must contend with limited metal availability in order to survive in the human body. Iron and zinc are essential trace elements, however, both are toxic at elevated concentrations [3, 4]. As the sequestration of iron is part of the non-specific mammalian immune response metal uptake and regulation of metal ion concentrations are central to host-pathogen interactions [5]. Metal ion homeostasis in prokaryotes is generally maintained at the level of transcription by various metal dependent regulator proteins [3, 5, 6]. The *M. tuberculosis* genome contains genes encoding for four of these regulators belonging to two different families [7]. IdeR (for Iron-dependent Regulator) and SirR (for staphylococcal iron regulatory repressor) belong to the Diphtheria toxin Repressor (DtxR) family. FurA and FurB are members of the second, the ferric uptake regulator (Fur) family.

IdeR is an essential protein in *M. tuberculosis* and plays the central role as an iron-dependent regulator which controls a large cohort of genes encoding proteins required for iron-uptake and storage [8, 9]. A series of crystal structures of holo-IdeR and IdeR in complex with DNA have shown that this protein undergoes a conformational change upon  $\text{Fe}^{2+}$ -binding and most likely binds its target as a double dimer [10-12]. SirR has originally been identified in *Staphylococcus epidermidis* [13] as a 25 kDa protein with approximately 33% sequence identity to IdeR. In *S. epidermidis* SirR (SirR<sub>SE</sub>) controls the three-gene operon *sitABC* encoding for an ABC transporter system that has been suggested to be involved in iron-uptake [13, 14]. Furthermore a pseudo-palindromic region in the operator/promoter sequence of *sitABC* has been shown to be specifically recognized by  $\text{Fe}^{2+}$ - or  $\text{Mn}^{2+}$ -activated SirR<sub>SE</sub> [13]. The biological role of the SirR homologue in *M. tuberculosis*, however, has yet to be determined.

Fur was originally identified in *E. coli* where it acts as global regulator influencing the expression of close to 100 genes [15, 16]. Members of the Fur family affect the expression of an array of genes, not only associated with iron acquisition and storage, but also those involved in intermediary metabolism as well as responses to acidic environments and oxidative stress. Moreover, they directly or indirectly control the expression of a variety of genes associated with virulence in Gram-positive and negative bacteria [6, 16-19]. In the initial working model it was suggested that Fur is activated by ferric iron and then binds to its operator, a 19 base pair pseudo-palindrome called Fur-Box [16, 20]. Binding to the operator region of a regulated gene blocks access of the RNA-polymerase thereby repressing the transcription of the downstream gene(s). More recent studies, however, found that the


regulation exerted by members of the Fur family is far more intricate than initially perceived. Several homologues, including Fur from *E. coli* (Fur<sub>EC</sub>) contain at least one tightly bound Zn<sup>2+</sup>-ion that is believed to serve a structural rather than a regulatory role [21-23]. Moreover, certain members of the Fur family are activated by divalent transition metals other than Fe<sup>2+</sup>, which include zinc, nickel and manganese. Accordingly, such members have now been named Zur [24], Nur [25] and Mur [26] which connotes that these family members are capable of selectively binding to a specific divalent transition metal other than Fe<sup>2+</sup> to activate their DNA binding function. Furthermore, Fur can also act as a positive regulatory factor, rather than as a classical repressor. While different mechanisms for this type of regulation have been proposed, in a number of bacterial pathogens, Fur homologues directly repress the expression of non-coding small RNAs named RHyhB in *E. coli* [27] and *V. cholerae* [28], PrrF in *P. aeruginosa* [29] and NrrF in *Neisseria meningitidis* [30] that in turn repress the expression of certain genes. Under iron-starvation conditions RyhB is rapidly expressed and stabilized by the RNA-chaperone Hfq. The single-stranded RNA in the RNA-Hfq complex then pairs with the complementary messenger RNA which becomes susceptible to rapid degradation [31].

## 2. Discussion

### 2.1. Biological role of FurA and FurB in *M. tuberculosis*

The *furA* gene is located immediately upstream of the *katG* gene encoding a catalase-peroxidase. KatG is involved in the oxidative stress response and a significant virulence factor of *M. tuberculosis* [32]. FurA is co-expressed along with KatG and it auto-represses its expression by binding to a unique sequence upstream of the *furA* gene [33]. Although there are indications that this protein is also involved in the regulation of other genes [32], the biological role of FurA, in contrast to the role of most members of the family appears to be more specialized (Figure 1). FurA could represent the metal-dependent peroxide sensor similar to the peroxide regulon repressor PerR that has been extensively characterized in *B. subtilis* [34]. Both proteins share a sequence identity of approximately 28% (Figure 2). PerR<sub>BS</sub> adopts a similar fold to Fur<sub>PA</sub> and contains one structural Zn<sup>2+</sup>- and one regulatory Fe<sup>2+</sup>/Mn<sup>2+</sup>-binding site [35]. Once activated and bound to its target DNA-sequence the regulator senses oxidative stress by iron-catalyzed histidine oxidation that leads to the loss of DNA-binding activity [36]. FurA is of particular biomedical importance as it controls a protein central to the contemporary TB therapy. One of the more efficacious therapeutics for this disease is Isoniazid (isonicotinic acid hydrazide = INH), which is able to traverse the complex lipid membrane of *M. tuberculosis*, entirely by passive diffusion. While unmodified INH is not toxic to the pathogen it becomes activated by the mycobacterial catalase KatG, which modifies INH into a range of reactive intermediates including NAD<sup>+</sup> and NADP<sup>+</sup> adducts. These then act as potent inhibitors of the NADPH-dependent enoyl acyl carrier protein reductase (InhA) of the fatty acid synthase type II [37, 38]. InhA is an essential enzyme in the mycolic acid synthesis [39]. INH-resistant *M. tuberculosis* strains isolated from patients predominantly have mutations in the *katG* gene leading to a catalase-peroxidase with reduced or abolished catalytic activity [38]. In addition, mutations were also found in the *inhA* gene as well as various other genes including some in the *furA* gene [38, 40]. FurA has attracted increased interest as a potentially novel drug target for the development of inhibitors that could enhance KatG levels possibly boosting INH potency.

**Figure 1.** Schematic representation of the regulatory functions of *M. tuberculosis* FurA and FurB, respectively. Full arrows refer to proven repression, dashed arrows refer to proposed activation pathways. Note, that the structural  $Zn^{2+}$ -site present in both FurA and FurB has been omitted for clarity.



The biological role of *M. tuberculosis* FurB has only recently been examined. Elegant biological and biochemical studies [41, 42] as well as biophysical and structural analyses [43] strongly suggested that FurB represents the authentic Zinc uptake regulator (Zur) in *M. tuberculosis* (Figure 1). The *furB* gene is co-transcribed with its upstream gene (Rv2358), which encodes another zinc-dependent regulator. In the absence of  $Zn^{2+}$  the gene product of Rv2359 represses the expression of both genes [41]. FurB is responsible for repressing at least 32 genes a number of which have been implicated with zinc homeostasis. For instance, the triplet gene cluster Rv2059-Rv2061c encodes for two components of an ABC-transport system, Rv2059 shows homology with the TroA superfamily, Rv2060 is similar to a membrane protein part of an ABC-type  $Mn^{2+}/Zn^{2+}$  transport system [42, 43]. However, since the two reading frames of Rv2059 and 2060 overlap it is not clear if functional proteins are expressed from these genes [42]. In addition, FurB controls five genes encoding ribosomal proteins three of which containing a putative zinc-binding motif. Based on comparative genomics data these ribosomal proteins have also been suggested to be involved in zinc homeostasis [44]. While FurB plays an important biological role as the genuine zinc-dependent repressor no phenotype difference was observed between wild-type and a *M. tuberculosis* strain in which the *zur* gene was deleted [42].

Although FurA and FurB belong to the same protein family they only share a sequence identity of approximately 25%. Thus, while it is likely that they have a similar overall structure, certain key elements in their respective architectures are likely to differ considerably. For instance, some of the distinct structural details discriminate in divalent metal binding while others are critical to the recognition of DNA sequence motifs in the operator of regulated genes.

**Figure 2.** Sequence alignment of Fur<sub>EC</sub> with FurA, PerR<sub>BS</sub> together with Zur<sub>EC</sub> and FurB (performed with ClustalW2 [45]). Residues of the putative regulatory binding site identified in the crystal structure of FurB are depicted in green, residues of the putative structural binding site are red [43]. The proposed binding residues for the second, regulator binding site in PerR<sub>BS</sub> His37, Asp85, His91, His 93 and Asp104 are shown in blue [46]. Secondary structure assignment is based on the FurB crystal structure: blue rods refers to  $\alpha$ -helix, and violet arrows to  $\beta$ -strands. The last line shows sequence conservation: '\*' denote conserved residues, ':' and ':' indicate similar residues in the alignment.



## 2.2. Crystal structure of FurB

The crystal structure of FurB in complex with Zn<sup>2+</sup> determined at a resolution of 2.7 Å revealed the familiar two-domain structure with the N-terminal DNA-binding domain composed of a three-helix bundle followed by a short antiparallel  $\beta$ -sheet, and the C-terminal metal-binding and dimerization domain [43]. Although the overall fold is similar to the first crystal structure of another member of the Fur-family (Fur from *P. aeruginosa*, Fur<sub>PA</sub> [47]) there are noteworthy differences in domain orientation as well as metal binding sites. Overall, the individual domains of FurB are very similar

with the exception that the DNA-binding domain of FurB lacks the N-terminal helix that is present in Fur<sub>PA</sub> and Fur<sub>EC</sub> [48]. However, the relative orientations of their two domains are very different (Figure 3a-c). The FurB homodimer adopts a much more open conformation where the DNA-binding domains are further separated and the DNA-recognition helices (helix 3, residues 45-58 in FurB in Figure 2) are almost collinear to each other. While the amino-acid sequence of the DNA binding helix is well conserved within the Fur family there are sufficient differences to enable DNA sequence specific recognition. The mobility of DNA-binding domains with respect to the dimer interface is also highlighted by the crystal structure of apo-PerR<sub>BS</sub> which shows a similar fold but a distinctively different domain orientation [46]. It should be noted that such domain motions can also be caused by packing effects as have been observed previously in the crystal structures of apo- and holo-DtxR [49].

**Figure 3.** Ribbon diagrams of the crystal structure of Fur homologues. All three proteins are dimers with the N-terminal DNA-binding domain depicted in green and the C-terminal dimerization domain shown in red. All Zn<sup>2+</sup>-ions are shown as golden spheres. (a) FurB in complex with Zn<sup>2+</sup> solved at 2.7 Å resolution, the putative regulatory sites are labeled Zn1 and Zn1', the structural sites are labeled Zn2 and Zn2', respectively. The labels for the third site were omitted for clarity [43]. (b) Fur<sub>PA</sub> complexed with Zn<sup>2+</sup> [47]. (c) apo-PerR<sub>BS</sub> with its putative structural Zn<sup>2+</sup>-site [46]. All Figures were produced with PyMol [50].





Figure 3. Cont.




### 2.3. Metal binding sites and metal selectivity

Three  $\text{Zn}^{2+}$ -binding sites were identified in the crystal structure of FurB, two of which were further characterized by EXAFS measurements in solution. The biological role of the third  $\text{Zn}^{2+}$ -site is not clear and it seems likely that this site may represent a crystallization artifact. The first metal site depicted in Figure 4a, revealed a  $\text{Zn}^{2+}$ -ion tetrahedrally coordinated by Asp61 (corresponding to Thr69 in Fur<sub>EC</sub>), Cys75 (Thr83), His80 and His82, respectively (His88 and His90). Further spectroscopic analysis showed that this site can be readily exchanged against  $\text{Co}^{2+}$ -ions with the same coordination sphere [43], hence suggesting it represents the lower-affinity regulatory site used to switch the DNA-

binding capability of FurB to *on*. It should be noted that this site corresponds roughly to the originally denoted putative structural site in Fur<sub>PA</sub> [47]. However, more recent data indicate that this assignment based on crystallographic and spectroscopic data may not represent the biologically relevant state [18].

**Figure 4.** Close-up of the two metal binding sites in the crystal structure of FurB: (a) putative regulatory Zn<sup>2+</sup>-site. (b) putative structural Zn<sup>2+</sup>-site. The residue numbers refer to the sequence given in Figure 1.





The sequence comparison shows that the residues of this site depicted in green in the sequence alignment (Figure 2) are only partially conserved, presumably because the different members of the Fur family have evolved to recognize different metals, possibly at different concentrations, thereby fine-tuning the regulatory networks in each organism. It is noteworthy that both Asp 61 and Cys75 in FurB are not conserved in FurA, rather they are changed to arginine residues, which cannot serve as ligands for divalent transition metals. Thus, it is likely that in FurA the framework of the overall Fur fold provides a chemically different metal environment better suited for  $\text{Fe}^{2+}$  than  $\text{Zn}^{2+}$ . Compared to zinc ions, ferric iron prefers octahedral coordination with O,N ligands rather than tetrahedral coordination with S-containing ligands [51]. Considering that the FurA sequence shows a number of histidine and aspartic acid residues in the vicinity a preferred  $\text{Fe}^{2+}$ -binding site can be realized without major structural changes. The residues that have been suggested to constitute the second, regulatory binding site in PerR<sub>BS</sub> (depicted in italic and blue/green in Figure 2) are in fact highly conserved in FurA (His37, Asp85, His91, His93 and Asp104, numbering according to PerR<sub>BS</sub>). Thus, it is possible, that FurA employs a similar mechanism to function as a peroxide sensor in *M. tuberculosis*.

The second metal binding site in FurB depicted in Figure 4b represents a regular tetrahedral coordination of the four cysteine residues Cys85, Cys88, Cys125 and Cys128.  $\text{ZnS}_4$  clusters in this arrangement have been observed in various proteins and are typical for structural  $\text{Zn}^{2+}$ -sites [51, 52]. The notion of a structural  $\text{Zn}^{2+}$ -site that ties the N-terminus to the core of the dimer was further supported by a series of experiments [43]. First of all it was not possible to completely remove the  $\text{Zn}^{2+}$  from FurB without precipitating the protein sample presumably due to (partial) unfolding. Secondly, microPIXE analysis [53] showed approximately one zinc atom per monomer and thirdly, the same tetrahedral  $\text{ZnS}_4$  environment was observed in EXAFS studies performed using a protein solution that had not been incubated with any additional metal. Moreover, a similar structural  $\text{Zn}^{2+}$  site was also revealed in the recent crystal structure of apo-PerR<sub>BS</sub> [46]. Even though the sequence alignment appears to indicate that all cysteine residues in this cluster are highly conserved between Fur<sub>EC</sub>, FurA, Zur<sub>EC</sub> and FurB (depicted in red in Figure 2) it should be noted that a series of mutational and spectroscopic experiments point to slightly different  $\text{Zn}^{2+}$  surroundings for the structural sites ranging from  $\text{Zn}(\text{S})_2(\text{O/N})_2$  in Fur<sub>EC</sub> to  $\text{Zn}(\text{S})_3(\text{O/N})$  in Zur<sub>EC</sub> [21, 54]. This metal site may be structurally less well conserved than deduced from sequence alone. However, preliminary EXAFS analysis on apo-FurA confirms a single  $\text{Zn}^{2+}$ -ion tightly bound in a structural binding site. (Lucarelli, Pohl, Meyer-Klaucke, unpublished data). It is therefore likely that FurA and FurB share the same structural  $\text{Zn}^{2+}$ -spot.

### 3. Conclusions

The past several years have seen significant progress in understanding the control of gene expression by metal dependent transcriptional regulators in *M. tuberculosis* and other bacterial pathogens. FurB has been identified as the Zinc uptake regulator (Zur) [42], while FurA controls the production of the catalase-peroxidase KatG, an essential enzyme in INH therapy of TB treatment [32]. FurA could thus represent the functional homologue of the peroxide sensor PerR. In general the mechanisms governing gene repression are reasonably well understood. By contrast, the full cohort of genes (activated and repressed) controlled by these Fur family members, and other metal-dependent

regulatory proteins have not yet been identified, nor have the mechanisms associated with their activation been fully elucidated. Although metal binding and activation mechanism have now been examined in greater detail for a significant number of Fur family members, including FurA and FurB, their exact mode of DNA-binding and in particular the structural basis of specific target recognition, remains unknown. Further knowledge about the architecture and function of Fur-operator complexes will certainly enhance future structurally driven drug design efforts.

### Acknowledgements

We would like to thank Drs. Alain Baulard (Institute Pasteur, Lille), Santina Russo and Anuschka Pauluhn (Paul Scherrer Institut) for their ongoing support and many valuable discussions. We are grateful to the anonymous reviewers for their comments that significantly improved the manuscript. This work was generously supported by a short-term fellowship to DL from the Boehringer Ingelheim Fonds and in part by a grant from the National Institutes of Health (NIAID R37AI15940) to MLV.

### References

1. WHO. *Report on tuberculosis: global tuberculosis control-surveillance, planning, financing*. World Health Organization: Geneva, Switzerland, **2008**.
2. Zignol, M.; Hosseini, M.S.; Wright, A.; Weezenbeek, C.L.; Nunn, P.; Watt, C.J.; Williams, B.G.; Dye, C. Global incidence of multidrug-resistant tuberculosis. *J. Infect. Dis.* **2006**, *194*, 479-485.
3. Nelson, N. Metal ion transporters and homeostasis. *EMBO J.* **1999**, *18*, 4361-4371.
4. Blencowe, D.K.; Morby, A.P. Zn(II) metabolism in prokaryotes. *FEMS Microbiol. Rev.* **2003**, *27*, 291-311.
5. Agranoff, D.; Krishna, S. Metal ion transport and regulation in Mycobacterium tuberculosis. *Front. Biosci.* **2004**, *9*, 2996-3006.
6. Braun, V. Iron uptake mechanisms and their regulation in pathogenic bacteria. *Int. J. Med. Microbiol.* **2001**, *291*, 67-79.
7. Cole, S.T.; Brosch, R.; Parkhill, J.; Garnier, T.; Churcher, C.; Harris, D.; Gordon, S.V.; Eiglmeier, K.; Gas, S.; Barry, C.E. 3<sup>rd</sup>; *et al.* Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. *Nature* **1998**, *393*, 537-544.
8. Rodriguez, G.M.; Smith, I. Mechanisms of iron regulation in mycobacteria: role in physiology and virulence. *Mol. Microbiol.* **2003**, *47*, 1485-1494.
9. Rodriguez, G. Control of iron metabolism in Mycobacterium tuberculosis. *Trends. Microbiol.* **2006**, *14*, 320-327.
10. Pohl, E.; Holmes, R.K.; Hol, W.G. Crystal structure of the iron-dependent regulator (IdeR) from Mycobacterium tuberculosis shows both metal binding sites fully occupied. *J. Mol. Biol.* **1999**, *285*, 1145-1156.
11. Wisedchaisri, G.; Holmes, R.K.; Hol, W.G. Crystal structure of an IdeR-DNA complex reveals a conformational change in activated IdeR for base-specific interactions. *J. Mol. Biol.* **2004**, *342*, 1155-1169.
12. Feese, M.D.; Ingason, B.P.; Goranson-Siekierke, J.; Holmes, R.K.; Hol, W.G. Crystal structure of the iron-dependent regulator from Mycobacterium tuberculosis at 2.0-Å resolution reveals the Src

- homology domain 3-like fold and metal binding function of the third domain. *J. Biol. Chem.* **2001**, *276*, 5959-5965.
13. Hill, P.J.; Cockayne, A.; Landers, P.; Morrissey, J.A.; Sims, C.M.; Williams, P. SirR, a novel iron-dependent repressor in *Staphylococcus epidermidis*. *Infect. Immun.* **1998**, *66*, 4123-4129.
  14. Morrissey, J.; Cockayne, A.; Hill, P.; Williams, P. Molecular cloning and analysis of a putative siderophore ABC transporter from *Staphylococcus aureus*. *Infect. Immun.* **2001**, *68*, 6281-6288.
  15. Hantke, K. Regulation of ferric iron transport in *Escherichia coli* K12: isolation of a constitutive mutant. *Mol. Gen. Genet.* **1981**, *182*, 288-292.
  16. Hantke, K. Iron and metal regulation in bacteria. *Curr. Opin. Microbiol.* **2001**, *4*(2), 172-177.
  17. Vasil, M.L.; Ochsner, U.A. The response of *Pseudomonas aeruginosa* to iron: genetics, biochemistry and virulence. *Mol. Microbiol.* **1999**, *34*, 399-413.
  18. Lee, J.W.; Helmann, J.D. Functional specialization within the Fur family of metalloregulators. *Biometals.* **2007**, *20*, 485-499.
  19. Vasil, M.L. How we learnt about iron acquisition in *Pseudomonas aeruginosa*: a series of very fortunate events. *Biometals.* **2007**, *20*, 587-601.
  20. Bagg, A.; Neilands, J.B. Molecular mechanism of regulation of siderophore-mediated iron assimilation. *Microbiol. Rev.* **1987**, *51*, 509-518.
  21. Jacquamet, L.; Aberdam, D.; Adrait, A.; Hazemann, J.L.; Latour, J.M.; Michaud-Soret, I. X-ray absorption spectroscopy of a new zinc site in the fur protein from *Escherichia coli*. *Biochemistry* **1998**, *37*, 2564-2571.
  22. Althaus, E.W.; Outten, C.E.; Olson, K.E.; Cao, H.; O'Halloran, T.V. The ferric uptake regulation (Fur) repressor is a zinc metalloprotein. *Biochemistry* **1999**, *38*, 6559-6569.
  23. Zheleznova, E.E.; Crosa, J.H.; Brennan, R.G. Characterization of the DNA- and metal-binding properties of *Vibrio anguillarum* fur reveals conservation of a structural Zn(2+) ion. *J. Bacteriol.* **2000**, *182*, 6264-6267.
  24. Patzer, S.I.; Hantke, K. The ZnuABC high-affinity zinc uptake system and its regulator Zur in *Escherichia coli*. *Mol. Microbiol.* **1998**, *28*, 1199-1210.
  25. Ahn, B.E.; Cha, J.; Lee, E.J.; Han, A.R.; Thompson, C.J.; Roe, J.H. Nur, a nickel-responsive regulator of the Fur family, regulates superoxide dismutases and nickel transport in *Streptomyces coelicolor*. *Mol. Microbiol.* **2006**, *59*, 1848-1858.
  26. Diaz-Mireles, E.; Wexler, M.; Sawers, G.; Bellini, D.; Todd, J.D.; Johnston, A.W. The Fur-like protein Mur of *Rhizobium leguminosarum* is a Mn(2+)-responsive transcriptional regulator. *Microbiology* **2004**, *150*, 1447-1456.
  27. Masse, E.; Gottesman, S. A small RNA regulates the expression of genes involved in iron metabolism in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 4620-4625.
  28. Davis, B.M.; Quinones, M.; Pratt, J.; Ding, Y.; Waldor, M.K. Characterization of the small untranslated RNA RyhB and its regulon in *Vibrio cholerae*. *J. Bacteriol.* **2005**, *187*, 4005-4014.

29. Wilderman, P.J.; Sowa, N.A.; FitzGerald, D.J.; FitzGerald, P.C.; Gottesman, S.; Ochsner, U.A.; Vasil, M.L. Identification of tandem duplicate regulatory small RNAs in *Pseudomonas aeruginosa* involved in iron homeostasis. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 9792-9797.
30. Mellin, J.R.; Goswami, S.; Grogan, S.; Tjaden, B.; Genco, C.A. A novel fur-and iron-regulated small RNA, NrrF, is required for indirect fur-mediated regulation of the *sdhA* and *sdhC* genes in *Neisseria meningitidis*. *J. Bacteriol.* **2007**, *189*, 3686-3694.
31. Massé, E.; Salvaila, H.; Desnoyers, G.; Arguin, M. Small RNAs controlling iron metabolism. *Curr. Opin. Microbiol.* **2007**, *10*, 140-145.
32. Pym, A.S.; Domenech, P.; Honore, N.; Song, J.; Deretic, V.; Cole, S.T. Regulation of catalase-peroxidase (KatG) expression, isoniazid sensitivity and virulence by *furA* of *Mycobacterium tuberculosis*. *Mol. Microbiol.* **2001**, *40*, 879-889.
33. Sala, C.; Forti, F.; Di, Florio E.; Canneva, F.; Milano, A.; Riccardi, G.; Ghisotti, D. *Mycobacterium tuberculosis* FurA autoregulates its own expression. *J. Bacteriol.* **2003**, *185*, 5357-5362.
34. Mongkolsuk, S.; Helmann, J.D. Regulation of inducible peroxide stress responses. *Mol. Microbiol.* **2002**, *45*, 9-15.
35. Lee, J.W.; Helmann, J.D. Biochemical characterization of the structural Zn<sup>2+</sup> site in the *Bacillus subtilis* peroxide sensor PerR. *J. Biol. Chem.* **2006**, *281*, 23567-23578.
36. Lee, J.W.; Helmann, J.D. The PerR transcription factor senses H<sub>2</sub>O<sub>2</sub> by metal-catalysed histidine oxidation. *Nature* **2006**, *440*, 363-367.
37. Marrakchi, H.; Laneelle, G.; Quemard, A. *InhA*, a target of the antituberculous drug isoniazid, is involved in a mycobacterial fatty acid elongation system, FAS-II. *Microbiology* **2000**, *146*, 289-296.
38. Vilchèze, C.; Jacobs, W.R.J. The mechanism of isoniazid killing: clarity through the scope of genetics. *Annu. Rev. Microbiol.* **2007**, *61*, 35-50.
39. Timmins, G.S.; Deretic, V. Mechanisms of action of isoniazid. *Mol. Microbiol.* **2006**, *62*, 1220-1227.
40. Zhang, M.; Yue, J.; Yang, Y.P.; Zhang, H.M.; Lei, J.Q.; Jin, R.L.; Zhang, X.L.; Wang, H.H. Detection of mutations associated with isoniazid resistance in *Mycobacterium tuberculosis* isolates from China. *J. Clin. Microbiol.* **2005**, *43*, 5477-5482.
41. Canneva, F.; Branzoni, M.; Riccardi, G.; Provvedi, R.; Milano, A. Rv2358 and FurB: two transcriptional regulators from *Mycobacterium tuberculosis* which respond to zinc. *J. Bacteriol.* **2005**, *187*, 5837-5840.
42. Maciag, A.; Dainese, E.; Rodriguez, G.M.; Milano, A.; Provvedi, R.; Pasca, M.R.; Smith, I.; Palu, G.; Riccardi, G.; Manganeli, R. Global analysis of *Mycobacterium tuberculosis* Zur (FurB) regulon. *J. Bacteriol.* **2006**, *189*, 730-740.
43. Lucarelli, D.; Russo, S.; Garman, E.; Milano, A.; Meyer-Klaucke, W.; Pohl, E. Crystal structure and function of the zinc uptake regulator FurB from *Mycobacterium tuberculosis*. *J. Biol. Chem.* **2007**, *282*, 9914-9922.

44. Panina, E.; Mironov, A.; Gelfand, M. Comparative genomics of bacterial zinc regulons: enhanced ion transport, pathogenesis, and rearrangement of ribosomal proteins. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 9912-9917.
45. Larkin, M.; Blackshields, G.; Brown, N.; Chenna, R.; McGettigan, P.; McWilliam, H.; Valentin, F.; Wallace, I.; Wilm, A.; Lopez, R.; *et al.* ClustalW and ClustalX version 2. *Bioinformatics* **2007**, *23*, 2947-2948.
46. Traoré, D.A.; El Ghazouani, A.; Ilango, S.; Dupuy, J.; Jacquamet, L.; Ferrer, J.L.; Caux-Thang, C.; Duarte, V.; Latour, J.M. Crystal structure of the apo-PerR-Zn protein from *Bacillus subtilis*. *Mol. Microbiol.* **2006**, *61*, 1211-1219.
47. Pohl, E.; Haller, J.C.; Mijovilovich, A.; Meyer-Klaucke, W.; Garman, E.; Vasil, M.L. Architecture of a protein central to iron homeostasis: crystal structure and spectroscopic analysis of the ferric uptake regulator. *Mol. Microbiol.* **2003**, *47*, 903-915.
48. Pecqueu, L.; D'Autréaux, B.; Dupuy, J.; Nicolet, Y.; Jacquamet, L.; Brutscher, B.; Michaud-Soret, I.; Bersch, B. Structural changes of *Escherichia coli* ferric uptake regulator during metal-dependent dimerization and activation explored by NMR and X-ray crystallography. *J. Biol. Chem.* **2006**, *281*, 21286-21295.
49. Pohl, E.; Holmes, R.K.; Hol, W.G. Motion of the DNA-binding domain with respect to the core of the diphtheria toxin repressor (DtxR) revealed in the crystal structures of apo- and holo-DtxR. *J. Biol. Chem.* **1998**, *273*, 22420-22427.
50. DeLano, W.L. The PyMOL graphics systems, 2002; <http://www.pymol.org>.
51. Harding, M.M. Geometry of metal-ligand interactions in proteins. *Acta Crystallogr. D. Biol. Crystallogr.* **2001**, *57*, 401-411.
52. Auld, D.S. Zinc coordination sphere in biochemical zinc sites. *Biometals.* **2001**, *14*, 271-313.
53. Garman, E.F.; Grime, G.W. Elemental analysis of proteins by microPIXE. *Prog. Biophys. Mol. Biol.* **2005**, *89*, 173-205.
54. Outten, C.E.; Tobin, D.A.; Penner-Hahn, J.E.; O'Halloran, T.V. Characterization of the metal receptor sites in *Escherichia coli* Zur, an ultrasensitive zinc(II) metalloregulatory protein. *Biochemistry* **2001**, *40*, 10417-10423.

© 2008 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).